

Nonparametric Bayesian Statistics: Part II

Tamara Broderick

ITT Career Development Assistant Professor
Electrical Engineering & Computer Science
MIT

Nonparametric Bayes: Part II

- Last time:

Nonparametric Bayes: Part II

- Last time:
 - Understand what it means to have a growing number of parameters

Nonparametric Bayes: Part II

- Last time:
 - Understand what it means to have a growing number of parameters; understand having an infinite # parameters

Nonparametric Bayes: Part II

- Last time:
 - Understand what it means to have a growing number of parameters; understand having an infinite # parameters
 - GEM

Nonparametric Bayes: Part II

- Last time:
 - Understand what it means to have a growing number of parameters; understand having an infinite # parameters
 - GEM
 - `www.tamarabroderick.com/tutorial_2015_summer_npb.html`

Nonparametric Bayes: Part II

- Last time:
 - Understand what it means to have a growing number of parameters; understand having an infinite # parameters
 - GEM
 - `www.tamarabroderick.com/tutorial_2015_summer_npb.html`
- This time:

Nonparametric Bayes: Part II

- Last time:
 - Understand what it means to have a growing number of parameters; understand having an infinite # parameters
 - GEM
 - `www.tamarabroderick.com/tutorial_2015_summer_npb.html`
- This time:
 - Understand what it means to have a growing/infinite number of parameters

Nonparametric Bayes: Part II

- Last time:
 - Understand what it means to have a growing number of parameters; understand having an infinite # parameters
 - GEM
 - `www.tamarabroderick.com/tutorial_2015_summer_npb.html`
- This time:
 - Understand what it means to have a growing/infinite number of parameters
 - Dirichlet process

Nonparametric Bayes: Part II

- Last time:
 - Understand what it means to have a growing number of parameters; understand having an infinite # parameters
 - GEM
 - `www.tamarabroderick.com/tutorial_2015_summer_npb.html`
- This time:
 - Understand what it means to have a growing/infinite number of parameters
 - Dirichlet process
 - Dirichlet process mixture model

Nonparametric Bayes: Part II

- Last time:
 - Understand what it means to have a growing number of parameters; understand having an infinite # parameters
 - GEM
 - `www.tamarabroderick.com/tutorial_2015_summer_npb.html`
- This time:
 - Understand what it means to have a growing/infinite number of parameters
 - Dirichlet process
 - Dirichlet process mixture model
 - A finite representation of an infinite process

Recall: Part I

Recall: Part I

- **Dirichlet process (DP) stick-breaking**

Recall: Part I

- **Dirichlet process (DP) stick-breaking**
- Griffiths-Engen-McCloskey (**GEM**) distribution:

Recall: Part I

- **Dirichlet process (DP) stick-breaking**
- Griffiths-Engen-McCloskey (**GEM**) distribution:
$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

Recall: Part I

- **Dirichlet process (DP) stick-breaking**
- Griffiths-Engen-McCloskey (**GEM**) distribution:
$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

Recall: Part I

- **Dirichlet process (DP) stick-breaking**

- Griffiths-Engen-McCloskey (**GEM**) distribution:

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$



Recall: Part I

- **Dirichlet process (DP) stick-breaking**

- Griffiths-Engen-McCloskey (**GEM**) distribution:

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$




Recall: Part I

- **Dirichlet process (DP) stick-breaking**

- Griffiths-Engen-McCloskey (**GEM**) distribution:

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$




Recall: Part I

- **Dirichlet process (DP) stick-breaking**

- Griffiths-Engen-McCloskey (**GEM**) distribution:

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$




Recall: Part I

- **Dirichlet process (DP) stick-breaking**

- Griffiths-Engen-McCloskey (**GEM**) distribution:

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$




Recall: Part I

- **Dirichlet process (DP) stick-breaking**

- Griffiths-Engen-McCloskey (**GEM**) distribution:

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$




Recall: Part I

- **Dirichlet process (DP) stick-breaking**

- Griffiths-Engen-McCloskey (**GEM**) distribution:

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$



$$V_k \stackrel{iid}{\sim} \text{Beta}(1, \alpha)$$

$$\rho_k = \left[\prod_{j=1}^{k-1} (1 - V_j) \right] V_k$$


...

Recall: Part I

- **Dirichlet process (DP) stick-breaking**

- Griffiths-Engen-McCloskey (**GEM**) distribution:

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$



$$V_k \stackrel{iid}{\sim} \text{Beta}(1, \alpha)$$

$$\rho_k = \left[\prod_{j=1}^{k-1} (1 - V_j) \right] V_k$$

...

[demo]


Distributions

Distributions

- Beta \rightarrow random distribution over 1, 2


Distributions

- Beta \rightarrow random distribution over 1, 2




Distributions

- Beta \rightarrow random distribution over 1, 2
- Dirichlet \rightarrow random distribution over $1, 2, \dots, K$




Distributions

- Beta \rightarrow random distribution over 1, 2
- Dirichlet \rightarrow random distribution over 1, 2, \dots , K




Distributions

- Beta \rightarrow random distribution over 1, 2
- Dirichlet \rightarrow random distribution over 1, 2, \dots , K
- GEM / Dirichlet stick-breaking \rightarrow random distribution over 1, 2, \dots




Distributions

- Beta \rightarrow random distribution over 1, 2
- Dirichlet \rightarrow random distribution over 1, 2, \dots , K
- GEM / Dirichlet stick-breaking \rightarrow random distribution over 1, 2, \dots



Distributions


- Beta \rightarrow random distribution over 1, 2
- Dirichlet \rightarrow random distribution over 1, 2, \dots , K
- GEM / Dirichlet stick-breaking \rightarrow random distribution over 1, 2, \dots



$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

Distributions


- Beta \rightarrow random distribution over 1, 2
- Dirichlet \rightarrow random distribution over 1, 2, \dots , K
- GEM / Dirichlet stick-breaking \rightarrow random distribution over 1, 2, \dots



$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$
$$\phi_k \stackrel{iid}{\sim} G_0$$

Distributions

- Beta \rightarrow random distribution over 1, 2
- Dirichlet \rightarrow random distribution over 1, 2, \dots , K
- GEM / Dirichlet stick-breaking \rightarrow random distribution over 1, 2, \dots




$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\phi_k \stackrel{iid}{\sim} G_0$$

$$G = \sum_{k=1}^{\infty} \rho_k \delta_{\phi_k}$$

Distributions

- Beta \rightarrow random distribution over 1, 2
- Dirichlet \rightarrow random distribution over 1, 2, \dots , K
- GEM / Dirichlet stick-breaking \rightarrow random distribution over 1, 2, \dots



$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\phi_k \stackrel{iid}{\sim} G_0$$

$$G = \sum_{k=1}^{\infty} \rho_k \delta_{\phi_k}$$


Distributions

- Beta \rightarrow random distribution over 1, 2
- Dirichlet \rightarrow random distribution over $1, 2, \dots, K$
- GEM / Dirichlet stick-breaking \rightarrow random distribution over $1, 2, \dots$

- **Dirichlet process** \rightarrow random distribution over Φ :
 $\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$

$$\phi_k \stackrel{iid}{\sim} G_0$$

$$G = \sum_{k=1}^{\infty} \rho_k \delta_{\phi_k}$$



Dirichlet process mixture model

Dirichlet process mixture model

- Gaussian mixture model

Dirichlet process mixture model


- Gaussian mixture model

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

Dirichlet process mixture model

- Gaussian mixture model

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$




Dirichlet process mixture model

- Gaussian mixture model

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\mu_k \stackrel{iid}{\sim} \mathcal{N}(\mu_0, \Sigma_0), k = 1, 2, \dots$$





Dirichlet process mixture model

- Gaussian mixture model

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\mu_k \stackrel{iid}{\sim} \mathcal{N}(\mu_0, \Sigma_0), k = 1, 2, \dots$$





Dirichlet process mixture model

- Gaussian mixture model

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\mu_k \stackrel{iid}{\sim} \mathcal{N}(\mu_0, \Sigma_0), k = 1, 2, \dots$$

- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\mu_k}$





Dirichlet process mixture model

- Gaussian mixture model

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\mu_k \stackrel{iid}{\sim} \mathcal{N}(\mu_0, \Sigma_0), k = 1, 2, \dots$$

- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\mu_k} \stackrel{d}{=} \text{DP}(\alpha, \mathcal{N}(\mu_0, \Sigma_0))$



Dirichlet process mixture model



- Gaussian mixture model

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\mu_k \stackrel{iid}{\sim} \mathcal{N}(\mu_0, \Sigma_0), k = 1, 2, \dots$$

- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\mu_k} \stackrel{d}{=} \text{DP}(\alpha, \mathcal{N}(\mu_0, \Sigma_0))$

$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$





Dirichlet process mixture model

- Gaussian mixture model

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\mu_k \stackrel{iid}{\sim} \mathcal{N}(\mu_0, \Sigma_0), k = 1, 2, \dots$$

- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\mu_k} \stackrel{d}{=} \text{DP}(\alpha, \mathcal{N}(\mu_0, \Sigma_0))$



$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$

$$\mu_n^* = \mu_{z_n}$$


Dirichlet process mixture model

- Gaussian mixture model

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$



$$\mu_k \stackrel{iid}{\sim} \mathcal{N}(\mu_0, \Sigma_0), k = 1, 2, \dots$$

- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\mu_k} \stackrel{d}{=} \text{DP}(\alpha, \mathcal{N}(\mu_0, \Sigma_0))$



$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$

$$\mu_n^* = \mu_{z_n}$$




Dirichlet process mixture model

- Gaussian mixture model

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\mu_k \stackrel{iid}{\sim} \mathcal{N}(\mu_0, \Sigma_0), k = 1, 2, \dots$$



- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\mu_k} \stackrel{d}{=} \text{DP}(\alpha, \mathcal{N}(\mu_0, \Sigma_0))$



$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$

$$\mu_n^* = \mu_{z_n}$$

- i.e. $\mu_n^* \stackrel{iid}{\sim} G$




Dirichlet process mixture model

- Gaussian mixture model

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\mu_k \stackrel{iid}{\sim} \mathcal{N}(\mu_0, \Sigma_0), k = 1, 2, \dots$$


- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\mu_k} \stackrel{d}{=} \text{DP}(\alpha, \mathcal{N}(\mu_0, \Sigma_0))$




$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$

$$\mu_n^* = \mu_{z_n}$$

- i.e. $\mu_n^* \stackrel{iid}{\sim} G$



$$x_n \stackrel{indep}{\sim} \mathcal{N}(\mu_n^*, \Sigma)$$




Dirichlet process mixture model

- Gaussian mixture model

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\mu_k \stackrel{iid}{\sim} \mathcal{N}(\mu_0, \Sigma_0), k = 1, 2, \dots$$


- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\mu_k} \stackrel{d}{=} \text{DP}(\alpha, \mathcal{N}(\mu_0, \Sigma_0))$




$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$

$$\mu_n^* = \mu_{z_n}$$

- i.e. $\mu_n^* \stackrel{iid}{\sim} G$



$$x_n \stackrel{indep}{\sim} \mathcal{N}(\mu_n^*, \Sigma)$$




Dirichlet process mixture model

- Gaussian mixture model

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\mu_k \stackrel{iid}{\sim} \mathcal{N}(\mu_0, \Sigma_0), k = 1, 2, \dots$$


- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\mu_k} \stackrel{d}{=} \text{DP}(\alpha, \mathcal{N}(\mu_0, \Sigma_0))$



$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$


$$\mu_n^* = \mu_{z_n}$$

- i.e. $\mu_n^* \stackrel{iid}{\sim} G$



$$x_n \stackrel{indep}{\sim} \mathcal{N}(\mu_n^*, \Sigma)$$

[demo]




Dirichlet process mixture model

- More generally

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\mu_k \stackrel{iid}{\sim} \mathcal{N}(\mu_0, \Sigma_0), k = 1, 2, \dots$$


- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\mu_k} \stackrel{d}{=} \text{DP}(\alpha, \mathcal{N}(\mu_0, \Sigma_0))$




$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$

$$\mu_n^* = \mu_{z_n}$$

- i.e. $\mu_n^* \stackrel{iid}{\sim} G$



$$x_n \stackrel{indep}{\sim} \mathcal{N}(\mu_n^*, \Sigma)$$



Dirichlet process mixture model

- More generally

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\phi_k \stackrel{iid}{\sim} G_0 \quad k = 1, 2, \dots$$




- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\mu_k} \stackrel{d}{=} \text{DP}(\alpha, \mathcal{N}(\mu_0, \Sigma_0))$

$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$

$$\mu_n^* = \mu_{z_n}$$

- i.e. $\mu_n^* \stackrel{iid}{\sim} G$

$$x_n \stackrel{indep}{\sim} \mathcal{N}(\mu_n^*, \Sigma)$$



Dirichlet process mixture model

- More generally

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\phi_k \stackrel{iid}{\sim} G_0 \quad k = 1, 2, \dots$$




- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\mu_k} \stackrel{d}{=} \text{DP}(\alpha, \mathcal{N}(\mu_0, \Sigma_0))$

$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$

$$\mu_n^* = \mu_{z_n}$$

- i.e. $\mu_n^* \stackrel{iid}{\sim} G$

$$x_n \stackrel{indep}{\sim} \mathcal{N}(\mu_n^*, \Sigma)$$



Dirichlet process mixture model

- More generally

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\phi_k \stackrel{iid}{\sim} G_0 \quad k = 1, 2, \dots$$




- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\mu_k} \stackrel{d}{=} \text{DP}(\alpha, \mathcal{N}(\mu_0, \Sigma_0))$

$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$

$$\mu_n^* = \mu_{z_n}$$

- i.e. $\mu_n^* \stackrel{iid}{\sim} G$

$$x_n \stackrel{indep}{\sim} \mathcal{N}(\mu_n^*, \Sigma)$$



Dirichlet process mixture model

- More generally

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\phi_k \stackrel{iid}{\sim} G_0 \quad k = 1, 2, \dots$$




- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\phi_k} \stackrel{d}{=} \text{DP}(\alpha, \mathcal{N}(\mu_0, \Sigma_0))$

$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$

$$\mu_n^* = \mu_{z_n}$$

- i.e. $\mu_n^* \stackrel{iid}{\sim} G$

$$x_n \stackrel{indep}{\sim} \mathcal{N}(\mu_n^*, \Sigma)$$



Dirichlet process mixture model

- More generally

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\phi_k \stackrel{iid}{\sim} G_0 \quad k = 1, 2, \dots$$




- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\phi_k} \stackrel{d}{=} \text{DP}(\alpha, G_0)$

$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$

$$\mu_n^* = \mu_{z_n}$$

- i.e. $\mu_n^* \stackrel{iid}{\sim} G$

$$x_n \stackrel{indep}{\sim} \mathcal{N}(\mu_n^*, \Sigma)$$



Dirichlet process mixture model

- More generally

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\phi_k \stackrel{iid}{\sim} G_0 \quad k = 1, 2, \dots$$




- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\phi_k} \stackrel{d}{=} \text{DP}(\alpha, G_0)$

$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$

$$\theta_n = \phi_{z_n}$$

- i.e. $\mu_n^* \stackrel{iid}{\sim} G$

$$x_n \stackrel{indep}{\sim} \mathcal{N}(\mu_n^*, \Sigma)$$



Dirichlet process mixture model

- More generally

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\phi_k \stackrel{iid}{\sim} G_0 \quad k = 1, 2, \dots$$




- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\phi_k} \stackrel{d}{=} \text{DP}(\alpha, G_0)$

$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$

$$\theta_n = \phi_{z_n}$$

- i.e. $\theta_n \stackrel{iid}{\sim} G$

$$x_n \stackrel{indep}{\sim} \mathcal{N}(\mu_n^*, \Sigma)$$



Dirichlet process mixture model

- More generally

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\phi_k \stackrel{iid}{\sim} G_0 \quad k = 1, 2, \dots$$




- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\phi_k} \stackrel{d}{=} \text{DP}(\alpha, G_0)$

$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$

$$\theta_n = \phi_{z_n}$$

- i.e. $\theta_n \stackrel{iid}{\sim} G$

$$x_n \stackrel{indep}{\sim} F(\theta_n)$$



Dirichlet process mixture model

- More generally

$$\rho = (\rho_1, \rho_2, \dots) \sim \text{GEM}(\alpha)$$

$$\phi_k \stackrel{iid}{\sim} G_0 \quad k = 1, 2, \dots$$




- i.e. $G = \sum_{k=1}^{\infty} \rho_k \delta_{\phi_k} \stackrel{d}{=} \text{DP}(\alpha, G_0)$

$$z_n \stackrel{iid}{\sim} \text{Categorical}(\rho)$$

$$\theta_n = \phi_{z_n}$$


- i.e. $\theta_n \stackrel{iid}{\sim} G$

$$x_n \stackrel{indep}{\sim} F(\theta_n)$$




[Antoniak 1974; Ferguson 1983; West, Müller, Escobar 1994;
Escobar, West 1995; MacEachern, Müller 1998]

DP or not DP, that is the question





DP or not DP, that is the question

- GEM: 





DP or not DP, that is the question

- GEM: 
- Compare to:



DP or not DP, that is the question

- GEM: 
- Compare to:
 - Finite (small K) mixture model




DP or not DP, that is the question


- GEM: 
- Compare to:
 - Finite (small K) mixture model



- Finite (large K) mixture model



DP or not DP, that is the question



- GEM: 
- Compare to:
 - Finite (small K) mixture model



- Finite (large K) mixture model




- Time series



Marginal cluster assignments


$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$



Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$




Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$

$$p(z_n = 1 | z_1, \dots, z_{n-1})$$




Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$

$$p(z_n = 1 | z_1, \dots, z_{n-1}) \\ = \int p(z_n = 1, \rho_1 | z_1, \dots, z_{n-1}) d\rho_1$$




Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$

$$p(z_n = 1 | z_1, \dots, z_{n-1})$$

$$= \int p(z_n = 1 | \rho_1) p(\rho_1 | z_1, \dots, z_{n-1}) d\rho_1$$



Marginal cluster assignments


- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$

$$p(z_n = 1 | z_1, \dots, z_{n-1})$$

$$= \int p(z_n = 1 | \rho_1) p(\rho_1 | z_1, \dots, z_{n-1}) d\rho_1$$

$$= \int \rho_1 \text{Beta}(\rho_1 | a_{1,n}, a_{2,n}) d\rho_1$$



Marginal cluster assignments

- Integrate out the frequencies


$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$

$$p(z_n = 1 | z_1, \dots, z_{n-1})$$

$$= \int p(z_n = 1 | \rho_1) p(\rho_1 | z_1, \dots, z_{n-1}) d\rho_1$$

$$= \int \rho_1 \text{Beta}(\rho_1 | a_{1,n}, a_{2,n}) d\rho_1$$

$$a_{1,n} := a_1 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 1\}, a_{2,n} = a_2 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 2\}$$



Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$


$$p(z_n = 1 | z_1, \dots, z_{n-1})$$

$$= \int p(z_n = 1 | \rho_1) p(\rho_1 | z_1, \dots, z_{n-1}) d\rho_1$$

$$= \int \rho_1 \text{Beta}(\rho_1 | a_{1,n}, a_{2,n}) d\rho_1$$

$$a_{1,n} := a_1 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 1\}, a_{2,n} = a_2 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 2\}$$


$$= \int \rho_1 \frac{\Gamma(a_{1,n} + a_{2,n})}{\Gamma(a_{1,n})\Gamma(a_{2,n})} \rho_1^{a_{1,n}-1} (1 - \rho_1)^{a_{2,n}-1} d\rho_1$$



Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$



$$p(z_n = 1 | z_1, \dots, z_{n-1})$$

$$= \int p(z_n = 1 | \rho_1) p(\rho_1 | z_1, \dots, z_{n-1}) d\rho_1$$

$$= \int \rho_1 \text{Beta}(\rho_1 | a_{1,n}, a_{2,n}) d\rho_1$$

$$a_{1,n} := a_1 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 1\}, a_{2,n} = a_2 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 2\}$$


$$= \int \rho_1 \frac{\Gamma(a_{1,n} + a_{2,n})}{\Gamma(a_{1,n})\Gamma(a_{2,n})} \rho_1^{a_{1,n}-1} (1 - \rho_1)^{a_{2,n}-1} d\rho_1$$

$$= \frac{\Gamma(a_{1,n} + a_{2,n})}{\Gamma(a_{1,n})\Gamma(a_{2,n})} \frac{\Gamma(a_{1,n} + 1)\Gamma(a_{2,n})}{\Gamma(a_{1,n} + a_{2,n} + 1)}$$

Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$



$$p(z_n = 1 | z_1, \dots, z_{n-1})$$

$$= \int p(z_n = 1 | \rho_1) p(\rho_1 | z_1, \dots, z_{n-1}) d\rho_1$$

$$= \int \rho_1 \text{Beta}(\rho_1 | a_{1,n}, a_{2,n}) d\rho_1$$

$$a_{1,n} := a_1 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 1\}, a_{2,n} = a_2 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 2\}$$

$$= \int \rho_1 \frac{\Gamma(a_{1,n} + a_{2,n})}{\Gamma(a_{1,n})\Gamma(a_{2,n})} \rho_1^{a_{1,n}-1} (1 - \rho_1)^{a_{2,n}-1} d\rho_1$$

$$= \frac{\Gamma(a_{1,n} + a_{2,n})}{\Gamma(a_{1,n})\Gamma(a_{2,n})} \frac{\Gamma(a_{1,n} + 1)\Gamma(a_{2,n})}{\Gamma(a_{1,n} + a_{2,n} + 1)}$$

$$= \frac{a_{1,n}}{a_{1,n} + a_{2,n}}$$


Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$

$$p(z_n = 1 | z_1, \dots, z_{n-1}) = \frac{a_{1,n}}{a_{1,n} + a_{2,n}}$$

$$a_{1,n} := a_1 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 1\}, a_{2,n} = a_2 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 2\}$$




Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$

$$p(z_n = 1 | z_1, \dots, z_{n-1}) = \frac{a_{1,n}}{a_{1,n} + a_{2,n}}$$



$$a_{1,n} := a_1 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 1\}, a_{2,n} := a_2 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 2\}$$


- Pólya urn

Marginal cluster assignments

- Integrate out the frequencies


$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$

$$p(z_n = 1 | z_1, \dots, z_{n-1}) = \frac{a_{1,n}}{a_{1,n} + a_{2,n}}$$



$$a_{1,n} := a_1 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 1\}, a_{2,n} := a_2 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 2\}$$

- Pólya urn




Marginal cluster assignments

- Integrate out the frequencies


$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$

$$p(z_n = 1 | z_1, \dots, z_{n-1}) = \frac{a_{1,n}}{a_{1,n} + a_{2,n}}$$



$$a_{1,n} := a_1 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 1\}, a_{2,n} = a_2 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 2\}$$

- Pólya urn




Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$


$$p(z_n = 1 | z_1, \dots, z_{n-1}) = \frac{a_{1,n}}{a_{1,n} + a_{2,n}}$$



$$a_{1,n} := a_1 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 1\}, a_{2,n} := a_2 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 2\}$$

- Pólya urn

- Choose any ball with equal probability




Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$


$$p(z_n = 1 | z_1, \dots, z_{n-1}) = \frac{a_{1,n}}{a_{1,n} + a_{2,n}}$$



$$a_{1,n} := a_1 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 1\}, a_{2,n} = a_2 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 2\}$$

- Pólya urn

- Choose any ball with equal probability
- Replace and add ball of same color




Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$


$$p(z_n = 1 | z_1, \dots, z_{n-1}) = \frac{a_{1,n}}{a_{1,n} + a_{2,n}}$$



$$a_{1,n} := a_1 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 1\}, a_{2,n} = a_2 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 2\}$$

- Pólya urn

- Choose any ball with equal probability
- Replace and add ball of same color




$$\lim_{n \rightarrow \infty} \frac{\# \text{ orange}}{\# \text{ total}}$$

Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$


$$p(z_n = 1 | z_1, \dots, z_{n-1}) = \frac{a_{1,n}}{a_{1,n} + a_{2,n}}$$



$$a_{1,n} := a_1 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 1\}, a_{2,n} = a_2 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 2\}$$

- Pólya urn

- Choose any ball with equal probability
- Replace and add ball of same color




$$\lim_{n \rightarrow \infty} \frac{\# \text{ orange}}{\# \text{ total}} = \rho_{\text{orange}}$$

Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$


$$p(z_n = 1 | z_1, \dots, z_{n-1}) = \frac{a_{1,n}}{a_{1,n} + a_{2,n}}$$



$$a_{1,n} := a_1 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 1\}, a_{2,n} = a_2 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 2\}$$

- Pólya urn

- Choose any ball with equal probability
- Replace and add ball of same color




$$\lim_{n \rightarrow \infty} \frac{\# \text{ orange}}{\# \text{ total}} = \rho_{\text{orange}} \stackrel{d}{=} \text{Beta}(a_{\text{orange}}, a_{\text{green}})$$

Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$


$$p(z_n = 1 | z_1, \dots, z_{n-1}) = \frac{a_{1,n}}{a_{1,n} + a_{2,n}}$$



$$a_{1,n} := a_1 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 1\}, a_{2,n} = a_2 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 2\}$$

- Pólya urn

- Choose any ball with equal probability
- Replace and add ball of same color




$$\lim_{n \rightarrow \infty} \frac{\# \text{ orange}}{\# \text{ total}} = \rho_{\text{orange}} \stackrel{d}{=} \text{Beta}(a_{\text{orange}}, a_{\text{green}})$$

Marginal cluster assignments

- Integrate out the frequencies

$$\rho_1 \sim \text{Beta}(a_1, a_2), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_1, \rho_2)$$


$$p(z_n = 1 | z_1, \dots, z_{n-1}) = \frac{a_{1,n}}{a_{1,n} + a_{2,n}}$$



$$a_{1,n} := a_1 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 1\}, a_{2,n} = a_2 + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = 2\}$$

- Pólya urn


- Choose any ball with prob proportional to its mass
- Replace and add ball of same color



$$\lim_{n \rightarrow \infty} \frac{\# \text{ orange}}{\# \text{ total}} = \rho_{\text{orange}} \stackrel{d}{=} \text{Beta}(a_{\text{orange}}, a_{\text{green}})$$

Marginal cluster assignments


- Integrate out the frequencies



Marginal cluster assignments

- Integrate out the frequencies

$$\rho_{1:K} \sim \text{Dirichlet}(a_{1:K}), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_{1:K})$$




Marginal cluster assignments

- Integrate out the frequencies

$$\rho_{1:K} \sim \text{Dirichlet}(a_{1:K}), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_{1:K})$$

$$p(z_n = k | z_1, \dots, z_{n-1}) = \frac{a_{k,n}}{\sum_{j=1}^K a_{j,n}}$$




Marginal cluster assignments

- Integrate out the frequencies

$$\rho_{1:K} \sim \text{Dirichlet}(a_{1:K}), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_{1:K})$$

$$p(z_n = k | z_1, \dots, z_{n-1}) = \frac{a_{k,n}}{\sum_{j=1}^K a_{j,n}}$$

$$a_{k,n} := a_k + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = k\}$$



Marginal cluster assignments


- Integrate out the frequencies

$$\rho_{1:K} \sim \text{Dirichlet}(a_{1:K}), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_{1:K})$$

$$p(z_n = k | z_1, \dots, z_{n-1}) = \frac{a_{k,n}}{\sum_{j=1}^K a_{j,n}}$$

$$a_{k,n} := a_k + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = k\}$$

- multivariate Pólya urn



Marginal cluster assignments


- Integrate out the frequencies

$$\rho_{1:K} \sim \text{Dirichlet}(a_{1:K}), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_{1:K})$$

$$p(z_n = k | z_1, \dots, z_{n-1}) = \frac{a_{k,n}}{\sum_{j=1}^K a_{j,n}}$$

$$a_{k,n} := a_k + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = k\}$$

- multivariate Pólya urn



Marginal cluster assignments


- Integrate out the frequencies

$$\rho_{1:K} \sim \text{Dirichlet}(a_{1:K}), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_{1:K})$$

$$p(z_n = k | z_1, \dots, z_{n-1}) = \frac{a_{k,n}}{\sum_{j=1}^K a_{j,n}}$$

$$a_{k,n} := a_k + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = k\}$$

- multivariate Pólya urn
 - Choose any ball with prob proportional to its mass



Marginal cluster assignments

- Integrate out the frequencies


$$\rho_{1:K} \sim \text{Dirichlet}(a_{1:K}), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_{1:K})$$

$$p(z_n = k | z_1, \dots, z_{n-1}) = \frac{a_{k,n}}{\sum_{j=1}^K a_{j,n}}$$

$$a_{k,n} := a_k + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = k\}$$

- multivariate Pólya urn

- Choose any ball with prob proportional to its mass
- Replace and add ball of same color



Marginal cluster assignments

- Integrate out the frequencies


$$\rho_{1:K} \sim \text{Dirichlet}(a_{1:K}), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_{1:K})$$

$$p(z_n = k | z_1, \dots, z_{n-1}) = \frac{a_{k,n}}{\sum_{j=1}^K a_{j,n}}$$

$$a_{k,n} := a_k + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = k\}$$

- multivariate Pólya urn

- Choose any ball with prob proportional to its mass
- Replace and add ball of same color



$$\lim_{n \rightarrow \infty} \frac{(\# \text{ orange}, \# \text{ green}, \# \text{ red}, \# \text{ yellow})}{\# \text{ total}}$$

Marginal cluster assignments

- Integrate out the frequencies


$$\rho_{1:K} \sim \text{Dirichlet}(a_{1:K}), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_{1:K})$$

$$p(z_n = k | z_1, \dots, z_{n-1}) = \frac{a_{k,n}}{\sum_{j=1}^K a_{j,n}}$$

$$a_{k,n} := a_k + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = k\}$$

- multivariate Pólya urn

- Choose any ball with prob proportional to its mass
- Replace and add ball of same color



$$\lim_{n \rightarrow \infty} \frac{(\# \text{ orange}, \# \text{ green}, \# \text{ red}, \# \text{ yellow})}{\# \text{ total}} \rightarrow (\rho_{\text{orange}}, \rho_{\text{green}}, \rho_{\text{red}}, \rho_{\text{yellow}})$$

Marginal cluster assignments

- Integrate out the frequencies


$$\rho_{1:K} \sim \text{Dirichlet}(a_{1:K}), z_n \stackrel{iid}{\sim} \text{Cat}(\rho_{1:K})$$

$$p(z_n = k | z_1, \dots, z_{n-1}) = \frac{a_{k,n}}{\sum_{j=1}^K a_{j,n}}$$

$$a_{k,n} := a_k + \sum_{m=1}^{n-1} \mathbf{1}\{z_m = k\}$$

- multivariate Pólya urn

- Choose any ball with prob proportional to its mass
- Replace and add ball of same color



$$\lim_{n \rightarrow \infty} \frac{(\# \text{ orange}, \# \text{ green}, \# \text{ red}, \# \text{ yellow})}{\# \text{ total}}$$

$$\rightarrow (\rho_{\text{orange}}, \rho_{\text{green}}, \rho_{\text{red}}, \rho_{\text{yellow}})$$


$$\stackrel{d}{=} \text{Dirichlet}(a_{\text{orange}}, a_{\text{green}}, a_{\text{red}}, a_{\text{yellow}})$$

Marginal cluster assignments

- Hoppe urn / Blackwell-MacQueen urn


Marginal cluster assignments

- Hoppe urn / Blackwell-MacQueen urn




Marginal cluster assignments

- Hoppe urn / Blackwell-MacQueen urn



Marginal cluster assignments


- Hoppe urn / Blackwell-MacQueen urn



- Choose ball with prob proportional to its mass

Marginal cluster assignments


- Hoppe urn / Blackwell-MacQueen urn



- Choose ball with prob proportional to its mass
 - If black, replace and add ball of new color

Marginal cluster assignments


- Hoppe urn / Blackwell-MacQueen urn



- Choose ball with prob proportional to its mass
 - If black, replace and add ball of new color
 - Else, replace and add ball of same color

Marginal cluster assignments

- Hoppe urn / Blackwell-MacQueen urn




- Choose ball with prob proportional to its mass
 - If black, replace and add ball of new color
 - Else, replace and add ball of same color

Step 0




Marginal cluster assignments

- Hoppe urn / Blackwell-MacQueen urn




- Choose ball with prob proportional to its mass
 - If black, replace and add ball of new color
 - Else, replace and add ball of same color




Marginal cluster assignments

- Hoppe urn / Blackwell-MacQueen urn




- Choose ball with prob proportional to its mass
 - If black, replace and add ball of new color
 - Else, replace and add ball of same color




Marginal cluster assignments

- Hoppe urn / Blackwell-MacQueen urn




- Choose ball with prob proportional to its mass
 - If black, replace and add ball of new color
 - Else, replace and add ball of same color




Marginal cluster assignments

- Hoppe urn / Blackwell-MacQueen urn




- Choose ball with prob proportional to its mass
 - If black, replace and add ball of new color
 - Else, replace and add ball of same color




Marginal cluster assignments

- Hoppe urn / Blackwell-MacQueen urn




- Choose ball with prob proportional to its mass
 - If black, replace and add ball of new color
 - Else, replace and add ball of same color




Marginal cluster assignments

- Hoppe urn / Blackwell-MacQueen urn



- Choose ball with prob proportional to its mass
 - If black, replace and add ball of new color
 - Else, replace and add ball of same color




$(\# \text{orange}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$


- not orange: $(\# \text{green}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$

Marginal cluster assignments

- Hoppe urn / Blackwell-MacQueen urn



- Choose ball with prob proportional to its mass
 - If black, replace and add ball of new color
 - Else, replace and add ball of same color




$(\# \text{orange}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$


- not orange: $(\# \text{green}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$
- not orange, green: $(\# \text{red}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$

Marginal cluster assignments

- Hoppe urn / Blackwell-MacQueen urn



- Choose ball with prob proportional to its mass
 - If black, replace and add ball of new color
 - Else, replace and add ball of same color




$(\# \text{orange}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$


- not orange: $(\# \text{green}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$
- not orange, green: $(\# \text{red}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$

Marginal cluster assignments

- Hoppe urn / Blackwell-MacQueen urn



- Choose ball with prob proportional to its mass
 - If black, replace and add ball of new color
 - Else, replace and add ball of same color




$(\# \text{orange}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$


- not orange: $(\# \text{green}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$
- not orange, green: $(\# \text{red}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$

Marginal cluster assignments

- Hoppe urn / Blackwell-MacQueen urn



- Choose ball with prob proportional to its mass
 - If black, replace and add ball of new color
 - Else, replace and add ball of same color




$(\# \text{orange}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$


- not orange: $(\# \text{green}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$
- not orange, green: $(\# \text{red}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$

Marginal cluster assignments

- Hoppe urn / Blackwell-MacQueen urn




- Choose ball with prob proportional to its mass
 - If black, replace and add ball of new color
 - Else, replace and add ball of same color




$(\# \text{orange}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$

- not orange: $(\# \text{green}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$
- not orange, green: $(\# \text{red}, \# \text{other}) = \text{PolyaUrn}(1, \alpha)$

Chinese restaurant process




Chinese restaurant process




- Same thing we just did

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process



- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process



- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α
- Marginal for the Categorical likelihood with GEM prior

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α
- Marginal for the Categorical likelihood with GEM prior
 $z_1 = z_2 = z_7 = z_8 = 1, z_3 = z_5 = z_6 = 2, z_4 = 3$

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α
- Marginal for the Categorical likelihood with GEM prior
$$z_1 = z_2 = z_7 = z_8 = 1, z_3 = z_5 = z_6 = 2, z_4 = 3$$
$$\Rightarrow \Pi_8 = \{\{1, 2, 7, 8\}, \{3, 5, 6\}, \{4\}\}$$

Chinese restaurant process




- Same thing we just did
- Each customer walks into the restaurant
 - Sits at existing table with prob proportional to # people there
 - Forms new table with prob proportional to α
- Marginal for the Categorical likelihood with GEM prior
$$z_1 = z_2 = z_7 = z_8 = 1, z_3 = z_5 = z_6 = 2, z_4 = 3$$
$$\Rightarrow \Pi_8 = \{\{1, 2, 7, 8\}, \{3, 5, 6\}, \{4\}\}$$
- *Partition of [8]*: set of mutually exclusive & exhaustive sets of $[8] = \{1, \dots, 8\}$

Chinese restaurant process



- Probability of this seating:


Chinese restaurant process



- Probability of this seating:

$$\frac{\alpha}{\alpha}$$


Chinese restaurant process



- Probability of this seating:

$$\frac{\alpha}{\alpha} \cdot \frac{1}{\alpha + 1}$$


Chinese restaurant process



- Probability of this seating:

$$\frac{\alpha}{\alpha} \cdot \frac{1}{\alpha + 1} \cdot \frac{\alpha}{\alpha + 2}$$


Chinese restaurant process



- Probability of this seating:

$$\frac{\alpha}{\alpha} \cdot \frac{1}{\alpha + 1} \cdot \frac{\alpha}{\alpha + 2} \cdot \frac{\alpha}{\alpha + 3}$$


Chinese restaurant process



- Probability of this seating:

$$\frac{\alpha}{\alpha} \cdot \frac{1}{\alpha + 1} \cdot \frac{\alpha}{\alpha + 2} \cdot \frac{\alpha}{\alpha + 3} \cdot \frac{1}{\alpha + 4}$$


Chinese restaurant process



- Probability of this seating:

$$\frac{\alpha}{\alpha} \cdot \frac{1}{\alpha + 1} \cdot \frac{\alpha}{\alpha + 2} \cdot \frac{\alpha}{\alpha + 3} \cdot \frac{1}{\alpha + 4} \cdot \frac{2}{\alpha + 5}$$


Chinese restaurant process



- Probability of this seating:

$$\frac{\alpha}{\alpha} \cdot \frac{1}{\alpha + 1} \cdot \frac{\alpha}{\alpha + 2} \cdot \frac{\alpha}{\alpha + 3} \cdot \frac{1}{\alpha + 4} \cdot \frac{2}{\alpha + 5} \cdot \frac{2}{\alpha + 6}$$


Chinese restaurant process



- Probability of this seating:

$$\frac{\alpha}{\alpha} \cdot \frac{1}{\alpha + 1} \cdot \frac{\alpha}{\alpha + 2} \cdot \frac{\alpha}{\alpha + 3} \cdot \frac{1}{\alpha + 4} \cdot \frac{2}{\alpha + 5} \cdot \frac{2}{\alpha + 6} \cdot \frac{3}{\alpha + 7}$$

Exercises



- If I run the Chinese restaurant process (CRP) for N customers, what is the probability of the customers sitting at K_N tables where the k th table to form has n_k ($n_k > 0$) people at it?
- What is the expected number of clusters generated by a $\text{CRP}(\alpha)$ after N data points?
- What is the distribution of the number of clusters under a $\text{CRP}(\alpha)$ after N data points? (consider simulations and/or theory)

References (Part II), Page 1

DJ Aldous. *Exchangeability and related topics*. Springer, 1983.

CE Antoniak. Mixtures of Dirichlet processes with applications to Bayesian nonparametric problems. *The Annals of Statistics*, 1974.

D Blackwell and JB MacQueen. Ferguson distributions via Pólya urn schemes. *The Annals of Statistics*, 1973.

S Engen. A note on the geometric series as a species frequency model. *Biometrika*, 1975.

MD Escobar and M West. Bayesian density estimation and inference using mixtures. *Journal of the American Statistical Association*, 1995.

W Ewens. Population genetics theory -- the past and the future. *Mathematical and Statistical Developments of Evolutionary Theory*, 1987.

TS Ferguson. A Bayesian analysis of some nonparametric problems. *The Annals of Statistics*, 1973.

TS Ferguson. Bayesian density estimation by mixtures of normal distributions. *Recent Advances in Statistics*, 1983.

FM Hoppe. Pólya-like urns and the Ewens' sampling formula. *Journal of Mathematical Biology*, 1984.

H Ishwaran and LF James. Gibbs sampling methods for stick-breaking priors. *Journal of the American Statistical Association*, 2001.

SN MacEachern and P Müller. Estimating mixtures of Dirichlet process models. *Journal of Computational and Graphical Statistics*, 1998.

JW McCloskey. A model for the distribution of individuals by species in an environment. Ph.D. thesis, Michigan State University, 1965.

References (Part II), Page 2

GP Patil and C Taillie. Diversity as a concept and its implications for random communities. *Bulletin of the International Statistical Institute*, 1977.

J Sethuraman. A constructive definition of Dirichlet priors. *Statistica Sinica*, 1994.

M West, P Müller, and MD Escobar. Hierarchical Priors and Mixture Models, With Application in Regression and Density Estimation. *Aspects of Uncertainty*, 1994.