

# STAT 7200

## Introduction to Advanced Probability

### Lecture 4

Taylor R. Brown

## 1 Probability Triples

- Extension Theorem
  - Constructing Probability Triples
  - Semialgebra
  - Algebra
  - Extension Theorem
  - Outer Measure  $P^*$
  - Outer Measure  $P^*$  is Countably Subadditive
  - $\mathcal{M}$ : The Measurable Sets
  - $\mathcal{M}$  and  $P^*$

“A First Look at Rigorous Probability Theory” (Jeffrey Rosenthal)  
Sections 2.1, 2.2, 2.3

# Constructing Probability Triples

Here we outline the major steps in constructing (complicated) probability triples on a sample space such as  $\Omega = [0, 1]$  or  $\mathbb{R}$ .

# Constructing Probability Triples

Here we outline the major steps in constructing (complicated) probability triples on a sample space such as  $\Omega = [0, 1]$  or  $\mathbb{R}$ .

- 1) Define a “probability measure”  $P$  on  $\mathcal{J}$ , which is a collection of subsets of  $\Omega$  that forms a *semialgebra*. For instance,  $\mathcal{J}$  can be chosen as the collection of all intervals.

# Constructing Probability Triples

Here we outline the major steps in constructing (complicated) probability triples on a sample space such as  $\Omega = [0, 1]$  or  $\mathbb{R}$ .

- 1) Define a “probability measure”  $P$  on  $\mathcal{J}$ , which is a collection of subsets of  $\Omega$  that forms a *semialgebra*. For instance,  $\mathcal{J}$  can be chosen as the collection of all intervals.
- 2) Construct a new function  $P^*$  as an extension of  $P$  over *\*all\** subsets of  $\Omega$ .  $P^*$  is usually not a proper probability measure for all the subsets of  $\Omega$ .

# Constructing Probability Triples

Here we outline the major steps in constructing (complicated) probability triples on a sample space such as  $\Omega = [0, 1]$  or  $\mathbb{R}$ .

- 1) Define a “probability measure”  $P$  on  $\mathcal{J}$ , which is a collection of subsets of  $\Omega$  that forms a *semialgebra*. For instance,  $\mathcal{J}$  can be chosen as the collection of all intervals.
- 2) Construct a new function  $P^*$  as an extension of  $P$  over *\*all\** subsets of  $\Omega$ .  $P^*$  is usually not a proper probability measure for all the subsets of  $\Omega$ .
- 3) Based on  $P^*$ , we construct a new collection of subsets, denoted as  $\mathcal{M}$ .  $\mathcal{M}$  stands for “measurable.”

# Constructing Probability Triples

Here we outline the major steps in constructing (complicated) probability triples on a sample space such as  $\Omega = [0, 1]$  or  $\mathbb{R}$ .

- 1) Define a “probability measure”  $P$  on  $\mathcal{J}$ , which is a collection of subsets of  $\Omega$  that forms a *semialgebra*. For instance,  $\mathcal{J}$  can be chosen as the collection of all intervals.
- 2) Construct a new function  $P^*$  as an extension of  $P$  over *\*all\** subsets of  $\Omega$ .  $P^*$  is usually not a proper probability measure for all the subsets of  $\Omega$ .
- 3) Based on  $P^*$ , we construct a new collection of subsets, denoted as  $\mathcal{M}$ .  $\mathcal{M}$  stands for “measurable.”
- 4)  $\mathcal{M}$  is a  $\sigma$ -algebra and  $P^*$  is a proper probability measure on  $\mathcal{M}$ .

# Constructing Probability Triples

Here we outline the major steps in constructing (complicated) probability triples on a sample space such as  $\Omega = [0, 1]$  or  $\mathbb{R}$ .

- 1) Define a “probability measure”  $P$  on  $\mathcal{J}$ , which is a collection of subsets of  $\Omega$  that forms a *semialgebra*. For instance,  $\mathcal{J}$  can be chosen as the collection of all intervals.
- 2) Construct a new function  $P^*$  as an extension of  $P$  over *\*all\** subsets of  $\Omega$ .  $P^*$  is usually not a proper probability measure for all the subsets of  $\Omega$ .
- 3) Based on  $P^*$ , we construct a new collection of subsets, denoted as  $\mathcal{M}$ .  $\mathcal{M}$  stands for “measurable.”
- 4)  $\mathcal{M}$  is a  $\sigma$ -algebra and  $P^*$  is a proper probability measure on  $\mathcal{M}$ .
- 5)  $\mathcal{J} \subseteq \mathcal{M}$  and the extension from  $P$  to  $P^*$  is “unique.”



# Constructing Probability Triples

Here we outline the major steps in constructing (complicated) probability triples on a sample space such as  $\Omega = [0, 1]$  or  $\mathbb{R}$ .

- 1) Define a “probability measure”  $P$  on  $\mathcal{J}$ , which is a collection of subsets of  $\Omega$  that forms a *semialgebra*. For instance,  $\mathcal{J}$  can be chosen as the collection of all intervals.
- 2) Construct a new function  $P^*$  as an extension of  $P$  over *\*all\** subsets of  $\Omega$ .  $P^*$  is usually not a proper probability measure for all the subsets of  $\Omega$ .
- 3) Based on  $P^*$ , we construct a new collection of subsets, denoted as  $\mathcal{M}$ .  $\mathcal{M}$  stands for “measurable.”
- 4)  $\mathcal{M}$  is a  $\sigma$ -algebra and  $P^*$  is a proper probability measure on  $\mathcal{M}$ .
- 5)  $\mathcal{J} \subseteq \mathcal{M}$  and the extension from  $P$  to  $P^*$  is “unique.”

The above steps lead to a probability triple  $\{\Omega, \mathcal{M}, P^*\}$ .

# Semialgebra

- **Semialgebra**

Let  $\mathcal{J}$  be a collection of subsets of  $\Omega$ .  $\mathcal{J}$  is a semialgebra if

a)  $\emptyset, \Omega \in \mathcal{J}$ .

b) If  $A_1, A_2, \dots, A_k \in \mathcal{J}$ , then  $\bigcap_{i=1}^k A_i \in \mathcal{J}$ . (Closed under finite intersections)

c) If  $A \in \mathcal{J}$ , then there is a pairwise disjoint sequence of sets  $B_1, B_2, \dots, B_m \in \mathcal{J}$  such that  $A^c = \bigcup_{i=1}^m B_i$ .

# Semialgebra

- **Semialgebra**

Let  $\mathcal{J}$  be a collection of subsets of  $\Omega$ .  $\mathcal{J}$  is a semialgebra if

a)  $\emptyset, \Omega \in \mathcal{J}$ .

b) If  $A_1, A_2, \dots, A_k \in \mathcal{J}$ , then  $\bigcap_{i=1}^k A_i \in \mathcal{J}$ . (Closed under finite intersections)

c) If  $A \in \mathcal{J}$ , then there is a pairwise disjoint sequence of sets  $B_1, B_2, \dots, B_m \in \mathcal{J}$  such that  $A^c = \bigcup_{i=1}^m B_i$ .

## Proposition 1 (Exercise 2.2.3)

$\mathcal{J} = \{ \text{All "intervals" contained in } [0,1] \text{ (or } \mathbb{R}) \}$  is a semialgebra.

# Semialgebra

- **Semialgebra**

Let  $\mathcal{J}$  be a collection of subsets of  $\Omega$ .  $\mathcal{J}$  is a semialgebra if

a)  $\emptyset, \Omega \in \mathcal{J}$ .

b) If  $A_1, A_2, \dots, A_k \in \mathcal{J}$ , then  $\bigcap_{i=1}^k A_i \in \mathcal{J}$ . (Closed under finite intersections)

c) If  $A \in \mathcal{J}$ , then there is a pairwise disjoint sequence of sets  $B_1, B_2, \dots, B_m \in \mathcal{J}$  such that  $A^c = \bigcup_{i=1}^m B_i$ .

## Proposition 1 (Exercise 2.2.3)

$\mathcal{J} = \{ \text{All "intervals" contained in } [0,1] \text{ (or } \mathbb{R}) \}$  is a semialgebra.

- **Remark** The notion of an “interval” includes singletons and open/closed/half-open/empty intervals.

# Algebra

- **Algebra**

Let  $\mathcal{B}_0$  be a collection of subsets of  $\Omega$ .  $\mathcal{B}_0$  is an **algebra** if

a)  $\emptyset \in \mathcal{B}_0$ .

b) If  $A \in \mathcal{B}_0$ , then  $A^c \in \mathcal{B}_0$ . (Closed under complement)

c) If  $A_1, \dots, A_m \in \mathcal{B}_0$ , then  $\bigcup_{i=1}^m A_i \in \mathcal{B}_0$ .

# Algebra

- **Algebra**

Let  $\mathcal{B}_0$  be a collection of subsets of  $\Omega$ .  $\mathcal{B}_0$  is an **algebra** if

a)  $\emptyset \in \mathcal{B}_0$ .

b) If  $A \in \mathcal{B}_0$ , then  $A^c \in \mathcal{B}_0$ . (Closed under complement)

c) If  $A_1, \dots, A_m \in \mathcal{B}_0$ , then  $\bigcup_{i=1}^m A_i \in \mathcal{B}_0$ .

- **Remark** it's nearly the same as the definition for a  $\sigma$ -algebra, except the union in c) is *\*finite\**.

# Algebra

- **Algebra**

Let  $\mathcal{B}_0$  be a collection of subsets of  $\Omega$ .  $\mathcal{B}_0$  is an **algebra** if

a)  $\emptyset \in \mathcal{B}_0$ .

b) If  $A \in \mathcal{B}_0$ , then  $A^c \in \mathcal{B}_0$ . (Closed under complement)

c) If  $A_1, \dots, A_m \in \mathcal{B}_0$ , then  $\bigcup_{i=1}^m A_i \in \mathcal{B}_0$ .

- **Remark** it's nearly the same as the definition for a  $\sigma$ -algebra, except the union in c) is \*finite\*.

- **Remark** b), c) and De Morgan's law together imply further that algebras are closed under finite intersections, as well.

# Algebra

- **Algebra**

Let  $\mathcal{B}_0$  be a collection of subsets of  $\Omega$ .  $\mathcal{B}_0$  is an **algebra** if

a)  $\emptyset \in \mathcal{B}_0$ .

b) If  $A \in \mathcal{B}_0$ , then  $A^c \in \mathcal{B}_0$ . (Closed under complement)

c) If  $A_1, \dots, A_m \in \mathcal{B}_0$ , then  $\bigcup_{i=1}^m A_i \in \mathcal{B}_0$ .

- **Remark** it's nearly the same as the definition for a  $\sigma$ -algebra, except the union in c) is \*finite\*.

- **Remark** b), c) and De Morgan's law together imply further that algebras are closed under finite intersections, as well.

## Proposition 2 (Exercise 2.2.5)

$\mathcal{B}_0 = \{ \text{All finite unions of "intervals" in } [0,1] \text{ (or } \mathbb{R}) \}$  is an algebra.



# Extension Theorem

## Theorem 3

The Extension Theorem *Let  $\mathcal{J}$  be a semialgebra of subsets of  $\Omega$  and  $P : \mathcal{J} \rightarrow [0, 1]$  such that:*

a)  $P(\emptyset) = 0, P(\Omega) = 1.$

b)  $P(\bigcup_{i=1}^k A_i) \geq \sum_{i=1}^k P(A_i)$  whenever  $A_1, \dots, A_k \in \mathcal{J}$ ,  $\bigcup_{i=1}^k A_i \in \mathcal{J}$ , and  $A_1, \dots, A_k$  are pairwise disjoint (finite superadditivity).

c)  $P(A) \leq \sum_n P(A_n)$  whenever  $A, A_1, A_2, \dots \in \mathcal{J}$ , and  $A \subseteq \bigcup_n A_n$  (countable monotonicity).

*Then there is a  $\sigma$ -algebra  $\mathcal{M} \supseteq \mathcal{J}$  and a countably-additive probability measure  $P^*$  on  $\mathcal{M}$  such that  $P^*(A) = P(A)$  for all  $A \in \mathcal{J}$ .*

## Outer Measure $P^*$

- **Remarks:**  $P^*$  is a function defined over *\*all\** subsets of  $\Omega$  (but its definition makes use of only sets from  $\mathcal{J}$ ) but it isn't necessarily a probability measure if you're looking at all of these subsets.

## Outer Measure $P^*$

- **Remarks:**  $P^*$  is a function defined over *\*all\** subsets of  $\Omega$  (but its definition makes use of only sets from  $\mathcal{J}$ ) but it isn't necessarily a probability measure if you're looking at all of these subsets.
- For any  $A \subseteq \Omega$

$$P^*(A) := \inf_{A_1, A_2, \dots \in \mathcal{J}, A \subseteq \bigcup_i A_i} \sum_i P(A_i)$$

## Outer Measure $P^*$

- **Remarks:**  $P^*$  is a function defined over *\*all\** subsets of  $\Omega$  (but its definition makes use of only sets from  $\mathcal{J}$ ) but it isn't necessarily a probability measure if you're looking at all of these subsets.
- For any  $A \subseteq \Omega$

$$P^*(A) := \inf_{A_1, A_2, \dots \in \mathcal{J}, A \subseteq \bigcup_i A_i} \sum_i P(A_i)$$

### Lemma 4

*Outer measure satisfies the following properties:*

- a)  $P^*(\emptyset) = 0$ .
- b)  $P^*(A) \leq P^*(B)$  if  $A \subseteq B$ . (Monotonicity)
- c)  $P^*(A) = P(A)$  if  $A \in \mathcal{J}$ . ( $P^*$  is an extension of  $P$ )

# Outer Measure $P^*$ is Countably Subadditive

## Lemma 5 (2.3.6.)

*Outer measure  $P^*$  is countably subadditive:*

$$P^*(\bigcup_{n=1}^{\infty} B_n) \leq \sum_{n=1}^{\infty} P^*(B_n) \text{ for any } B_1, B_2, \dots \in \Omega$$

# Outer Measure $P^*$ is Countably Subadditive

## Lemma 5 (2.3.6.)

*Outer measure  $P^*$  is countably subadditive:*

$$P^*(\bigcup_{n=1}^{\infty} B_n) \leq \sum_{n=1}^{\infty} P^*(B_n) \text{ for any } B_1, B_2, \dots \in \Omega$$

**Proof:** The key is  $P^*(A)$  is defined as the infimum of the countable unions of sets in  $\mathcal{J}$  that “cover”  $A$ .

# Outer Measure $P^*$ is Countably Subadditive

## Lemma 5 (2.3.6.)

*Outer measure  $P^*$  is countably subadditive:*

$$P^*(\bigcup_{n=1}^{\infty} B_n) \leq \sum_{n=1}^{\infty} P^*(B_n) \text{ for any } B_1, B_2, \dots \in \Omega$$

**Proof:** The key is  $P^*(A)$  is defined as the infimum of the countable unions of sets in  $\mathcal{J}$  that “cover”  $A$ .

- 1) Given  $\varepsilon > 0$ , for each  $B_n$ , there must be a sequence  $\{C_{nk}\}_{k=1}^{\infty}$ , s.t.  $C_{nk} \in \mathcal{J}$ ,  $B_n \subseteq \bigcup_k C_{nk}$  and  $\sum_k P(C_{nk}) < P^*(B_n) + \varepsilon/2^n$  (small typo in book)

# Outer Measure $P^*$ is Countably Subadditive

## Lemma 5 (2.3.6.)

*Outer measure  $P^*$  is countably subadditive:*

$$P^*(\bigcup_{n=1}^{\infty} B_n) \leq \sum_{n=1}^{\infty} P^*(B_n) \text{ for any } B_1, B_2, \dots \in \mathcal{J}$$

**Proof:** The key is  $P^*(A)$  is defined as the infimum of the countable unions of sets in  $\mathcal{J}$  that “cover”  $A$ .

- 1) Given  $\varepsilon > 0$ , for each  $B_n$ , there must be a sequence  $\{C_{nk}\}_{k=1}^{\infty}$ , s.t.  $C_{nk} \in \mathcal{J}$ ,  $B_n \subseteq \bigcup_k C_{nk}$  and  $\sum_k P(C_{nk}) < P^*(B_n) + \varepsilon/2^n$  (small typo in book)
- 2) Since  $\bigcup_{n=1}^{\infty} B_n \subseteq \bigcup_{n,k} C_{n,k}$ ,  
$$P^*(\bigcup_{n=1}^{\infty} B_n) \leq \sum_{n,k} P(C_{n,k}) < \sum_n P^*(B_n) + \varepsilon.$$
- 3) As  $\varepsilon$  is an arbitrary positive constant, we must have  
$$P^*(\bigcup_{n=1}^{\infty} B_n) \leq \sum_{n=1}^{\infty} P^*(B_n).$$



## $\mathcal{M}$ : The Measurable Sets

- Outer measure cannot always be a probability measure over *all* subsets of  $\Omega$  (recall Proposition 1.2.6). Define a refined collection of subsets using  $P^*$ :

$$\mathcal{M} = \{A \subseteq \Omega : P^*(A \cap E) + P^*(A^c \cap E) = P^*(E) \text{ for all } E \subseteq \Omega\}$$

## $\mathcal{M}$ : The Measurable Sets

- Outer measure cannot always be a probability measure over *\*all\** subsets of  $\Omega$  (recall Proposition 1.2.6). Define a refined collection of subsets using  $P^*$ :

$$\mathcal{M} = \{A \subseteq \Omega : P^*(A \cap E) + P^*(A^c \cap E) = P^*(E) \text{ for all } E \subseteq \Omega\}$$

### Lemma 6

*We have the following results regarding  $\mathcal{M}$ :*

*a)  $\emptyset \in \mathcal{M}, \Omega \in \mathcal{M}$*

*b) If  $A \in \mathcal{M}$ , then  $A^c \in \mathcal{M}$  (Closed under complement)*

## $\mathcal{M}$ : The Measurable Sets

- Outer measure cannot always be a probability measure over *\*all\** subsets of  $\Omega$  (recall Proposition 1.2.6). Define a refined collection of subsets using  $P^*$ :

$$\mathcal{M} = \{A \subseteq \Omega : P^*(A \cap E) + P^*(A^c \cap E) = P^*(E) \text{ for all } E \subseteq \Omega\}$$

### Lemma 6

*We have the following results regarding  $\mathcal{M}$ :*

- a)  $\emptyset \in \mathcal{M}, \Omega \in \mathcal{M}$*
- b) If  $A \in \mathcal{M}$ , then  $A^c \in \mathcal{M}$  (Closed under complement)*

- Remark:** We often need to verify that a given set  $A \in \mathcal{M}$ . By the countable subadditivity of our outer measure, we always have  $P^*(E) \leq P^*(A \cap E) + P^*(A^c \cap E)$  for all  $E \subseteq \Omega$ . If it's easier, we only need need to verify  $P^*(E) \geq P^*(A \cap E) + P^*(A^c \cap E)$  for all  $E \subseteq \Omega$ . This would can be achieved by using the finite superadditivity of  $P$ .

# $\mathcal{M}$ and $P^*$

- We still need to show that  $\mathcal{M}$  is a  $\sigma$ -algebra and  $P^*$  is a proper probability measure over  $\mathcal{M}$ . A review of our results so far:

# $\mathcal{M}$ and $P^*$

- We still need to show that  $\mathcal{M}$  is a  $\sigma$ -algebra and  $P^*$  is a proper probability measure over  $\mathcal{M}$ . A review of our results so far:
- $\mathcal{M} = \{A : A \subseteq \Omega, P^*(A \cap E) + P^*(A^c \cap E) = P^*(E) \text{ for all } E \subseteq \Omega\}$ .  
We already have  $\emptyset \in \mathcal{M}, \Omega \in \mathcal{M}$  and  $\mathcal{M}$  is closed under complement. So we still need to prove that  $\mathcal{M}$  is closed under countable unions.

# $\mathcal{M}$ and $P^*$

- We still need to show that  $\mathcal{M}$  is a  $\sigma$ -algebra and  $P^*$  is a proper probability measure over  $\mathcal{M}$ . A review of our results so far:
- $\mathcal{M} = \{A : A \subseteq \Omega, P^*(A \cap E) + P^*(A^c \cap E) = P^*(E) \text{ for all } E \subseteq \Omega\}$ . We already have  $\emptyset \in \mathcal{M}, \Omega \in \mathcal{M}$  and  $\mathcal{M}$  is closed under complement. So we still need to prove that  $\mathcal{M}$  is closed under countable unions.
- Regarding  $P^*$ , for any  $A \in \mathcal{M}$ , first,  $P^*(A) \geq 0$ ; second, in the definition of  $\mathcal{M}$ , by choosing  $E = \Omega$ , we have  $P^*(A) = 1 - P^*(A^c)$ . So we still need to show that, on  $\mathcal{M}$ ,  $P^*$  is countably additive.

# $\mathcal{M}$ and $P^*$

- We still need to show that  $\mathcal{M}$  is a  $\sigma$ -algebra and  $P^*$  is a proper probability measure over  $\mathcal{M}$ . A review of our results so far:
- $\mathcal{M} = \{A : A \subseteq \Omega, P^*(A \cap E) + P^*(A^c \cap E) = P^*(E) \text{ for all } E \subseteq \Omega\}$ . We already have  $\emptyset \in \mathcal{M}, \Omega \in \mathcal{M}$  and  $\mathcal{M}$  is closed under complement. So we still need to prove that  $\mathcal{M}$  is closed under countable unions.
- Regarding  $P^*$ , for any  $A \in \mathcal{M}$ , first,  $P^*(A) \geq 0$ ; second, in the definition of  $\mathcal{M}$ , by choosing  $E = \Omega$ , we have  $P^*(A) = 1 - P^*(A^c)$ . So we still need to show that, on  $\mathcal{M}$ ,  $P^*$  is countably additive.
- Countable additivity of  $P^*$  is shown in the next lecture, as well as the fact that  $\mathcal{M}$  is a  $\sigma$ -field.