

STAT 7200

Introduction to Advanced Probability

Lecture 4

Taylor R. Brown

1 Probability Triples

- Extension Theorem
 - Constructing Probability Triples
 - Semialgebra
 - Algebra
 - Extension Theorem
 - Outer Measure P^*
 - Outer Measure P^* is Countably Subadditive
 - \mathcal{M} : The Measurable Sets
 - \mathcal{M} and P^*

“A First Look at Rigorous Probability Theory” (Jeffrey Rosenthal)
Sections 2.1, 2.2, 2.3

Constructing Probability Triples

Here we outline the major steps in constructing (complicated) probability triples on a sample space such as $\Omega = [0, 1]$ or \mathbb{R} .

Constructing Probability Triples

Here we outline the major steps in constructing (complicated) probability triples on a sample space such as $\Omega = [0, 1]$ or \mathbb{R} .

- 1) Define a “probability measure” P on \mathcal{J} , which is a collection of subsets of Ω that forms a *semialgebra*. For instance, \mathcal{J} can be chosen as the collection of all intervals.

Constructing Probability Triples

Here we outline the major steps in constructing (complicated) probability triples on a sample space such as $\Omega = [0, 1]$ or \mathbb{R} .

- 1) Define a “probability measure” P on \mathcal{J} , which is a collection of subsets of Ω that forms a *semialgebra*. For instance, \mathcal{J} can be chosen as the collection of all intervals.
- 2) Construct a new function P^* as an extension of P over **all** subsets of Ω . P^* is usually not a proper probability measure for all the subsets of Ω .

Constructing Probability Triples

Here we outline the major steps in constructing (complicated) probability triples on a sample space such as $\Omega = [0, 1]$ or \mathbb{R} .

- 1) Define a “probability measure” P on \mathcal{J} , which is a collection of subsets of Ω that forms a *semialgebra*. For instance, \mathcal{J} can be chosen as the collection of all intervals.
- 2) Construct a new function P^* as an extension of P over **all** subsets of Ω . P^* is usually not a proper probability measure for all the subsets of Ω .
- 3) Based on P^* , we construct a new collection of subsets, denoted as \mathcal{M} . \mathcal{M} stands for “measurable.”

Constructing Probability Triples

Here we outline the major steps in constructing (complicated) probability triples on a sample space such as $\Omega = [0, 1]$ or \mathbb{R} .

- 1) Define a “probability measure” P on \mathcal{J} , which is a collection of subsets of Ω that forms a *semialgebra*. For instance, \mathcal{J} can be chosen as the collection of all intervals.
- 2) Construct a new function P^* as an extension of P over **all** subsets of Ω . P^* is usually not a proper probability measure for all the subsets of Ω .
- 3) Based on P^* , we construct a new collection of subsets, denoted as \mathcal{M} . \mathcal{M} stands for “measurable.”
- 4) \mathcal{M} is a σ -algebra and P^* is a proper probability measure on \mathcal{M} .

Constructing Probability Triples

Here we outline the major steps in constructing (complicated) probability triples on a sample space such as $\Omega = [0, 1]$ or \mathbb{R} .

- 1) Define a “probability measure” P on \mathcal{J} , which is a collection of subsets of Ω that forms a *semialgebra*. For instance, \mathcal{J} can be chosen as the collection of all intervals.
- 2) Construct a new function P^* as an extension of P over **all** subsets of Ω . P^* is usually not a proper probability measure for all the subsets of Ω .
- 3) Based on P^* , we construct a new collection of subsets, denoted as \mathcal{M} . \mathcal{M} stands for “measurable.”
- 4) \mathcal{M} is a σ -algebra and P^* is a proper probability measure on \mathcal{M} .
- 5) $\mathcal{J} \subseteq \mathcal{M}$ and the extension from P to P^* is “unique.”

Constructing Probability Triples

Here we outline the major steps in constructing (complicated) probability triples on a sample space such as $\Omega = [0, 1]$ or \mathbb{R} .

- 1) Define a “probability measure” P on \mathcal{J} , which is a collection of subsets of Ω that forms a *semialgebra*. For instance, \mathcal{J} can be chosen as the collection of all intervals.
- 2) Construct a new function P^* as an extension of P over **all** subsets of Ω . P^* is usually not a proper probability measure for all the subsets of Ω .
- 3) Based on P^* , we construct a new collection of subsets, denoted as \mathcal{M} . \mathcal{M} stands for “measurable.”
- 4) \mathcal{M} is a σ -algebra and P^* is a proper probability measure on \mathcal{M} .
- 5) $\mathcal{J} \subseteq \mathcal{M}$ and the extension from P to P^* is “unique.”

The above steps lead to a probability triple $\{\Omega, \mathcal{M}, P^*\}$.

Semialgebra

- **Semialgebra**

Let \mathcal{J} be a collection of subsets of Ω . \mathcal{J} is a semialgebra if

a) $\emptyset, \Omega \in \mathcal{J}$.

b) If $A_1, A_2, \dots, A_k \in \mathcal{J}$, then $\bigcap_{i=1}^k A_i \in \mathcal{J}$. (Closed under finite intersections)

c) If $A \in \mathcal{J}$, then there is a pairwise disjoint sequence of sets $B_1, B_2, \dots, B_m \in \mathcal{J}$ such that $A^c = \bigcup_{i=1}^m B_i$.

Semialgebra

- **Semialgebra**

Let \mathcal{J} be a collection of subsets of Ω . \mathcal{J} is a semialgebra if

a) $\emptyset, \Omega \in \mathcal{J}$.

b) If $A_1, A_2, \dots, A_k \in \mathcal{J}$, then $\bigcap_{i=1}^k A_i \in \mathcal{J}$. (Closed under finite intersections)

c) If $A \in \mathcal{J}$, then there is a pairwise disjoint sequence of sets $B_1, B_2, \dots, B_m \in \mathcal{J}$ such that $A^c = \bigcup_{i=1}^m B_i$.

Proposition 1 (Exercise 2.2.3)

$\mathcal{J} = \{ \text{All "intervals" contained in } [0,1] \text{ (or } \mathbb{R}) \}$ is a semialgebra.

Semialgebra

- **Semialgebra**

Let \mathcal{J} be a collection of subsets of Ω . \mathcal{J} is a semialgebra if

a) $\emptyset, \Omega \in \mathcal{J}$.

b) If $A_1, A_2, \dots, A_k \in \mathcal{J}$, then $\bigcap_{i=1}^k A_i \in \mathcal{J}$. (Closed under finite intersections)

c) If $A \in \mathcal{J}$, then there is a pairwise disjoint sequence of sets $B_1, B_2, \dots, B_m \in \mathcal{J}$ such that $A^c = \bigcup_{i=1}^m B_i$.

Proposition 1 (Exercise 2.2.3)

$\mathcal{J} = \{ \text{All "intervals" contained in } [0,1] \text{ (or } \mathbb{R}) \}$ is a semialgebra.

- **Remark** The notion of an “interval” includes singletons and open/closed/half-open/empty intervals.

Algebra

- **Algebra**

Let \mathcal{B}_0 be a collection of subsets of Ω . \mathcal{B}_0 is an **algebra** if

a) $\emptyset \in \mathcal{B}_0$.

b) If $A \in \mathcal{B}_0$, then $A^c \in \mathcal{B}_0$. (Closed under complement)

c) If $A_1, \dots, A_m \in \mathcal{B}_0$, then $\bigcup_{i=1}^m A_i \in \mathcal{B}_0$.

Algebra

- **Algebra**

Let \mathcal{B}_0 be a collection of subsets of Ω . \mathcal{B}_0 is an **algebra** if

a) $\emptyset \in \mathcal{B}_0$.

b) If $A \in \mathcal{B}_0$, then $A^c \in \mathcal{B}_0$. (Closed under complement)

c) If $A_1, \dots, A_m \in \mathcal{B}_0$, then $\bigcup_{i=1}^m A_i \in \mathcal{B}_0$.

- **Remark** it's nearly the same as the definition for a σ -algebra, except the union in c) is **finite**.

Algebra

- **Algebra**

Let \mathcal{B}_0 be a collection of subsets of Ω . \mathcal{B}_0 is an **algebra** if

a) $\emptyset \in \mathcal{B}_0$.

b) If $A \in \mathcal{B}_0$, then $A^c \in \mathcal{B}_0$. (Closed under complement)

c) If $A_1, \dots, A_m \in \mathcal{B}_0$, then $\bigcup_{i=1}^m A_i \in \mathcal{B}_0$.

- **Remark** it's nearly the same as the definition for a σ -algebra, except the union in c) is *finite*.

- **Remark** b), c) and De Morgan's law together imply further that algebras are closed under finite intersections, as well.

Algebra

- **Algebra**

Let \mathcal{B}_0 be a collection of subsets of Ω . \mathcal{B}_0 is an **algebra** if

a) $\emptyset \in \mathcal{B}_0$.

b) If $A \in \mathcal{B}_0$, then $A^c \in \mathcal{B}_0$. (Closed under complement)

c) If $A_1, \dots, A_m \in \mathcal{B}_0$, then $\bigcup_{i=1}^m A_i \in \mathcal{B}_0$.

- **Remark** it's nearly the same as the definition for a σ -algebra, except the union in c) is *finite*.

- **Remark** b), c) and De Morgan's law together imply further that algebras are closed under finite intersections, as well.

Proposition 2 (Exercise 2.2.5)

$\mathcal{B}_0 = \{ \text{All finite unions of "intervals" in } [0,1] \text{ (or } \mathbb{R}) \}$ is an algebra.

Extension Theorem

Theorem 3

The Extension Theorem *Let \mathcal{J} be a semialgebra of subsets of Ω and $P : \mathcal{J} \rightarrow [0, 1]$ such that:*

a) $P(\emptyset) = 0, P(\Omega) = 1.$

b) $P(\bigcup_{i=1}^k A_i) \geq \sum_{i=1}^k P(A_i)$ whenever $A_1, \dots, A_k \in \mathcal{J}$, $\bigcup_{i=1}^k A_i \in \mathcal{J}$, and A_1, \dots, A_k are pairwise disjoint (finite superadditivity).

c) $P(A) \leq \sum_n P(A_n)$ whenever $A, A_1, A_2, \dots \in \mathcal{J}$, and $A \subseteq \bigcup_n A_n$ (countable monotonicity).

Then there is a σ -algebra $\mathcal{M} \supseteq \mathcal{J}$ and a countably-additive probability measure P^ on \mathcal{M} such that $P^*(A) = P(A)$ for all $A \in \mathcal{J}$.*

Outer Measure P^*

- **Remarks:** P^* is a function defined over **all** subsets of Ω (but its definition makes use of only sets from \mathcal{J}) but it isn't necessarily a probability measure if you're looking at all of these subsets.

Outer Measure P^*

- **Remarks:** P^* is a function defined over **all** subsets of Ω (but its definition makes use of only sets from \mathcal{J}) but it isn't necessarily a probability measure if you're looking at all of these subsets.
- For any $A \subseteq \Omega$

$$P^*(A) := \inf_{A_1, A_2, \dots \in \mathcal{J}, A \subseteq \bigcup_i A_i} \sum_i P(A_i)$$

Outer Measure P^*

- **Remarks:** P^* is a function defined over **all** subsets of Ω (but its definition makes use of only sets from \mathcal{J}) but it isn't necessarily a probability measure if you're looking at all of these subsets.
- For any $A \subseteq \Omega$

$$P^*(A) := \inf_{A_1, A_2, \dots \in \mathcal{J}, A \subseteq \bigcup_i A_i} \sum_i P(A_i)$$

Lemma 4

Outer measure satisfies the following properties:

- a) $P^*(\emptyset) = 0$.
- b) $P^*(A) \leq P^*(B)$ if $A \subseteq B$. (Monotonicity)
- c) $P^*(A) = P(A)$ if $A \in \mathcal{J}$. (P^* is an extension of P)

Outer Measure P^* is Countably Subadditive

Lemma 5 (2.3.6.)

Outer measure P^ is countably subadditive:*

$$P^*(\bigcup_{n=1}^{\infty} B_n) \leq \sum_{n=1}^{\infty} P^*(B_n) \text{ for any } B_1, B_2, \dots \subseteq \Omega$$

Outer Measure P^* is Countably Subadditive

Lemma 5 (2.3.6.)

Outer measure P^ is countably subadditive:*

$$P^*(\bigcup_{n=1}^{\infty} B_n) \leq \sum_{n=1}^{\infty} P^*(B_n) \text{ for any } B_1, B_2, \dots \subseteq \Omega$$

Proof: The key is $P^*(A)$ is defined as the infimum of the countable unions of sets in \mathcal{J} that “cover” A .

Outer Measure P^* is Countably Subadditive

Lemma 5 (2.3.6.)

Outer measure P^ is countably subadditive:*

$$P^*(\bigcup_{n=1}^{\infty} B_n) \leq \sum_{n=1}^{\infty} P^*(B_n) \text{ for any } B_1, B_2, \dots \subseteq \Omega$$

Proof: The key is $P^*(A)$ is defined as the infimum of the countable unions of sets in \mathcal{J} that “cover” A .

- 1) Given $\varepsilon > 0$, for each B_n , there must be a sequence $\{C_{nk}\}_{k=1}^{\infty}$, s.t. $C_{nk} \in \mathcal{J}$, $B_n \subseteq \bigcup_k C_{nk}$ and $\sum_k P(C_{nk}) < P^*(B_n) + \varepsilon/2^n$ (small typo in book)

Outer Measure P^* is Countably Subadditive

Lemma 5 (2.3.6.)

Outer measure P^ is countably subadditive:*

$$P^*(\bigcup_{n=1}^{\infty} B_n) \leq \sum_{n=1}^{\infty} P^*(B_n) \text{ for any } B_1, B_2, \dots \subseteq \Omega$$

Proof: The key is $P^*(A)$ is defined as the infimum of the countable unions of sets in \mathcal{J} that “cover” A .

- 1) Given $\varepsilon > 0$, for each B_n , there must be a sequence $\{C_{nk}\}_{k=1}^{\infty}$, s.t. $C_{nk} \in \mathcal{J}$, $B_n \subseteq \bigcup_k C_{nk}$ and $\sum_k P(C_{nk}) < P^*(B_n) + \varepsilon/2^n$ (small typo in book)
- 2) Since $\bigcup_{n=1}^{\infty} B_n \subseteq \bigcup_{n,k} C_{n,k}$,
$$P^*(\bigcup_{n=1}^{\infty} B_n) \leq \sum_{n,k} P(C_{n,k}) < \sum_n P^*(B_n) + \varepsilon.$$
- 3) As ε is an arbitrary positive constant, we must have
$$P^*(\bigcup_{n=1}^{\infty} B_n) \leq \sum_{n=1}^{\infty} P^*(B_n).$$

\mathcal{M} : The Measurable Sets

- Outer measure cannot always be a probability measure over **all** subsets of Ω (recall Proposition 1.2.6). Define a refined collection of subsets using P^* :

$$\mathcal{M} = \{A \subseteq \Omega : P^*(A \cap E) + P^*(A^c \cap E) = P^*(E) \text{ for all } E \subseteq \Omega\}$$

\mathcal{M} : The Measurable Sets

- Outer measure cannot always be a probability measure over **all** subsets of Ω (recall Proposition 1.2.6). Define a refined collection of subsets using P^* :

$$\mathcal{M} = \{A \subseteq \Omega : P^*(A \cap E) + P^*(A^c \cap E) = P^*(E) \text{ for all } E \subseteq \Omega\}$$

Lemma 6

We have the following results regarding \mathcal{M} :

- a) $\emptyset \in \mathcal{M}, \Omega \in \mathcal{M}$
- b) *If $A \in \mathcal{M}$, then $A^c \in \mathcal{M}$ (Closed under complement)*

\mathcal{M} : The Measurable Sets

- Outer measure cannot always be a probability measure over **all** subsets of Ω (recall Proposition 1.2.6). Define a refined collection of subsets using P^* :

$$\mathcal{M} = \{A \subseteq \Omega : P^*(A \cap E) + P^*(A^c \cap E) = P^*(E) \text{ for all } E \subseteq \Omega\}$$

Lemma 6

We have the following results regarding \mathcal{M} :

- a) $\emptyset \in \mathcal{M}, \Omega \in \mathcal{M}$*
- b) If $A \in \mathcal{M}$, then $A^c \in \mathcal{M}$ (Closed under complement)*

- Remark:** We often need to verify that a given set $A \in \mathcal{M}$. By the countable subadditivity of our outer measure, we always have $P^*(E) \leq P^*(A \cap E) + P^*(A^c \cap E)$ for all $E \subseteq \Omega$. If it's easier, we only need need to verify $P^*(E) \geq P^*(A \cap E) + P^*(A^c \cap E)$ for all $E \subseteq \Omega$. This would can be achieved by using the finite superadditivity of P .

\mathcal{M} and P^*

- We still need to show that \mathcal{M} is a σ -algebra and P^* is a proper probability measure over \mathcal{M} . A review of our results so far:

\mathcal{M} and P^*

- We still need to show that \mathcal{M} is a σ -algebra and P^* is a proper probability measure over \mathcal{M} . A review of our results so far:
- $\mathcal{M} = \{A : A \subseteq \Omega, P^*(A \cap E) + P^*(A^c \cap E) = P^*(E) \text{ for all } E \subseteq \Omega\}$. We already have $\emptyset \in \mathcal{M}, \Omega \in \mathcal{M}$ and \mathcal{M} is closed under complement. So we still need to prove that \mathcal{M} is closed under countable unions.

\mathcal{M} and P^*

- We still need to show that \mathcal{M} is a σ -algebra and P^* is a proper probability measure over \mathcal{M} . A review of our results so far:
- $\mathcal{M} = \{A : A \subseteq \Omega, P^*(A \cap E) + P^*(A^c \cap E) = P^*(E) \text{ for all } E \subseteq \Omega\}$. We already have $\emptyset \in \mathcal{M}, \Omega \in \mathcal{M}$ and \mathcal{M} is closed under complement. So we still need to prove that \mathcal{M} is closed under countable unions.
- Regarding P^* , for any $A \in \mathcal{M}$, first, $P^*(A) \geq 0$; second, in the definition of \mathcal{M} , by choosing $E = \Omega$, we have $P^*(A) = 1 - P^*(A^c)$. So we still need to show that, on \mathcal{M} , P^* is countably additive.

\mathcal{M} and P^*

- We still need to show that \mathcal{M} is a σ -algebra and P^* is a proper probability measure over \mathcal{M} . A review of our results so far:
- $\mathcal{M} = \{A : A \subseteq \Omega, P^*(A \cap E) + P^*(A^c \cap E) = P^*(E) \text{ for all } E \subseteq \Omega\}$. We already have $\emptyset \in \mathcal{M}, \Omega \in \mathcal{M}$ and \mathcal{M} is closed under complement. So we still need to prove that \mathcal{M} is closed under countable unions.
- Regarding P^* , for any $A \in \mathcal{M}$, first, $P^*(A) \geq 0$; second, in the definition of \mathcal{M} , by choosing $E = \Omega$, we have $P^*(A) = 1 - P^*(A^c)$. So we still need to show that, on \mathcal{M} , P^* is countably additive.
- Countable additivity of P^* is shown in the next lecture, as well as the fact that \mathcal{M} is a σ -field.