

## 4.4: The Gamma Distribution and Its Relatives

Taylor

University of Virginia

Normal distributions are probably the most popular but are defined for rvs that take on values on any part of the real line  $(-\infty, \infty)$ . Gamma rvs are defined for only positive numbers. Also, they include a lot of useful specific distributions.

# Definition

This is a special function; it isn't a pdf or cdf. We use it a lot whenever we work with integrals for gamma-ish pdfs.

## Definition

For  $\alpha > 0$ , the **gamma function**  $\Gamma(\alpha)$  is defined by

$$\Gamma(\alpha) = \int_0^{\infty} e^{-z} z^{\alpha-1} dz$$

Note:  $z$  is just a dummy variable. This is a function in  $\alpha$

# Properties

- 1  $\Gamma(\alpha) = (\alpha - 1)\Gamma(\alpha - 1)$  for  $\alpha > 1$
- 2  $\Gamma(\frac{1}{2}) = \sqrt{\pi}$

So it generalizes the factorial function for non-integral arguments, and it comes up with “special numbers” a lot.

Note: you can always just use wolfram alpha or some other symbolic computational thing to evaluate the gamma function

## Definition

A cts rv  $X$  is a **Gamma Distribution** if it has pdf

$$f(x; \alpha, \beta) = \frac{1}{\beta^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-\frac{x}{\beta}}$$

with  $x, \alpha$  and  $\beta$  all being positive.

it's MGF is

$$M_X(t) = (1 - \beta t)^{-\alpha}$$

It has simple-to-remember mean and variance

$$EX = \alpha\beta$$

$$VX = \alpha\beta^2$$

Also you can multiply gamma rvs by a constant and they're still gamma (you can't add constants though)

$$X \sim \text{Gamma}(\alpha, \beta) \rightarrow cX \sim \text{Gamma}(\alpha, c\beta)$$

# Proof

Sometimes “recognizing different gamma densities” is even easier than “recognizing gamma functions”

$$\begin{aligned} EX &= \int_0^{\infty} x \frac{1}{\beta^{\alpha} \Gamma(\alpha)} x^{\alpha-1} e^{-x/\beta} dx \\ &= \frac{1}{\beta^{\alpha} \Gamma(\alpha)} \int_0^{\infty} x^{(\alpha+1)-1} e^{-x/\beta} dx \\ &= \frac{1}{\beta^{\alpha} \Gamma(\alpha)} \frac{\beta^{\alpha+1} \Gamma(\alpha+1)}{1} \left[ \frac{1}{\beta^{\alpha+1} \Gamma(\alpha+1)} \int_0^{\infty} x^{(\alpha+1)-1} e^{-x/\beta} dx \right] \\ &= \frac{1}{\beta^{\alpha} \Gamma(\alpha)} \frac{\beta^{\alpha+1} \Gamma(\alpha+1)}{1} \\ &= \beta \frac{\Gamma(\alpha+1)}{\Gamma(\alpha)} \\ &= \beta \alpha \frac{\Gamma(\alpha)}{\Gamma(\alpha)} \end{aligned}$$

Let's start out with  $X \sim \text{Gamma}(\alpha, \beta)$ . We want to show that  $Y = cX$  follows a  $\text{Gamma}(\alpha, c\beta)$  distribution. This is easiest to do with mgfs..

$$M_Y(t) = E[e^{tY}] = E[e^{tcX}] = E[e^{(tc)X}] = M_X(tc) = (1 - \beta ct)^{-\alpha}$$



Another way to prove it:

$$F_Y(y) = P(Y \leq y) = P(cX \leq y) = P(X \leq y/c) = F_X(y/c)$$

So  $F_Y(y) = F_X(y/c)$ .

So  $f_Y(y) = f_X(y/c) \frac{1}{c}$  (by the chain rule)

## Some special cases

A **chi-squared distribution** with parameter  $\nu$  is the same as a  $\text{Gamma}(\nu/2, 2)$

So its density is  $f(x; \nu) = \frac{1}{2^{\nu/2}\Gamma(\nu/2)} e^{-x/2} x^{\nu/2-1}$

A **exponential distribution** with parameter  $\lambda$  is the same as a  $\text{Gamma}(1, \frac{1}{\lambda})$

So its density is  $f(y; \lambda) = \lambda e^{-y\lambda}$

# Exercise

This is similar to example 4.30 on page 199. Let  $T \sim \text{Exponential}(\lambda)$  denote the waiting time until a call arrives (in days). Find its cdf. Then use its cdf to find the probability that we wait more than 2 days for a call (use  $\lambda = .5$ ).

# Exercise

$$P(T \leq t) = \int_0^t \lambda e^{-\lambda x} dx = [-e^{-\lambda x}]_{x=0}^{x=t} = -e^{-\lambda t} + 1$$

So  $F_T(t) = 1 - e^{-\lambda t}$ . Then

$$P(T > 2) = 1 - P(T \leq 2) = 1 - [1 - e^{-\lambda 2}] = e^{-\lambda 2}$$

So if we use  $\lambda = \frac{1}{2}$ , then  $P(T > 2) = \frac{1}{e}$