

# Unit 19: Spectral Density for Causal ARMA Processes

Taylor R. Brown

Department of Statistics, University of Virginia

Fall 2020

## Readings for Unit 19

Textbook chapter 4.2 (from page 176).

# Last Unit

- 1 Spectral Density: Fourier Transformation of Autocovariance.
- 2 Properties of Spectral Density.

# Motivation

We generalize the spectral density for causal ARMA processes.

# 1 Autocovariance Generating Function

## 2 Rational Spectrum

# Causal ARMA Process

A zero-mean causal ARMA process can be written as

$$x_t = \sum_{j=0}^{\infty} \psi_j w_{t-j} = \psi(B)w_t.$$

where  $\psi(B) = \sum_{j=0}^{\infty} \psi_j B^j$ . Therefore, the autocovariance function is

$$\begin{aligned} \gamma(h) &= E(x_t x_{t+h}) \\ &= E\left(\sum_{j=0}^{\infty} \psi_j w_{t-j} \sum_{i=0}^{\infty} \psi_i w_{t+h-i}\right) \\ &= \sigma_w^2 \sum_{j=0}^{\infty} \psi_j \psi_{j+h}. \end{aligned}$$

# Spectral Density of Causal ARMA models

Therefore the spectral density for a causal ARMA process can be expressed as

$$\begin{aligned} f(\omega) &= \sum_{h=-\infty}^{\infty} \gamma(h) e^{-2\pi i \omega h} \\ &= \\ &= \\ &= \end{aligned} \tag{1}$$

# Example: $MA(q)$



# Example: AR(p)

1 Autocovariance Generating Function

2 Rational Spectrum

# Autocovariance Generating Function: ARMA(p,q)

From (1), we have

$$f(\omega) = \sigma_w^2 \left| \frac{\theta(e^{-2\pi i\omega})}{\phi(e^{-2\pi i\omega})} \right|^2.$$

This is also called the **rational spectrum** of an ARMA(p,q).

# Zeroes and Poles

Recall that every degree  $p$  polynomial  $a(z)$  can be factorized as

$$a(z) = a_p(z - z_1)(z - z_2) \cdots (z - z_p)$$

where  $z_1, \dots, z_p \in \mathbb{C}$  are the roots.

# Zeroes and Poles

For the MA and AR polynomials,

$$\theta(z) = \theta_q(z - z_1)(z - z_2) \cdots (z - z_q)$$

and

$$\phi(z) = \phi_p(z - p_1)(z - p_2) \cdots (z - p_p).$$

$z_1, \dots, z_q$  are called **zeroes**.  $p_1, \dots, p_p$  are called **poles**.

# Rational Spectrum

Therefore, the rational spectrum as expressed in (1) can be re-written as


$$\begin{aligned} f(\omega) &= \sigma_w^2 \left| \frac{\theta_q \prod_{j=1}^q (e^{-2\pi i\omega} - z_j)}{\phi_p \prod_{j=1}^p (e^{-2\pi i\omega} - p_j)} \right|^2 \\ &= \sigma_w^2 \frac{\theta_q^2 \prod_{j=1}^q |e^{-2\pi i\omega} - z_j|^2}{\phi_p^2 \prod_{j=1}^p |e^{-2\pi i\omega} - p_j|^2}. \end{aligned} \quad (2)$$

# Examples: AR(1)

Consider  $\phi > 0$ .

# Examples: AR(1)

**Power spectrum of AR(1) with  $\phi=0.9$**






# Examples: MA(1)

Consider  $\theta > 0$ .

## Examples: MA(1)

**Power spectrum of MA(1) with  $\theta=0.5$**



## Examples: AR(2)

Suppose we have the following AR(2) model:

$x_t = x_{t-1} - 0.9x_{t-2} + w_t$ , where  $\sigma_w^2 = 1$ . The roots (poles) of the AR polynomial  $\phi(z) = 0.9z^2 - z + 1$  are  $p_1, p_2 = 0.555 \pm i0.8958$ . Note that:

## Examples: AR(2)

Using the representation (2), the spectral density is

$$f(\omega) = \frac{1}{\phi_2^2 |e^{-2\pi i\omega} - p_1|^2 |e^{-2\pi i\omega} - p_2|^2}.$$

The peaks of the spectral density for this process occurs when  $e^{-2\pi i\omega}$  is near  $1.054e^{-2\pi i0.16165}$ .

# Examples: AR(2)

**Power spectrum of AR(2) with  $\phi = c(-1, 0.9)$**

