


P. 2-11



Since $OA = OB = OC$, we know $\angle OAC = \angle OCA$ and $\angle OBC = \angle OCB$. And since ABC is an triangle, $\angle OAC + \angle OBC + \angle ACB = \pi$, we can simply find that $\angle ACB = \pi/2$, so that an angle inscribed in a semicircle is a right angle.

P. 2-17

a)

$$|\mathbf{E}| = |\mathbf{a}_R(25/R^2)| = |\mathbf{a}_R| \cdot \frac{25}{3^2 + 4^2 + 5^2} = \frac{1}{2}.$$

$$E_x = |\mathbf{E}| \cdot \frac{x}{R} = \frac{1}{2} \cdot \frac{-3}{\sqrt{3^2 + 4^2 + 5^2}} = -\frac{3\sqrt{2}}{20}.$$

b)

$$\cos \theta = \frac{\mathbf{E} \cdot \mathbf{B}}{|\mathbf{E}||\mathbf{B}|} = \frac{-3 \cdot 2 + 4 \cdot -2 - 5 \cdot 1}{\sqrt{3^2 + 4^2 + 5^2} \cdot \sqrt{2^2 + 2^2 + 1^2}} = -\frac{19\sqrt{2}}{30},$$

$$\theta = \arccos -\frac{19\sqrt{2}}{30} \approx 2.681 \text{ rad.}$$

P. 2-21

$$\int_{P_1}^{P_2} \mathbf{E} \cdot d\mathbf{l} = \int_{P_1}^{P_2} (\mathbf{a}_x y + \mathbf{a}_y x)(\mathbf{a}_x dx + \mathbf{a}_y dy) = \int_{P_1}^{P_2} (y dx + x dy)$$

a)

$$\int_{P_1}^{P_2} \mathbf{E} \cdot d\mathbf{l} = \int_{P_1}^{P_2} (y \cdot 4y dy + 2y^2 \cdot dy) = \int_1^2 6y^2 dy = 14.$$

b)

$$x = 6y - 4,$$

$$\int_{P_1}^{P_2} \mathbf{E} \cdot d\mathbf{l} = \int_{P_1}^{P_2} [y \cdot 6dy + (6y - 4) \cdot dy] = \int_1^2 (12y - 4) dy = 14.$$

P. 2-26

a)

$$\nabla \cdot f_1(\mathbf{R}) = \frac{1}{R^2} \cdot \frac{\partial}{\partial R}(R^n \cdot R^2) = (n+2)R^{n-1}.$$

b)

$$\nabla \cdot f_2(\mathbf{R}) = \frac{1}{R^2} \cdot \frac{\partial}{\partial R}(k/R^2 \cdot R^2) = 0.$$

P. 2-29

$$\nabla \cdot \mathbf{A} = \frac{1}{r} \cdot \frac{\partial}{\partial r}(r^2 \cdot r) + \frac{\partial}{\partial z}(2z) = 3r + 2.$$

$$\int_V \nabla \cdot \mathbf{A} dV = \int_V (3r + 2) dV = \int_0^4 \int_0^{2\pi} \int_0^5 (3r + 2) r dr d\theta dz = 1200\pi.$$

$$\oint_S \mathbf{A} dS = \int_0^4 \int_0^{2\pi} (\mathbf{a}_r r^2) r d\theta dz \mathbf{a}_r + \int_0^{2\pi} \int_0^5 2z \mathbf{a}_z r dr d\theta \mathbf{a}_z = 1000\pi + 200\pi = 1200\pi.$$

So

$$\int_V \nabla \cdot \mathbf{A} dV = \oint_S \mathbf{A} dS.$$

P. 2-33

$$\nabla \cdot (\mathbf{E} \times \mathbf{H}) = \frac{\partial}{\partial x}(E_y H_z - E_z H_y) + \frac{\partial}{\partial y}(E_z H_x - E_x H_z) + \frac{\partial}{\partial z}(E_x H_y - E_y H_x),$$

$$\mathbf{H} \cdot (\nabla \times \mathbf{E}) = H_x \left(\frac{\partial E_z}{\partial y} - \frac{\partial E_y}{\partial z} \right) + H_y \left(\frac{\partial E_x}{\partial z} - \frac{\partial E_z}{\partial x} \right) + H_z \left(\frac{\partial E_y}{\partial x} - \frac{\partial E_x}{\partial y} \right),$$

$$\mathbf{E} \cdot (\nabla \times \mathbf{H}) = E_x \left(\frac{\partial H_z}{\partial y} - \frac{\partial H_y}{\partial z} \right) + E_y \left(\frac{\partial H_x}{\partial z} - \frac{\partial H_z}{\partial x} \right) + E_z \left(\frac{\partial H_y}{\partial x} - \frac{\partial H_x}{\partial y} \right).$$

Since

$$\frac{\partial}{\partial x} AB = A \frac{\partial B}{\partial x} + B \frac{\partial A}{\partial x},$$

we can find that

$$\nabla \cdot (\mathbf{E} \times \mathbf{H}) = \mathbf{H} \cdot (\nabla \times \mathbf{E}) - \mathbf{E} \cdot (\nabla \times \mathbf{H}).$$

P. 2-35

$$\Delta s_u = R^2 \sin \theta \Delta \theta \Delta \phi,$$

$$\oint_{C_u} \mathbf{A} \cdot d\mathbf{l} = A_\theta \cdot (R, \theta, \phi - \Delta \phi / 2) R \Delta \theta - A_\theta \cdot (R, \theta, \phi + \Delta \phi / 2) R \Delta \theta +$$

$$A_\phi \cdot (R, \theta + \Delta \theta / 2, \phi) R \Delta \phi \sin(\theta + \Delta \theta / 2) - A_\phi \cdot (R, \theta - \Delta \theta / 2, \phi) R \Delta \phi \sin(\theta - \Delta \theta / 2)$$

$$= -\frac{\partial A_\theta}{\partial \phi} R \Delta \phi \Delta \theta + \frac{\partial A_\phi \sin \theta}{\partial \theta} R \Delta \phi \Delta \theta.$$

$$(\nabla \times \mathbf{A})_u = \lim_{\Delta s_u \rightarrow 0} \frac{1}{\Delta s_u} \oint_{C_u} \mathbf{A} \cdot d\mathbf{l} = \lim_{\Delta s_u \rightarrow 0} \frac{1}{R^2 \sin \theta \Delta \theta \Delta \phi} \cdot \left(-\frac{\partial A_\theta}{\partial \phi} + \frac{\partial A_\phi \sin \theta}{\partial \theta} \right) R \Delta \phi \Delta \theta$$

$$= \frac{1}{R \sin \theta} \cdot \left(-\frac{\partial A_\theta}{\partial \phi} + \frac{\partial A_\phi \sin \theta}{\partial \theta} \right).$$

P. 2-39

a)

$$\nabla \times \mathbf{F} = \mathbf{a}_x(c_3 + 3) + \mathbf{a}_y(c_1 - 1) + \mathbf{a}_z c_2 = \mathbf{0},$$

$$c_1 = 1, c_2 = 0, c_3 = -3.$$

b)

$$\nabla \cdot \mathbf{F} = 1 + 0 + c_4 = 0,$$


$$c_4 = -1.$$

c)

$$\mathbf{F} = -\nabla V = \mathbf{a}_x(x + z) + \mathbf{a}_y(-3z) + \mathbf{a}_z(x - 3y - z),$$

$$V = -\frac{1}{2}x^2 - xz + 3yz + \frac{1}{2}z^2 + C.$$

P. 3-8




$$dQ = bd\theta\rho_l,$$

$$dE = \frac{dQ}{4\pi\epsilon_0 b^2} = \frac{\rho_l}{4\pi\epsilon_0 b} d\theta,$$

$$|\mathbf{E}| = \int_0^\pi dE \sin\theta = \frac{\rho_l}{4\pi\epsilon_0 b} \int_0^\pi \sin\theta d\theta = \frac{\rho_l}{2\pi\epsilon_0 b}.$$

The direction is downwards.

P. 3-9



$$dQ = \rho dl,$$

$$dE = \frac{dQ}{4\pi\epsilon_0(l^2 + L^2/12)} = \frac{\rho}{4\pi\epsilon_0} \cdot \frac{dl}{l^2 + L^2/12},$$

$$|\mathbf{E}_1| = \int_{-L/2}^{L/2} dE \cdot \sqrt{\frac{L^2/12}{l^2 + L^2/12}} = \frac{3\rho_{l1}}{2\pi\epsilon_0 L},$$

$$|\mathbf{E}_2| = |\mathbf{E}_3| = \frac{1}{2}|\mathbf{E}_1| = \frac{3\rho_{l1}}{4\pi\epsilon_0 L}.$$

$$|\mathbf{E}| = |\mathbf{E}_1| - \frac{1}{2}|\mathbf{E}_2| - \frac{1}{2}|\mathbf{E}_3| = \frac{3\rho_{l1}}{4\pi\epsilon_0 L}.$$

The direction is upwards.

P. 3-12

a) For $0 < r < a$, $\mathbf{E} = \mathbf{0}$.

For $a < r < b$,

$$2\pi aL \cdot \frac{\rho_{sa}}{\epsilon_0} = 2\pi rL \cdot E,$$

$$\mathbf{E} = \frac{a\rho_{sa}}{\epsilon_0 r} \mathbf{a}_r.$$

For $b < r$,

$$2\pi aL \cdot \frac{\rho_{sa}}{\epsilon_0} + 2\pi bL \cdot \frac{\rho_{sb}}{\epsilon_0} = 2\pi rL \cdot E,$$

$$\mathbf{E} = \frac{a\rho_{sa} + b\rho_{sb}}{\epsilon_0 r} \mathbf{a}_r.$$

b)

$$\frac{a\rho_{sa} + b\rho_{sb}}{\epsilon_0 r} = 0,$$

$$a = -\frac{\rho_{sb}}{\rho_{sa}} b.$$

P. 3-13

$$W = - \int \mathbf{E} q d\mathbf{l} = 2\mu \int (\mathbf{a}_x y + \mathbf{a}_y x)(\mathbf{a}_x dx + \mathbf{a}_y dy) = 2\mu \int y dx + x dy.$$

a)

$$W = 2\mu \int y dx + x dy = 2\mu C \int_2^8 4y^2 dy + 2y^2 dy = 28\mu.$$

b)

$$x = 6y - 4,$$

$$W = 2\mu \int y dx + x dy = 2\mu C \int_2^8 6y dy + (6y - 4) dy = 28\mu.$$

P. 3-16

a)

$$dQ = \rho_l dl,$$

$$dV = \frac{dQ}{4\pi\epsilon_0 \sqrt{l^2 + y^2}} = \frac{\rho_l}{4\pi\epsilon_0} \cdot \frac{dl}{\sqrt{l^2 + y^2}},$$

$$V = \int_{-L/2}^{L/2} dV = \frac{\rho_l}{4\pi\epsilon_0} \int_{-L/2}^{L/2} \frac{1}{\sqrt{l^2 + y^2}} dl = \frac{\rho_l}{2\pi\epsilon_0} \operatorname{arcsinh} \frac{L}{2y}.$$

b)

$$dE = \frac{dQ}{4\pi\epsilon_0(l^2 + y^2)} = \frac{\rho_l}{4\pi\epsilon_0} \cdot \frac{dl}{l^2 + y^2},$$

$$\mathbf{E} = \mathbf{a}_y \int_{-L/2}^{L/2} dE \cdot \sqrt{\frac{y^2}{l^2 + y^2}} = \frac{\rho_l}{2\pi\epsilon_0} \cdot \frac{y}{\sqrt{L^2 + 4y^2}} \mathbf{a}_y.$$

c)

$$-\nabla V = -\frac{dV}{dy} \mathbf{a}_y = \frac{\rho_l}{4\pi\epsilon_0} \cdot \frac{L}{\sqrt{L^2 + 4y^2}} \mathbf{a}_y = \mathbf{E}.$$

P. 3-19

$$dQ = \frac{Q}{h} dz,$$

$$dV = \frac{dQ}{4\pi\epsilon_0 \sqrt{b^2 + z^2}} = \frac{Q}{4\pi\epsilon_0 h \sqrt{b^2 + z^2}} dz.$$

a)

$$V = \int_{z-h}^z dV = \frac{Q}{4\pi\epsilon_0 h} \int_{z-h}^z \frac{1}{\sqrt{b^2 + z^2}} dz = \frac{Q}{4\pi\epsilon_0 h} \left[\operatorname{arcsinh} \frac{z}{b} - \operatorname{arcsinh} \frac{z-h}{b} \right].$$

$$\mathbf{E} = -\nabla V = -\frac{dV}{dz} \mathbf{a}_z = -\frac{Q}{4\pi\epsilon_0 h} \left[\frac{1}{\sqrt{b^2 + z^2}} - \frac{1}{\sqrt{b^2 + (z-h)^2}} \right] \mathbf{a}_z.$$

b)

$$V = \int_0^z dV + \int_0^{h-z} dV = \frac{Q}{4\pi\epsilon_0 h} \left[\operatorname{arcsinh} \frac{z}{b} + \operatorname{arcsinh} \frac{h-z}{b} \right].$$

$$\mathbf{E} = -\nabla V = -\frac{dV}{dz} \mathbf{a}_z = -\frac{Q}{4\pi\epsilon_0 h} \left[\frac{1}{\sqrt{b^2 + z^2}} - \frac{1}{\sqrt{b^2 + (h-z)^2}} \right] \mathbf{a}_z.$$

P. 3-22

a)

$$\rho_{ps} = \mathbf{P} \cdot \mathbf{a}_n|_{n=L/2} = \frac{1}{2} P_0 L.$$

$$\rho_p = -\nabla \cdot \mathbf{P} = -3P_0.$$

b)

$$Q_s = \oint_S \rho_{ps} dS = \frac{1}{2} P_0 L \cdot 6L^2 = 3P_0 L^3,$$

$$Q_v = \int_v \rho_p dV = \int_{-L/2}^{L/2} \int_{-L/2}^{L/2} \int_{-L/2}^{L/2} \rho_p dz dy dx = -3P_0 L^3,$$

$$Q = Q_s + Q_v = 0.$$

P. 3-23

Let $\mathbf{P} = \mathbf{a}_p P_0$, $\theta = \langle \mathbf{P}, \mathbf{a}_n \rangle$,

$$\rho_{ps}(\theta) = \mathbf{P} \cdot \mathbf{a}_n = P_0 \cos \theta,$$

$$dE_\theta = dv \cdot \frac{\rho_{ps}}{4\pi\epsilon_0 R^2} \cdot \cos \theta = 2\pi R^2 \sin \theta d\theta \cdot \frac{P_0 \cos \theta}{4\pi\epsilon_0 R^2} \cdot \cos \theta = \frac{P_0 \sin \theta \cos \theta^2}{2\epsilon_0} d\theta.$$

$$|\mathbf{E}| = \int dE_\theta = \int_0^\pi \frac{P_0 \sin \theta \cos \theta^2}{2\epsilon_0} d\theta = \frac{P_0}{3\epsilon_0},$$

$$\mathbf{E} = \mathbf{a}_p \frac{P_0}{3\epsilon_0} = \frac{\mathbf{P}}{3\epsilon_0}.$$

P. 3-25

$$E_{2t} = E_{1t} = \mathbf{a}_x 2y - \mathbf{a}_y 3x.$$

Since $\rho_s = 0$,

$$\epsilon_{r1} E_{1n} = \epsilon_{r2} E_{2n},$$

$$E_{2n} = \frac{\epsilon_{r1}}{\epsilon_{r2}} E_{1n} = \frac{2}{3} \cdot \mathbf{a}_z 5 = \mathbf{a}_z \frac{10}{3}.$$

$$E_2 = E_{2t} + E_{2n} = \mathbf{a}_x 2y - \mathbf{a}_y 3x + \mathbf{a}_z \frac{10}{3}.$$

$$\mathbf{D}_2 = \epsilon_2 \mathbf{E}_2 = 3\epsilon_0 \left(\mathbf{a}_x 2y - \mathbf{a}_y 3x + \mathbf{a}_z \frac{10}{3} \right).$$

P. 3-28

Obviously, \mathbf{E}_3 is parallel to \mathbf{E}_2 , so we only need to find ϵ_{r2} so that \mathbf{E}_2 is parallel to the x-axis.

$$E_{1t} = E_{2t} = -3.$$

$$\epsilon_{r1} E_{1n} = \epsilon_{r2} E_{2n},$$

$$E_{2n} = \frac{\varepsilon_{r1}}{\varepsilon_{r2}} E_{1n} = \frac{1}{\varepsilon_{r2}} \cdot 5 = \frac{5}{\varepsilon_{r2}}.$$

$$E_{2t} \cos \theta + E_{2n} \cos \theta = 0,$$

$$-3 + \frac{5}{\varepsilon_{r2}} = 0,$$

$$\varepsilon_{r2} = \frac{5}{3}.$$

P. 3-32

$$C_1 = \frac{2\pi\varepsilon L}{\ln \frac{b}{r_i}} = \frac{2\pi\varepsilon_0\varepsilon_{r1} L}{\ln \frac{b}{r_i}},$$

$$C_2 = \frac{2\pi\varepsilon L}{\ln \frac{r_o}{b}} = \frac{2\pi\varepsilon_0\varepsilon_{r2} L}{\ln \frac{r_o}{b}},$$

$$C = \frac{1}{\frac{1}{C_1} + \frac{1}{C_2}} = \frac{2\pi\varepsilon_0 L}{\frac{1}{\varepsilon_{r1}} \ln \frac{b}{r_i} + \frac{1}{\varepsilon_{r3}} \ln \frac{r_o}{b}},$$

$$\frac{C}{L} = \frac{2\pi\varepsilon_0}{\frac{1}{\varepsilon_{r1}} \ln \frac{b}{r_i} + \frac{1}{\varepsilon_{r3}} \ln \frac{r_o}{b}}.$$

P. 3-43

$$dW_e = Vdq = \frac{q}{C} dq,$$

$$W_e = \int dW_e = \int_0^Q \frac{q}{C} dq = \frac{Q^2}{2C}.$$

Since $Q = CV$, we can also get

$$W_e = \frac{1}{2} CV^2 = \frac{1}{2} QV.$$