

VE320 Homework 1

Liu Yihao 515370910207

Ex. 1

$$E = hv = \frac{hc}{\lambda} \implies \lambda = \frac{hc}{E}$$

For gold,

$$\lambda = \frac{hc}{E} = \frac{6.63 \times 10^{-34} \text{ J} \cdot \text{s} \cdot 3 \times 10^8 \text{ m/s}}{4.90 \text{ eV}} = 2.54 \times 10^{-7} \text{ m}$$

For Cesium,

$$\lambda = \frac{hc}{E} = \frac{6.63 \times 10^{-34} \text{ J} \cdot \text{s} \cdot 3 \times 10^8 \text{ m/s}}{1.90 \text{ eV}} = 6.54 \times 10^{-7} \text{ m}$$

Ex. 2

$$E_{avg} = \frac{3}{2}kT = 1.5 \cdot 0.0259 \text{ eV} = 3.885 \times 10^{-2} \text{ eV}$$

$$p_{avg} = \sqrt{2mE_{avg}} = \sqrt{2 \cdot 9.11 \times 10^{-31} \text{ kg} \cdot 3.885 \times 10^{-2} \text{ eV}} = 1.064 \times 10^{-25} \text{ kg} \cdot \text{m/s}$$

$$\lambda = \frac{h}{p} = \frac{6.63 \times 10^{-34}}{1.064 \times 10^{-25} \text{ kg} \cdot \text{m/s}} = 6.23 \times 10^{-9} \text{ m} = 62.3 \text{ \AA}$$

Ex. 3

(a)

$$p = \frac{h}{\lambda} = \frac{6.63 \times 10^{-34} \text{ J} \cdot \text{s}}{8.5 \times 10^{-9} \text{ m}} = 7.80 \times 10^{-26} \text{ kg} \cdot \text{m/s}$$

$$v = \frac{p}{m} = \frac{7.80 \times 10^{-26} \text{ kg} \cdot \text{m/s}}{9.11 \times 10^{-31} \text{ kg}} = 8.56 \times 10^4 \text{ m/s}$$

$$E = \frac{1}{2}mv^2 = \frac{1}{2} \cdot 9.11 \times 10^{-31} \text{ kg} \cdot (8.56 \times 10^4 \text{ m/s})^2 = 3.34 \times 10^{-21} \text{ J} = 2.09 \times 10^{-2} \text{ eV}$$

(b)

$$v = 8 \times 10^5 \text{ cm/s} = 8 \times 10^2 \text{ m/s}$$

$$E = \frac{1}{2}mv^2 = \frac{1}{2} \cdot 9.11 \times 10^{-31} \text{ kg} \cdot (8 \times 10^2 \text{ m/s})^2 = 2.92 \times 10^{-25} \text{ J} = 1.83 \times 10^{-4} \text{ eV}$$

$$p = mv = 9.11 \times 10^{-31} \text{ kg} \cdot 8 \times 10^2 \text{ m/s} = 7.29 \times 10^{-27} \text{ kg} \cdot \text{m/s}$$

$$\lambda = \frac{h}{p} = \frac{6.63 \times 10^{-34} \text{ J} \cdot \text{s}}{7.29 \times 10^{-27} \text{ kg} \cdot \text{m/s}} = 9.09 \times 10^{-8} \text{ m} = 909 \text{ \AA}$$

Ex. 4

$$\int_{-1}^{+3} |\Phi(x, t)|^2 dx = \int_{-1}^{+3} A^2 \cos^2\left(\frac{\pi x}{2}\right) dx = A^2 \cdot \left(\frac{x}{2} + \frac{\sin(\pi x)}{2\pi}\right) \Big|_{-1}^{+3} = 2A^2 = 1$$

$$A = \pm \frac{\sqrt{2}}{2}$$

Ex. 5

(a) (i)

$$v_p = \frac{\omega}{k} = \frac{8 \times 10^{12} \text{ rad/s}}{8 \times 10^8 \text{ m}^{-1}} = 10^4 \text{ m/s}$$

$$\lambda = \frac{2\pi}{k} = \frac{2 \cdot 3.14}{8 \times 10^8 \text{ m}^{-1}} = 7.85 \times 10^{-9} \text{ m} = 78.5 \text{ \AA}$$

(ii)

$$p = \hbar k = 1.05 \times 10^{-34} \text{ J} \cdot \text{s} \cdot 8 \times 10^8 \text{ m}^{-1} = 8.40 \times 10^{-26} \text{ kg} \cdot \text{m/s}$$

$$v = \frac{p}{m} = \frac{8.40 \times 10^{-26} \text{ kg} \cdot \text{m/s}}{9.11 \times 10^{-31} \text{ kg}} = 9.22 \times 10^4 \text{ m/s}$$

$$E = \frac{1}{2}mv^2 = \frac{1}{2} \cdot 9.11 \times 10^{-31} \text{ kg} \cdot (9.22 \times 10^4 \text{ m/s})^2 = 3.87 \times 10^{-21} \text{ J}$$

(b) (i)

$$v_p = \frac{\omega}{k} = \frac{1.5 \times 10^{12} \text{ rad/s}}{-1.5 \times 10^9 \text{ m}^{-1}} = -10^4 \text{ m/s}$$

$$\lambda = \frac{2\pi}{|k|} = \frac{2 \cdot 3.14}{1.5 \times 10^9 \text{ m}^{-1}} = 4.19 \times 10^{-9} \text{ m} = 41.9 \text{ \AA}$$

(ii)

$$p = \hbar|k| = 1.05 \times 10^{-34} \text{ J} \cdot \text{s} \cdot 1.5 \times 10^9 \text{ m}^{-1} = 1.58 \times 10^{-25} \text{ kg} \cdot \text{m/s}$$

$$v = \frac{p}{m} = \frac{1.58 \times 10^{-25} \text{ kg} \cdot \text{m/s}}{9.11 \times 10^{-31} \text{ kg}} = 1.73 \times 10^5 \text{ m/s}$$

$$E = \frac{1}{2}mv^2 = \frac{1}{2} \cdot 9.11 \times 10^{-31} \text{ kg} \cdot (1.73 \times 10^5 \text{ m/s})^2 = 1.36 \times 10^{-20} \text{ J}$$

Ex. 6

(a)

$$E_n = \frac{\hbar^2 n^2 \pi^2}{2ma^2} = \frac{\hbar^2 n^2}{8ma^2} = n^2 \cdot \frac{(6.63 \times 10^{-43} \text{ J} \cdot \text{s})}{8 \cdot 9.11 \times 10^{-31} \text{ kg} \cdot (10^{-9} \text{ m})^2} = n^2 \cdot 6.03 \times 10^{-20} \text{ J} = 0.377n^2 \text{ eV}$$

$$E_1 = 0.377 \text{ eV}$$

$$E_2 = 1.508 \text{ eV}$$

$$E_3 = 3.393 \text{ eV}$$

(b)

$$\Delta E = 5 \cdot 6.03 \times 10^{-20} \text{ J} = 3.015 \times 10^{-19} \text{ J}$$

$$\lambda = \frac{hc}{\Delta E} = \frac{6.63 \times 10^{-34} \text{ J} \cdot \text{s} \cdot 3 \times 10^8 \text{ m/s}}{3.015 \times 10^{-19} \text{ J}} = 6.60 \times 10^{-7} \text{ m}$$

Ex. 7

$$\psi_1(x) = A_1 e^{jk_1 x} + B_1 e^{-jk_1 x}$$

$$\psi_2(x) = A_2 e^{jk_2 x} + B_2 e^{-jk_2 x}$$

$$\psi_3(x) = A_3 e^{jk_3 x}$$

At $x = 0$, the function must be continuous, so $\psi_1(0) = \psi_2(0)$, $\frac{\partial \psi_1}{\partial x}(0) = \frac{\partial \psi_2}{\partial x}(0)$,

$$A_1 + B_1 = A_2 + B_2$$

$$k_1 A_1 - k_1 B_1 = k_2 A_2 - k_2 B_2$$

At $x = a$, the function must be continuous, so $\psi_2(a) = \psi_3(a)$, $\frac{\partial \psi_2}{\partial x}(a) = \frac{\partial \psi_3}{\partial x}(a)$,

$$A_2 e^{jk_2 a} + B_2 e^{-jk_2 a} = A_3 e^{jk_3 a}$$

$$k_2 A_2 e^{jk_2 a} - k_2 B_2 e^{-jk_2 a} = k_3 A_3 e^{jk_3 a}$$

Since $k_2 a = 2n\pi$, so $e^{jk_2 a} = e^{-jk_2 a} = 1$, we get

$$A_2 + B_2 = A_3 e^{jk_3 a}$$

$$k_2 A_2 - k_2 B_2 = k_3 A_3 e^{jk_3 a}$$

Use the equations, we can get

$$A_1 + B_1 = A_3 e^{jk_3 a}$$

$$A_1 - B_1 = \frac{k_3}{k_1} A_3 e^{jk_3 a}$$

$$2A_1 = \frac{k_3 + k_1}{k_1} A_3 e^{jk_3 a}$$

$$\frac{A_3 \cdot A_3^*}{A_1 \cdot A_1^*} = \frac{4k_1^2}{(k_1 + k_3)^2}$$

So the transmission coefficient is

$$T = \frac{v_t \cdot A_3 \cdot A_3^*}{v_i \cdot A_1 \cdot A_1^*} = \frac{k_3 \cdot A_3 \cdot A_3^*}{k_1 \cdot A_1 \cdot A_1^*} = \frac{4k_1 k_3}{(k_1 + k_3)^2}$$

Ex. 8

(a) For Region I,

$$\frac{\partial^2 \psi_1(x)}{\partial x^2} - \frac{2m(V_0 - E)}{\hbar^2} \psi_1(x) = 0$$

$$\psi_1(x) = A_1 e^{k_1 x}, k_1 = \sqrt{\frac{2m(V_0 - E)}{\hbar^2}}$$

For Region II,

$$\frac{\partial^2 \psi_2(x)}{\partial x^2} + \frac{2mE}{\hbar^2} \psi_2(x) = 0$$

$$\psi_2(x) = A_2 \sin(k_2 x) + B_2 \cos(k_2 x), k_2 = \sqrt{\frac{2mE}{\hbar^2}}$$

For Region III,

$$\psi_3(x) = 0$$

(b) At $x = 0$, the function must be continuous, so $\psi_1(0) = \psi_2(0)$, $\frac{\partial\psi_1}{\partial x}(0) = \frac{\partial\psi_2}{\partial x}(0)$,

$$A_1 = B_2$$

$$k_1 A_1 = k_2 A_2$$

which means

$$\frac{A_1}{A_2} = \frac{k_2}{k_1}$$

At $x = a$, the function must be continuous, so $\psi_2(a) = \psi_3(a)$, $\frac{\partial\psi_2}{\partial x}(a) = \frac{\partial\psi_3}{\partial x}(a)$,

$$A_2 \sin(k_2 a) + B_2 \cos(k_2 a) = 0$$

$$A_2 \cos(k_2 a) - B_2 \sin(k_2 a) = 0$$

which means

$$\frac{B_2}{A_2} = -\tan(k_2 a)$$

(c)

$$\frac{A_1}{A_2} = \frac{k_2}{k_1} = -\tan(k_2 a)$$

$$\sqrt{\frac{2mE}{2m(V_0 - E)}} = -\tan\left(\sqrt{\frac{2mE}{\hbar^2}}\right)$$

This equation can be solved so that we can find the exact values of E , so the energy levels of the electrons are quantized.