

Dietary Reference Intakes (DRI)

The Dietary Reference Intakes (DRI) include two sets of values that serve as goals for nutrient intake—Recommended Dietary Allowances (RDA) and Adequate Intakes (AI). The RDA reflect the average daily amount of a nutrient considered adequate to meet the needs of most healthy people. If there is insufficient evidence to determine an RDA, an AI is set. AI are more tentative than RDA, but both may be used as goals for nutrient intakes. (Chapter 2 provides more details.)

Estimated Energy Requirements (EER), Recommended Dietary Allowances (RDA), and Adequate Intakes (AI) for Water, Energy, and the Energy Nutrients

AGE (YR)	REFERENCE BMI (kg/m ²)	REFERENCE HEIGHT cm (in)	REFERENCE WEIGHT kg (lb)	WATER ^a AI (U/day)	ENERGY EER ^b (Cal/day)	CARBOHYDRATE RDA (g/day)	TOTAL FIBER AI (g/day)	TOTAL FAT AI (g/day)	LINOLEIC ACID ^c AI (g/day)	LINOLENIC ACID ^c AI (g/day)	PROTEIN RDA (g/day) ^d	PROTEIN RDA (g/kg/day)
MALES												
0–0.5	—	62 (24)	6 (13)	0.7 ^e	570	60	—	31	4.4	0.5	9.1	1.52
0.5–1	—	71 (28)	9 (20)	0.8 ^f	743	95	—	30	4.6	0.5	11	1.20
1–3 ^g	—	86 (34)	12 (27)	1.3	1046	130	19	—	7	0.7	13	1.05
4–8 ^g	15.3	115 (45)	20 (44)	1.7	1742	130	25	—	10	0.9	19	0.95
9–13	17.2	144 (57)	36 (79)	2.4	2279	130	31	—	12	1.2	34	0.95
14–18	20.5	174 (68)	61 (134)	3.3	3152	130	38	—	16	1.6	52	0.85
19–30	22.5	177 (70) ⁱ	70 (154) ^j	3.7	3067 ^k	130	38	—	17	1.6	56	0.80
31–50	22.5 ^l	177 (70) ⁱ	70 (154) ^j	3.7	3067 ^k	130	30	—	14	1.6	56	0.80
>50	22.5 ^l	177 (70) ⁱ	70 (154) ^j	3.7	3067 ^k	130	—	—	—	—	—	—
FEMALES												
0–0.5	—	62 (24)	6 (13)	0.7 ^e	520	60	—	31	4.4	0.5	9.1	1.52
0.5–1	—	71 (28)	9 (20)	0.8 ^f	676	95	—	30	4.6	0.5	11	1.20
1–3 ^g	—	86 (34)	12 (27)	1.3	992	130	19	—	7	0.7	13	1.05
4–8 ^g	15.3	115 (45)	20 (44)	1.7	1642	130	25	—	10	0.9	19	0.95
9–13	17.4	144 (57)	37 (81)	2.1	2071	130	26	—	10	1.0	34	0.95
14–18	20.4	163 (64)	54 (119)	2.3	2368	130	26	—	11	1.1	46	0.85
19–30	21.5	163 (64)	57 (126)	2.7	2403 ^l	130	25	—	12	1.1	46	0.80
31–50	21.5 ^l	163 (64) ⁱ	57 (126) ^j	2.7	2403 ^l	130	25	—	12	1.1	46	0.80
>50	21.5 ^l	163 (64) ⁱ	57 (126) ^j	2.7	2403 ^l	130	21	—	11	1.1	46	0.80
PREGNANCY												
1st trimester	—	—	—	3.0	+0	175	28	—	13	1.4	46	0.80
2nd trimester	—	—	—	3.0	+340	175	28	—	13	1.4	71	1.10
3rd trimester	—	—	—	3.0	+452	175	28	—	13	1.4	71	1.10
LACTATION												
1st 6 months	—	—	—	3.8	+330	210	29	—	13	1.3	71	1.30
2nd 6 months	—	—	—	3.8	+400	210	29	—	13	1.3	71	1.30

NOTE: For all nutrients, values for infants are AI. Dashes indicate that values have not been determined.

^aThe water AI includes drinking water, water in beverages, and water in foods; in general, drinking water and other beverages contribute about 70 to 80 percent, and foods, the remainder. Conversion factors: 1 L = 33.8 fluid oz; 1 L = 1.06 qt; 1 cup = 8 fluid oz.

^bThe Estimated Energy Requirement (EER) represents the average dietary energy intake that will maintain energy balance in a healthy person of a given gender, age, weight, height, and physical activity level. The values listed are based on an "active" person at the reference height and weight and at the midpoint ages for each group until age 19. Chapter 9 and Appendix H provide equations and tables to determine estimated energy requirements.

^cThe linoleic acid referred to in this table and text is the omega-6 fatty acid known as alpha-linoleic acid.

^dThe values listed are based on reference body weights.

In addition to the values that serve as goals for nutrient intakes (presented in the tables on these two pages), the DRI include a set of values called Tolerable Upper Intake Levels (UL). The UL represent the maximum amount of a nutrient that appears safe for most healthy people to consume on a regular basis. Turn the page for a listing of the UL for selected vitamins and minerals.

^eAssumed to be from human milk.

^fAssumed to be from human milk and complementary foods and beverages. This includes approximately 0.6 L (~2½ cups) as total fluid including formula, juices, and drinking water.

^gFor energy, the age groups for young children are 1–2 years and 3–8 years.

^hFor males, subtract 10 calories per day for each year of age above 19.

ⁱBecause weight need not change as adults age if activity is maintained, reference weights for adults 19 through 30 years are applied to all adult age groups.

^jFor females, subtract 7 calories per day for each year of age above 19.

SOURCE: Adapted from the Dietary Reference Intakes series, National Academies Press. Copyright 1997, 1998, 2000, 2001, 2002, 2004, 2005, 2011 by the National Academies of Sciences.

Recommended Dietary Allowances (RDA) and Adequate Intakes (AI) for Vitamins

AGE (YR)	Thiamin RDA (mg/day)	Riboflavin RDA (mg/day)	Niacin RDA (mg/day)	Biotin RDA (mg/day) ^a	Pantothenic Acid AI (µg/day)	Vitamin B ₆ RDA (mg/day)	Folate RDA (µg/day)	Vitamin B ₁₂ RDA (µg/day)	Choline AI (mg/day)	Vitamin C RDA (mg/day)	Vitamin A RDA (µg/day) ^c	Vitamin D RDA (IU/day) ^d	Vitamin E RDA (mg/day) ^e	Vitamin K AI (µg/day)
INFANTS														
0-0.5	0.2	0.3	2	5	1.7	0.1	65	0.4	125	40	400	400 (10 µg)	4	2.0
0.5-1	0.3	0.4	4	6	1.8	0.3	80	0.5	150	50	500	400 (10 µg)	5	2.5
CHILDREN														
1-3	0.5	0.5	6	8	2	0.5	150	0.9	200	15	300	600 (15 µg)	6	30
4-8	0.6	0.6	8	12	3	0.6	200	1.2	250	25	400	600 (15 µg)	7	55
MALES														
9-13	0.9	0.9	12	20	4	1.0	300	1.8	375	45	600	600 (15 µg)	11	60
14-18	1.2	1.3	16	25	5	1.3	400	2.4	550	75	900	600 (15 µg)	15	75
19-30	1.2	1.3	16	30	5	1.3	400	2.4	550	90	900	600 (15 µg)	15	120
31-50	1.2	1.3	16	30	5	1.3	400	2.4	550	90	900	600 (15 µg)	15	120
51-70	1.2	1.3	16	30	5	1.7	400	2.4	550	90	900	600 (15 µg)	15	120
>70	1.2	1.3	16	30	5	1.7	400	2.4	550	90	900	800 (20 µg)	15	120
FEMALES														
9-13	0.9	0.9	12	20	4	1.0	300	1.8	375	45	600	600 (15 µg)	11	60
14-18	1.0	1.0	14	25	5	1.2	400	2.4	400	65	700	600 (15 µg)	15	75
19-30	1.1	1.1	14	30	5	1.3	400	2.4	425	75	700	600 (15 µg)	15	90
31-50	1.1	1.1	14	30	5	1.3	400	2.4	425	75	700	600 (15 µg)	15	90
51-70	1.1	1.1	14	30	5	1.5	400	2.4	425	75	700	600 (15 µg)	15	90
>70	1.1	1.1	14	30	5	1.5	400	2.4	425	75	700	800 (20 µg)	15	90
PREGNANCY														
≤18	1.4	1.4	18	30	6	1.9	600	2.6	450	80	750	600 (15 µg)	15	75
19-30	1.4	1.4	18	30	6	1.9	600	2.6	450	85	770	600 (15 µg)	15	90
31-50	1.4	1.4	18	30	6	1.9	600	2.6	450	85	770	600 (15 µg)	15	90
LACTATION														
≤18	1.4	1.6	17	35	7	2.0	500	2.8	550	115	1200	600 (15 µg)	19	75
19-30	1.4	1.6	17	35	7	2.0	500	2.8	550	120	1300	600 (15 µg)	19	90
31-50	1.4	1.6	17	35	7	2.0	500	2.8	550	120	1300	600 (15 µg)	19	90

NOTE: For all nutrients, values for infants are AI. The table on page Y defines units of nutrient measure.

^aNiacin recommendations are expressed as niacin equivalents (NE), except for recommendations for infants younger than 6 months, which are expressed as preformed niacin.

^bFolate recommendations are expressed as dietary folate equivalents (DFE).

^cVitamin A recommendations are expressed as retinol activity equivalents (RAE).

^dVitamin D recommendations are expressed as cholecalciferol and assume an absence of adequate exposure to sunlight. Pregnant or lactating girls ages 14-18 also need 15 micrograms vitamin D per day.

^eVitamin E recommendations are expressed as α-tocopherol.

Recommended Dietary Allowances (RDA) and Adequate Intakes (AI) for Minerals

AGE (YR)	Sodium AI (mg/day)	Chloride AI (mg/day)	Potassium AI (mg/day)	Calcium RDA (mg/day)	Phosphorus RDA (mg/day)	Magnesium RDA (mg/day)	Iron RDA (mg/day)	Zinc RDA (mg/day)	Iodine RDA (µg/day)	Selenium RDA (µg/day)	Copper RDA (µg/day)	Manganese AI (mg/day) ^f	Fluoride AI (mg/day)	Chromium AI (µg/day)	Molybdenum RDA (µg/day)
INFANTS															
0-0.5	120	180	400	200	100	30	0.27	2	110	15	200	0.003	0.01	0.2	2
0.5-1	370	570	700	260	275	75	11	3	130	20	220	0.6	0.5	5.5	3
CHILDREN															
1-3	1000	1500	3000	700	460	80	7	3	90	20	340	1.2	0.7	11	17
4-8	1200	1900	3800	1000-	500	130	10	5	90	30	440	1.5	1.0	15	22
MALES															
9-13	1500	2300	4500	1300	1250	240	8	8	120	40	700	1.9	2	25	34
14-18	1500	2300	4700	1300	1250	410	11	11	150	55	890	2.2	3	35	43
19-30	1500	2300	4700	1000	700	400	8	11	150	55	900	2.3	4	35	45
31-50	1500	2300	4700	1000	700	420	8	11	150	55	900	2.3	4	35	45
51-70	1300	2000	4700	1000	700	420	8	11	150	55	900	2.3	4	30	45
>70	1200	1800	4700	1200	700	420	8	11	150	55	900	2.3	4	30	45
FEMALES															
9-13	1500	2300	4500	1300	1250	240	8	8	120	40	700	1.6	2	21	34
14-18	1500	2300	4700	1300	1250	360	15	9	150	55	890	1.6	3	24	43
19-30	1500	2300	4700	1000	700	310	18	8	150	55	900	1.8	3	25	45
31-50	1500	2300	4700	1000	700	320	18	8	150	55	900	1.8	3	25	45
51-70	1300	2000	4700	1200	700	320	8	8	150	55	900	1.8	3	20	45
>70	1200	1800	4700	1200	700	320	8	8	150	55	900	1.8	3	20	45
PREGNANCY															
≤18	1500	2300	4700	1300	1250	400	27	12	220	60	1000	2.0	3	29	50
19-30	1500	2300	4700	1000	700	350	27	11	220	60	1000	2.0	3	30	50
31-50	1500	2300	4700	1000	700	360	27	11	220	60	1000	2.0	3	30	50
LACTATION															
≤18	1500	2300	5100	1300	1250	360	10	13	290	70	1300	2.6	3	44	50
19-30	1500	2300	5100	1000	700	310	9	12	290	70	1300	2.6	3	45	50
31-50	1500	2300	5100	1000	700	320	9	12	290	70	1300	2.6	3	45	50

NOTE: For all nutrients, values for infants are AI.

Tolerable Upper Intake Levels (UL) for Vitamins

AGE (YR)	NIAKIN (mg/day) ^a	VITAMIN B ₆ (mg/day)	FOLATE (μg/day) ^a	CHOLINE (mg/day)	VITAMIN C (mg/day)	VITAMIN A (μg/day) ^b	VITAMIN D (IU/day)	VITAMIN E (mg/day) ^c
INFANTS								
0-0.5	—	—	—	—	—	600	1000 (25 μg)	—
0.5-1	—	—	—	—	—	600	1500 (38 μg)	—
CHILDREN								
1-3	10	30	300	1000	400	600	2500 (63 μg)	200
4-8	15	40	400	1000	650	900	3000 (75 μg)	300
9-13	20	60	600	2000	1200	1700	4000 (100 μg)	600
ADOLESCENTS								
14-18	30	80	800	3000	1800	2800	4000 (100 μg)	800
ADULTS								
19-50	35	100	1000	3500	2000	3000	4000 (100 μg)	1000
>70	35	100	1000	3500	2000	3000	4000 (100 μg)	1000
PREGNANCY								
≤18	30	80	800	3000	1800	2800	4000 (100 μg)	800
19-50	35	100	1000	3500	2000	3000	4000 (100 μg)	1000
LACTATION								
≤18	30	80	800	3000	1800	2800	4000 (100 μg)	800
19-50	35	100	1000	3500	2000	3000	4000 (100 μg)	1000

^aThe UL for niacin and folate apply to synthetic forms obtained from supplements, fortified foods, or a combination of the two.

^bThe UL for vitamin A applies to the preformed vitamin only.

^cThe UL for vitamin E applies to any form of supplemental α-tocopherol, fortified foods, or a combination of the two.

Tolerable Upper Intake Levels (UL) for Minerals

AGE (YR)	SODIUM (mg/day)	CHLORIDE (mg/day)	CALCIUM (mg/day)	PHOSPHORUS (mg/day)	MAGNESIUM (mg/day) ^d	IRON (mg/day)	ZINC (mg/day)	IODINE (μg/day)	SELENIUM (μg/day)	COPPER (μg/day)	MANGANESE (mg/day)	FLUORIDE (mg/day)	MOLYBDENUM (μg/day)	BORON (mg/day)	NICKEL (mg/day)	VANADIUM (mg/day)
INFANTS																
0-0.5	—	—	1000	—	—	40	4	—	45	—	—	0.7	—	—	—	—
0.5-1	—	—	1500	—	—	40	5	—	60	—	—	0.9	—	—	—	—
CHILDREN																
1-3	1500	2300	2500	3000	65	40	7	200	90	1000	2	1.3	300	3	0.2	—
4-8	1900	2900	2500	3000	110	40	12	300	150	3000	3	2.2	600	6	0.3	—
9-13	2200	3400	3000	4000	350	40	23	600	280	5000	6	10	1100	11	0.6	—
ADOLESCENTS																
14-18	2300	3600	3000	4000	350	45	34	900	400	8000	9	10	1700	17	1.0	—
ADULTS																
19-50	2300	3600	2500	4000	350	45	40	1100	400	10,000	11	10	2000	20	1.0	1.8
51-70	2300	3600	2000	4000	350	45	40	1100	400	10,000	11	10	2000	20	1.0	1.8
>70	2300	3600	2000	3000	350	45	40	1100	400	10,000	11	10	2000	20	1.0	1.8
PREGNANCY																
≤18	2300	3600	3000	3500	350	45	34	900	400	8000	9	10	1700	17	1.0	—
19-50	2300	3600	2500	3500	350	45	40	1100	400	10,000	11	10	2000	20	1.0	—
LACTATION																
≤18	2300	3600	3000	4000	350	45	34	900	400	8000	9	10	1700	17	1.0	—
19-50	2300	3600	2500	4000	350	45	40	1100	400	10,000	11	10	2000	20	1.0	—

^dThe UL for magnesium applies to synthetic forms obtained from supplements or drugs only.

SOURCE: Adapted with permission from the Dietary Reference Intakes series, National Academies Press. Copyright 1997, 1998, 2000, 2001, 2002, 2005, 2011 by the National Academies of Sciences.

NOTE: An Upper Limit was not established for vitamins and minerals not listed and for those age groups listed with a dash (—) because of a lack of data, not because these nutrients are safe to consume at any level of intake. All nutrients can have adverse effects when intakes are excessive.

Daily Values for Food Labels

The Daily Values are standard values developed by the Food and Drug Administration (FDA) for use on food labels. The values are based on 2,000 calories a day for adults and children over 4 years old. Chapter 2 provides more details.

NUTRIENT	AMOUNT	NUTRIENT	AMOUNT
Protein ^a	50 g	Vitamin K	80 µg
Thiamin	1.5 mg	Calcium	1000 mg
Riboflavin	1.7 mg	Iron	18 mg
Niacin	20 mg NE	Zinc	15 mg
Biotin	300 µg	Iodine	150 µg
Pantothenic acid	10 mg	Copper	2 mg
Vitamin B ₆	2 mg	Chromium	120 µg
Folate	400 µg	Selenium	70 µg
Vitamin B ₁₂	6 µg	Molybdenum	75 µg
Vitamin C	60 mg	Manganese	2 mg
Vitamin A	5000 IU ^b	Chloride	3400 mg
Vitamin D	400 IU ^b	Magnesium	400 mg
Vitamin E	30 IU ^b	Phosphorus	1000 mg

^aThe Daily Values for protein vary for different groups of people: pregnant women, 60 g; nursing mothers, 65 g; infants under 1 year, 14 g; children 1 to 4 years, 16 g.

^bEquivalent values for nutrients expressed as IU are: vitamin A, 1,500 RAE (assumes a mixture of 40% retinol and 60% beta-carotene); vitamin D, 10 µg; vitamin E, 20 mg.

FOOD COMPONENT	AMOUNT	CALCULATION FACTORS
Fat	65 g	30% of calories
Saturated fat	20 g	10% of calories
Cholesterol	300 mg	Same regardless of calories
Carbohydrate (total)	300 g	60% of calories
Fiber	25 g	11.5 g per 1000 calories
Protein	50 g	10% of calories
Sodium	2400 mg	Same regardless of calories
Potassium	3500 mg	Same regardless of calories