

Gauge-invariant expectation values of the energy of a molecule in an electromagnetic field

Anirban Mandal and Katharine L. C. Hunt

Department of Chemistry, Michigan State University, East Lansing, MI 48824

For a molecule interacting with an electromagnetic field, we show that the Hamiltonian can be separated into a molecular Hamiltonian and a field Hamiltonian, both with *gauge-invariant* expectation values.¹ The expectation value of the molecular Hamiltonian gives physically meaningful result for the energy of a molecule in a time-dependent field. The usual partitioning of the full Hamiltonian results in gauge dependence of both the molecular Hamiltonian and the field Hamiltonian.^{2,3} With the usual partitioning of the Hamiltonian, gauge dependence arises even in the absence of an applied field, as we have shown explicitly by considering a gauge transformation from zero applied field and zero external potentials to zero applied field, but non-zero vector and scalar potentials.^{1,4} By repartitioning the full Hamiltonian, we resolve this problem and also remove the gauge-dependence from the Hamiltonian for a molecule in a non-zero applied field.¹ We treat the electromagnetic field classically and treat the molecule quantum mechanically, but nonrelativistically. We construct the full Hamiltonian using a Lagrange multiplier method originally suggested by Fermi⁵ and later applied by Goldman,⁶ partition this Hamiltonian into a molecular term H_m and a field term H_f , and then show that both H_m and H_f have gauge-invariant expectation values.¹ Any gauge may be chosen for the calculation; but following our partitioning, the expectation values of the molecular Hamiltonian are identical to those obtained in the Coulomb gauge. As a corollary of this result, the power absorbed by a molecule from a time-dependent, applied electromagnetic field is equal to the time-derivative of the nonadiabatic term in the molecular energy, in any gauge.^{7,8}

1. A. Mandal and K. L. C. Hunt, *J. Chem. Phys.* **144**, 044109 (2016).
2. H. A. Kramers, *Quantum Mechanics* (North Holland, Amsterdam, 1957), pp. 268 – 269.
3. C. Cohen-Tannoudji, B. Diu, and F. Laloë, *Quantum Mechanics* (Wiley, New York, 1977), Vol. 1, pp. 315 – 326.
4. D. H. Kobe and S. M. Golshan, *J. Phys. A: Math. Gen.* **20**, 2813 (1987).
5. E. Fermi, *Rev. Mod. Phys.* **4**, 87 (1932).
6. T. Goldman, *Phys. Rev. D* **15**, 1063 (1977).
7. A. Mandal and K. L. C. Hunt, *J. Chem. Phys.* **137**, 164109 (2012).
8. A. Mandal and K. L. C. Hunt, *J. Chem. Phys.* **143**, 034102 (2015).