

# On the Fibrations Underlying Optimization and Elimination

Thomas C. Fraser<sup>1,2</sup> and Tobias Fritz<sup>1</sup>

<sup>1</sup>*Perimeter Institute for Theoretical Physics, Waterloo, Ontario, Canada, N2L 2Y5*

<sup>2</sup>*Dept. of Physics and Astronomy, University of Waterloo, Waterloo, Ontario, Canada,  
N2L 3G1*

August 26, 2019

## Abstract

*As of July 25, 2019:* The theory of fibrations and fibered categories appears to be a natural place to discuss the theory of various optimization and elimination problems, including resolution in logic, linear and non-linear quantifier elimination, polytope projection, lattice optimization over various spaces, etc. These notes aim to investigate that claim and furthermore attempts to determine any and all structural similarities between the various cases.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Category Theory Terminology</b>	<b>2</b>
2.1	Cartesian Arrows . . . . .	2
2.2	Fibrations, Fibered Categories, and Cleavages . . . . .	3
2.3	Pseudo-Functors, Splitting Cleavages . . . . .	4
2.4	Nearby Fibrations: Opfibrations and *-fibrations . . . . .	4
2.5	Hom-Functors . . . . .	5
2.6	Adjoint Functors . . . . .	5
2.7	Beck-Chevalley Conditions . . . . .	6
2.8	Slice and Coslice Categories . . . . .	6
2.9	Functors of Monoidal Categories . . . . .	7
2.10	Frobenius Reciprocity . . . . .	7
<b>3</b>	<b>Case Studies of Interest</b>	<b>8</b>
3.1	Polyhedra and Affine Maps . . . . .	8
3.2	The Beck-Chevalley Condition for the Polyhedral Fibration . . . . .	9
3.3	The Subobject Fibration . . . . .	9
3.4	The Beck-Chevalley Condition for Any Subobject Fibration . . . . .	10
3.5	The Category of Convex Cones and Linear Maps . . . . .	12
3.6	Subset Projection . . . . .	12
3.7	Optimization of real-valued functions . . . . .	14

# Notation Proposals

- Aff the category of affine spaces (?) and affine maps between them
- Vect the category of vector spaces and the linear maps between them
- Poly the category of polyhedra and the affine maps between them
- Cone the category of cones and the linear maps between them

## 1 Introduction

## 2 Category Theory Terminology

The following unordered list of categorical concepts are anticipated to be utilized:

- adjunctions
- fibered categories
- cleavages
- pseudo functors (and if cleavages are splitting, functors)
- Beck-Chevalley condition
- Frobenius reciprocity (and functors of monoidal categories)

**Tobias:** Cleavages are not really important because for any two different choices of cleavage, the resulting pullback functors are naturally isomorphic. So cleavages are just a technical tool relevant for proving the equivalence between fibred cats and pseudofunctors, but not relevant in practice

**TC:** The above comment makes sense. Overall there are isomorphisms lurking behind every corner: first, there are natural isomorphisms present when considering the equivalence between pseudo-functors and "cleaved" fibered categories, and second, whenever the cartesian arrows are indeed pullbacks, they are unique up to unique isomorphism and thus entire cleavages are unique up to unique isomorphisms. For a discussion see [Vis04] at the end of Section 3.1.3. starting on page 50.

### 2.1 Cartesian Arrows

**Definition 2.1.** Let  $P : \mathcal{E} \rightarrow \mathcal{B}$  be a functor between categories  $\mathcal{E}$  and  $\mathcal{B}$ . An arrow  $\phi : \alpha \rightarrow \beta$  of  $\mathcal{E}$  is *cartesian* with respect to  $P$  (sometimes *P-cartesian*) if for every arrow  $\psi : \gamma \rightarrow \beta$  sharing a codomain with  $\phi$ , and for every arrow  $g : P(\gamma) \rightarrow P(\alpha)$  in  $\mathcal{B}$  satisfying  $g \circ P(\phi) = P(\psi)$ , there exists a unique arrow  $\theta : \gamma \rightarrow \alpha$  in  $\mathcal{E}$  satisfying  $\phi \circ \theta = \psi$  and  $P(\theta) = g$ .

$$\begin{array}{ccccc}
 & & \forall \psi & & \\
 & \gamma & \xrightarrow{\quad} & \beta & \\
 & \downarrow & \searrow \exists! \theta & \downarrow \phi & \\
 & P(\gamma) & \xrightarrow{P(\psi)} & P(\alpha) & \xrightarrow{P(\phi)} P(\beta) \\
 & \downarrow & \downarrow & \downarrow & \\
 & P(\gamma) & \xrightarrow{\forall g} & P(\alpha) & \xrightarrow{P(\phi)} P(\beta)
 \end{array}
 \tag{1}$$

**Corollary 2.0.1.** *A cartesian morphism  $\phi : \alpha \rightarrow \beta$  in  $\mathcal{E}$  with respect to a functor  $P : \mathcal{E} \rightarrow \mathcal{B}$  establishes an isomorphism of categories  $\mathcal{E}/\phi \cong \mathcal{E}/\beta \times_{\mathcal{B}/P(\beta)} \mathcal{B}/P(\phi)$  [Lur09, Section 2.4.1]<sup>1</sup>*

$$\mathcal{E}/\phi \cong \mathcal{E}/\beta \times_{\mathcal{B}/P(\beta)} \mathcal{B}/P(\phi) \quad (2)$$

where  $\mathcal{E}/\beta \times_{\mathcal{B}/P(\beta)} \mathcal{B}/P(\phi)$  is the pullback of functors.

$$\begin{array}{ccc} \mathcal{E}/\phi & \xrightarrow{P/\phi} & \mathcal{B}/P(\phi) \\ \downarrow \cong & \searrow & \downarrow \text{cod} \\ \mathcal{E}/\beta \times_{\mathcal{B}/P(\beta)} \mathcal{B}/P(\phi) & \longrightarrow & \mathcal{B}/P(\phi) \\ \downarrow \perp & & \downarrow \text{cod} \\ \mathcal{E}/\beta & \xrightarrow{P/\beta} & \mathcal{B}/P(\beta) \end{array} \quad (3)$$

The pullback category  $\mathcal{E}/\beta \times_{\mathcal{B}/P(\beta)} \mathcal{B}/P(\phi)$  has morphisms associated with diagrams of  $\mathcal{B}$  with the following format:

$$\begin{array}{ccc} & P(\gamma) & \\ f \swarrow & \downarrow P(\chi) & \searrow P(\omega) \\ & P(\delta) & \\ g \swarrow & \downarrow P(\psi) & \searrow \\ P(\alpha) & \xrightarrow{P(\phi)} & P(\beta) \end{array} \quad (4) \quad \text{eq:pullback\_}$$

Evidently, if  $\phi : \alpha \rightarrow \beta$  is cartesian, then there exists unique morphisms  $\zeta : \gamma \rightarrow \alpha$  and  $\eta : \delta \rightarrow \alpha$  such that  $P(\zeta) = f$  and  $P(\eta) = g$  and the following diagram of  $\mathcal{E}$  commutes:

$$\begin{array}{ccc} & \gamma & \\ \zeta \swarrow & \downarrow \chi & \searrow \omega \\ & \delta & \\ \eta \swarrow & \downarrow \psi & \searrow \\ \alpha & \xrightarrow{\phi} & \beta \end{array} \quad (5) \quad \text{eq:cartesian}$$

Intuitively, if  $\phi$  is cartesian, then in order to determine the category  $\mathcal{E}/\phi$  over  $\phi$ , it is sufficient to specify  $\mathcal{E}/\beta \times_{\mathcal{B}/P(\beta)} \mathcal{B}/P(\phi)$ .

## 2.2 Fibrations, Fibered Categories, and Cleavages

**Definition 2.2.** A *fibered category over  $\mathcal{B}$*  is a category  $\mathcal{E}$  associated to the domain of a functor, referred to as the *fibration*,  $P : \mathcal{E} \rightarrow \mathcal{B}$  with the property that for every morphism  $f : a \rightarrow b$  of  $\mathcal{B}$  and object  $\beta$  such that  $P(\beta) = b$ , there exists a cartesian arrow  $\phi : \alpha \rightarrow \beta$  with  $P(\phi) = f$ .

**Lemma 2.1.** A fibration  $P : \mathcal{E} \rightarrow \mathcal{B}$  is a faithful functor if and only if its fibers are thin.

*Proof.* Recall that if  $P : \mathcal{E} \rightarrow \mathcal{B}$  is a faithful functor, then by definition every pair of parallel arrows  $\phi, \psi : \alpha \rightarrow \beta$  in  $\mathcal{E}$  satisfies

$$P(\phi) = P(\psi) : P(\alpha) \rightarrow P(\beta) \implies \phi = \psi. \quad (6) \quad \text{eq:faithful}$$

<sup>1</sup>This formulation is also discussed here: <https://ncatlab.org/nlab/show/Cartesian+morphism#CartInOrdCatReformulation>.

$\Rightarrow$  : Assuming  $P : \mathcal{E} \rightarrow \mathcal{B}$  is faithful functor, consider an arbitrary pair of parallel arrows  $\phi, \psi : \alpha \rightarrow \beta$  in an arbitrary fiber  $\mathcal{E}_x$  over  $x$ ; i.e.  $P(\phi) = P(\psi) = \text{id}_x$ . In such cases, faithfulness of  $P$  (Eq. [6](#)) guarantees that  $\phi = \psi$  and thus  $\mathcal{E}_x$  is a thin category.

$\Leftarrow$  : If the fiber  $\mathcal{E}_x$  for every object  $x$  in  $\mathcal{B}$  is a thin category, then clearly  $P : \mathcal{E} \rightarrow \mathcal{B}$  must be faithful when restricted to an individual fiber. The non-trivial case is to consider an arbitrary pair of parallel morphisms  $\phi, \psi : \alpha \rightarrow \beta$  not belonging to any fibers of  $\mathcal{E}$ . Denote  $a := P(\alpha)$  and  $b := P(\beta)$  and suppose  $f := P(\phi) = P(\psi) : a \rightarrow b$ . Then, because  $\mathcal{E}$  is a fibered category, there exists a cartesian arrow  $\zeta : \gamma \rightarrow \beta$ , such that  $P(\zeta) = f$  (note that  $a = P(\alpha) = P(\gamma)$  but  $\gamma$  is not necessarily equal to  $\alpha$ ). Since  $\zeta$  is a cartesian arrow, there exists a unique arrows  $\mu, \nu : \alpha \rightarrow \gamma$  completing the top edges of the following diagram:

$$\begin{array}{ccccc}
 \alpha & \xrightarrow{\mu} & \gamma & \xleftarrow{\nu} & \alpha \\
 \downarrow \psi & & \downarrow \zeta & & \downarrow \phi \\
 & & \beta & & \\
 \downarrow & & \downarrow & & \downarrow \\
 a & \xrightarrow{f} & b & \xleftarrow{f} & a
 \end{array}$$

(7)

However,  $P(\nu) = \text{id}_a = P(\mu)$  and therefore  $\mu$  and  $\nu$  are parallel arrows in the fiber  $\mathcal{E}_a$  and therefore  $\mu = \nu$  because  $\mathcal{E}_a$  is assumed thin. Therefore,  $\psi = \zeta \circ \mu = \zeta \circ \nu = \phi$  and thus  $P$  is a faithful functor.  $\square$

**Definition 2.3.** A *cleavage* for a fibration  $P : \mathcal{E} \rightarrow \mathcal{B}$  is an assignment to each morphism  $f : a \rightarrow b$  of  $\mathcal{B}$  and object  $\beta$  in  $\mathcal{E}_b$  (i.e.  $P(\beta) = b$ ), a unique cartesian morphism  $\kappa_f(\beta)$  of  $\mathcal{E}$  such that  $P(\kappa_f(\beta)) = f$ .

Given a cleavage for a fibration, the cartesianness of morphisms within a cleavage permits one to establish functors between the fibers of the fibration. This concept is visualized in the following figure:

(8)

## 2.3 Pseudo-Functors, Splitting Cleavages

Pages 47-48 of [\[Vis04\]](#) explicate the notions of pseudo-functors and their equivalence to fibrations with cleavages. Moreover if the cleavage is splitting, the induced pseudo-functor is in fact a functor.

## 2.4 Nearby Fibrations: Opfibrations and \*-fibrations

Given a functor  $P : \mathcal{E} \rightarrow \mathcal{B}$ , it can be considered as a fibration in many different ways. For example, if  $P^{\text{op}} : \mathcal{E}^{\text{op}} \rightarrow \mathcal{B}^{\text{op}}$  is a fibration, then  $P$  is said to be an *opfibration*.

## 2.5 Hom-Functors

For a locally small category  $\mathcal{C}$ , the hom-functor of  $\mathcal{C}$  is a functor  $\text{Hom}_{\mathcal{C}} : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathbf{Set}$  constructed in the following manner. Given objects  $a, b, c, \dots \in \mathcal{C}_0$  of  $\mathcal{C}$ , the hom-functor  $\text{Hom}_{\mathcal{C}}$  maps a pair of objects  $(a, b) \in (\mathcal{C}^{\text{op}} \times \mathcal{C})_0 = \mathcal{C}_0 \times \mathcal{C}_0 = \mathcal{C}_0^2$  into the set<sup>2</sup> of morphisms  $\mathcal{C}_1$  of  $\mathcal{C}$  with source  $a$  and target  $b$ . Therefore,  $\text{Hom}_{\mathcal{C}}(a, b)$  is the set of morphisms in  $\mathcal{C}$  of type  $a \rightarrow b$ . Given morphisms  $g^{\text{op}} \in \text{Hom}_{\mathcal{C}^{\text{op}}}(a, c)$  and  $h \in \text{Hom}_{\mathcal{C}}(b, d)$ , the hom-functor  $\text{Hom}_{\mathcal{C}}$  constructs a function

$$\text{Hom}_{\mathcal{C}}(g^{\text{op}}, h) : \text{Hom}_{\mathcal{C}}(a, b) \rightarrow \text{Hom}_{\mathcal{C}}(c, d)$$

which takes a morphism  $f : a \rightarrow b \in \text{Hom}_{\mathcal{C}}(a, b)$  and produces the morphism  $h \circ f \circ g : c \rightarrow d \in \text{Hom}_{\mathcal{C}}(c, d)$ . Graphically,

$$\text{Hom}_{\mathcal{C}}(g^{\text{op}}, h) \left( a \xrightarrow{f} b \right) = c \xrightarrow{g} a \xrightarrow{f} b \xrightarrow{h} d$$

## 2.6 Adjoint Functors

Given two categories  $\mathcal{C}$  and  $\mathcal{D}$ , a pair of functors  $L : \mathcal{C} \rightarrow \mathcal{D}, R : \mathcal{D} \rightarrow \mathcal{C}$  are called an *adjoint pair*, denoted  $L \dashv R$  or

$$\begin{array}{ccc} & L & \\ \mathcal{C} & \xrightleftharpoons{\perp} & \mathcal{D} \\ & R & \end{array}$$

if there exists a natural isomorphism  $\alpha$  between the following pair of hom-functors of type  $\mathcal{C}^{\text{op}} \times \mathcal{D} \rightarrow \mathbf{Set}$ :

$$\text{Hom}_{\mathcal{D}}(L^{\text{op}}(-), -) \xrightarrow{\alpha} \text{Hom}_{\mathcal{C}}(-, R(-))$$

This relationship can be depicted graphically as 2-cell (and its inverse) in  $\mathbf{Cat}$ ,

$$\begin{array}{ccc} \mathcal{C}^{\text{op}} \times \mathcal{D} & \xrightarrow{I_{\mathcal{C}^{\text{op}}} \times R} & \mathcal{C}^{\text{op}} \times \mathcal{C} \\ \downarrow L^{\text{op}} \times I_{\mathcal{D}} & \searrow \alpha & \downarrow \text{Hom}_{\mathcal{C}} \\ \mathcal{D}^{\text{op}} \times \mathcal{D} & \xrightarrow{\text{Hom}_{\mathcal{D}}} & \mathbf{Set} \end{array}$$

$\alpha^{-1}$

Concretely, the naturality of  $\alpha$  means that for every morphism  $(f^{\text{op}} : b \rightarrow a, g : c \rightarrow d) \in (\mathcal{C}^{\text{op}} \times \mathcal{D})_1$  the components  $\alpha_{(b,c)}$  and  $\alpha_{(a,d)}$  of  $\alpha$  make the following square commute:

$$\begin{array}{ccc} \text{Hom}_{\mathcal{D}}(L^{\text{op}}(b), c) & \xrightarrow{\text{Hom}_{\mathcal{D}}(L^{\text{op}}(f^{\text{op}}), g)} & \text{Hom}_{\mathcal{D}}(L^{\text{op}}(a), d) \\ \downarrow \alpha_{(b,c)} & & \downarrow \alpha_{(a,d)} \\ \text{Hom}_{\mathcal{C}}(b, R(c)) & \xrightarrow{\text{Hom}_{\mathcal{C}}(f^{\text{op}}, R(g))} & \text{Hom}_{\mathcal{C}}(a, R(d)) \end{array}$$

<sup>2</sup>The collection of morphisms of type  $a \rightarrow b$  forms a set because  $\mathcal{C}$  is locally small.

## 2.7 Beck-Chevalley Conditions

The Beck-Chevalley Conditions are conditions that may or may not be satisfied by a quadruplet of functors  $F, H, G, K$  which form a natural isomorphism  $\alpha : KF \Rightarrow HG$  square:

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{F} & \mathcal{B} \\ G \downarrow & \alpha \swarrow & \downarrow K \\ \mathcal{C} & \xrightarrow{H} & \mathcal{D} \end{array}$$

To define the *left* Beck-Chevalley condition, one needs functors  $F_L : \mathcal{B} \rightarrow \mathcal{A}$  and  $H_L : \mathcal{D} \rightarrow \mathcal{A}$  which are respectively left adjoint functors to  $F$  and  $H$ ,

$$\begin{array}{ccc} & F_L & \\ \mathcal{A} & \xleftarrow{\quad} & \mathcal{B} \\ & \perp & \\ & F & \end{array}, \quad \begin{array}{ccc} & H_L & \\ \mathcal{C} & \xleftarrow{\quad} & \mathcal{D} \\ & \perp & \\ & H & \end{array}.$$

Using these left adjoint functors, it becomes possible to construct a natural transformation  $\beta : KH_L \Rightarrow GF_L$  from  $\alpha$ <sup>3</sup>. Graphically,  $\beta$  can be identified as the outer cell of the following diagram:

$$\begin{array}{ccc} & F_L & \\ \mathcal{A} & \xleftarrow{\quad} & \mathcal{B} \\ & \perp & \\ & F & \\ G \downarrow & \alpha \swarrow & \downarrow K \\ \mathcal{C} & \xleftarrow{\quad} & \mathcal{D} \\ & \top & \\ & H_L & \end{array}, \quad \text{i.e.} \quad \begin{array}{ccc} & F_L & \\ \mathcal{A} & \xleftarrow{\quad} & \mathcal{B} \\ & \beta \swarrow & \downarrow K \\ G \downarrow & & \downarrow \\ \mathcal{C} & \xleftarrow{\quad} & \mathcal{D} \\ & H_L & \end{array}.$$

Although the natural transformation  $\alpha$  is assumed to be a natural isomorphism, the natural transformation  $\beta$  need not be; if  $\beta$  happens to be a natural isomorphism, then we say that the original square satisfies the *left* Beck-Chevalley condition<sup>4</sup>. The *right* Beck-Chevalley condition is defined analogously with functors  $F_R, H_R$  which are respectively right adjoints  $F \dashv F_R$  and  $H \dashv H_R$ .

## 2.8 Slice and Coslice Categories

Given a category  $\mathcal{C}$  and an object  $c \in \mathcal{C}_0$  of  $\mathcal{C}$ , the *slice category* (or *over category*)  $\mathcal{C}/c$  is the “stuff in  $\mathcal{C}$  that is on top of  $c$ ”. Specifically, the objects of  $\mathcal{C}/c$  are all the morphisms  $f \in \mathcal{C}_1$  from  $\mathcal{C}$  whose codomain is  $\text{cod}(f) = c$  (alternatively you could write  $(\mathcal{C}/c)_0 = \text{Hom}_{\mathcal{C}}(-, c)$ ). A morphism of  $\mathcal{C}/c$  between objects  $f : a \rightarrow c, g : b \rightarrow c \in (\mathcal{C}/c)_0$  is a commuting triangle completed by a third morphism  $h : a \rightarrow b \in \mathcal{C}_1$ :

$$\begin{array}{ccc} a & \xrightarrow{h} & b \\ g \searrow & & \swarrow f \\ & c & \end{array}$$

<sup>3</sup>The natural transformations  $\alpha$  and  $\beta$  are known as *mates* or *conjugates*.

<sup>4</sup>Are the left adjoints  $F_L, H_L$  unique? If not, it might be better to say the original square satisfies the left Beck-Chevalley condition with respect to  $F_L, H_L$ .

Composition of morphisms in  $\mathcal{C}/c$  is induced by the composition of morphisms in  $\mathcal{C}$ :

$$\left( \begin{array}{ccc} y & \xrightarrow{n} & z \\ f \searrow & & \nearrow h \\ & c & \end{array} \right) \circ_{\mathcal{C}/c} \left( \begin{array}{ccc} x & \xrightarrow{m} & y \\ g \searrow & & \nearrow f \\ & c & \end{array} \right) = \begin{array}{ccccc} x & \xrightarrow{m} & y & \xrightarrow{n} & z \\ & \searrow g & \downarrow f & \nearrow h & \\ & & c & & \end{array}$$

The assignment of an overcategory  $\mathcal{C}/c$  to each object  $c$  can be extended to a *slice functor*  $\mathcal{C}/(-) : \mathcal{C} \rightarrow \mathbf{Cat}$  in the following sense. For objects  $c \in \mathcal{C}_0$ , the slice functor takes  $c$  to the slice category  $\mathcal{C}/c$ ; for morphisms  $f : a \rightarrow b \in \mathcal{C}_1$ , the slice functor takes  $f$  to the functor  $\mathcal{C}/f : \mathcal{C}/a \rightarrow \mathcal{C}/b$  defined graphically; for every morphism of  $\mathcal{C}/a$  (commuting triangle in  $\mathcal{C}$  over  $a$ ), construct the morphism of  $\mathcal{C}/b$  (commuting triangle in  $\mathcal{C}$  over  $b$ ) as follows:

$$\begin{array}{ccc} l & \xrightarrow{m} & r \\ x \searrow & & \nearrow x' \\ & a & \\ f \downarrow & & \\ & b & \end{array} \quad \begin{array}{c} f \circ_{\mathcal{C}} x \\ \quad \quad \quad \\ f \circ_{\mathcal{C}} x' \end{array}$$

where the inner triangle is a morphism of  $\mathcal{C}/a$  and the outer triangle is a morphism of  $\mathcal{C}/b$  given by the functor  $\mathcal{C}/f$ .

Given a category  $\mathcal{C}$  and an object  $c \in \mathcal{C}_0$  of  $\mathcal{C}$  the *coslice category* (or *under category*)  $c/\mathcal{C}$  is the “stuff in  $\mathcal{C}$  that is underneath  $c$ ”. Specifically, the objects of  $c/\mathcal{C}$  are all the morphisms  $f \in \mathcal{C}_1$  from  $\mathcal{C}$  whose domain is  $\text{dom}(f) = c$  (alternatively you could write  $(c/\mathcal{C})_0 = \text{Hom}_{\mathcal{C}}(c, -)$ ). A morphism of  $c/\mathcal{C}$  between objects  $f : c \rightarrow a, g : c \rightarrow b \in (c/\mathcal{C})_0$  is a commuting triangle completed by a third morphism  $h : a \rightarrow b \in \mathcal{C}_1$ :

$$\begin{array}{ccc} & c & \\ g \swarrow & & \searrow f \\ a & \xrightarrow{h} & b \end{array}$$

Everything about coslice categories is defined as expected analogously to that of a slice categories.

**TODO:** determine how the details of the Grothendieck construction transform the slice (pseudo-)functor  $\mathcal{C}/(-) : \mathcal{C} \rightarrow \mathbf{Cat}$  into the codomain fibration

## 2.9 Functors of Monoidal Categories

[TODO]

## 2.10 Frobenius Reciprocity


[TODO]

### 3 Case Studies of Interest

#### 3.1 Polyhedra and Affine Maps


One of the primary motivating examples for this project is the theory of *(finite) convex polyhedra* and the affine maps between them. Following Boyd and Vandenberghe [BV04], a *polyhedron*<sup>5,6</sup>  $P$  is the intersection of a finite number of *halfspaces* of some ambient vector space  $V \cong \mathbb{R}^n$ . A *halfspace*  $H \subseteq \mathbb{R}^n$  is a subset of a vector space (of dimension  $n$ ) which is the solution set of a linear inequality constraint over canonical coordinates  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ :

$$H = \{x \in V \mid a^\top x = \sum_{i=1}^n a_i x_i \geq b\} \quad (9)$$



As previously mentioned, a polyhedra is the intersection of finitely many halfspaces and therefore corresponds to

$$P = \{x \in V \mid \bigwedge_{j=1}^k (a_j^\top x \geq b_j)\} \quad (10)$$



$$\begin{array}{ccc} f^*(\mathbb{R}_+^k) & \xrightarrow{f'} & \mathbb{R}_+^k \\ \downarrow j & \lrcorner & \downarrow i \\ \mathbb{R}^n & \xrightarrow{f} & \mathbb{R}^k \end{array} \quad (11)$$

eq:pullback\_

<sup>5</sup>The term polytope will be reserved for the context of *bounded polyhedron*. Note that the opposite convention is sometimes used by other authors as pointed out by [BV04].

<sup>6</sup>Alternative and sometimes inequivalent definitions for “polyhedra” do exist; oftentimes, these alternative definitions accommodate more general notions of polyhedra, such as non-convex polyhedra. Understanding the relationship between these various definitions, and the proposal of new ones, is a mathematical endeavour which dates back to antiquity and continues today [Gru03; Lak15].




eq:pullback\_of\_positive\_orthant

In Equation [11](#), the morphism  $j$  is simply the inclusion of the polyhedra  $f^*(\mathbb{R}_+^k)$  into its ambient vector space  $\mathbb{R}^n$  and the morphism  $f'$  is provided by the restriction of  $f$  onto  $f^*(\mathbb{R}_+^k)$ .


### 3.2 The Beck-Chevalley Condition for the Polyhedral Fibration

Consider a commuting square of affine maps between vector spaces in the base category of affine maps

**Definition 3.1.** The category  $\text{Poly}$  consists of polyhedra as objects and affine maps between them.



$$\begin{array}{ccccc}
 & \text{Poly}_Y & \xleftarrow{\pi_{Y,!}} & \text{Poly}_{X \otimes Y} & \\
 \nearrow \pi_{Y,!} & \downarrow \pi_{Y \otimes Z,!} & & \nearrow \pi_{X \otimes Y,!} & \\
 \text{Poly}_{Y \otimes Z} & \xleftarrow{\pi_{1,!}} & \text{Poly}_{X \otimes Y \otimes Z} & \xleftarrow{\pi_{X,!}} & \text{Poly}_X \\
 \downarrow \pi_{Z,!} & \downarrow \pi_{1,!} & \downarrow \pi_{X \otimes Z,!} & \downarrow \pi_{X,!} & \\
 & \text{Poly}_1 & \xleftarrow{\pi_{1,!}} & \text{Poly}_X & \\
 \downarrow \pi_{Z,!} & \nearrow \pi_{1,!} & & \nearrow \pi_{X,!} & \\
 \text{Poly}_Z & \xleftarrow{\pi_{Z,!}} & \text{Poly}_{X \otimes Z} & & 
 \end{array} \tag{12}$$




### 3.3 The Subobject Fibration

Given any category  $\mathcal{B}$ , and an object  $X$  of  $\mathcal{B}$ , a *subobject* of  $X$  is an isomorphism class of monomorphisms  $f : Y \hookrightarrow X$ ; two monomorphisms  $f : Y \hookrightarrow X, g : Z \hookrightarrow X$  with shared codomain  $X$  are isomorphic if there

exists an isomorphism  $h : Y \xrightarrow{\sim} Z$  such that  $f = g \circ h$ . The individual monomorphisms of a subobject class can be equipped with a preorder relation  $(f : Y \hookrightarrow Z) \leq (g : Z \hookrightarrow X)$  if there exists  $k : Y \rightarrow Z$  such that  $f = g \circ k$ :

$$\begin{array}{ccc} Y & \xrightarrow{k} & Z \\ f \swarrow & & \searrow g \\ & X & \end{array} \quad (13)$$

Note that if such a  $k : Y \rightarrow Z$  exists, then it is unique because  $g$  is a monomorphism; if  $k' : Y \rightarrow Z$  satisfied  $f = g \circ k'$  as well, then  $g \circ k = g \circ k'$  which implies  $k = k'$ . Moreover, this preorder relation on the individual monomorphisms extends to a poset  $\mathbf{Sub}_{\mathcal{B}}(X)$  between the subobjects. For example, if  $\mathcal{B}$  is the category  $\mathbf{Set}$ , the subobjects of  $X$  are subsets of  $X$  and thus  $\mathbf{Sub}_{\mathbf{Set}}(X) \cong \mathbf{P}(X)$ . Even more concretely, if  $X$  is the set  $\mathbb{R}^2$ , an exemplary diagram of  $\mathbf{Sub}_{\mathbf{Set}}(\mathbb{R}^2)$  is



If the category  $\mathcal{B}$  has pullbacks, the posetal categories  $\mathbf{Sub}_{\mathcal{B}}(X)$  for varying objects  $X$  of  $\mathcal{B}$  can be “stitched together” to form an enveloping category denoted  $\mathbf{Sub}(\mathcal{B})$ . The objects of  $\mathbf{Sub}(\mathcal{B})$  are the objects of  $\mathbf{Sub}_{\mathcal{B}}(X)$  for various  $X$ . The morphisms of  $\mathbf{Sub}(\mathcal{B})$  are defined using the morphisms of  $\mathcal{B}$ . Given an equivalence class of monomorphisms  $B = \{b_i : Z_i \hookrightarrow X\}_{i \in \mathcal{I}} \in \mathbf{Sub}_{\mathcal{B}}(X)$  with shared codomain  $X$ , and morphism  $f : Y \rightarrow X$  (not necessarily a monomorphism) with codomain  $X$ , the pullbacks  $f^*b_i$  of  $b_i$  along  $f$  for each  $i \in \mathcal{I}$  constitute an equivalence class of monomorphisms  $f^*B := \{f^*b_i : Y \times_X Z_i \hookrightarrow Y\}_{i \in \mathcal{I}} \in \mathbf{Sub}_{\mathcal{B}}(Y)$ .<sup>7,8</sup>

$$\begin{array}{ccc} Y \times_X Z_i & \xrightarrow{b_i^* f} & Z_i \\ f^* b_i \downarrow & \lrcorner & \downarrow b_i \\ Y & \xrightarrow{f} & X \end{array} \quad (14)$$

To every morphism  $f : Y \rightarrow X$  of  $\mathcal{B}$ , and object  $B \in \mathbf{Sub}_{\mathcal{B}}(X)$ , denote  $\kappa_f(B) : f^*B \rightarrow B$  where  $f^*B \in \mathbf{Sub}_{\mathcal{B}}(Y)$ . Finally, a morphism of  $\mathbf{Sub}(\mathcal{B})$  between  $A \in \mathbf{Sub}_{\mathcal{B}}(Y)$  and  $B \in \mathbf{Sub}_{\mathcal{B}}(X)$  is the formal sequence  $A \xrightarrow{\leq} f^*(B) \xrightarrow{\kappa_f(B)} B$ . The projection functor  $P_{\mathcal{B}} : \mathbf{Sub}(\mathcal{B}) \rightarrow \mathcal{B}$  defines the *subobject fibration*<sup>9</sup>.

**TODO:** figure out the relationship between a subobject fibration and the codomain fibration via the notion of *subterminal objects*.

### 3.4 The Beck-Chevalley Condition for Any Subobject Fibration

Given any category  $\mathcal{B}$  with all pullbacks, the functor  $P_{\mathcal{B}} : \mathbf{Sub}_{\mathcal{B}} \rightarrow \mathcal{B}$  which sends subobjects  $[\psi]$  to their shared codomains, constitutes a bifibration. In particular, given a morphism  $f : a \rightarrow b$  of  $\mathcal{B}$ , there is an induced adjoint pair of functors between the subobject fibers:

$$\begin{array}{ccc} & f_! & \\ \text{Sub}_{\mathcal{B}}(a) & \xrightarrow{\quad} & \text{Sub}_{\mathcal{B}}(b) \\ & f^* & \end{array} \quad (15)$$

<sup>7</sup>Evidently, this relies on the fact that pullbacks preserve monomorphisms; i.e.  $b : Z \hookrightarrow X$  is a monomorphism, then the pullback  $f^*b : Y \times_X Z \hookrightarrow Y$  is also.

<sup>8</sup>The isomorphisms connecting the monomorphisms of  $f^*B$  are also given by pullback of the isomorphisms connecting the monomorphisms of  $B$ .

<sup>9</sup>Note that while the fibres  $\mathbf{Sub}_{\mathcal{B}}(X)$  are thin, the total category  $\mathbf{Sub}(\mathcal{B})$  is not necessarily thin.

Specifically, left adjoint functor  $f_!$  acting on a subobject  $[\psi] : \mathbf{Sub}_{\mathcal{B}}(a)$  (where  $[\psi]$  is the equivalence class of monomorphisms into  $a$  containing  $\psi : a' \hookrightarrow a$ ) is given by post-composition  $f_!([\psi]) = [f\psi] \in \mathbf{Sub}_{\mathcal{B}}(b)$  in  $\mathcal{B}$ . The right adjoint functor  $f^*$  acting on a subobject  $[\phi] \in \mathbf{Sub}_{\mathcal{B}}(b)$  is given by pullback of  $a \xrightarrow{f} b \xleftarrow{\phi} b'$  in  $\mathcal{B}$ :

$$\begin{array}{ccc} f^*b' & \longrightarrow & b' \\ f^*\phi \downarrow & \lrcorner & \downarrow \phi \\ a & \xrightarrow{f} & b \end{array} \quad (16)$$

Generally speaking, it is important to determine how diagrams in the base category  $\mathcal{B}$  behave when lifted to the total category  $\mathbf{Sub}_{\mathcal{B}}$  under the associated pseudo-functors. For example, given a pullback square in base category  $\mathcal{B}$ ,

$$\begin{array}{ccc} a & \xrightarrow{f} & b \\ h \downarrow & \lrcorner & \downarrow g \\ c & \xrightarrow{k} & d \end{array} \quad (17)$$

There exists a natural isomorphism  $\alpha$  between the functors  $f^*g^*$  and  $h^*k^*$ :

$$\begin{array}{ccc} \mathbf{Sub}_{\mathcal{B}}(a) & \xleftarrow{f} & \mathbf{Sub}_{\mathcal{B}}(b) \\ h \uparrow & \swarrow \alpha & \uparrow g \\ \mathbf{Sub}_{\mathcal{B}}(c) & \xleftarrow{k} & \mathbf{Sub}_{\mathcal{B}}(d) \end{array} \quad (18)$$

The existence of components  $\alpha_{[\psi]}$  for each subobject class containing the monomorphism  $\psi : p \hookrightarrow d$  can be determined by examining the following diagram:

$$\begin{array}{ccccc} & & f^*g^*\psi & \longrightarrow & f^*g^*p \\ & & \downarrow & \lrcorner & \downarrow \\ & & a & \xrightarrow{f} & b \xleftarrow{g^*\psi} g^*p \\ & \nearrow h^*k^*\psi & \downarrow h & \lrcorner & \downarrow g \\ & & c & \xrightarrow{k} & d \xleftarrow{\psi} p \\ & & \downarrow k^*\psi & \lrcorner & \downarrow \\ h^*k^*p & \longrightarrow & k^*p & \longrightarrow & p \end{array} \quad (19)$$

Using the composition of pullback squares, it is clear that *both*  $f^*g^*p$  and  $h^*k^*p$  are pullbacks of  $a \xrightarrow{gf=kh} d \xleftarrow{\psi} p$  and therefore, they are unique up to a unique isomorphism which will be denoted  $\beta$ . It is important to notice that this does not depend on  $a$  being a pullback itself.

$$\begin{array}{ccc} & \beta & \\ & \swarrow \beta^{-1} & \searrow \\ h^*k^*p & & f^*g^*p \\ \downarrow h^*k^*\psi & \lrcorner & \downarrow \\ a & \xrightarrow{gf=kh} d \xleftarrow{\psi} p \end{array} \quad (20)$$

These isomorphisms actually establish that  $h^*k^*\psi \simeq f^*g^*\psi$  and therefore they belong to the same equivalence class  $[h^*k^*\psi] = [f^*g^*\psi]$  for each  $\psi$ . Therefore, it becomes clear that the natural isomorphism is indeed the identity natural transformation  $\text{id}_{f^*g^*} = \text{id}_{h^*k^*} = \alpha : f^*g^* \rightarrow h^*k^*$ .<sup>10</sup>

<sup>10</sup>Ultimately, the reason for  $\alpha = \text{id}$  is due to considering  $\mathbf{Sub}_{\mathcal{B}}(a)$ , the *poset* of subobjects of  $a$ , instead of  $\mathbf{Monog}(a)$  the *preorder* of monomorphisms into  $a$ .

Alternatively, one can consider whether or not there exists natural transformations (or natural isomorphisms) of the form  $\alpha : h_! f^* \rightarrow k^* g_!$ . Remembering that the left adjoints  $f_!$  are specified by post-composition  $f_![\psi] \rightarrow [f\psi]$ , this question can be answered by considering the following diagram:

$$\begin{array}{ccccc}
 & & hf^*\psi & & \\
 & & \curvearrowright & & \\
 k^*gp & \xrightarrow{k^*g\psi} & c & \xleftarrow{h} & a & \xleftarrow{f^*\psi} & f^*p \\
 \downarrow & \lrcorner & \downarrow k & \lrcorner & \downarrow f & \lrcorner & \downarrow \\
 p & \xrightarrow{g\psi} & d & \xleftarrow{g} & b & \xleftarrow{\psi} & p \\
 & \curvearrowright & & \curvearrowright & & \curvearrowright & 
 \end{array} \tag{21}$$

Similarly,  $k^*gp$  and  $f^*p$  are *both* pullbacks of  $p \xrightarrow{g\psi} d \xleftarrow{k} c$  and therefore, we obtain an analogous statement: since  $hf^*\psi \simeq k^*g\psi$ , we have  $[hf^*\psi] = [k^*g\psi]$  and therefore  $h_! f^* = k^* g_!$ . The key difference in this case is that this argument *does* rely on  $a$  being the pullback of a pullback square.

In summary, the subobject fibration *satisfies* the Beck-Chevalley condition. Moreover, since none of the above argument relies on the fact that the objects of total category are represented by *monomorphisms* in the base, just that they correspond to *morphisms* in the base, it is clear that any codomain bifibration also satisfies the Beck-Chevalley condition.

### 3.5 The Category of Convex Cones and Linear Maps

Given any  $\mathbb{R}$ -vector space  $V$ , a (*closed*) *cone*  $C \subseteq V$  is a subset of  $V$  such that for any elements  $c_1, c_2 \in C$  and for any positive coefficients  $\gamma_1, \gamma_2 \geq 0$ ,  $\gamma_1 c_1 + \gamma_2 c_2 \in C$ . A finite cone  $C \subseteq V$  admits a halfspace representation in terms a finite number of linear constraints:

$$C = \{x \in V \mid \bigwedge_{i=1}^K (a_i \cdot x \geq 0)\} \tag{22}$$

Alternatively, a Cone can be expressed in terms of the pullback of the positive orthant

$$\mathbb{R}_+^n := \{v \in \mathbb{R}^n \mid \forall i \in [n] : v_i \geq 0\} \tag{23}$$

by a linear transformation  $f : V \rightarrow \mathbb{R}^n$  into  $\mathbb{R}^n$ .

$$\begin{array}{ccc}
 f^*(\mathbb{R}_+^n) & \longrightarrow & \mathbb{R}_+^n \\
 \downarrow \lrcorner & & \downarrow i_+ \\
 V & \xrightarrow{f} & \mathbb{R}^n
 \end{array} \tag{24}$$

$$f^*(\mathbb{R}_+^n) \cong \{v \in V \mid f(v) \in \mathbb{R}_+^n\} \tag{25}$$

Given a cone  $f^*(\mathbb{R}_+^n) \subseteq V$  associated with a finite set of  $n$  linear expressions  $f : V \rightarrow \mathbb{R}^n$ , and a linear transformation  $g : V \rightarrow W$ ,

### 3.6 Subset Projection

A prototypical example wherein an adjoint triple

$$f_!, \exists_f \dashv f^*, f^{-1} \dashv f^!, \forall_f$$

arises is that of functions  $f : X \rightarrow Y$  between sets  $X$  and  $Y$ . The inverse image functor  $f^* : \mathcal{P}Y \rightarrow \mathcal{P}X$  is defined on a subset  $T \subseteq Y$


$$f^*(T) = \{x \in X : f(x) \in T\},$$

and is functorial in the sense that if  $T \subseteq T' \subseteq Y$  then  $f^*(T) \subseteq f^*(T') \subseteq f^*(T)$ . The adjoint functors  $\exists_f, \forall_f : \mathcal{P}X \rightarrow \mathcal{P}Y$  are defined on  $S \subseteq X$  as

$$\begin{aligned}\exists_f(S) &= \{y \in Y : \exists x \in f^*(y) : x \in S\} \\ \forall_f(S) &= \{y \in Y : \forall x \in f^*(y) : x \in S\}\end{aligned}$$

form an adjoint triple in the sense that  $\exists_f \dashv f^* \dashv \forall_f$ :

$$\begin{aligned}\exists_f \dashv f^* : \quad \exists_f(S) \subseteq T &\iff S \subseteq f^*(T) \\ f^* \dashv \forall_f : \quad f^*(T) \subseteq R &\iff T \subseteq \forall_f(R)\end{aligned}$$




Consider a pair of sets  $A$  and  $B$  and a subset  $S \subseteq A \times B$  of their cartesian product. The projection morphisms associated with  $A \times B$  are  $p : A \times B \rightarrow A$  and  $q : A \times B \rightarrow B$ . The projection of the subset  $S$  onto  $A$  is then the subset  $X \subseteq A$  defined by:

$$X = \{a \in A \mid \exists s \in S, p(s) = a\}$$

$$S \subseteq p^*(X) \iff \exists_p(S) \subseteq X \tag{26}$$

### 3.7 Optimization of real-valued functions



### Comments on selected references

This section is temporary and reserved for recording comments toward various references.

- [vistoli2004notes](#) [\[Vis04\]](#)
- [street1974fibrations](#) [\[Str74\]](#)
- [koudenburg2018categorical](#) [\[Kou18\]](#)
- [brown2009algebraic](#) [\[BS09\]](#)
- [lurie2009higher](#) [\[Lur09\]](#)
- [shulman2008framed](#) [\[Shu08\]](#)
- [boyd2004convex](#) [\[BV04\]](#)
- [bogart2013hom](#) [\[BCG13\]](#)
- [gubeladze2016affine](#) [\[Gub16\]](#)
- [fausk2003isomorphisms](#) [\[FHM03\]](#)
- [hofstra2011dialectica](#) [\[Hof11\]](#)

- ~~ponto2012duality~~ Ponto and Shulman [~~PS12~~]
- ~~mac2013categories~~ Mac Lane [~~Mac13~~]
- ~~ziegler2012lectures~~ Ziegler [~~Zie12~~]
- [Spectrahedron](https://www.youtube.com/watch?v=AevFRN5sx0U) are interesting semi-algebraic sets. (<https://www.youtube.com/watch?v=AevFRN5sx0U>).

## References

- |   |  |
|---|--|
| <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">bogart2013hom</div> <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">009algebraic</div> <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">oyd2004convex</div> <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">isomorphisms</div> <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">baum2003your</div> <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">ze2016affine</div> <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">11dialectica</div> <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">8categorical</div> <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">os2015proofs</div> <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">ie2009higher</div> <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">13categories</div> <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">o2012duality</div> <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">an2008framed</div> <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">74fibrations</div> <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">oli2004notes</div> <div style="border: 1px solid black; padding: 2px; margin-bottom: 5px;">2012lectures</div> | <div style="margin-bottom: 10px;">[BCG13] Tristram Bogart, Mark Contois, and Joseph Gubeladze. “Hom-polytopes”. In: <i>Mathematische Zeitschrift</i> 273.3-4 (2013), pp. 1267–1296.</div> <div style="margin-bottom: 10px;">[BS09] Ronald Brown and Rafael Sivera. “Algebraic colimit calculations in homotopy theory using fibred and cofibred categories”. In: <i>Theory and Applications of Categories</i> 22.8 (2009), pp. 222–251.</div> <div style="margin-bottom: 10px;">[BV04] Stephen Boyd and Lieven Vandenbergh. <i>Convex optimization</i>. Cambridge university press, 2004.</div> <div style="margin-bottom: 10px;">[FHM03] Halvard Fausk, Po Hu, and J Peter May. “Isomorphisms between left and right adjoints”. In: <i>Theory Appl. Categ</i> 11.4 (2003), pp. 107–131.</div> <div style="margin-bottom: 10px;">[Grü03] Branko Grünbaum. “Are your polyhedra the same as my polyhedra?” In: <i>Discrete and computational geometry</i>. Springer, 2003, pp. 461–488.</div> <div style="margin-bottom: 10px;">[Gub16] Joseph Gubeladze. “Affine-compact functors”. In: <i>Advances in Geometry</i> (2016).</div> <div style="margin-bottom: 10px;">[Hof11] Pieter Hofstra. “The dialectica monad and its cousins”. In: <i>Models, logics, and higherdimensional categories: A tribute to the work of Mihály Makkai</i> 53 (2011), pp. 107–139.</div> <div style="margin-bottom: 10px;">[Kou18] Seerp Roald Koudenburg. “A categorical approach to the maximum theorem”. In: <i>Journal of Pure and Applied Algebra</i> 222.8 (2018), pp. 2099–2142.</div> <div style="margin-bottom: 10px;">[Lak15] Imre Lakatos. <i>Proofs and Refutations: The Logic of Mathematical Discovery (Cambridge Philosophy Classics)</i>. Cambridge University Press, 2015. ISBN: 1107534054.</div> <div style="margin-bottom: 10px;">[Lur09] Jacob Lurie. <i>Higher Topos Theory (AM-170)</i>. Vol. 189. Princeton University Press, 2009.</div> <div style="margin-bottom: 10px;">[Mac13] Saunders Mac Lane. <i>Categories for the working mathematician</i>. Vol. 5. Springer Science &amp; Business Media, 2013.</div> <div style="margin-bottom: 10px;">[PS12] Kate Ponto and Michael Shulman. “Duality and traces for indexed monoidal categories”. In: <i>Theory and Applications of Categories</i> 26.23 (2012), pp. 582–659.</div> <div style="margin-bottom: 10px;">[Shu08] Michael Shulman. “Framed bicategories and monoidal fibrations”. In: <i>Theory and applications of categories</i> 20.18 (2008), pp. 650–738.</div> <div style="margin-bottom: 10px;">[Str74] Ross Street. “Fibrations and Yoneda’s lemma in a 2-category”. In: <i>Category seminar</i>. Springer. 1974, pp. 104–133.</div> <div style="margin-bottom: 10px;">[Vis04] Angelo Vistoli. “Notes on Grothendieck topologies, fibered categories and descent theory”. In: <i>arXiv preprint math/0412512</i> (2004).</div> <div style="margin-bottom: 10px;">[Zie12] Günter M Ziegler. <i>Lectures on polytopes</i>. Vol. 152. Springer Science &amp; Business Media, 2012.</div> |
|---|--|