

On the Kronecker Product of S_n Characters

YOAV DVIR*

*Department of Theoretical Mathematics, The Weizmann Institute of Science,
Rehovot 76100, Israel*

Communicated by Gordon James

Received February 15, 1991

Let $\chi_\nu, \chi_\lambda, \chi_\mu$ be irreducible S_n characters and assume χ_μ appears in the Kronecker product $\chi_\lambda \otimes \chi_\nu$ with maximal first part ρ_1 . Then $\rho_1 = |\lambda \cap \mu| = \sum \min(\lambda_i, \mu_i)$. A similar result holds for the maximal first column. We also give a recursive formula for $\chi_\lambda \otimes \chi_\mu$. As an application, we show that if $n = \lambda_1 + \mu_1 - \rho_1$, then $\langle \chi_\lambda \otimes \chi_\mu, \chi_\nu \rangle_{S_n} = \langle \chi_{(\lambda_2, \lambda_3, \dots)} \hat{\otimes} \chi_{(\mu_2, \mu_3, \dots)}, \chi_{(\rho_2, \rho_3, \dots)} \rangle_{S_{n-\rho_1}}$, where $\hat{\otimes}$ denotes the outer tensor product. These results are applied to study the character $\sum \chi_\lambda \otimes \chi_\lambda$ where λ runs through the partitions with no more than k parts. This character is closely related to the polynomial identities of the algebra of $k \times k$ matrices. © 1993 Academic Press, Inc.

0. INTRODUCTION

Throughout this paper the base field is \mathbb{C} . Let S_n denote the symmetric group. The partition $\lambda = (\lambda_1, \lambda_2, \dots)$ of n ($\lambda \vdash n$) corresponds to the irreducible S_n character $\lambda \leftrightarrow \chi_\lambda$. Given $\lambda, \mu \vdash n$ then $\langle \chi_\mu, \chi_\lambda \rangle_{S_n} = \delta_{\mu, \lambda}$ defines an inner product: $\langle \chi, \chi_\lambda \rangle_{S_n}$ is the coefficient of χ_λ in the S_n character χ . Let λ' denote the conjugate partition of λ . Given an S_n character χ , we have

$$\chi = \sum_{\lambda \vdash n} \langle \chi, \chi_\lambda \rangle_{S_n} \cdot \chi_\lambda.$$

Define (the conjugate character)

$$\chi' = \sum_{\lambda \vdash n} \langle \chi, \chi_\lambda \rangle_{S_n} \cdot \chi_{\lambda'}.$$

Define the width of χ as follows: $w(\chi_\lambda) = w(\lambda) = w(\lambda_1, \lambda_2, \dots) = \lambda_1$, and

* This work was done under the supervision of Professor Amitai Regev as a part of the Ph.D. thesis in the Department of Theoretical Mathematics, The Weizmann Institute of Science.

$w(\chi) = \max\{w(\lambda') \mid \langle \chi, \chi_\lambda \rangle_{S_n} \neq 0\}$. Define the height $h(\chi) = w(\chi')$. Obviously

$$h(\chi) = \max\{w(\lambda') \mid \langle \chi, \chi_\lambda \rangle_{S_n} \neq 0\}.$$

Let $\chi_\lambda \otimes \chi_\mu$ be the inner (Kronecker) products, and let

$$\chi_\lambda \otimes \chi_\mu = \sum_{\rho \vdash n} c(\lambda, \mu, \rho) \chi_\rho \quad (\text{i.e., } c(\lambda, \mu, \rho) = \langle \chi_\lambda \otimes \chi_\mu, \chi_\rho \rangle_{S_n})$$

denote the decomposition of the inner product as a sum of irreducible characters. The known algorithms for calculating (i.e., decomposing) $\chi_\lambda \otimes \chi_\mu$ [Gar-Rem; Jam-Ker, 2.9] are rather ineffective in the general case. The problem of finding effective ways for calculating, and even estimating $\chi_\lambda \otimes \chi_\mu$, is, probably, the last major problem in the ordinary representation theory of S_n .

The inequality $h(\chi_\lambda \otimes \chi_\mu) \leq h(\chi_\lambda) \cdot h(\chi_\mu)$ was proved in [Reg 1], and was crucial for the results there. The main result here is a precise formula which expresses $h(\chi_\lambda \otimes \chi_\mu)$ as the area of the intersection of the young diagrams of λ and of μ' , the conjugate of μ . This is

THEOREM (1.6 below). *Write $\lambda \cap \mu = (\min(\lambda_1, \mu_1), \min(\lambda_2, \mu_2), \dots)$, so $|\lambda \cap \mu|$ is the area of the intersection of the diagrams of λ and of μ . Then*

$$(a) \quad w(\chi_\lambda \otimes \chi_\mu) = |\lambda \cap \mu|$$

$$(b) \quad h(\chi_\lambda \otimes \chi_\mu) = |\lambda \cap \mu'|.$$

Note that $|\lambda \cap \mu'| \leq \lambda'_1 \cdot \mu'_1 = h(\lambda) \cdot h(\mu)$. Thus 1.6(b) here replaces the estimate in [Reg 1] (1.7 here) by a precise formula. In 1.8 we extend part of 1.6 to skew characters.

In Section 2 we establish a recursive formula, of interest in its own, for computing $\langle \chi_\lambda \otimes \chi_\mu, \chi_\rho \rangle_{S_n}$. The recursion in that formula is on n and on $n - \rho_1$. In the special case that $\rho_1 = |\lambda \cap \mu|$ we obtain

THEOREM (2.4 below). *Given partitions α, β , let α/β denote the skew partition with corresponding skew character $\chi_{\alpha/\beta}$ [Mac]. Let $\lambda, \mu, \rho \vdash n$, $\rho_1 = |\lambda \cap \mu|$. Then*

$$\langle \chi_\lambda \otimes \chi_\mu, \chi_\rho \rangle_{S_n} = \langle \chi_{\lambda/\lambda \cap \mu} \otimes \chi_{\mu/\lambda \cap \mu}, \chi_{(\rho_2, \rho_3, \dots)} \rangle_{S_{n-\rho_1}}.$$

In section 3 we apply these results to decompose $\chi_\lambda \otimes \chi_\mu$ in some special cases. Let $\lambda = (\lambda_1, \lambda_2, \dots)$. Then denote $\theta(\lambda) = (\lambda_2, \lambda_3, \dots)$ and $d(\lambda) = |\theta(\lambda)| = n - \lambda_1$. Let $\alpha \vdash n$, $\beta \vdash m$. Then $\chi_\alpha \hat{\otimes} \chi_\beta$ denotes the outer product of χ_α and χ_β . The decomposition of $\chi_\alpha \hat{\otimes} \chi_\beta$ is given by the Littlewood-Richardson Rule [Jam-Ker]. Our main result in Section 3 is

THEOREM (3.3 below). *Given λ, μ, ρ partitions of n such that $d(\lambda) + d(\mu) = d(\rho)$ (or equivalently $n = \lambda_1 + \mu_1 - \rho_1$) then*

$$\langle \chi_\lambda \otimes \chi_\mu, \chi_\rho \rangle_{S_n} = \langle \chi_{\theta(\lambda)} \hat{\otimes} \chi_{\theta(\mu)}, \chi_{\theta(\rho)} \rangle_{S_{d(\rho)}}.$$

In Section 4 we apply the above results to study the character

$$\psi(k, n) = \sum_{\lambda \in H(k, 0; n)} \chi_\lambda \otimes \chi_\lambda$$

and its restrictions

$$\left(\sum_{\lambda \in H(k, 0; n)} \chi_\lambda \otimes \chi_\lambda \right)_{S_m},$$

where $m \leq n$ and

$$H(k, 0; n) = \{ \lambda \vdash n \mid \lambda_{k+1} = 0 \}.$$

When $m = n$ and $m = n - 1$, the corresponding characters describe the trace identities of the $k \times k$ matrices, which are closely related to the polynomial identities of such matrices. Write:

$$\left(\sum_{\lambda \in H(k, 0; n)} \chi_\lambda \otimes \chi_\lambda \right)_{S_m} = \sum_{\rho \in H(k^2, 0; m)} a(\rho, k, n) \chi_\rho.$$

The calculation of the coefficients $a(\rho, k, n)$ when $m = n$ and $m = n - 1$ is a major open problem in the theory of polynomial identities. We show (4.3 below) that if $\rho = (\rho_1, \dots, \rho_{k^2})$ is a partition such that $|\rho| \leq n$ and $0 \leq t \leq \rho_{k^2}$ then

$$a((\rho_1, \dots, \rho_{k^2}), k, n) = a((\rho_1 - t, \dots, \rho_{k^2} - t, 0), k, n - k^2 t),$$

i.e., $k^2 \times t$ rectangles can be removed.

In the special case that $\rho = \phi$ is a $k^2 \times t$ rectangle

$$\phi = (\underbrace{t, t, \dots, t}_{k^2})$$

(and also for

$$\phi = (t + 1, \underbrace{t, \dots, t}_{k^2 - 1})$$

we have the following intriguing formula (4.5 below):

$$a(\phi, k, n) = \sum_{\tilde{\lambda} \in H(k, 0; n - k^2 t)} (\deg \chi_{\tilde{\lambda}})^2.$$

Note that

$$\deg \psi(k, n) = \sum_{\lambda \in H(k, 0; n)} (\deg \chi_\lambda)^2.$$

The asymptotics and the generating functions of these sums were studied in [Reg 2, Bec-Reg].

1. THE WIDTH AND HEIGHT OF $\chi_\lambda \otimes \chi_\mu$

All groups here are finite. In addition, the following definitions and notations will hold.

Let $F(G)$ be the ring of class functions of G into \mathbb{C} , $CH(G)$ the subring generated by the characters of G , $\text{Irr}(G)$ the irreducible characters.

Denote by $\langle, \rangle_G: F(G) \times F(G) \rightarrow \mathbb{C}$ the natural bilinear form defined by $\langle f_1, f_2 \rangle_G = \delta_{f_1, f_2}$ for all $f_1, f_2 \in \text{Irr}(G)$.

Let H, G be finite groups, $H \subseteq G$. For any $f \in F(H)$, f^G will denote the induced class function and for any $f \in F(G)$, f_H will denote the restricted (to H) class function.

For groups G_1, G_2 , $f_i \in F(G_i)$, define $f_1 \times f_2 \in F(G_1 \times G_2)$ by $(f_1 \times f_2)(g_1, g_2) = f_1(g_1) \cdot f_2(g_2)$. We denote the inner (or the Kronecker) product of $f_1, f_2 \in F(G)$ by $f_1 \otimes f_2$. Denote by $f \hat{\otimes} g$ the outer product of $f \in F(S_m)$ and $g \in F(S_n)$ [Mac].

Following [Mac], define the skew character $\chi_{\lambda/\mu}$ of the skew tableau λ/μ as follows: let $n, m \in \mathbb{N} \cup \{0\}$, $\lambda \vdash n$, and $\mu \vdash m$; then $\chi_{\lambda/\mu} = \sum_{\rho \vdash n-m} \langle \chi_\lambda, \chi_\mu \hat{\otimes} \chi_\rho \rangle_{S_n} \chi_\rho$.

Throughout we will assume S_0 is the group of one element, with $\chi_{(0)} = \chi_{(0, 0, \dots)}$ the unique trivial character of S_0 . In this case, for any $\lambda \vdash n$, $\chi_\lambda \hat{\otimes} \chi_{(0)} = \chi_\lambda$ (by definition).

DEFINITION 1.1. Let $\lambda \vdash n$, $\mu \vdash m$, $n \geq m$, and $S_m \subseteq S_n$ in the natural embedding. Define $k(\lambda, \mu) = \langle \chi_\lambda, (\chi_\mu)^{S_n} \rangle_{S_n}$. By Frobenius Reciprocity:

$$k(\lambda, \mu) = \langle (\chi_\lambda)_{S_m}, \chi_\mu \rangle_{S_m}.$$

Remarks 1.2. Let $n \geq m$, $\lambda \vdash n$, $\mu \vdash m$.

(a) It is well known that $k(\lambda, \mu) > 0$ if and only if $\mu \subseteq \lambda$ [Mac, 5.7, 5.12]. Actually, $k(\lambda, \mu)$ is the degree of $\chi_{\lambda/\mu}$.

In later sections we shall also need:

(b) Let $\alpha \vdash m - m$ and $\langle \chi_\alpha \hat{\otimes} \chi_\mu, \chi_\lambda \rangle_{S_n} \neq 0$. Then $\alpha, \mu \subseteq \lambda$ (by the Littlewood–Richardson Rule).

(c) $(\chi_\lambda)_{S_m \times S_{n-m}} = \sum_{\theta \vdash m} \chi_\theta \times \chi_{\lambda/\theta}$ [Jam-Ker, 2.3.12].

Clearly we can also write:

$$(1.3) \quad (\chi_\lambda)_{S_m} = \sum_{\substack{\alpha \vdash m \\ \alpha \subseteq \lambda}} k(\lambda, \alpha) \chi_\alpha$$

$$(1.4) \quad (\chi_\mu)^{S_n} = \sum_{\substack{\beta \vdash n \\ \beta \supseteq \mu}} k(\beta, \mu) \chi_\beta.$$

PROPOSITION 1.5. *Let $(m) \vdash m$, $\lambda, \mu \vdash n$, and $m \leq n$. Then:*

$$\sum_{\substack{\beta \vdash n \\ \beta \supseteq (m)}} k(\beta, (m)) \langle \chi_\beta, \chi_\lambda \otimes \chi_\mu \rangle_{S_n} = \sum_{\substack{\alpha \vdash m \\ \alpha \subseteq \lambda \cap \mu}} k(\lambda, \alpha) k(\mu, \alpha).$$

Proof. Let

$$p = \langle (\chi_{(m)})^{S_n}, \chi_\lambda \otimes \chi_\mu \rangle_{S_n}$$

and

$$q = \langle \chi_{(m)}, (\chi_\lambda)_{S_m} \otimes (\chi_\mu)_{S_m} \rangle_{S_m}.$$

By Frobenius Reciprocity (and $(\chi_\lambda \otimes \chi_\mu)_{S_m} = (\chi_\lambda)_{S_m} \otimes (\chi_\mu)_{S_m}$) $p = q$. By (1.3)

$$q = \left\langle \chi_{(m)}, \left(\sum_{\substack{\alpha \vdash m \\ \alpha \subseteq \lambda}} k(\lambda, \alpha) \chi_\alpha \right) \otimes \left(\sum_{\substack{\gamma \vdash m \\ \gamma \subseteq \mu}} k(\mu, \gamma) \chi_\gamma \right) \right\rangle_{S_m},$$

and by (1.4)

$$p = \left\langle \sum_{\substack{\beta \vdash n \\ \beta \supseteq (m)}} k(\beta, (m)) \chi_\beta, \chi_\lambda \otimes \chi_\mu \right\rangle_{S_n}.$$

It easily follows from basic properties of character tables that for any $\alpha, \gamma \vdash m$, $\langle \chi_\alpha \otimes \chi_\gamma, \chi_{(m)} \rangle_{S_m} = \delta_{\alpha\gamma}$. Thus

$$q = \sum_{\substack{\alpha \vdash m \\ \alpha \subseteq \lambda \cap \mu}} k(\lambda, \alpha) k(\mu, \alpha)$$

and the statement follows. ■

The main result in this section is (recall $w(\chi)$, $h(\chi)$, and $\lambda \cap \mu$ from Sect. 0):

THEOREM 1.6. For any partitions λ, μ of n the following holds:

- (a) $w(\chi_\lambda \otimes \chi_\mu) = |\lambda \cap \mu|$
 (b) $h(\chi_\lambda \otimes \chi_\mu) = |\lambda \cap \mu'|.$

Proof. (a). We show first that $w(\chi_\lambda \otimes \chi_\mu) \leq |\lambda \cap \mu|$. In order to do so, apply 1.5 with $|\lambda \cap \mu| = m - 1$,

$$\begin{aligned} & \sum_{\substack{\beta \vdash n \\ \beta \supseteq (|\lambda \cap \mu| + 1)}} k(\beta, |\lambda \cap \mu| + 1) \langle \chi_\lambda \otimes \chi_\mu, \chi_\beta \rangle_{S_n} \\ &= \sum_{\substack{\alpha \vdash |\lambda \cap \mu| + 1 \\ \alpha \subseteq \lambda \cap \mu}} k(\lambda, \alpha) k(\mu, \alpha) = 0, \end{aligned}$$

since clearly, $\{\alpha \mid \alpha \vdash |\lambda \cap \mu| + 1 \text{ and } \alpha \subseteq \lambda \cap \mu\}$ is empty.

For all $\beta = (\beta_1, \beta_2, \dots)$ such that $(|\lambda \cap \mu| + 1) \subseteq \beta$ (i.e., $\beta_1 \geq |\lambda \cap \mu| + 1$), $k(\beta, (|\lambda \cap \mu| + 1)) > 0$ (Remark 1.2(a)); hence $\langle \chi_\lambda \otimes \chi_\mu, \chi_\beta \rangle_{S_n} = 0$. By the definition of w , this implies: $w(\chi_\lambda \otimes \chi_\mu) \leq |\lambda \cap \mu|$.

Next we show that $w(\chi_\lambda \otimes \chi_\mu) \geq |\lambda \cap \mu|$.

Again apply 1.5, with $m = |\lambda \cap \mu|$:

$$\begin{aligned} & \sum_{\substack{\beta \vdash n \\ \beta \supseteq (|\lambda \cap \mu|)}} k(\beta, (|\lambda \cap \mu|)) \langle \chi_\lambda \otimes \chi_\mu, \chi_\beta \rangle_{S_n} \\ &= \sum_{\substack{\alpha \vdash |\lambda \cap \mu| \\ \alpha \subseteq \lambda \cap \mu}} k(\lambda, \alpha) k(\mu, \alpha) = k(\lambda, \lambda \cap \mu) k(\mu, \lambda \cap \mu) > 0. \end{aligned}$$

Hence there exists $\beta = (\beta_1, \beta_2, \dots) \vdash n$ with $\beta_1 \geq |\lambda \cap \mu|$ (i.e., $w(\beta) \geq |\lambda \cap \mu|$) such that $\langle \chi_\beta, \chi_\lambda \otimes \chi_\mu \rangle_{S_n} \neq 0$ so:

$$w(\chi_\lambda \otimes \chi_\mu) \geq |\lambda \cap \mu|.$$

The proof of (a) is now complete.

(b) Since $\chi_\lambda \otimes \chi_{\mu'} = (\chi_\lambda \otimes \chi_\mu)'$, therefore

$$h(\chi_\lambda \otimes \chi_\mu) = w((\chi_\lambda \otimes \chi_\mu)') = w(\chi_\lambda \otimes \chi_{\mu'}) = |\lambda \cap \mu'| \quad (\text{by 1.6(a)}). \quad \blacksquare$$

COROLLARY 1.7 [Reg 1]. Let $\lambda, \mu \vdash n$, $h(\lambda) = k$, and $h(\mu) = l$. Then

$$h(\chi_\lambda \otimes \chi_\mu) \leq kl.$$

Proof. Obviously, $|\lambda \cap \mu'| \leq kl$, so 1.7 follows from 1.6(b). \blacksquare

Let χ_λ be an irreducible character appearing in the skew character $\chi_{\lambda/\eta}$. Then by definition of skew characters and 1.2(b), $\alpha \subseteq \lambda$. Thus, we obviously have the following corollary, which extends part of 1.6 to skew characters:

COROLLARY 1.8. Let $m_1, m_2, l \geq 0$, $n_1 = m_1 + l$, $n_2 = m_2 + l$, $\lambda \vdash n_1$, $\eta \vdash m_1$, $\mu \vdash n_2$ and $\xi \vdash m_2$. Then

- (a) $w(\chi_{\lambda/\eta} \otimes \chi_{\mu/\xi}) \leq |\lambda \cap \mu|$.
- (b) $h(\chi_{\lambda/\eta} \otimes \chi_{\mu/\xi}) \leq |\lambda \cap \mu'|$.

2. A RECURSIVE FORMULA FOR $\chi_\lambda \otimes \chi_\mu$

Theorems 2.3, 2.4, and 2.4' below give recursive formulas for the coefficients $c(\lambda, \mu, \rho)$ (see Sect. 0). A major tool here is the well-known Young's Rule [Jam-Ker].

PROPOSITION 2.1. Let $m, l \geq 0$ and $\theta \vdash l$. Define

$$Y(\theta, m) = \{\eta \mid \eta \vdash m + l, \eta_i \geq \theta_i \geq \eta_{i+1} \text{ for all } i \geq 1\}.$$

Then

- (a) Young's Rule can be written as follows: $\chi_{(m)} \hat{\otimes} \chi_\theta = \sum_{\eta \in Y(\theta, m)} \chi_\eta$.
- (b) $\langle \chi_{(m)} \times \chi_\theta, (\chi_\gamma)_{S_m \times S_l} \rangle_{S_m \times S_l} = \begin{cases} 1, & \gamma \in Y(\theta, m) \\ 0, & \text{otherwise.} \end{cases}$
- (c) Let $m \geq \theta_1$, and define $\rho \vdash m + l$ as follows: $\rho_1 = m$, $\rho_i = \theta_{i-1}$ ($i \geq 2$); then $\rho \in Y(\theta, m)$. Also, if $\eta \in Y(\theta, m)$ and $\eta \neq \rho$ then $\eta_1 > m$.

Proof. (a) See [Jam-Ker, 2.3.14].

(b) Applying Frobenius Reciprocity, we have

$$\begin{aligned} \langle \chi_{(m)} \times \chi_\theta, (\chi_\gamma)_{S_m \times S_l} \rangle_{S_m \times S_l} &= \langle (\chi_{(m)} \times \chi_\theta)^{S_m + l}, \chi_\gamma \rangle_{S_{m+l}} \\ &= \langle \chi_{(m)} \hat{\otimes} \chi_\theta, \chi_\gamma \rangle_{S_{m+l}} \end{aligned}$$

(Young's Rule)

$$= \left\langle \sum_{\eta \in Y(\theta, m)} \chi_\eta, \chi_\gamma \right\rangle_{S_{m+l}} = \begin{cases} 1, & \gamma \in Y(\theta, m); \\ 0, & \text{otherwise.} \end{cases}$$

(c) Obviously, $\rho \in Y(\theta, m)$. Let $\eta \in Y(\theta, m)$, $\eta \neq \rho$, and assume $\eta_1 \leq m$. Since $\eta \in Y(\theta, m)$, therefore $\eta_i \leq \theta_{i-1} = \rho_i$ ($i \geq 2$), by the definition of $Y(\theta, m)$. Thus $\eta_i \leq \rho_i$ ($i \geq 1$). But $\eta, \rho \vdash m + l$; hence $\eta = \rho$, contradiction. ■

Recall the coefficients $c(\lambda, \mu, \rho)$ (see Sect. 0). We shall need the following technical

PROPOSITION 2.2. Let $m, l \geq 0$, $\lambda, \mu \vdash m+l$, $\theta \vdash l$. Then

- (a) $(\chi_\lambda \otimes \chi_\mu)_{S_m \times S_l}, \chi(m) \times \chi_\theta \rangle_{S_m \times S_l} = \sum_{\rho \in Y(\theta, m)} c(\lambda, \mu, \rho)$, and also
 (b) $\langle (\chi_\lambda \otimes \chi_\mu)_{S_m \times S_l}, \chi(m) \times \chi_\theta \rangle_{S_m \times S_l} = \sum_{\alpha \vdash m} \langle \chi_{\lambda/\alpha} \otimes \chi_{\mu/\alpha}, \chi_\theta \rangle_{S_l}$.

Proof. (a)

$$\begin{aligned} & \langle (\chi_\lambda \otimes \chi_\mu)_{S_m \times S_l}, \chi(m) \times \chi_\theta \rangle_{S_m \times S_l} \\ &= \left\langle \left(\sum_{\rho \vdash m+l} c(\lambda, \mu, \rho) \chi_\rho \right)_{S_m \times S_l}, \chi(m) \times \chi_\theta \right\rangle_{S_m \times S_l} \quad (\text{by linearity}) \\ &= \sum_{\rho \vdash m+l} c(\lambda, \mu, \rho) \langle (\chi_\rho)_{S_m \times S_l}, \chi(m) \times \chi_\theta \rangle_{S_m \times S_l} \quad (\text{by 2.1(b)}) \\ &= \sum_{\rho \in Y(\theta, m)} c(\lambda, \mu, \rho). \end{aligned}$$

(b) The following is well known: Let H, G be finite groups, $H \subseteq G$, and $f_1, f_2 \in F(G)$. Then $(f_1 \otimes f_2)_H = f_{1_H} \otimes f_{2_H}$. Therefore

$$\begin{aligned} & \langle (\chi_\lambda \otimes \chi_\mu)_{S_m \times S_l}, \chi(m) \times \chi_\theta \rangle_{S_m \times S_l} \\ &= \langle (\chi_\lambda)_{S_m \times S_l} \otimes (\chi_\mu)_{S_m \times S_l}, \chi(m) \times \chi_\theta \rangle_{S_m \times S_l} \quad (\text{by 1.2(c)}) \\ &= \left\langle \left(\sum_{\alpha \vdash m} \chi_\alpha \times \chi_{\lambda/\alpha} \right) \otimes \left(\sum_{\beta \vdash m} \chi_\beta \times \chi_{\mu/\beta} \right), \chi(m) \times \chi_\theta \right\rangle_{S_m \times S_l} \\ & \quad (\text{by linearity}) \\ &= \sum_{\alpha, \beta \vdash m} \langle \chi_\alpha \times \chi_{\lambda/\alpha} \otimes (\chi_\beta \times \chi_{\mu/\beta}), \chi(m) \times \chi_\theta \rangle_{S_m \times S_l} = D. \end{aligned}$$

For any finite groups H, G , if $f_1, f_2 \in F(H)$, $g_1, g_2 \in F(G)$ then $(f_1 \times g_1) \otimes (f_2 \times g_2) = (f_1 \otimes f_2) \times (g_1 \otimes g_2)$. Also, $\langle f_1 \times g_1, f_2 \times g_2 \rangle_{H \times G} = \langle f_1, f_2 \rangle_H \cdot \langle g_1, g_2 \rangle_G$. Thus

$$\begin{aligned} D &= \sum_{\alpha, \beta \vdash m} \langle (\chi_\alpha \otimes \chi_\beta) \times (\chi_{\lambda/\alpha} \otimes \chi_{\mu/\beta}), \chi(m) \times \chi_\theta \rangle_{S_m \times S_l} \\ &= \sum_{\alpha, \beta \vdash m} \langle \chi_\alpha \otimes \chi_\beta, \chi(m) \rangle_{S_m} \langle \chi_{\lambda/\alpha} \otimes \chi_{\mu/\beta}, \chi_\theta \rangle_{S_l} \end{aligned}$$

Note that $\chi(m)$ is the identity character of S_m ; hence $\langle \chi_\alpha \otimes \chi_\beta, \chi(m) \rangle = \delta_{\alpha, \beta}$. Thus

$$\begin{aligned} D &= \langle (\chi_\lambda \otimes \chi_\mu)_{S_m \times S_l}, \chi(m) \otimes \chi_\theta \rangle_{S_m \times S_l} \\ &= \sum_{\alpha \vdash m} \langle \chi_{\lambda/\alpha} \otimes \chi_{\mu/\alpha}, \chi_\theta \rangle_{S_l}. \quad \blacksquare \end{aligned}$$

We can now give a recursion formula for the coefficients $c(\lambda, \mu, \rho)$.

THEOREM 2.3. *Let $\lambda, \mu, \rho \vdash m + l = n$, such that $m = \rho_1$. Define $\theta \vdash l$ as follows: $\theta_i = \rho_{i+1}$, $i \geq 1$. Then*

$$c(\lambda, \mu, \rho) = \sum_{\substack{\alpha \vdash m \\ \alpha \subseteq \lambda \cap \mu}} \langle \chi_{\lambda/\alpha} \otimes \chi_{\mu/\alpha}, \chi_\theta \rangle_{S_l} - \sum_{\substack{\eta \in Y(\theta, m) \\ \eta \neq \rho \\ \eta_1 \leq |\lambda \cap \mu|}} c(\lambda, \mu, \eta).$$

Note that this is indeed a recursive formula: the skew characters can be decomposed into irreducible ones of S_l [Gar-Rem, Mac]; thus the problem (in the first summand) is reduced to smaller symmetric groups. Also, in the second summand, $\eta_1 > \rho_1$ (by 2.1.c); hence $n - \eta_1 < n - \rho_1$ and the recursion is on the invariant $I(\lambda, \mu, \rho) = n - \rho_1$ (note that when $I(\lambda, \mu, \rho) = 0$, $\rho = (n)$, and $c(\lambda, \mu, (n)) = \delta_{\lambda\mu}$).

Proof. Denote $L = \sum_{\eta \in Y(\theta, m)} c(\lambda, \mu, \eta)$ and $R = \sum_{\alpha \vdash m} \langle \chi_{\lambda/\alpha} \otimes \chi_{\mu/\alpha}, \chi_\theta \rangle_{S_l}$. By 1.6,

$$L = \sum_{\substack{\eta \in Y(\theta, m) \\ \eta_1 \leq |\lambda \cap \mu|}} c(\lambda, \mu, \eta)$$

and it easily follows from 1.2(a) that

$$R = \sum_{\substack{\alpha \vdash m \\ \alpha \subseteq \lambda \cap \mu}} \langle \chi_{\lambda/\alpha} \otimes \chi_{\mu/\alpha}, \chi_\theta \rangle_{S_l}.$$

By 2.2 $L = R$; thus

$$\sum_{\substack{\eta \in Y(\theta, m) \\ \eta_1 \leq |\lambda \cap \mu|}} c(\lambda, \mu, \eta) = \sum_{\substack{\alpha \vdash m \\ \alpha \subseteq \lambda \cap \mu}} \langle \chi_{\lambda/\alpha} \otimes \chi_{\mu/\alpha}, \chi_\theta \rangle_{S_l}.$$

By the definition of $Y(\theta, m)$ it follows that $\rho \in Y(\theta, m)$, and the formula follows. ■

In the case $\rho_1 = |\lambda \cap \mu|$ the formula is particularly efficient:

THEOREM 2.4. *Let $\lambda, \mu, \rho \vdash m + l$, $\rho_1 = |\lambda \cap \mu| = m$, and define $\theta = \theta(\rho) \vdash l$ by $\theta_i = \rho_{i+1}$, $i \geq 1$. Then*

$$c(\lambda, \mu, \rho) = \langle \chi_{\lambda/\lambda \cap \mu} \otimes \chi_{\mu/\lambda \cap \mu}, \chi_\theta \rangle_{S_l}.$$

Proof. If $\alpha \vdash |\lambda \cap \mu|$ and $\alpha \subseteq \lambda \cap \mu$ then $\alpha = \lambda \cap \mu$. For any $\eta \in Y(\theta, m)$ with $\eta \neq \rho$, $\eta_1 > \rho_1 = |\lambda \cap \mu|$ (by 2.1(c)) and therefore $c(\lambda, \mu, \eta) = 0$ (by 1.6(a)). The proof now follows from 2.3. ■

There is a conjugate to 2.4:

THEOREM 2.4'. *Let $\lambda, \mu, \rho \vdash m+l$, $h(\rho) = |\lambda \cap \mu'| = m$, and define $\zeta = \zeta(\rho) \vdash n$ by $\zeta_i = \rho_i - 1$, $i \geq 1$, and $\rho_i \geq 1$. Then*

$$c(\lambda, \mu, \rho) = \langle \chi_{\lambda/\lambda \cap \mu'} \otimes \chi_{\mu/\lambda' \cap \mu}, \chi_{\zeta} \rangle_{S_l}.$$

Proof. We make use of the following well-known properties of conjugation:

Let χ, ψ be S_n characters, then $\langle \chi, \psi \rangle_{S_n} = \langle \chi', \psi' \rangle_{S_n}$ and $\chi \otimes \psi' = (\chi \otimes \psi)'$. Thus $c(\lambda, \mu, \rho) = c(\lambda, \mu', \rho')$. Finally let $\alpha \vdash m \leq n$. Then $(\chi_{\lambda/\alpha})' = \chi_{\lambda'/\alpha'}$. Hence

$$\begin{aligned} c(\lambda, \mu, \rho) &= c(\lambda, \mu', \rho') \quad (\text{by 2.4}) \\ &= \langle \chi_{\lambda/\lambda \cap \mu'} \otimes \chi_{\mu'/\lambda \cap \mu'}, \chi_{\zeta'} \rangle_{S_{n-m}} \\ &= \langle (\chi_{\lambda/\lambda \cap \mu'} \otimes \chi_{\mu'/\lambda \cap \mu'})', \chi_{\zeta} \rangle_{S_{n-m}} \\ &= \langle \chi_{\lambda/\lambda \cap \mu'} \otimes (\chi_{\mu'/\lambda \cap \mu'})', \chi_{\zeta} \rangle_{S_{n-m}} \\ &= \langle \chi_{\lambda/\lambda \cap \mu'} \otimes \chi_{\mu/\lambda' \cap \mu}, \chi_{\zeta} \rangle_{S_{n-m}}. \quad \blacksquare \end{aligned}$$

3. APPLICATIONS: EVALUATING $\langle \chi_{\lambda} \otimes \chi_{\mu}, \chi_{\rho} \rangle_{S_n}$ WHEN $d(\lambda) + d(\mu) = d(\rho)$

Let $\alpha = (\alpha_1, \alpha_2, \dots)$ be a partition, and define $d(\alpha) = \sum_{i \geq 2} \alpha_i = |\alpha| - \alpha_1$. In this section we consider triples of partitions $\lambda, \mu, \rho \vdash n$ for which

$$d(\lambda) + d(\mu) = d(\rho).$$

For such partitions we reduce the calculation of $c(\lambda, \mu, \rho)$ to that of outer products (given by the Littlewood–Richardson rule). This is 3.3 below.

Given $\alpha \vdash n$ we denote $\theta(\alpha) = (\alpha_2, \alpha_3, \dots)$; $\theta(\alpha) \vdash d(\alpha)$.

The proof of 3.3 requires the following two lemmas.

LEMMA 3.1. *Let $\lambda, \mu, \rho \vdash n$ be any partitions. Then*

- (a) $n - |\lambda_1 - \rho_1| \geq |\lambda \cap \rho|$.
- (b) *If in addition $d(\lambda) + d(\mu) = d(\rho)$ then*
 - (b.1) $\mu_1 = n - \lambda_1 + \rho_1$.
 - (b.2) $\lambda_1 \geq \rho_1$.
 - (b.3) $\mu_1 = |\lambda \cap \rho|$ *if and only if* $\theta(\lambda) \subseteq \theta(\rho)$.

Proof. (a) Without loss of generality we may assume that $\lambda_1 \geq \rho_1$. Now $n = \lambda_1 + \lambda_2 + \dots$; hence

$$\begin{aligned} n - |\lambda_1 - \rho_1| &= \rho_1 + \lambda_2 + \lambda_3 + \dots \\ &\geq \min(\rho_1, \lambda_1) + \min(\rho_2, \lambda_2) + \dots = |\lambda \cap \rho|. \end{aligned}$$

(b.1) By assumption $d(\lambda) + d(\mu) = d(\rho)$, so $(n - \lambda_1) + (n - \mu_1) = (n - \rho_1)$; hence $\mu_1 = n - \lambda_1 + \rho_1$.

(b.2) Since $\mu_1 = n - \lambda_1 + \rho_1$ and $n \geq \mu_1$, therefore $\lambda_1 \geq \rho_1$.

(b.3) Assume first that $\mu_1 = |\lambda \cap \rho|$, so (b.1) $|\lambda \cap \rho| = n - \lambda_1 + \rho_1 = \rho_1 + \sum_{i \geq 2} \lambda_i$. Also $|\lambda \cap \rho| = \sum_{i \geq 1} \min(\lambda_i, \rho_i) = \rho_1 + \sum_{i \geq 2} \min(\lambda_i, \rho_i)$ (by (b.2)). Thus $\sum_{i \geq 2} \lambda_i = \sum_{i \geq 2} \min(\lambda_i, \rho_i)$; hence $\theta(\lambda) \subseteq \theta(\rho)$.

Next, assume $\theta(\lambda) \subseteq \theta(\rho)$, or equivalently, $\lambda_i \leq \rho_i$ ($i \geq 2$). By (b.2) $\lambda_1 \geq \rho_1$, so $|\lambda \cap \rho| = \sum_{i \geq 1} \min(\lambda_i, \rho_i) = \rho_1 + \sum_{i \geq 2} \lambda_i = \rho_1 + n - \lambda_1 = \mu_1$ (b.1). ■

LEMMA 3.2. As in 3.1(b), let $\lambda, \mu, \rho \vdash n$ with $d(\lambda) + d(\mu) = d(\rho)$, and assume in addition that $\langle \chi_\lambda \otimes \chi_\mu, \chi_\rho \rangle_{S_n} \neq 0$. Then

$$\mu_1 = |\lambda \cap \rho|.$$

Proof. Since $\langle \chi_\lambda \otimes \chi_\rho, \chi_\mu \rangle_{S_n} = \langle \chi_\lambda \otimes \chi_\mu, \chi_\rho \rangle_{S_n} \neq 0$, by 1.6(a)

$$\begin{aligned} \mu_1 &\leq |\lambda \cap \rho| \\ &\leq n - |\lambda_1 - \rho_1| \quad (\text{by 3.1(a)}) \\ &= n - \lambda_1 + \rho_1 \quad (\text{by 3.1(b.2)}) \\ &= \mu_1 \quad (\text{by 3.1(b.1)}). \end{aligned}$$

Thus $\mu_1 = |\lambda \cap \rho|$. ■

THEOREM 3.3. Let $\lambda, \mu, \rho \vdash n$ and $d(\lambda) + d(\mu) = d(\rho)$. Then

$$\langle \chi_\lambda \otimes \chi_\mu, \chi_\rho \rangle_{S_n} = \langle \chi_{\theta(\lambda)} \hat{\otimes} \chi_{\theta(\mu)}, \chi_{\theta(\rho)} \rangle_{S_{d(\rho)}}.$$

Proof.

Case 1. $\mu_1 = |\lambda \cap \rho|$.

Denote $\eta = \lambda \cap \rho$ and apply 2.4:

$$\begin{aligned} \langle \chi_\lambda \otimes \chi_\mu, \chi_\rho \rangle_{S_n} &= \langle \chi_\lambda \otimes \chi_\rho, \chi_\mu \rangle_{S_n} \\ &= \langle \chi_{\lambda/\eta} \otimes \chi_{\rho/\eta}, \chi_{\theta(\mu)} \rangle_{S_{n-\mu_1}}. \end{aligned}$$

Now $\theta(\lambda) \subseteq \theta(\rho)$ (3.1(b.3)), so $\lambda_i \leq \rho_i$ and $\eta_i = \min(\lambda_i, \rho_i) = \lambda_i$ ($i \geq 2$); hence $\theta(\eta) = \theta(\lambda)$. Also, by 3.1(b.2) $\lambda_1 \geq \rho_1$ and therefore $\eta_1 = \rho_1$. Thus $\rho/\eta = \theta(\rho)/\theta(\lambda)$ so $\chi_{\rho/\eta} = \chi_{\theta(\rho)/\theta(\lambda)}$. Also, $\lambda/\eta = (\lambda_1 - \rho_1)$ so $\chi_{\lambda/\eta} = \chi_{(\lambda_1 - \rho_1)}$. It follows that

$$\begin{aligned} \langle \chi_{\lambda/\eta} \otimes \chi_{\rho/\eta}, \chi_{\theta(\mu)} \rangle_{S_{n-\mu_1}} &= \langle \chi_{(\lambda_1 - \rho_1)} \otimes \chi_{\theta(\rho)/\theta(\lambda)}, \chi_{\theta(\mu)} \rangle_{S_{n-\mu_1}} \\ &= \langle \chi_{\theta(\rho)/\theta(\lambda)}, \chi_{\theta(\mu)} \rangle_{S_{n-\mu_1}} \\ &\quad (\chi_{(\lambda_1 - \rho_1)} \text{ is the identity character}) \\ &= \langle \chi_{\theta(\lambda)} \hat{\otimes} \chi_{\theta(\mu)}, \chi_{\theta(\rho)} \rangle_{S_{n-\rho_1}} \\ &\quad (\text{by definition of skew characters; see Sect. 1}). \end{aligned}$$

Case 2. $\mu_1 \neq |\lambda \cap \rho|$.

We shall show that

$$\langle \chi_\lambda \otimes \chi_\mu, \chi_\rho \rangle_{S_n} = \langle \chi_{\theta(\lambda)} \hat{\otimes} \chi_{\theta(\mu)}, \chi_{\theta(\rho)} \rangle_{S_{\theta(\rho)}} = 0.$$

By 3.2

$$\langle \chi_\lambda \otimes \chi_\mu, \chi_\rho \rangle_{S_n} = 0.$$

By 3.1(b.3)

$$\theta(\lambda) \not\supseteq \theta(\rho);$$

hence, by 1.2(b),

$$\langle \chi_{\theta(\lambda)} \hat{\otimes} \chi_{\theta(\mu)}, \chi_{\theta(\rho)} \rangle_{S_{\theta(\rho)}} = 0. \quad \blacksquare$$

EXAMPLE.

$$\begin{aligned} &\langle \chi_{(7,3,2)} \otimes \chi_{(9,2,1)}, \chi_{(4,4,2,1^2)} \rangle_{S_{12}} \quad (\text{by Theorem 3.3}) \\ &= \langle \chi_{(3,2)} \hat{\otimes} \chi_{(2,1)}, \chi_{(4,2,1^2)} \rangle_{S_8} \\ &= 2 \quad (\text{by [Jam-Ker, p. 439]}). \end{aligned}$$

4. APPLICATIONS: REMOVING HIGHEST RECTANGLES IN SOME CHARACTERS

DEFINITION 4.1. (a) Let $H(k, l; n) = \{\lambda \vdash n \mid \text{for all } i > k, \lambda_i \leq l\}$.

(b) Let $m \leq n$. Then denote

$$\psi(k, l; n, m) = \left(\sum_{\lambda \in H(k, l; n)} \chi_\lambda \otimes \chi_\lambda \right)_{S_m}.$$

In this paper we only study the case $l=0$. When $l=0$ and $m=n, n-1$, these are the cocharacters describing the various "trace identities" of the $k \times k$ matrices. These are very close to the cocharacters of the polynomial identities (P.I.'s) of the $k \times k$ matrices. For that reason, the analysis of these characters is of special interest in the theory of P.I. algebras. When $k=2$, $\psi(2, 0; n, n-1)$ was calculated by Procesi (see [Reg 3]). No further results are known for $k \geq 3$. We calculate below (4.5) a rather intriguing formula for the multiplicities of highest rectangles, for any k, n , and m .

By 1.7 we can write:

DEFINITION 4.2. Let $n \geq m$. The equation

$$\psi(k, 0; n, m) = \left(\sum_{\lambda \in H(k, 0; n)} \chi_\lambda \otimes \chi_\lambda \right)_{S_m} = \sum_{\rho \in H(k^2, 0; m)} a(\rho, k, n) \chi_\rho$$

defines the coefficients $a(\rho, k, n)$. Thus $a(\rho, k, n) = \langle \psi(k, 0; n, |\rho|), \chi_\rho \rangle_{S_{|\rho|}}$. We shall prove

PROPOSITION 4.3. Let $0 \leq m \leq n$ and $\rho \vdash m$. Let $t \geq 0$ and assume ρ contains a $k^2 \times t$ rectangle: $\rho = (\hat{\rho} + t, \dots, \hat{\rho}_{k^2} + t)$, $\hat{\rho}_{k^2} \geq 0$. Finally let $a(\rho, k, n)$, $a(\hat{\rho}, k, n - k^2 t)$ be given by 4.2. Then $a(\rho, k, n) = a(\hat{\rho}, k, n - k^2 t)$.

The proof of 4.3 will follow from the following

LEMMA 4.4. Let $\lambda \in H(k, 0; n)$, $\mu \in H(l, 0; n)$. By 1.7, all components of $\chi_\lambda \otimes \chi_\mu$ are in $H(kl, 0; n)$. Let $\eta \in H(kl, 0; n)$ and let $0 \leq t \leq \eta_{kl}$.

(a) Assume $\lambda_k \geq lt$ and $\mu_l \geq kt$. Write $\hat{\lambda} = (\hat{\lambda}_1 + lt, \dots, \hat{\lambda}_k + lt)$, $\hat{\mu} = (\hat{\mu}_1 + kt, \dots, \hat{\mu}_l + kt)$, and $\eta = (\hat{\eta}_1 + t, \dots, \hat{\eta}_{kl} + t)$. Then

$$\langle \chi_\lambda \otimes \chi_\mu, \chi_\eta \rangle_{S_n} = \langle \chi_{\hat{\lambda}} \otimes \chi_{\hat{\mu}}, \chi_{\hat{\eta}} \rangle_{S_{n-kl}}.$$

(b) Let $t \geq 1$. If either $\lambda_k < lt$ or $\mu_l < kt$, then

$$\langle \chi_\lambda \otimes \chi_\mu, \chi_\eta \rangle_{S_n} = 0.$$

Proof. (a) Induction on $t \geq 0$. Trivial for $t=0$. Assume true for $t-1 \geq 0$, and apply 2.4': notice that

$$\lambda \cap \mu' = (\underbrace{l, l, \dots, l}_k)$$

hence $\lambda/\lambda \cap \mu' = (\lambda_1 - l, \lambda_2 - l, \dots, \lambda_k - l)$. Similarly, $\mu/\lambda' \cap \mu = (\mu_1 - k,$

$\mu_2 - k, \dots, \mu_l - k$). In the notation of 2.4', $\zeta = \zeta(\eta) = (\eta_1 - 1, \eta_2 - 1, \dots, \eta_{kl} - 1)$. Thus, by 2.4'

$$\begin{aligned} & \langle \chi_\lambda \otimes \chi_\mu, \chi_\eta \rangle_{S_n} \\ &= \langle \chi_{(\lambda_1 - l, \lambda_2 - l, \dots)} \otimes \chi_{(\mu_1 - k, \mu_2 - k, \dots)}, \chi_{(\eta_1 - 1, \eta_2 - 1, \dots, \eta_{kl} - 1)} \rangle_{S_{n - kl}}. \end{aligned}$$

By induction the proof is now complete.

(b) Here $|\lambda \cap \mu'| < kl$ but $h(\eta) = kl$; hence by 1.6(b), $\langle \chi_\lambda \otimes \chi_\mu, \chi_\eta \rangle_{S_n} = 0$. ■

Proof of 4.3. Apply 4.4 with $k = l$. Let

$$\alpha = (\underbrace{kt, kt, \dots, kt}_k) \vdash k^2 t \quad \text{and} \quad \beta = (\underbrace{t, t, \dots, t}_{k^2}) \vdash k^2 t.$$

Thus

$$\begin{aligned} a(\rho, k, n) &= \langle \psi(k, 0; n, m), \chi_\rho \rangle_{S_m} \quad (\text{by Frobenius Reciprocity}) \\ &= \left\langle \sum_{\lambda \in H(k, 0; n)} \chi_\lambda \otimes \chi_\lambda, (\chi_\rho)^{S_n} \right\rangle_{S_n} \quad (\text{by 1.4}) \\ &= \left\langle \sum_{\lambda \in H(k, 0; n)} \chi_\lambda \otimes \chi_\lambda, \sum_{\substack{\eta \supseteq n \\ \eta \supseteq \rho}} k(\eta, \rho) \chi_\eta \right\rangle_{S_n} \quad (\text{by 1.7}) \\ &= \sum_{\lambda \in H(k, 0; n)} \sum_{\substack{\eta \in H(k^2, 0; n) \\ \eta \supseteq \rho}} k(\eta, \rho) \langle \chi_\lambda \otimes \chi_\lambda, \chi_\eta \rangle_{S_n} \quad (\text{by 4.4 with } l = k) \\ &= \sum_{\substack{\lambda \in H(k, 0; n) \\ \lambda \supseteq \alpha}} \sum_{\substack{\eta \in H(k^2, 0; n) \\ \eta \supseteq \beta}} k(\eta, \rho) \langle \chi_{\lambda/\alpha} \otimes \chi_{\lambda/\alpha}, \chi_{\eta/\beta} \rangle_{S_n}. \end{aligned}$$

Obviously, $\lambda \leftrightarrow \lambda/\alpha = \hat{\lambda}$ is a bijection between $\{\lambda \in H(k, 0; n) \mid \alpha \subseteq \lambda\}$ and $H(k, 0; n - k^2 t)$. Similarly, $\eta \leftrightarrow \eta/\beta = \hat{\eta}$ is a bijection between $\{\eta \in H(k^2, 0; n) \mid \rho \subseteq \eta\}$ and $\{\hat{\eta} \in H(k^2, 0, n - k^2 t) \mid (\rho/\beta) \subseteq \hat{\eta}\}$. Finally $k(\eta, \rho)$, which is the degree of $\chi_{\eta/\rho}$, depends only on the shape η/ρ ; hence $k(\eta, \rho) = k(\hat{\eta}, \hat{\rho})$. Thus

$$\begin{aligned} & a(\rho, k, n) \\ &= \sum_{\hat{\lambda} \in H(k, 0; n - k^2 t)} \sum_{\substack{\hat{\eta} \in H(k^2, 0; n - k^2 t) \\ \hat{\eta} \supseteq \hat{\rho}}} k(\hat{\eta}, \hat{\rho}) \langle \chi_{\hat{\lambda}} \otimes \chi_{\hat{\lambda}}, \chi_{\hat{\eta}} \rangle_{S_{n - k^2 t}} \\ & \quad (\text{by linearity and 1.6}) \\ &= \left\langle \sum_{\hat{\lambda} \in H(k, 0; n - k^2 t)} \chi_{\hat{\lambda}} \otimes \chi_{\hat{\lambda}}, \sum_{\substack{\hat{\eta} \vdash n - k^2 t \\ \hat{\eta} \supseteq \hat{\rho}}} k(\hat{\eta}, \hat{\rho}) \chi_{\hat{\eta}} \right\rangle_{S_{n - k^2 t}} \quad (\text{by 1.4}) \end{aligned}$$

$$\begin{aligned}
 &= \left\langle \sum_{\hat{\lambda} \in H(k, 0; n - k^2 t)} \chi_{\hat{\lambda}} \otimes \chi_{\hat{\lambda}}, (\chi_{\hat{\rho}})^{S_{n - k^2 t}} \right\rangle_{S_{n - k^2 t}} \quad (\text{Frobenius Reciprocity}) \\
 &= \langle \psi(k, 0; n - k^2 t, m - k^2 t, m - k^2 t), \chi_{\hat{\rho}} \rangle_{S_{m - k^2 t}} = a(\hat{\rho}, k, n - k^2 t). \quad \blacksquare
 \end{aligned}$$

COROLLARY 4.5. Let $k \geq 1$, $t \geq 0$, $i \in \{0, 1\}$, and

$$\phi = (t + i, \underbrace{t, \dots, t}_{k^2 - 1}).$$

Let $a(\phi, k, n)$ be given by 4.2. Then

(a) $a(\phi, k, n) = \sum_{\hat{\lambda} \in H(k, 0; n - k^2 t)} (\deg(\chi_{\hat{\lambda}}))^2.$

(b) The case $m = |\phi| = n$ and $m = |\phi| = n - 1$ is of special importance (see 4.1). In that case

$$a(\phi, k, n) = \begin{cases} 2 & \text{if } k \geq 2, i = 1, \text{ and } |\phi| = n - 1; \\ 1 & \text{otherwise.} \end{cases}$$

Proof. (a) By 4.3, $a(\phi, k, n) = a((i), k, n - k^2 t)$. Thus

$$a(\phi, k, n) = a((i), k, n - k^2 t) = \langle \psi(k, 0; n - k^2 t, i), \chi_{(i)} \rangle_{S_i}$$

(since the trivial character is the only irreducible one of S_i , $i = 0, 1$)

$$= \deg(\psi(k, 0; n - k^2 t, i))$$

$$= \deg \left(\left(\sum_{\hat{\lambda} \in H(k, 0, n - k^2 t)} \chi_{\hat{\lambda}} \otimes \chi_{\hat{\lambda}} \right)_{S_i} \right) = \deg \left(\sum_{\hat{\lambda} \in H(k, 0, n - k^2 t)} \chi_{\hat{\lambda}} \otimes \chi_{\hat{\lambda}} \right)$$

$$= \sum_{\hat{\lambda} \in H(k, 0; n - k^2 t)} (\deg(\chi_{\hat{\lambda}}))^2.$$

(b) Easy to check from (a). \blacksquare

Remark 4.6. Let $k \geq 1$, $n \geq 0$, and denote $S_k^{(\beta)}(n) = \sum_{\hat{\lambda} \in H(k, 0, n)} (\deg(\chi_{\hat{\lambda}}))^\beta.$

(a) By 4.1, $S_k^{(2)}(n) = \deg(\psi(k, 0; n, m))$, and by 4.5, $a(\phi, k, n) = S_k^{(2)}(n - k^2 t).$

(b) Explicit values of $S_k^{(\beta)}(n)$ are known only in few special cases [Reg 2, Introduction], such as: $S_2^{(2)}(n) = (1/(n+1))\binom{2n}{n}$, and if $k \geq n$ then $S_k^{(2)}(n) = n!$.

(c) The asymptotic behavior of $S_k^{(\beta)}(n)$ is given in [Reg 2]. The algebraicity of the related generating functions is studied in [Bec-Reg]. For example, it is shown there that when $k \geq 3$ is odd, $\sum_{n \geq 0} S_k^{(2)}(n) t^n$ is not algebraic.

ACKNOWLEDGMENT

I wish to thank Professor Amitai Regev for introducing me to the subject of this work and for the patience and helpful guidance he gave me.

REFERENCES

- [Bec-Reg] W. BECKNER AND A. REGEV, Asymptotics and algebraicity of some generating functions, *Adv. Math.* **65** (1987), 1–15.
- [Gar-Rem] A. GARSIA AND J. REMMEL, Shuffles of permutations and the Kronecker product, *Graphs Combin.* **1** (1985), 217–263.
- [Jam-Ker] G. JAMES AND A. KERBER, “The Representation Theory of the Symmetric Group” (G. C. Rota, Ed.), Encyclopedia of Mathematics and Its Application, Vol. 16, Cambridge Univ. Press, Cambridge, 1984.
- [Mac] I. G. MACDONALD, “Symmetric Functions and Hall Polynomials,” Oxford Univ. Press (Clarendon), Oxford, 1979.
- [Reg1] A. REGEV, The Kronecker product of S_n characters and an $A \otimes B$ theorem for Capelli identities, *J. Algebra* **66** (1980), 505–510.
- [Reg2] A. REGEV, Asymptotic values for degrees associated with strips of Young diagrams, *Adv. Math.* **41** (1981), 115–136.
- [Reg3] A. REGEV, A combinatorial proof of an identity of processi of S_n -characters, *J. Linear Multilinear Algebra* **21**, No. 1 (1987), 29–40.