

The tetrahedral Horn problem and asymptotics of $U(n)$ $6j$ symbols

Anton Alekseev¹, Matthias Christandl², and Thomas C. Fraser²

¹Department of Mathematics, University of Geneva, 7-9 rue du Conseil Général, 1211 Geneva 4, Switzerland

²Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, 2100 Denmark

June 13, 2025

Abstract

Horn's problem is concerned with characterizing the eigenvalues (a, b, c) of a Hermitian matrices (A, B, C) satisfying the constraint $A + B = C$ and has deep connections to tensor product invariants, Littlewood-Richardson coefficients, geometric invariant theory and the intersection theory of Schubert varieties. This paper concerns a generalization of Horn's problem we call the *tetrahedral Horn problem* that aims to characterize the eigenvalues (a, b, c, d, e, f) of Hermitian matrices (A, B, C, D, E, F) forming the edges of a tetrahedron in the space of Hermitian matrices and thus satisfying the constraints $A + B = C$, $B + D = F$, $D + C = E$ and $A + F = E$. We derive new inequalities satisfied by the Schur-polynomials of such eigenvalues and, using eigenvalue estimation techniques from quantum information theory, prove their satisfaction up to degree k implies the existence of approximate solutions with error $O(\sqrt{\frac{\log k}{k}})$. Moreover, the existence of such a matrices is connected to the semiclassical asymptotics of the $6j$ -symbols for the unitary group $U(n)$, which are maps between multiplicity spaces which encode the associativity constraint $(V_\alpha \otimes V_\beta) \otimes V_\delta \cong V_\alpha \otimes (V_\beta \otimes V_\delta)$ for unitary irreducible representations. Using our techniques, we prove the decay rate of the norms of such $6j$ -symbols is either inverse-polynomial or exponential depending on whether there exists such tetrahedra of Hermitian matrices.

Contents

1	Introduction	2
2	The tetrahedral Horn problem	7
2.1	Eigenvalue problems for matrix sums	7
2.2	Deviation Measures	9
2.3	Low dimensional solutions	9
2.4	Symmetries of tetrahedral eigenvalues	10
3	Inequalities for tetrahedral eigenvalues	14
3.1	Schur-Weyl duality	15
3.2	The binomial theorem	16
3.3	Inequalities for Horn's original problem	19
3.4	The $6j$ symbol for $U(n)$	19
3.5	Inequalities for the tetrahedral Horn problem	20
3.6	The large n limit & an entropic inequality	23
4	Distance bounds via eigenvalue concentration	24
4.1	Eigenvalue estimation & separation	25
4.2	Results for random matrices	26
4.3	Distance bound for the original Horn problem	27

4.4	Distance for the tetrahedral Horn problem	28
4.5	Asymptotics of the $6j$ symbols	30
A	Review of previous work of eigenvalues of sums	32
A.1	Horn's inequalities	32
A.2	A simple quadratic inequality	33
A.3	Ptolemy & Cayley-Menger inequalities	33
A.4	Strongly subadditive/Submodular inequalities	34
A.5	Seesaw reciprocity & eigenvalues of blocks	35
A.6	The Courtade-Sra determinant inequality	38
A.7	Some conjectures	39

1 Introduction

Horn's problem: The problem of determining the set of eigenvalues (a, b, c) of Hermitian matrices (A, B, C) satisfying $A + B = C$ has been investigated for over a century [Wey12; Fan49; Lid53; Wie55; TF71; TF70] and has come to be known as *Horn's problem* after Horn who, in 1962, conjectured a solution to the problem in the form of an inductively defined, finite set of homogeneous linear inequalities. In 1998, Klyachko provided a solution to Horn's problem [Kly98] which, when combined with a proof of the so-called saturation conjecture, due to Knutson and Tao [KT99], constitutes a proof of Horn's original conjecture [Ful98]. These linear inequalities, now known as the *Horn inequalities* describe the convex polytope of admissible values for (a, b, c) . Recent developments on Horn's problem include the development of the geometric method by Sherman [She15] and Berline, Vergne, and Walter [BVW18] originally pioneered by the geometric proofs due to Belkale [Bel02; Bel05], as well the characterization and computation of the induced distribution over the eigenvalues γ of the sum $C = A + B$ if A and B are sampled uniformly and independently from the respective sets of Hermitian matrices with eigenvalues a and b by Coquereaux, McSwiggen, and Zuber in series of papers [CZ18; CZ18; CMZ19; Zub18]. For a pedagogical and historical introduction to Horn's problem, we recommend the review articles of Bhatia [Bha01], Knutson [Knu00], and Fulton [Ful00] or alternatively [NP06, Appendix D]. See also the article by Daftuar and Hayden for an account of applications to topics in quantum information theory [DH05].

In parallel to Horn's problem, numerous variations and extensions of Horn's problem have been explored by various authors. For example, numerous authors have considered the admissible eigenvalues of unitary matrices and their product [AW97; Bel08; BK16], or the admissible singular values of matrices and their product [Spe05; TT71]. In 2000, it was shown by [Kly00] that *logarithms* of the admissible singular values of non-singular matrices and their product are identical to the admissible eigenvalues of Hermitian matrices and their sum, thus proving a conjecture due to Thompson [Kly00; AMW01].

The tetrahedral Horn problem: In this paper, we consider a generalization of Horn's problem investigated previously only briefly in [CSW18]. While Horn's problem aims to characterize the eigenvalues (a, b, c) of a triple of $n \times n$ Hermitian matrices (A, B, C) which are subjected to the constraint $A + B = C$, the generalization we consider here aims to characterize the possible eigenvalues (a, b, c, d, e, f) of a *sextuple* of $n \times n$ matrices (A, B, C, D, E, F) subject to four constraints:

$$A + B = C, \quad B + D = F, \quad D + C = E, \quad \text{and} \quad A + F = E. \quad \text{\texttt{\{eq:tet_cond\}}} \quad (1)$$

These constraints arise naturally from building a tetrahedral arrangement of $n \times n$ Hermitian matrices inside the vector space of such Hermitian matrices with edges labelled according to Fig. 1. For this reason, we refer to this eigenvalue problem as the *tetrahedral* Horn problem and we say a sextuple of eigenvalues, (a, b, c, d, e, f) , is *tetrahedral* if such matrices exist, and not tetrahedral otherwise. Beyond the superficial connection between the imposed constraints and the resulting shape formed in the vector space of Hermitian matrices (as in Fig. 1), in the special case of 2×2 Hermitian matrices, this connection is much deeper. It is well-known that the space of admissible eigenvalues for a triangular arrangement of 2×2 traceless Hermitian matrices is equivalent to the moduli space of Euclidean *triangles* in $\mathbb{R}^3$ [HK96; FKL10; Knu00]. This correspondence generalizes to the tetrahedral case in the sense that the space of admissible eigenvalues




Figure 1: (Left) Horn's problem subjects three Hermitian matrices, (A, B, C) , to the constraint $A + B = C$ forming a triangle in the vector space of Hermitian matrices. (Right) The *tetrahedral* Horn problem subjects six Hermitian matrices, (A, B, C, D, E, F) , to a quadruple of constraints forming a tetrahedron in the vector space of Hermitian matrices. fig:tetrahedron_labelling_scheme

for a tetrahedral arrangements of 2×2 traceless Hermitian matrices is equivalent to the moduli space of Euclidean *tetrahedra* in $\mathbb{R}^3$. While these correspondences are straightforward to prove and also well-known, this geometric knowledge about the solution space leads to a complete solution to the tetrahedral Horn's problem for 2×2 matrices; specifically, the moduli space of Euclidean tetrahedra in $\mathbb{R}^3$ is described by the conjunction of the triangle inequalities for each of the four faces (these inequalities coincide with the solution to Horn's original problem) together with the positivity of so-called Cayley-Menger determinants which are polynomial inequalities in the side-lengths (a, b, c, d, e, f) [SJS18]. Even for $n = 2$, we remark that there is a strong qualitative difference between the tetrahedral Horn problem and the original Horn problem. In particular, while the Horn's problem is completely characterized by a *finite* number of *linear* inequalities in the eigenvalues (indeed the solution space is an instance of a moment polytope known as *Horn's polytope*), the solution space for the tetrahedral Horn's problem is not convex nor a polytope.

Our contributions: Regarding the tetrahedral Horn problem, we derive a new family of inequalities constraining the set of tetrahedral eigenvalues. These inequalities are expressed in terms of Schur polynomials $s_\lambda(x)$ of the involved eigenvalues which are homogeneous symmetric polynomials indexed by Young diagrams λ . More accurately, we find our inequalities are best articulated in terms of a simple rescaling of the Schur polynomials, $\varphi_\lambda(x) := s_\lambda(x)/H_\lambda$, where H_λ is a positive integer equal to the product of the hook-lengths of all boxes in the Young diagram λ . These inequalities are presented in the following theorem. thm:iff_tet

Theorem 1. *Let (a, b, c, d, e, f) be a sextuple of non-negative eigenvalues of length n satisfying the trace conditions. Then there exists Hermitian matrices (A, B, C, D, E, F) with satisfying $A + B = C$, $B + D = F$, $C + D = E$ and $A + F = E$ with eigenvalues (a, b, c, d, e, f) if and only if for all triples of Young diagrams (α, β, δ) ,*

$$\varphi_\alpha(a)^{\frac{2}{3}} \varphi_\beta(b) \varphi_\delta(d)^{\frac{2}{3}} \leq \sum_{\gamma, \epsilon, \phi} \left\| \begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix} \right\|_\infty \varphi_\phi(f)^{\frac{1}{3}} \varphi_\gamma(c)^{\frac{1}{3}} \varphi_\epsilon(e)^{\frac{1}{3}}. \quad (2) \quad \text{eq:nec_tet}$$

The expression $\begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix}$ appearing here is an operator known as the $6j$ -symbol for the group of unitary $n \times n$ matrices $U(n)$ and its significance will be explained shortly. While Theorem 1 provides a complete characterization of the set of tetrahedral eigenvalues, it does so using an infinite number of inequalities. Nevertheless, we prove the finite subset of such inequalities of fixed degree $k = |\alpha| + |\beta| + |\delta|$ is sufficient to characterize the set of sextuples of eigenvalues, (a, b, c, d, e, f) , which are *approximately* tetrahedral in the sense that their distance, $D\left(\begin{smallmatrix} a & b & c \\ d & e & f \end{smallmatrix}\right)$, to the set of tetrahedral eigenvalues is small. thm:approx_tet

Theorem 2. *If (a, b, c, d, e, f) are non-negative eigenvalues satisfying the trace conditions and also Eq. (2) for all triples (α, β, δ) such that $|\alpha| + |\beta| + |\delta| = k$ for some positive integer k , then*

$$\frac{1}{\text{Tr}[e]} D\left(\begin{smallmatrix} a & b & c \\ d & e & f \end{smallmatrix}\right) \leq O\left(n \sqrt{\frac{\log k}{k}}\right), \quad (3)$$

where $D\left(\begin{smallmatrix} a & b & c \\ d & e & f \end{smallmatrix}\right) \geq 0$ with equality holding if and only if (a, b, c, d, e, f) are tetrahedral.

Note this theorem actually implies the “if” portion of Theorem 1 as one can take $k \rightarrow \infty$. Along the way, we derive an analogous set of inequalities for the original Horn problem along with an accompanying distance bound (see Proposition 53).

Eigenvalue estimation: Our approach to proving the above theorems relies on the concept of *eigenvalue estimation*, a probabilistic technique developed in the context of quantum information theory for estimating the eigenvalues of positive semidefinite matrices which model the state of quantum systems [KW01; Chr08; OW16; HM02; CHW07; Hay17]. These techniques rely heavily on the concept of Schur-Weyl duality which describes the decomposition of the k -th tensor power of the n -dimensional complex vector space $\mathbb{C}^n$, which supports representations of both the symmetric group S_k by permutation of tensor factors and the group of unitary matrices $U(n)$ (or more generally invertible complex matrices $GL(n, \mathbb{C})$), into irreducible invariant subspaces indexed by Young diagrams $\lambda = (\lambda_1, \dots, \lambda_n)$ with k boxes and at most n rows. Given a positive semidefinite matrix $X \geq 0$ with eigenvalues $x = (x_1, \dots, x_n)$, one can analogously decompose the k -th tensor power of X into the same irreducible invariant subspaces indexed by λ and then investigate the aptly named Schur-Weyl distribution, $p_x(\lambda) := \text{Tr}[P_\lambda X^{\otimes k}] / \text{Tr}[X^{\otimes k}]$, which measures the relative proportion of the decomposed pieces of $X^{\otimes k}$ and depends only on the eigenvalues x of X . The concept of eigenvalue estimation relies on the fact that the Schur-Weyl distribution concentrates strongly around those Young diagrams λ which are proportional to eigenvalues x and that this concentration strengthens as the sizes of the Young diagrams grow to infinity.

6j-symbols: Given a complex reductive group G with unitary irreducible representations (irreps) V_λ labelled by highest weights λ along with tensor-product multiplicity spaces, $M_{\alpha, \beta}^\gamma := \text{Hom}_G(V_\gamma, V_\alpha \otimes V_\beta)$, the 6j-symbols are maps between these multiplicity spaces which encode the associativity constraint $(V_\alpha \otimes V_\beta) \otimes V_\gamma \cong V_\alpha \otimes (V_\beta \otimes V_\gamma)$ between a triple of irreps, V_α, V_β , and V_δ of G . Specifically, a 6j-symbol depends on six irreps, $V_\alpha, V_\beta, V_\gamma, V_\delta, V_\epsilon$, and V_ϕ , and is the linear map between multiplicity spaces of the type

$$\left\{ \begin{matrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{matrix} \right\} : M_{\alpha, \beta}^\gamma \otimes M_{\delta, \gamma}^\epsilon \rightarrow M_{\alpha, \phi}^\epsilon \otimes M_{\beta, \delta}^\phi,$$

corresponding to the block-wise components of the canonical isomorphism from $\bigotimes_\gamma M_{\alpha, \beta}^\gamma \otimes M_{\delta, \gamma}^\epsilon$ to $\bigotimes_\phi M_{\alpha, \beta}^\gamma \otimes M_{\delta, \gamma}^\epsilon$. Whenever the group G is multiplicity-free, in the sense that the dimensions of all multiplicity spaces, $m_{\alpha, \beta}^\gamma := \dim M_{\alpha, \beta}^\gamma$, are either zero or one, the corresponding 6j-symbol is merely a scalar quantity.

Wigner’s 6j-symbol for SU(2): The group SU(2) is multiplicity-free and the 6j-symbols for SU(2) are, depending on the normalization and/or phase convention, are also referred to as Wigner 6j-symbols or Wigner 6j-coefficients [Wig93] or the Racah W -coefficients [Rac42]. The 6j-symbols for SU(2) play a foundational role in the quantum theory of angular momentum, modelling the recoupling theory of three angular momenta, and thus they are used frequently in calculations involving the scattering of multiple particles in nuclear, atomic and molecular physics [VMK88; Edm96; Aqu+12]. Beyond the direct application to the theory of angular momenta, the 6j-symbols for SU(2) also constitute a primitive object in the theory of spin networks [Pen71; BS10], representing a minimal non-trivial closed spin network. General spin networks are networks of trivalent vertices and connections labelled by irreps (spins) of SU(2) which provide a discrete/combinatorial and manifestly invariant model with applications to gauge theories on graphs or manifolds [Bae96], invariants of knots and manifolds [Res87; TV92], topological quantum field theory [DJN93], quantum computing [MR05], quantum change-point discrimination [AH11], and various aspects of proposed models for quantum gravity [Bae99; BC98; Bar03; Rov04; Mou84].

Throughout this paper, by the *asymptotics* of 6j-symbols, we are referring to their behaviour in the regime wherein the six highest weights, $\lambda \in \{\alpha, \beta, \gamma, \delta, \epsilon, \phi\}$, are becoming larger and larger in a suitable sense. The asymptotics of the 6j-symbol for SU(2), wherein the irreps of dimension $\ell + 1$ may be identified either by integers $\ell \in \{0, 1, \dots\}$ or half-integers $j = \frac{\ell}{2} \in \{0, \frac{1}{2}, 1, \frac{3}{2}, \dots\}$, were initially investigated by Wigner [Wig60, Chapter 27]. In the context of the quantum theory of angular momenta, the limit of large ℓ , or equivalently large spins j , may also be interpreted as the limit wherein Planck’s constant tends to zero, $\hbar \propto \frac{1}{k} \rightarrow 0$, which suppresses the effects of quantizing angular momentum; therefore, these asymptotic limits are also known as *semiclassical* limits. Given three particles with spins $\frac{a}{2}$, $\frac{b}{2}$ and $\frac{d}{2}$ and total spin $\frac{e}{2}$, the 6j-symbol, $\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\}$, can be interpreted as the quantum mechanical amplitude to observe the first and second

particles to have combined spin $\frac{c}{2}$ given the second and third have combined spin $\frac{f}{2}$ [Wig60; Rob99]. Wigner considered the probability for this event to occur in the asymptotic limit of large spins, i.e., the square of the $6j$ -symbol, and conjectured the estimate:

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\}^2 \approx \frac{1}{3\pi V}$$

where V is the volume of the Euclidean tetrahedron with side-lengths (a, b, c, d, e, f) . Building upon the work of Wigner, Ponzano and Regge conjectured a formula for the amplitude of the $6j$ -symbol itself in the limit of large spins [PR68]. Later, using techniques from geometric quantization theory, Roberts rigorously proved an asymptotic formula which is equivalent to the conjecture of Ponzano and Regge (up to minor differences) [Rob99].

Theorem 3 (Roberts). *If (a, b, c, d, e, f) are integer side-lengths of a Euclidean tetrahedron with volume V and corresponding exterior dihedral angles $(\theta_a, \theta_b, \theta_c, \theta_d, \theta_e, \theta_f)$, then*

$$\left\{ \begin{matrix} ka & kb & kc \\ kd & ke & kf \end{matrix} \right\} \stackrel{k \rightarrow \infty}{\sim} \sqrt{\frac{2}{3\pi V k^3}} \cos \left(\sum_{\ell} (k\ell + 1) \frac{\theta_{\ell}}{2} + \frac{\pi}{4} \right),$$

where the sum is over the six sides ℓ . Otherwise, if there does not exist a Euclidean tetrahedron with side-lengths (a, b, c, d, e, f) , then $\left\{ \begin{matrix} ka & kb & kc \\ kd & ke & kf \end{matrix} \right\}$ decays to zero at an exponential rate as $k \rightarrow \infty$.

It is worth emphasizing that Chen, Ismail, and Simeonov [CIS99] obtained a rigorous proof of the Ponzano and Regge asymptotics (with minor correction due to [Wat99]) slightly before Chen, Ismail, and Simeonov using estimates from the theory of hypergeometric functions. Additional proofs of this asymptotic correspondence between $6j$ -symbols for $SU(2)$ and Euclidean tetrahedra have emerged in recent years including the elementary proof by Gurau which applies Stirling's approximation directly to Racah's single-sum formula [Gur08] as well as group integral approach to evaluate so-called simple spin networks as Feynman graphs [FK00; FL03]. See also the paper by Bartlett and Nzaganya for an accessible presentation of these kinds of calculations [BN23]. Analogous asymptotic formulas have also been obtained for the $6j$ symbol for the quantum group $U_q(\mathfrak{sl}_2)$ for a k -dependent parameter q [TW06]. The asymptotics calculated by Taylor and Woodward encode (i) the existence of a spherical tetrahedron on the unit sphere with prescribed edge-lengths whenever $q = \exp(i\pi/r(k))$ is an $r(k)$ th root of unity or (ii) the existence of a hyperbolic tetrahedron with prescribed edge-lengths when $q = \exp(\pi/r(k))$ is positive real (here $r(k) = k(r - 2) + 2$ for $r > 2$ a fixed integer) [TW06].

More generally, that the asymptotics of representation-theoretic quantities (such as $6j$ -symbols) are controlled by the existence of some critical geometric object is explained by the theory of geometric quantization [Rob02]. Geometric quantization, broadly speaking, associates classical phase spaces modelled by symplectic manifolds to quantum phase spaces modelled by Hilbert spaces [Rob02; GS82; Kir04]. In particular, Kirillov's orbit principle [Kir04] proposes an association between (i) the coadjoint orbits of a compact Lie group G passing through an integral weight λ to (ii) the corresponding unitary irreps of G of highest weight λ . Furthermore, products of coadjoint orbits correspond to tensor products of irreps and moreover symplectic quotients of the resulting products of orbits are identified with invariant vectors in the resulting tensor products of irreps [Rob02; GS82]. For the aforementioned case of $SU(2)$, the coadjoint orbits can be identified the $SO(3)$ rotations of vectors in Euclidean $\mathbb{R}^3$ (i.e., spheres of fixed radius), and thus the asymptotics of representation-theoretic quantities for $SU(2)$ are controlled by the existence of corresponding geometric objects in Euclidean $\mathbb{R}^3$ with prescribed edge-lengths [Rob02].

$6j$ -symbols for $U(n)$: One application of the techniques developed in this paper is to provide estimates for the asymptotics of the $6j$ symbols for the group $U(n)$ (or equivalently for $SU(n)$) wherein the polynomial irreps V_{λ}^n are labelled by highest-weights which can also be identified with Young diagrams λ . One of the primary difficulties in proving statements about $6j$ symbols for $U(n)$ for $n \geq 3$ is the presence of non-trivial multiplicity spaces, meaning the dimensions of tensor-product multiplicity spaces for $U(n)$, also known as Littlewood-Richardson coefficients, can be greater than one. Because of these difficulties, over the years, considerable attention has been placed on the study of $U(n)$ (or $U_q(\mathfrak{sl}_n)$) $6j$ symbols which are multiplicity-free in the sense that the dimensions of each of the multiplicity spaces is either one or zero

(in the latter case the $6j$ symbol defined to be zero). For instance, tensor products involving symmetric representations, where $\lambda = (k)$, or conjugates of symmetric representations, $\lambda = (k^{(n-1)})^1$, are necessarily multiplicity free; Nawata, Pichai, and Zodinmawia conjectured closed form expressions for the $U_q(\mathfrak{sl}_n)$ $6j$ symbols when a, b, d and e are integers and either $(\alpha, \beta, \delta, \epsilon) = ((a), (b^{(n-1)}), (d), (e^{(n-1)}))$ and $a + d = b + e$ (Type I) or $(\alpha, \beta, \delta, \epsilon) = ((a), (b), (d^{(n-1)}), (e^{(n-1)}))$ and $a + b = d + e$ (Type II) as well as the cases which can be related to these by known symmetries of the corresponding $6j$ symbol [NPZ13]. Building upon this work, Alekseev, Morozov, and Sleptsov studied both the asymptotics of and symmetries in each of the Type I and Type II cases [AMS20]. More recently, Alcock-Zeilinger et al. have derived formulae for the family of $U(n)$ $6j$ symbols where at least one pair of non-adjacent edges, say (α, δ) correspond to a pair of fundamental/defining representations, i.e., $\alpha, \delta = (1)$ [Alc+23]. Shortly thereafter, Keppeler, Plätzer, and Sjødahl expanded upon these results to derive formulae for the family of $U(n)$ $6j$ symbols where either (i) at least one pair of non-adjacent edges consists of fundamental or adjoint representations, i.e., $\alpha, \delta \in \{(1), (2, 1^{(n-2)})\}$, or (ii) at least one vertex, say (α, β, ϕ) , consists entirely of fundamental or adjoint representations, i.e., $\alpha, \beta, \phi \in \{(1), (2, 1^{(n-2)})\}$ [KPS24]. As stated by Coquereaux, “apart from very few special cases (low dimension of irreps and no multiplicity), ... we are not aware of any published general result.” [Coq23].

Our approach bypasses the difficulties presented by multiplicities by focusing only on the *norms* of the $6j$ symbol for $U(n)$. Following the procedure provided by geometric quantization, one expects the asymptotics of the $6j$ -symbol for $U(n)$ to be controlled by existence of some kind of tetrahedral object in space of coadjoint orbits for $U(n)$. In particular, the coadjoint orbits of $U(n)$ can be identified with the orbits of $n \times n$ Hermitian matrices, X , under the conjugation action, $X \mapsto U^{-1}XU$, by unitary $n \times n$ matrices $U \in U(n)$; in other words, coadjoint orbits are associated to sets of $n \times n$ Hermitian matrices sharing a common set of eigenvalues $x = (x_1 \geq \dots \geq x_n)$. Using our methods we are able to prove the following inverse polynomial lower bound.

cor: asym_6j

Corollary 4. *If (a, b, c, d, e, f) are the eigenvalues of $n \times n$ of positive semidefinite matrices (A, B, C, D, E, F) satisfying $A + B = C$, $B + D = F$, $C + D = E$, and $A + F = E$, then there exists a sequence of Young diagrams $(\alpha_k, \beta_k, \gamma_k, \delta_k, \epsilon_k, \phi_k)$ converging in proportion to the eigenvalues (a, b, c, d, e, f) such that*

$$\left\| \begin{Bmatrix} \alpha_k & \beta_k & \gamma_k \\ \delta_k & \epsilon_k & \phi_k \end{Bmatrix} \right\| \geq \frac{1}{\text{poly}(k)}. \quad (4)$$

Otherwise, if (a, b, c, d, e, f) are not the eigenvalues of any such matrices, then for any sequence of Young diagrams converging in proportion to (a, b, c, d, e, f) , we have exponential decay in the value of $\left\| \begin{Bmatrix} \alpha_k & \beta_k & \gamma_k \\ \delta_k & \epsilon_k & \phi_k \end{Bmatrix} \right\|$.

Before continuing, it is worth emphasizing the similarities and differences between our result and the asymptotic formula proven by Roberts above [Rob99]. First and foremost, our result goes beyond that of Roberts in the sense that it applies to the higher-rank Lie group $U(n)$ for $n \geq 3$. Moreover, our result, even when restricted to the case for $SU(2)$, goes beyond the statement proven by Roberts in the sense our result is applicable to *irrational* eigenvalues/edge-lengths, whereas the former is only applicable to *integer*, or with a minor re-scaling, *rational* eigenvalues/edge-lengths. Nevertheless, one drawback of our result is that it merely promises the *existence* of some sequence of Young diagrams, $(\alpha_k, \beta_k, \gamma_k, \delta_k, \epsilon_k, \phi_k)$, converging to the eigenvalues (a, b, c, d, e, f) , whereas the result of Roberts provides a concrete statement about the particular sequence of *scaled* Young diagrams $(k\alpha, k\beta, k\gamma, k\delta, k\epsilon, k\phi)$.

Another drawback of our result is that it merely provides a polynomial lower bound on the value of the $\left\| \begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix} \right\|$ and does not provide an asymptotic formula which would serve as the proper generalization of the Ponzano and Regge formula above [PR68]. The Ponzano and Regge formula for $SU(2)$ was traditionally obtained using the method of *stationary phases* which relies on both an integral formula for the $6j$ -symbol with exponential integrand along with a geometric understanding of the *isolated* critical points for the exponential phase [PR68; Gur08]. In the case of $SU(2)$ with generic edge-lengths, there is a formulation involving exactly *two* isolated critical points corresponding the $SO(3)$ -orbit $[\tau]$ of a Euclidean tetrahedron τ with given edge-lengths and (ii) the $SO(3)$ -orbit $[\tau']$ of the Euclidean tetrahedron τ' which is the mirror-image of the tetrahedron τ with the same fixed side lengths. Generalizing this type of criticality analysis to $U(n)$ for $n \geq 3$ appears to be a significant challenge, and even if an analysis was performed, one would still

¹Here $k^{(n-1)}$ is shorthand for a string $k, k, \dots, k$ consisting of $n - 1$ repetitions of k .

need to compute a corresponding Hessian for each critical point and evaluate the total contribution [Rob02]. Perhaps another approach, such as the approach adopted by Taylor and Woodward (see also [CIS99]) to prove asymptotics for the $6j$ -symbol for $U_q(\mathfrak{sl}_2)$ using recursion relations [TW06] (see also the formulas for the $6j$ -symbols for $\mathfrak{gl}_n$ in terms of generalized hypergeometric functions due to [Art24] [Art23; Art24]).

Previous work: TC: Add a summary of all the strong subadditivity stuff. More recently, Alekseev et al. have investigated a few variants of the tetrahedral Horn problem concerning either the singular values of products of invertible matrices or the tropical singular values of concatenations of weighted planar networks [Ale+25]. These variants are termed *multiple* Horn problems in [Ale+25]; the tetrahedral Horn problem we consider here is termed the *additive* multiple Horn problem in the language of [Ale+25].

Quantum marginals & the $6j$ -symbol for the symmetric group S_k : The techniques used in this paper to derive bounds on the $6j$ symbol for the unitary group $U(n)$ are closely related to the techniques used in an earlier paper by Christandl, Şahinoğlu, and Walter which derived analogous bounds on the $6j$ symbol for the symmetric group S_k [CŞW18]. In that paper, the authors considered the problem of determining whether there exists a density matrix ρ_{ABC} with partial traces $\rho_{AB}, \rho_{BC}, \rho_A, \rho_B$, and ρ_C with a given sextuple of eigenvalues [CŞW18]. Such problems arise naturally in the context of quantum information theory and are typically referred to as quantum marginals problems [CŞW18; Fra22]. In particular, they proved the $6j$ symbols for the symmetric group S_k (referred to as the recoupling coefficients) in the limit as $k \rightarrow \infty$ either decay exponentially or can be bounded from below by a polynomial factor depending on whether there exists a density matrix with the prescribed marginal spectra [CŞW18]. Moreover, they also demonstrated how the tetrahedral Horn problem may be viewed as a special case of the quantum marginals problem and, in a complementary manner, furthermore demonstrated how, by restricting to the subgroup $S_k \subset U(k)$ of permutation matrices and applying Nishiyama’s restriction theorem [Nis00], the $6j$ symbols for unitary group $U(n)$ may be viewed as a special case of the $6j$ symbols for the symmetric group S_k [CŞW18]. This latter result obtained by [CŞW18] is a generalizes how the Littlewood-Richardson coefficients can be cast as a special case of the Kronecker coefficients [Lit58; Mur55; IP23]. The approach taken by this paper differs from that of [CŞW18] in that we prove the correspondence between solutions to the tetrahedral Horn problem on one hand and the asymptotics of the $6j$ symbol for $U(n)$ on the other hand *directly*, rather than first pass through the quantum marginals problem or through the $6j$ symbols for the symmetric group. We note that this more direct approach was initiated, in the $n = 2$ case, by Backens [Bac10].

Organization of the paper: In Section 2 we formally define the tetrahedral Horn problem, describe some of its symmetries, and describe its solution for 2×2 matrices. Then, in Section 3, we develop the mathematical machinery needed to prove the *necessity* of the inequalities in Theorem 1, specifically the binomial theorem for representations (Theorem 26). Afterwards, Section 4 is dedicated to proving the distance bound in Theorem 2 and thus the *sufficiency* of the inequalities in Theorem 1. The results in Section 4 rely on the results of Section 3, the aforementioned eigenvalue concentration bounds (Section 4.1), and some standard results from random matrix theory (Section 4.2). TC: For the sake of completeness, we provide a brief review of some previous works concerning the eigenvalues of sums of Hermitian matrices in Appendix A.

2 The tetrahedral Horn problem

sec:tet_horn_prob

2.1 Eigenvalue problems for matrix sums

Throughout this paper, we consider the relationship between $n \times n$ Hermitian matrices and their real-valued tuples of eigenvalues. To denote the set of $n \times n$ Hermitian matrices, we use

$$\mathcal{H}(n) := \{X \in M_n(\mathbb{C}) \mid X^\dagger = X\}, \quad (5)$$

and to denote the set of possible eigenvalues of $n \times n$ matrices, we use

$$\text{Eig}(n) := \{x = (x_1, \dots, x_n) \in \mathbb{R}^n \mid x_1 \geq \dots \geq x_n\}. \quad (6)$$

Let $\mathcal{H}_+(n)$ be the subset of positive semidefinite $n \times n$ Hermitian matrices and $\text{Eig}^+(n)$ the corresponding set of non-negative eigenvalues. Given a Hermitian matrix $X \in \mathcal{H}(n)$, $\text{eig}(X)$ denotes the eigenvalues of X . The set of Hermitian matrices with eigenvalues $x \in \text{Eig}(n)$ is denoted by

$$\mathcal{O}_x := \{X \in \mathcal{H}(n) \mid \text{eig}(X) = x\}, \quad (7)$$

and $[x]$ denotes the diagonal matrix with entries $(x_1, \dots, x_n)$ along the diagonal:

$$[x] := \text{diag}(x_1, \dots, x_n). \quad (8)$$

The set of Hermitian matrices with eigenvalues x can then be identified with the unitary orbit of the diagonal matrix $[x]$, i.e., $\mathcal{O}_x = \{U[x]U^{-1} \mid U \in \text{U}(n)\}$. Moreover, for any function of Hermitian matrices, f , which unitarily invariant in the sense that $f(X) = f(UXU^{-1})$ holds for all $X \in \mathcal{H}(n)$ and $U \in \text{U}(n)$, we let $f[x]$ be shorthand for $f([x])$. For example,

$$\text{Tr}[x] := \text{Tr}([x]), \quad \text{and} \quad \text{rank}[x] := \text{rank}([x]). \quad (9)$$

`prob:tri_problem`

Definition 5 (Trigonal Horn problem). A triple $(a, b, c) \in \text{Eig}(n)^3$ of candidate eigenvalues are said to be *trigonal* if there exists a corresponding triple $(A, B, C) \in \mathcal{H}(n)^3$ of $n \times n$ Hermitian matrices satisfying (i) the linear constraint

$$A + B = C, \quad (10)$$

`{eq:tri_linear_constraints}`

and (ii) the three eigenvalue constraints

$$\text{eig}(A) = a, \quad \text{eig}(B) = b, \quad \text{eig}(C) = c. \quad (11)$$

`{eq:tri_eigenvalue_constraints}`

The problem of determining the set of trigonal eigenvalues (a, b, c) is known as *Horn's problem*.

`prob:tet_problem`

Definition 6 (Tetrahedral Horn problem). A sextuple $(a, b, c, d, e, f) \in \text{Eig}(n)^6$ of candidate eigenvalues are said to be *tetrahedral* if there exists a corresponding sextuple $(A, B, C, D, E, F) \in \mathcal{H}(n)^6$ of $n \times n$ Hermitian matrices satisfying (i) the four linear constraints

$$A + B = C, \quad B + D = F, \quad C + D = E, \quad A + F = E, \quad (12)$$

`{eq:tet_linear_constraints}`

and (ii) the six eigenvalue constraints

$$\text{eig}(A) = a, \quad \text{eig}(B) = b, \quad \text{eig}(C) = c, \quad \text{eig}(D) = d, \quad \text{eig}(E) = e, \quad \text{eig}(F) = f. \quad (13)$$

`{eq:tet_eigenvalue_constraints}`

The problem of determining the set of tetrahedral eigenvalues (a, b, c, d, e, f) will be referred to as the *tetrahedral Horn problem*.

The tetrahedral Horn problem can be viewed as a generalization of the Horn's problem in a variety of ways. For instance, a triplet (a, b, c) of eigenvalues is trigonal if and only if the sextuple $(a, b, c, 0, c, b)$ is tetrahedral. More generally, it is possible to view the tetrahedral Horn problem as the conjunction of four separate instances of Horn's original problem subject to additional matching constraints. Specifically, a sextuple of eigenvalues (a, b, c, d, e, f) is tetrahedral only if each of the following four triplets are trigonal:

$$(a, b, c), \quad (b, d, f), \quad (c, d, e), \quad (a, f, e). \quad (14)$$

`{eq:four_faces}`

If each of the above triplets are trigonal, then there exists a set of twelve Hermitian matrices with the corresponding eigenvalues satisfying

$$A + B = C, \quad \tilde{B} + D = F, \quad \tilde{C} + \tilde{D} = E, \quad \tilde{A} + \tilde{F} = \tilde{E}. \quad (15)$$

`{eq:four_faces_matrices}`

The tetrahedral Horn problem goes beyond four separate instances of Horn's original problem in that it additionally requires the corresponding pairs of primed and unprimed Hermitian matrices above must coincide.

2.2 Deviation Measures

In order to quantify the degree to which a candidate collection of eigenvalues is compatible, either in the case of Horn's original problem or in the case of the tetrahedral Horn problem, it will be helpful to define some measures of deviation or distance which are non-negative and vanish if and only if the eigenvalues are compatible.

Proposition 7. *Let $1 \leq p \leq \infty$. Then for pairs of Hermitian matrices $X, Y \in \mathcal{H}(n)$ with eigenvalues $\text{eig}(X) = x = (x_1 \geq \dots \geq x_n)$ and $\text{eig}(Y) = y = (y_1 \geq \dots \geq y_n)$, one has*

$$\|X - Y\|_p \geq \|x - y\|_p. \quad (16)$$

Moreover, if $S \subseteq \mathcal{H}(n)$ is a unitarily invariant and closed subset of Hermitian matrices. Then

$$\min_{Y \in S} \|X - Y\|_p = \min_{y \in \text{eig}(S)} \|x - y\|_p, \quad (17)$$

In terms of Horn's original problem, we can consider the set S of Hermitian matrices which are of the form $A + B$ for some $(A, B) \in \mathcal{O}_a \times \mathcal{O}_b$.

Proposition 8. *For a given triple of eigenvalues, (a, b, c) , and $1 \leq p \leq \infty$, let*

$$D_p(a, b, c) := \min_{U, V \in \text{U}(n)} \|\text{eig}(U \cdot [a] + V \cdot [b]) - c\|_p. \quad (18)$$

Then $D_p(a, b, c) \geq 0$ with equality holding if and only if (a, b, c) are trigonal. Moreover,

$$D_p(a, b, c) = \min_{A, B, C \in \mathcal{O}_a \times \mathcal{O}_b \times \mathcal{O}_c} \|A + B - C\|_p. \quad (19)$$

For the tetrahedral Horn problem, there is some additional freedom in how one incorporates the various constraints. The measure of deviation which emerges naturally from our analysis is the following. {defn:tet_deviation}

Definition 9. For a given sextuple of eigenvalues, (a, b, c, d, e, f) , and $1 \leq p \leq \infty$, let

$$D \begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix} := \min_{U, V, W \in \text{U}(n)} \left(\|\text{eig}(U \cdot [a] + V \cdot [b]) - c\|_1^2 + \|\text{eig}(V \cdot [b] + W \cdot [d]) - f\|_1^2 + \|\text{eig}(U \cdot [a] + V \cdot [b] + W \cdot [d]) - e\|_1^2 \right)^{\frac{1}{2}}. \quad (20)$$

Then $D \begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix} \geq 0$ with equality holding if and only if (a, b, c, d, e, f) are trigonal.

2.3 Low dimensional solutions

In the original version of Horn's problem, the eigenvalues (a, b, c) of Hermitian matrices (A, B, C) satisfying $A + B = C$ necessarily must satisfy a basic equality condition known as the *trace condition*: {eq:tri_trace_cond}

$$\text{Tr}[a] + \text{Tr}[b] = \text{Tr}[c], \quad (21)$$

where $\text{Tr}[x] := \sum_i x_i$ is shorthand for the sum of the eigenvalues $x = (x_1 \geq \dots \geq x_n)$. This condition must hold because $A + B = C$ implies $\text{Tr}(C) = \text{Tr}(A + B) = \text{Tr}(A) + \text{Tr}(B)$ and $\text{Tr}(X) = \text{Tr}[x]$ for any Hermitian matrix X with eigenvalues x . For the tetrahedral Horn's problem, a set of tetrahedral eigenvalues (a, b, c, d, e, f) necessarily satisfy four *trace conditions*: {eq:tet_trace_conds}

$$\text{Tr}[a] + \text{Tr}[b] = \text{Tr}[c], \quad \text{Tr}[b] + \text{Tr}[d] = \text{Tr}[f], \quad \text{Tr}[d] + \text{Tr}[c] = \text{Tr}[e], \quad \text{Tr}[a] + \text{Tr}[f] = \text{Tr}[e]. \quad (22)$$

Furthermore, for the trivial case where $n = 1$, the above trace constraints are both necessary and sufficient for the tetrahedral Horn problem.

Example 10 (Tetrahedral Horn's problem for $n = 2$). Here we articulate the solution to the tetrahedral Horn's problem for 2×2 Hermitian matrices. We restrict our focus to the case of *traceless* 2×2 Hermitian matrices². In this context, the sextuple of eigenvalues (a, b, c, d, e, f) can be identified with a sextuple of non-negative real numbers $(a, b, c, d, e, f) \in \mathbb{R}_{\geq 0}^6$ such that the sextuple of traceless eigenvalues are

$$(a, -a), \quad (b, -b), \quad (c, -c), \quad (d, -d), \quad (e, -e), \quad (f, -f). \quad (\text{eq:traceless_two_dim_tetra}) \quad (23)$$

Moreover, each traceless 2×2 Hermitian matrix, X , may be identified with a three-dimensional vector $(x_1, x_2, x_3) \in \mathbb{R}^3$ using the standard Pauli-matrix basis $\{\sigma_1, \sigma_2, \sigma_3\}$ according to the formula:

$$X = x_1\sigma_1 + x_2\sigma_2 + x_3\sigma_3 = \begin{pmatrix} x_3 & x_1 - ix_2 \\ x_1 + ix_2 & -x_3 \end{pmatrix}. \quad (24)$$

Under this identification, the eigenvalues of X are determined by the Euclidean length of the vector (x_1, x_2, x_3) :

$$\text{eig}(X) = \text{eig} \begin{pmatrix} x_3 & x_1 - ix_2 \\ x_1 + ix_2 & -x_3 \end{pmatrix} = (\|x\|_2, -\|x\|_2), \quad \text{where,} \quad \|x\|_2 = \sqrt{x_1^2 + x_2^2 + x_3^2}. \quad (25)$$

Furthermore, if A is determined by the coordinates (a_1, a_2, a_3) and B by the coordinates (b_1, b_2, b_3) , then their sum $C = A + B$ is determined by the coordinates $(c_1, c_2, c_3) = (a_1 + b_1, a_2 + b_2, a_3 + b_3)$. Consequently, in this setting, the tetrahedral Horn's problem reduces to problem of describing the space of all possible edge-lengths (a, b, c, d, e, f) of a Euclidean tetrahedron in $\mathbb{R}^3$. This problem admits of a well-studied solution [Ric02; WD09], namely, the conjunction of a quadruplet of triangle-inequalities

$$|a - b| \leq c \leq a + b, \quad |b - d| \leq f \leq b + d, \quad |d - c| \leq e \leq d + c, \quad |a - f| \leq e \leq a + f, \quad (\text{eq:quadruplet_triangle_constraints}) \quad (26)$$

(which ensure the possibility of forming a Euclidean triangle for each of the four faces), together with a single non-linear inequality of the form

$$\text{CM}(a, b, c, d, e, f) = \frac{1}{288} \det \begin{pmatrix} 0 & a^2 & c^2 & e^2 & 1 \\ a^2 & 0 & b^2 & f^2 & 1 \\ c^2 & b^2 & 0 & d^2 & 1 \\ e^2 & f^2 & d^2 & 0 & 1 \\ 1 & 1 & 1 & 1 & 0 \end{pmatrix} \geq 0. \quad (\text{eq:volume_positive_constraint}) \quad (27)$$

The quantity $\text{CM}(a, b, c, d, e, f)$ is known as a Cayley-Menger determinant and, when $\text{CM}(a, b, c, d, e, f)$ is positive and when the triangle inequalities above are also satisfied, $\text{CM}(a, b, c, d, e, f)$ is always equal to *square* of the volume of the corresponding tetrahedron [SJS18, Equation 5.2]³. Figure 2 depicts a three-dimensional projection of the solution space described by Eq. (26) and Eq. (27).

2.4 Symmetries of tetrahedral eigenvalues

sec:symmetries

There are numerous types of symmetries present in the statement of Definition 6. The purpose of this section is to describe the scaling, inversion, tetrahedral, and shifting symmetries of the problem. These symmetries can, in many cases, be used to simplify the problem by selecting a canonical representative of the corresponding equivalence class; in other cases, it will be useful to leave Definition 6 as stated.

prop:scaling

Proposition 11 (Scaling). *Let (a, b, c, d, e, f) be a sextuple of candidate eigenvalues and let $s \in (0, \infty)$ be a positive real number. Then the scaled eigenvalues, (sa, sb, sc, sd, se, sf) , are tetrahedral if and only if the original, original eigenvalues (a, b, c, d, e, f) are tetrahedral.*

²In Section 2.4, Corollary 16, we will see that this restriction can be made without loss of generality.

³The triangle inequalities $|a - b| \leq c \leq a + b$ are also equivalent to the non-negativity of the *area*, $\text{CM}(a, b, c)$, of the triangle with side-lengths a, b and c where $\text{CM}(a, b, c)^2 = -\frac{1}{16} \det \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & a^2 \\ 1 & a^2 & 0 \end{pmatrix}$ is equivalent to Heron's formula.




Figure 2: A slice of the set of admissible side-lengths (a, b, c, d, e, f) for Euclidean tetrahedron along the coordinates $(a, b, d) = (5, 7, 6)$ (Left) The polyhedron of values (c, e, f) satisfying the triangle inequalities for each face Eq. (26). (Right) The set of all values (c, e, f) satisfying both Eq. (26) and Eq. (27). tetrahedron_multi_space_example

Proof. The statement follows from the fact that $\text{eig}(sX) = s \text{eig}(X)$ holds for all Hermitian matrices X and positive scalars s and that each of the constraints in Eq. (12) are homogeneous in the six Hermitian matrices (A, B, C, D, E, F) ; for example, we have $sA + sB - sC = 0$ if and only if $s(A + B - C) = 0$ if and only if $A + B - C = 0$. $\square$

One may wonder whether it is possible to extend the scaling symmetry to consider scaling by negative $s < 0$. Indeed, this can be accomplished, provided one accounts for the fact that negating a Hermitian matrices both negates and reverses the order of its eigenvalues. prop:inversion

Proposition 12 (Inversion). *Let (a, b, c, d, e, f) be a sextuple of candidate eigenvalues. Given any list of non-increasing eigenvalues $x = (x_1 \geq \dots \geq x_n)$, let x^* denote the inverted list of resorted and negated eigenvalues: $x^* = (-x_n \geq \dots \geq -x_1)$. Then (a, b, c, d, e, f) are tetrahedral if and only if $(a^*, b^*, c^*, d^*, e^*, f^*)$ are tetrahedral.*

Proof. The statement holds because (i) (A, B, C, D, E, F) satisfy Eq. (12) if and only if $(A', B', C', D', E', F') := (-A, -B, -C, -D, -E, -F)$ satisfy Eq. (12), and (ii) $\text{eig}(X) = x$ if and only if $\text{eig}(-X) = x^*$. $\square$

The inversion symmetry Proposition 12 is just one element belonging to a group of 48 symmetries consisting of inversions of eigenvalues, $x \rightarrow x^*$, or permutations of elements in the sextuple (a, b, c, d, e, f) . prop:tet_symmetry

Proposition 13 (Oriented). *The sextuple (a, b, c, d, e, f) of eigenvalues are tetrahedral if and only if any (and thus all) of the following sextuples of eigenvalues are tetrahedral:*

$$(a \ f \ e \ d^* \ c \ b) \quad (c \ e^* \ d^* \ f \ b \ a^*) \quad \text{{eq:gens_of_tet_sym}} \quad (28)$$

Moreover, these two operations generate a finite group of order 48 whose action, grouped by conjugacy classes, produces:

$$\begin{array}{cccc} (a \ b \ c \ d \ e \ f) & (a^* \ c \ b \ d \ f \ e) & (a \ c^* \ b^* \ d^* \ f^* \ e^*) & (a^* \ b^* \ c^* \ d^* \ e^* \ f^*) \\ (d \ b \ f \ a \ e \ c) & (b^* \ a^* \ c^* \ e \ d \ f) & (b \ a \ c \ e^* \ d^* \ f^*) & (d^* \ b^* \ f^* \ a^* \ e^* \ c^*) \\ (d^* \ e \ c \ a^* \ b \ f) & (f^* \ b \ d^* \ c^* \ e^* \ a^*) & (f \ b^* \ d \ c \ e \ a) & (d \ e^* \ c^* \ a \ b^* \ f^*) \\ (a \ e^* \ f^* \ d \ b^* \ c^*) & (e \ d^* \ c \ b^* \ a \ f^*) & (e^* \ d \ c^* \ b \ a^* \ f) & (a^* \ e \ f \ d^* \ b \ c) \\ (b \ f^* \ d^* \ e \ c \ a) & (c \ b^* \ a \ f \ e \ d) & (c^* \ b \ a^* \ f^* \ e^* \ d^*) & (a^* \ f^* \ e^* \ d \ c^* \ b^*) \\ (c \ a^* \ b \ f^* \ d^* \ e^*) & (a \ f \ e \ d^* \ c \ b) & (a^* \ f^* \ e^* \ d \ c^* \ b^*) & (b^* \ f \ d \ e^* \ c^* \ a^*) \\ (e \ c^* \ d \ b \ f \ a^*) & (b \ d \ f \ e^* \ a^* \ c^*) & (b^* \ d^* \ f^* \ e \ a \ c) & (c^* \ a \ b^* \ f \ d \ e) \\ (f \ a \ e \ c^* \ d \ b^*) & (d \ f^* \ b^* \ a^* \ c^* \ e^*) & (d^* \ f \ b \ a \ c \ e) & (e^* \ c \ d^* \ b^* \ f^* \ a) \\ (f^* \ d \ b^* \ c \ a \ e) & (f \ e^* \ a^* \ c \ b \ d^*) & (f^* \ e \ a \ c^* \ b^* \ d) & (f^* \ a^* \ e^* \ c \ d^* \ b) \\ (e^* \ f \ a^* \ b^* \ c^* \ d) & (e \ a \ f^* \ b \ d^* \ c) & (e \ a^* \ f \ b^* \ d \ c^*) & (f \ d^* \ b \ c^* \ a^* \ e^*) \\ (c^* \ d^* \ e^* \ f \ a^* \ b) & (d^* \ c^* \ e^* \ a \ f^* \ b^*) & (d \ c \ e \ a^* \ f \ b) & (e \ f^* \ a \ b \ c \ d^*) \\ (b^* \ c \ a \ e^* \ f^* \ d^*) & (c^* \ e \ d \ f^* \ b^* \ a) & (c \ e^* \ d^* \ f \ b \ a^*) & (c \ d \ e \ f^* \ a \ b^*) \\ & & & (b \ c^* \ a^* \ e \ f \ d) \end{array} \quad \text{{eq:all_of_tet_sym}} \quad (29)$$

Proof. To begin, notice that each of the operations on (a, b, c, d, e, f) appearing in Eq. (28) are invertible (indeed the first operation has order two, while the second operation has order four). The claim that these two operations generate a finite group of order 48 can be checked explicitly using computer software. Now suppose (a, b, c, d, e, f) are the eigenvalues of a sextuple of Hermitian matrices (A, B, C, D, E, F) satisfying Eq. (12). Then, for example, the sextuple $(A', B', C', D', E', F') = (A, F, E, -D, C, B)$ (which has the eigenvalues (a, f, e, d^*, c, b)) is also tetrahedral because

$$\begin{aligned} A' + B' - C' &= (A + F - E) = 0, \\ B' + D' - F' &= -(B + D - F) = 0, \\ C' + D' - E' &= -(C + D - E) = 0, \\ A' + F' - E' &= (A + B - C) = 0. \end{aligned} \tag{30}$$

Moreover, as the operation mapping (a, b, c, d, e, f) to (a^*, c, b, d, f, e) is invertible, one also has that if (a, b, c, d, e, f) is not tetrahedral, then neither is (a^*, c, b, d, f, e) . An analogous argument holds for the operation mapping (a, b, c, d, e, f) to (c, e^*, d^*, f, b, a^*) with (A', B', C', D', E', F') given by $(C, -E, -D, F, B, -A)$. $\square$

Remark 14. The group G of order 48 appearing in Proposition 13 corresponds to the subgroup of all possible signed permutations of 6 elements generated by permutations and negations of the elements of the sextuple (A, B, C, D, E, F) which preserves the set of linear constraints in Eq. (12). The group G is isomorphic to both the direct product $S_2 \times S_4$ and to the signed permutation group of three elements, i.e., $(S_2 \times S_2 \times S_2) \rtimes S_3$.

The first isomorphism, $G \simeq S_2 \times S_4$, can be understood as S_2 acting by the inversion of all eigenvalues,

$$(a, b, c, d, e, f) \leftrightarrow (a^*, b^*, c^*, d^*, e^*, f^*), \tag{31}$$

while S_4 acts by permutation of the *vertices* of the tetrahedron in Fig. 1: each vertex permutation induces a permutation of the six directed edges with the possibility of inverting the orientation of a subset of the edges (such edge orientation reversals can be accounted for a corresponding inversion of eigenvalues).

The second isomorphism, $G \simeq (S_2 \times S_2 \times S_2) \rtimes S_3$, can be understood by considering the triplet $(A, D), (B, E), (C, F)$ of non-adjacent pairs of edges in Fig. 1. Accordingly, it is helpful to arrange the sextuple (a, b, c, d, e, f) into a 2×3 grid using the following identification:

$$(a \ b \ c \ d \ e \ f) \cong \begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix}. \tag{32}$$

Then one can see that each S_2 factor in the direct product $S_2 \times S_2 \times S_2$ acts faithfully on the sextuple by both negating a certain subset of the columns and permuting the upper and lower rows leaving just one column unchanged.

$$\begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix} \leftrightarrow \begin{pmatrix} a & e^* & f^* \\ d & b^* & c^* \end{pmatrix}, \quad \begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix} \leftrightarrow \begin{pmatrix} d & b & e \\ a & c & f \end{pmatrix}, \quad \begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix} \leftrightarrow \begin{pmatrix} d^* & e & c \\ a^* & b & f \end{pmatrix}. \tag{33}$$

The symmetric group S_3 of order $3! = 6$ acts instead by permuting the columns and negating a subset of the entries in the top row, producing a subgroup of order 6 of the form:

$$\begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix} \quad \begin{pmatrix} b & c^* & a^* \\ e & f & d \end{pmatrix} \quad \begin{pmatrix} c^* & a & b^* \\ f & d & e \end{pmatrix} \\ \begin{pmatrix} a^* & c & b \\ d & f & e \end{pmatrix} \quad \begin{pmatrix} b^* & a^* & c^* \\ e & d & f \end{pmatrix} \quad \begin{pmatrix} c & b^* & a \\ f & e & d \end{pmatrix} \tag{34}$$

We further remark that these symmetries also corresponding to symmetries of the Wigner $6j$ operator $\begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix}$ in Definition 31 with the involution $\lambda \mapsto \lambda^*$ corresponding taking the dual representation $V_\lambda^n \mapsto (V_{\lambda^*}^n)^*$. In the case where $n = 2$, the involution map becomes the identity map (as every irreducible representation of $SU(2)$ is self-dual, or the reversal of a pair $(x, -x)$ of traceless eigenvalues is trivial), and thus the symmetry group collapses to a group of order $|S_4| = 4! = 24$.

prop:shifting

Proposition 15 (Shifting symmetry). *Let (a, b, c, d, e, f) be a sextuple of candidate eigenvalues and let $x, y, z \in \mathbb{R}$ be fixed. Define shifted eigenvalues $(\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}, \tilde{e}, \tilde{f})$ by*

$$\begin{aligned} \tilde{a}_i &:= a_i + x, & \tilde{b}_i &:= b_i + y, & \tilde{c}_i &:= c_i + x + y, & \text{\textit{\{eq:shifted_tet_eigenvalues\}}} \\ \tilde{d}_i &:= d_i + z, & \tilde{e}_i &:= e_i + x + y + z, & \tilde{f}_i &:= f_i + y + z. \end{aligned} \quad (35)$$

Then shifted eigenvalues $(\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}, \tilde{e}, \tilde{f})$ are tetrahedral if and only if the original eigenvalues (a, b, c, d, e, f) are tetrahedral.

Proof. Given a Hermitian matrix A with eigenvalues $\text{eig}(A) = a$ and a constant $w \in \mathbb{R}$, the eigenvalues of $A + wI$ (where I is the identity matrix) are

$$\text{eig}(A + wI) = (a_1 + w \geq \dots \geq a_n + w). \quad (36)$$

The statement then follows because the sextuple of matrices $(\tilde{A}, \tilde{B}, \tilde{C}, \tilde{D}, \tilde{E}, \tilde{F})$, where

$$\begin{aligned} \tilde{A} &:= A + xI, & \tilde{B} &:= B + yI, & \tilde{C} &:= C + (x + y)I, \\ \tilde{D} &:= D + zI, & \tilde{E} &:= E + (x + y + z)I, & \tilde{F} &:= F + (y + z)I, \end{aligned} \quad (37)$$

have the shifted eigenvalues $(\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}, \tilde{e}, \tilde{f})$ and satisfy the tetrahedral constraints in Eq. (12) if and only if (A, B, C, D, E, F) satisfy the tetrahedral constraints in Eq. (12). $\square$

As a corollary of the shifting symmetries (Proposition 15) one concludes that to solve the tetrahedral Horn's problem, in a certain sense, it suffices to solve the tetrahedral Horn's problem merely for traceless matrices (Corollary 16) or, alternatively, for positive definite or positive semidefinite matrices (Corollary 17). *cor:traceless*

Corollary 16 (Traceless eigenvalues). *Let (a, b, c, d, e, f) be a sextuple of candidate eigenvalues and for any list of n eigenvalues $x = (x_1 \geq \dots \geq x_n)$, let $x^0 := x - \text{Tr}[x]/n$ denote the traceless list of eigenvalues obtained by shifting x by $-\text{Tr}[x]/n$. Then (a, b, c, d, e, f) are tetrahedral if and only if (i) their traces $(\text{Tr}[a], \text{Tr}[b], \text{Tr}[c], \text{Tr}[d], \text{Tr}[e], \text{Tr}[f])$ are tetrahedral (they satisfy Eq. (22)) and (ii) $(a^0, b^0, c^0, d^0, e^0, f^0)$ are tetrahedral.*

Proof. This “only if” portion of the statement follows from both the tetrahedral trace conditions (Eq. (22)) and applications of the shifting symmetry Proposition 15 with (x, y, z) values

$$x = -n^{-1} \text{Tr}(A), \quad y = -n^{-1} \text{Tr}(B) \quad z = -n^{-1} \text{Tr}(D). \quad (38)$$

The “if” portion of the statement follows from the shifting symmetry applied to the negated (x, y, z) values. $\square$

The next corollary of the shifting symmetry proves that to solve the tetrahedral Horn's problem it suffices to solve the tetrahedral Horn's problem for non-negative (or positive) eigenvalues only. Indeed, one can simply take the shifting parameters (x, y, z) appearing in Proposition 15 to be sufficiently large. The following corollary establishes, in a certain sense, the minimum shifting required to ensure non-negative (or positive) eigenvalues for all six Hermitian matrices. Note that in subsequent sections we will exclusively consider candidate eigenvalues (a, b, c, d, e, f) that are either non-negative. *cor:psd*

Corollary 17 (Non-negativity). *If (a, b, c, d, e, f) are a tetrahedral sextuple of eigenvalues, then necessarily the smallest eigenvalues satisfy*

$$c_n \geq a_n + b_n, \quad f_n \geq b_n + d_n, \quad e_n \geq d_n + c_n, \quad e_n \geq a_n + f_n. \quad (39) \quad \text{\textit{\{eq:min_eig_ineqs\}}}$$

Moreover, if (a, b, c, d, e, f) are a sextuple of candidate eigenvalues satisfying Eq. (39), then for any $(x, y, z) \in \mathbb{R}^3$ satisfying

$$x \geq -a_n, \quad y \geq -b_n, \quad z \geq -d_n, \quad (40) \quad \text{\textit{\{eq:positive_shift\}}}$$

the resulting shifted eigenvalues $(\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}, \tilde{e}, \tilde{f})$ (given by Eq. (35)) are all non-negative.

Proof. That tetrahedral eigenvalues satisfy Eq. (39) follows from four applications of the well-known inequality $c_n \geq a_n + b_n$ for Horn's problem where $C = A + B$ [Bha01]. As a reminder, the smallest eigenvalue x_n of a matrix X is characterized by a minimization: $x_n = \min_{\|v\|=1} \langle v, Xv \rangle$ which produces

$$c_n = \min_{\|v\|=1} \langle v, Cv \rangle = \min_{\|v\|=1} (\langle v, Av \rangle + \langle v, Bv \rangle) \geq \min_{\|v\|=1} \langle v, Av \rangle + \min_{\|w\|=1} \langle w, Bw \rangle = a_n + b_n. \quad (41)$$

Four applications of this inequality produces Eq. (39). Now if (x, y, z) satisfy Eq. (40), we have for all non-negativity of the shifted eigenvalues, e.g.,

$$\forall j : \quad \tilde{a}_j = a_j + x \geq a_n + x \geq 0, \quad (42)$$

Similar statements hold for b, c, d, e and f . $\square$

Remark 18. Although this section listed a handful of symmetries of the tetrahedral Horn's problem, there remains the possibility of additional symmetries which are unknown at this time. For example, in the case 2×2 Hermitian matrices, an additional class of symmetries on the set of edge-lengths (a, b, c, e, d, f) of tetrahedra, known as the *Regge symmetries*, hold [AI19; Boa07; PR68, Theorem 1.1]. If there exists a tetrahedron with edge lengths (a, b, c, d, e, f) , then there also exists a tetrahedron with edge lengths

$$(s_1 - a, s_1 - b, c, s_1 - d, s_1 - e, f) \quad \text{where} \quad s_1 = \frac{a + b + d + e}{2}. \quad (43)$$

Combining this symmetry with the permutational symmetries given by Proposition 13 (and noting that for the eigenvalues of traceless 2×2 Hermitian matrices are invariant under negation) one concludes the co-existence of a family of six tetrahedra with edge lengths given by [VMK88, Chapter 9]:

$$\begin{aligned} (a, b, c, d, e, f) & \quad (s_1 - a, s_1 - b, c, s_1 - d, s_1 - e, f) \\ (a, s_3 - b, s_3 - c, d, s_3 - e, s_3 - f) & \quad (s_2 - a, b, s_2 - c, s_2 - d, e, s_2 - f) \\ (s_2 - d, s_1 - e, s_3 - f, s_2 - a, s_1 - b, s_3 - c) & \quad (s_1 - d, s_3 - e, s_2 - f, s_1 - a, s_3 - b, s_2 - c) \end{aligned} \quad (44)$$

where

$$s_1 = \frac{a + b + d + e}{2}, \quad s_2 = \frac{a + c + d + f}{2}, \quad s_3 = \frac{b + c + e + f}{2}. \quad (45)$$

Remark 19. The Regge symmetry of a tetrahedron can be understood as a consequence of an earlier discovery of a surprising symmetry of Wigner $6j$ symbol for $SU(2)$ originally due to Regge [Reg59]. The paper due to Boalch provides a comprehensive and modern treatment of the Regge symmetry for the $6j$ symbol for $SU(2)$ by viewing it as a consequence of a symmetry of multiplicity-free $6j$ -symbols for $SU(3)$ [Boa07]. That the Regge symmetry of a tetrahedron emerges from the Regge symmetry of the associated $6j$ symbols can be viewed as a consequence of the semiclassical limit of the latter [PR68; Rob99]. It has also been shown that the Regge symmetry of Euclidean tetrahedra is actually a scissors-congruence preserving both the volume and Dehn invariants [AI19]. At the moment, it remains unclear if there exists analogous symmetries for the generalized Wigner $6j$ symbol for $n \geq 3$ (although there are conjectured symmetries for the $6j$ symbols for $U_q(\mathfrak{sl}_n)$ which would follow from the eigenvalue hypothesis [MS18; LS23]). It could also be possible for additional symmetries to be present in the tetrahedral Horn problem for $n \geq 3$ which are *not* present in the $6j$ symbol for $U(n)$.

3 Inequalities for tetrahedral eigenvalues

sec:ineqs_from_schur_weyl

The purpose of this section is to develop the technical machinery used to prove numerous new inequalities constraining the set of tetrahedral eigenvalues. Our approach relies heavily on the theory of irreducible representations of the group $U(n)$ of unitary $n \times n$ matrices, or equivalently, the theory of polynomial irreducible representations of the group $GL(n, \mathbb{C})$ of invertible $n \times n$ matrices with complex entries. More specifically, our approach relies on the decomposition of tensor powers $X^{\otimes k}$ of $n \times n$ matrices provided by Schur-Weyl duality. In Section 3.1 we review this prerequisite representation theory and provide elementary results regarding the decomposition of tensor powers $X^{\otimes k}$ and then afterwards, in Section 4.1, we describe

how these decompositions of $X^{\otimes k}$ encode information about the eigenvalues x of X . Building upon this machinery, in Section 3.2, we state and prove a binomial theorem concerning various decompositions of tensors powers of matrix sums $(X + Y)^{\otimes k}$. Using the binomial theorem, in Section 3.3, we will derive new inequalities constraining the set of eigenvalue triples (α, β, γ) of positive semidefinite matrices (A, B, C) satisfying $A + B = C$ relevant for Horn's original eigenvalue problem. Afterwards, in Section 3.5, we again leverage the binomial theorem (or rather its multinomial generalization) to derive new inequalities constraining the set of eigenvalue sextuples (a, b, c, d, e, f) of positive semidefinite matrices (A, B, C, D, E, F) satisfying $A + B = C$, $B + D = F$, $C + D = E$, and $A + F = E$ relevant for the tetrahedral Horn problem.

3.1 Schur-Weyl duality

[sec:schur_weyl](#)

The purpose of this section is to review the representation theory of $U(n)$, or equivalently, the polynomial representation theory of $GL(n)$. The isomorphism classes of irreducible polynomial representations of $GL(n)$ are indexed by Young diagrams.

A Young diagram λ of size k is a collection of k boxes arranged in rows of sorted length and can be identified with its sequence of non-zero row-lengths, $\lambda \simeq (\lambda_1, \dots, \lambda_\ell)$, where λ_i counts the number of boxes in the i th row, $\lambda_1 \geq \dots \geq \lambda_\ell > 0$, and where $\sum_i \lambda_i = k$. The number of non-zero rows in λ is called the depth or length of λ and is denoted by $\ell(\lambda)$ and the size of λ is denoted by $|\lambda|$. As a shorthand, we write $\lambda \vdash_n k$ to denote a Young diagram λ of size $|\lambda| = k$ with depth at most $\ell(\lambda) \leq n$. For example, $\lambda = \mathbb{F}^{\square} \simeq (4, 2, 1)$ is a Young diagram of size $|\lambda| = 4 + 2 + 1 = 7$ and depth 3.

For each Young diagram ℓ of size k with depth at most n , there is an irreducible polynomial representation of $U(n)$ (or $GL(n, \mathbb{C})$), denoted by $(\pi_\lambda, V_\lambda^n)$, whose matrix coefficients are polynomials of homogeneous degree k in the entries of $U \in U(n)$. The dimension of the V_λ^n is given by Weyl's dimension formula for $U(n)$ [Hal08; IN66],

$$\dim(V_\lambda^n) = \frac{\Delta(\lambda + \rho)}{\Delta(\rho)} = \frac{\prod_{1 \leq i < j \leq n} (\lambda_i - i - \lambda_j + j)}{\prod_{1 \leq i < j \leq n} (j - i)}, \quad \text{\texttt{eq:weyl_dim}} \quad (46)$$

where $\rho \simeq (n-1, n-2, \dots, 1, 0)$ is known as the Weyl vector and where

$$\Delta(x_1, \dots, x_n) = \prod_{1 \leq i < j \leq n} (x_i - x_j). \quad (47)$$

is known as the Vandermonde determinant. The defining representation of $U(n)$, wherein $U \in U(n)$ acts via matrix-vector multiplication on $\mathbb{C}^n$, is associated to the Young diagram with one box, $\lambda = \square \simeq (1)$, and as dimension $\dim(V_\square^n) = n$ as $V_\square^n \simeq \mathbb{C}^n$.

The so-called Schur-Weyl decomposition of $(V_\square^n)^{\otimes k} \simeq (\mathbb{C}^n)^{\otimes k}$ into irreducible representations of $U(n)$ is as follows.

Proposition 20 (Schur-Weyl duality). *Let k and n be positive integers. Then the decomposition of $(\mathbb{C}^n)^{\otimes k}$ into irreducible subrepresentations of $S_k \times U(n)$, known as the Schur-Weyl decomposition, is given by*

$$(\mathbb{C}^n)^{\otimes k} \cong \bigoplus_{\lambda \vdash_n k} W_\lambda \otimes V_\lambda^n \quad \text{\texttt{eq:schur_weyl_decomp}} \quad (48)$$

where the direct summands are indexed by Young diagrams λ of size k and depth at most n rows, and where W_λ supports an irreducible representation of S_k and V_λ^n supports the irreducible representation of $U(n)$.

The dimension W_λ can be computed using the Frobenius dimension formula [GW00, Corollaries 9.1.5],

$$\dim(W_\lambda) = \frac{|\lambda|! \Delta(\lambda + \rho)}{\prod_{i=1}^n (\lambda_i + \rho)_i!}. \quad \text{\texttt{eq:frob_dim}} \quad (49)$$

While this formula is useful for comparisons with $\dim(V_\lambda^n)$, it is not obviously independent of n for $n \geq \ell(\lambda)$. Indeed, the right hand side of Eq. (49) does not depend on n for $n \geq \ell(\lambda)$ and coincides with the Hook-length formula [GW00, Corollary 9.1.6], in this case

$$\dim(W_\lambda) = \frac{|\lambda|!}{H_\lambda}, \quad H_\lambda = \prod_{b \in \lambda} h_\lambda(b), \quad \text{\texttt{eq:hook_dim}} \quad (50)$$

where the hook length, $h_\lambda(b)$, of b counts the number of boxes to the right and below b plus one for itself. We refer to H_λ as the hook-product for λ .

Now let $(\pi_\lambda, V_\lambda^n)$ be an irreducible representation of $U(n)$ labelled by the Young diagram λ with depth at most n . Throughout this paper, we let the $S_k \times U(n)$ equivariant isometry for the λ -isotypic subspace of $V_\square^{\otimes |\lambda|} \simeq (\mathbb{C}^n)^{\otimes |\lambda|}$ be denoted by

$$\iota_\lambda : W_\lambda \otimes V_\lambda^n \rightarrow V_\square^{\otimes |\lambda|}. \quad (51)$$

Then given an $n \times n$ matrix X , viewed as a linear operator on the defining representation $V_\square^n \simeq \mathbb{C}^n$, let $\pi_\lambda(X)$ be defined by

$$\pi_\lambda(X) := \frac{\text{Tr}_{W_\lambda}[\iota_\lambda^\dagger X^{\otimes |\lambda|} \iota_\lambda]}{\dim(W_\lambda)}.$$

Note that when X is invertible, i.e. $X \in \text{GL}(n, \mathbb{C})$, the operator $\pi_\lambda(X)$ as defined above coincides with the representation of X on V_λ^n , so this abuse of notation is permitted. For any positive integer k , the k -th tensor power of X then admits a decomposition as $X^{\otimes k} \simeq \bigoplus_{\lambda \vdash_n k} \pi_\lambda(X)^{\oplus W_\lambda}$, or more precisely,

$$X^{\otimes k} = \sum_{\lambda \vdash_n k} \iota_\lambda(\pi_\lambda(X) \otimes \text{id}_{W_\lambda}) \iota_\lambda^\dagger. \quad (52)$$

As a special case, when $\lambda = \square$, then $\pi_\square(X) = X$. Note also that the matrix coefficients of $\pi_\lambda(X)$ are homogeneous degree $|\lambda|$ polynomials in the matrix coefficients of X meaning for any scalar $s \in \mathbb{C}$,

$$\pi_\lambda(sX) = s^{|\lambda|} \pi_\lambda(X). \quad (53)$$

Additionally, if X, Y are two matrices, we have

$$\pi_\lambda(XY) = \pi_\lambda(X) \pi_\lambda(Y). \quad (54)$$

The corresponding character $\chi_\lambda(X) = \text{Tr}(\pi_\lambda(X))$ of a representation of a Hermitian matrix X with eigenvalues x is equal to

$$\chi_\lambda(X) := \text{Tr}(\pi_\lambda(X)) = \frac{\text{Tr}[P_\lambda X^{\otimes |\lambda|}]}{\dim(W_\lambda)} = s_\lambda(x),$$

where $P_\lambda := \iota_\lambda^\dagger \iota_\lambda$ is the orthogonal projection operator onto the λ -isotypic subspace, and where $s_\lambda(x) = s_\lambda(x_1, \dots, x_n)$ Schur-polynomial associated to λ .

There are various formulas for the Schur-polynomial $s_\lambda(x_1, \dots, x_n)$. Here we present the Jacobi's bialternant formula

$$s_\lambda(x) = \frac{\det[x_i^{\lambda_j + n - j}]}{\det[x_i^{n - j}]} = \frac{1}{\Delta(x)} \det[x_i^{(\lambda + \rho)_j}]. \quad (55)$$

3.2 The binomial theorem

[sec:binomial_theorem](#)

The purpose of this section is to state and prove the binomial theorem, which serves as the foundation for all subsequent results in this paper. The binomial theorem expresses any polynomial irreducible representation of a matrix sum $X + Y$ as a sum over the polynomial irreducible representations of the summands X and Y .

The theorem inherits its name from the non-commutative binomial expansion of the k -th tensor power of the sum $X + Y$,

$$(X + Y)^{\otimes k} = \sum_{\ell=0}^k \binom{k}{\ell} \left[\frac{1}{k!} \sum_{\sigma \in S_k} \sigma(X^{\otimes \ell} \otimes Y^{\otimes (k-\ell)}) \sigma^{-1} \right] \quad (56)$$

where $\sigma \in S_k$ acts on $(\mathbb{C}^n)^{\otimes k}$ by permutation of the k tensor factors. For an example, consider the case where $k = 3$:

$$\begin{aligned} (X + Y)^{\otimes 3} = & \underbrace{X \otimes X \otimes X}_{\ell=3} + \underbrace{X \otimes X \otimes Y + X \otimes Y \otimes X + Y \otimes X \otimes X}_{\ell=2} + \\ & + \underbrace{X \otimes Y \otimes Y + Y \otimes X \otimes Y + Y \otimes Y \otimes X}_{\ell=1} + \underbrace{Y \otimes Y \otimes Y}_{\ell=0}. \end{aligned} \quad (57)$$

As we have seen previously, the k -th tensor power of a matrix can also be decomposed into a sum over polynomial irreducible representations labelled by Young diagrams γ with sizes $|\gamma| = k$. Combining this formula with the non-commutative binomial expansion yields

$$(X + Y)^{\otimes k} = \sum_{\gamma \vdash k} \iota_\gamma(\pi_\gamma(X + Y) \otimes \text{id}_{W_\gamma}) \iota_\gamma^\dagger \quad (58)$$

where

$$\pi_\gamma(X + Y) = \frac{1}{\dim(W_\gamma)} \sum_{\ell=0}^k \binom{k}{\ell} \text{Tr}_{W_\gamma}[\iota_\gamma^\dagger(X^{\otimes \ell} \otimes Y^{\otimes(k-\ell)}) \iota_\gamma], \quad (59)$$

where we have eliminated the symmetrization operation over $\sigma \in S_k$ because the partial trace over W_γ is S_k -invariant. At this stage, for each summand indexed by $0 \leq \ell \leq k$, we can decompose $X^{\otimes \ell}$ into representations of X labelled by α with sizes $|\alpha| = \ell$ and $Y^{\otimes(k-\ell)}$ into representations of Y labelled by β with sizes $|\beta| = k - \ell$.

$$X^{\otimes \ell} = \sum_{\alpha \vdash \ell} \iota_\alpha(\pi_\alpha(X) \otimes \text{id}_{W_\alpha}) \iota_\alpha^\dagger, \quad Y^{\otimes(k-\ell)} = \sum_{\beta \vdash k-\ell} \iota_\beta(\pi_\beta(Y) \otimes \text{id}_{W_\beta}) \iota_\beta^\dagger. \quad (60)$$

Upon doing so, one encounters the composite operator $\iota_\gamma^\dagger(\iota_\alpha \otimes \iota_\beta)$ which maps from $(V_\alpha^n \otimes W_\alpha) \otimes (V_\beta^n \otimes W_\beta)$ to $(V_\gamma^n \otimes W_\gamma)$ in an $S_\ell \times S_{k-\ell}$ -equivariant and $U(n)$ -equivariant manner. By Schur's lemma and Schur-Weyl duality, it can be written as

$$\iota_\gamma^\dagger(\iota_\alpha \otimes \iota_\beta) = (\text{id}_{V_\gamma^n} \otimes \omega_{\alpha\beta}^\gamma)(\iota_{\alpha\beta}^\gamma \otimes \text{id}_{W_\alpha \otimes W_\beta})^\dagger \quad (61)$$

where $\iota_{\alpha\beta}^\gamma : C_{\alpha\beta}^\gamma \otimes V_\gamma^n \rightarrow V_\alpha^n \otimes V_\beta^n$ is the $U(n)$ -equivariant isometry for the γ -isotypic subspace of $V_\alpha^n \otimes V_\beta^n$ and $\omega_{\alpha\beta}^\gamma : C_{\alpha\beta}^\gamma \otimes W_\alpha \otimes W_\beta \rightarrow W_\gamma$ the $S_\ell \times S_{k-\ell}$ -equivariant isometry for the (α, β) -isotypic subspace of W_γ .⁴ Combining everything together, and rearranging we obtain the formula

$$\frac{\dim W_\gamma \pi_\gamma(X + Y)}{|\gamma|!} = \sum_{\alpha, \beta} \text{Tr}_{C_{\alpha\beta}^\gamma} \left[(\iota_{\alpha\beta}^\gamma)^\dagger \frac{\dim W_\alpha \pi_\alpha(X)}{|\alpha|!} \otimes \frac{\dim W_\beta \pi_\beta(Y)}{|\beta|!} (\iota_{\alpha\beta}^\gamma) \right], \quad (62)$$

where we leave the summation over all pairs (α, β) such that $|\alpha| = \ell$, $|\beta| = k - \ell$ for $|\gamma| = k$ implicit simply because the multiplicity space $C_{\alpha\beta}^\gamma$ is empty if $|\alpha| + |\beta| \neq |\gamma|$ and thus the corresponding summand is taken to be zero.

In the remaining of the paper, we will use the above formula, Eq. (62), to prove numerous results concerning the eigenvalues of Hermitian matrices and their sums. In order to simplify the presentation of our results considerably, we introduce the following definitions.

Definition 21 (Hook-rescaled representations). Given an $n \times n$ matrix X and a Young diagram λ , define an operator $\Phi_\lambda(X)$ acting on the representation space V_λ^n as:

$$\Phi_\lambda(X) := \frac{\pi_\lambda(X)}{H_\lambda}, \quad (63)$$

where $H_\lambda = \prod_{b \in \lambda} h_\lambda(b)$ is the product of the Hook-lengths of the boxes in the Young diagram λ . Furthermore, the trace of $\Phi_\lambda(X)$ be

$$\varphi_\lambda(X) = \text{Tr}[\Phi_\lambda(X)] = \frac{s_\lambda(X)}{H_\lambda}. \quad (64)$$

Remark 22. The operator $\Phi_\lambda(X)$ and trace $\varphi_\lambda(X)$ are simply rescalings of the representation matrix $\pi_\lambda(X)$ of X and the character $\chi_\lambda(X)$ of X respectively. This normalization scheme is such that summing the rescaled character over all Young diagrams λ with fixed size k or over all sizes yields respectively

$$\sum_{\lambda \vdash k} \varphi_\lambda(X) = \frac{\text{Tr}(X)^k}{k!}, \quad \sum_{\lambda} \varphi_\lambda(X) = \exp(\text{Tr}(X)). \quad (65)$$

⁴Schur-Weyl duality provides us with a isometric and bijective identification between the multiplicity spaces for V_γ^n inside $V_\alpha^n \otimes V_\beta^n$ with respect to $U(n)$ and $W_\alpha \otimes W_\beta$ inside W_γ with respect to $S_\ell \times S_{k-\ell}$, hence why we use, by abuse of notation, the symbol $C_{\alpha\beta}^\gamma$ in both cases.

Before stating the main result of this section, we first generalize the quantities $\Phi_\lambda(X)$ and $\varphi_\lambda(X)$ to consider arguments involving more than one matrix.

Definition 23 (Mixed hook-rescaled representations). Given a pair (X, Y) of $n \times n$ matrices and a triple of Young diagrams (α, β, γ) , define an operator $\Phi_\gamma^{\alpha\beta}(X, Y)$ acting on the representation space V_γ^n as

$$\Phi_\gamma^{\alpha\beta}(X, Y) := \text{Tr}_{C_{\alpha,\beta}^\gamma} [\iota_{\alpha,\beta}^{\gamma;\dagger}(\Phi_\alpha(X) \otimes \Phi_\beta(Y)) \iota_{\alpha,\beta}^\gamma], \quad (66)$$

where $\iota_{\alpha,\beta}^\gamma : C_{\alpha,\beta}^\gamma \otimes V_\gamma^n \rightarrow V_\alpha^n \otimes V_\beta^n$ is the $U(n)$ -equivariant isometry for the γ -isotypic subspace of $V_\alpha^n \otimes V_\beta^n$. Note the sum over Young diagrams α , and β is implicitly restricted to those satisfying $|\alpha| + |\beta| = |\gamma|$, since otherwise the multiplicity space $C_{\alpha,\beta}^\gamma$ is zero-dimensional and thus the summand is taken to be zero. Analogously, let the trace of $\Phi_\gamma^{\alpha\beta}(X, Y)$ be denoted by

$$\varphi_\gamma^{\alpha\beta}(X, Y) := \text{Tr}[\Phi_\gamma^{\alpha\beta}(X, Y)] = \text{Tr}[P_{\alpha,\beta}^\gamma(\Phi_\alpha(X) \otimes \Phi_\beta(Y))], \quad (67)$$

where $P_{\alpha,\beta}^\gamma = \iota_{\alpha,\beta}^\gamma \iota_{\alpha,\beta}^{\gamma;\dagger}$ is the projection operator onto the γ -isotypic subspace of $V_\alpha^n \otimes V_\beta^n$.

Remark 24. While $\Phi_\lambda(X)$ is a homogeneous polynomial in the matrix coefficients of X of degree $|\lambda|$ in the sense that $\Phi_\lambda(sX) = s^{|\lambda|} \Phi_\lambda(X)$ holds for any $s \in \mathbb{R}$, $\Phi_\gamma^{\alpha\beta}(X, Y)$ is a bihomogeneous polynomial in X and Y of degrees $|\alpha|$ and $|\beta|$ respectively in the sense that for any $s, t \in \mathbb{R}$,

$$\Phi_\gamma^{\alpha\beta}(sX, tY) = s^{|\alpha|} t^{|\beta|} \Phi_\gamma^{\alpha\beta}(X, Y). \quad (68)$$

Remark 25. The quantity $\Phi_\gamma^{\alpha\beta}(X, Y)$ generalizes $\Phi_\lambda(X)$ in the sense that one can take the triple $(\alpha, \beta, \gamma) = (\lambda, (), \lambda)$ (where $()$ denotes the Young diagram with zero boxes), in which case $V_{()}^n$ supports the trivial one-dimensional representation with $\Phi_{()}(Y) = 1$ and because $\dim(C_{\lambda,()}^\lambda) = 1$,

$$\forall Y : \quad \Phi_\lambda^{\lambda,()}(X, Y) = \Phi_\lambda(X). \quad (69)$$

With these definitions in place, we are in a position to state the main result of this section.

Theorem 26 (Binomial theorem). For all $n \times n$ matrices X, Y , and all Young diagrams γ ,

$$\Phi_\gamma(X + Y) = \sum_{\alpha,\beta} \Phi_\gamma^{\alpha\beta}(X, Y), \quad (70)$$

where the sum is implicitly taken over all Young diagrams such that $|\alpha| + |\beta| = |\gamma|$.

Proof. See preceding discussion. □

Corollary 27. Let X, Y be $n \times n$ matrices. Then the following identities hold:

$$\forall \alpha, \beta : \quad \sum_{\gamma} \varphi_\gamma^{\alpha\beta}(X, Y) = \varphi_\alpha(X) \varphi_\beta(Y), \quad (71)$$

$$\forall \gamma : \quad \sum_{\alpha,\beta} \varphi_\gamma^{\alpha\beta}(X, Y) = \varphi_\gamma(X + Y). \quad (72)$$

Proof. The first identity involving a summation over α and β is simply the result of taking the trace of both sides of Theorem 26. For the second identity, involving a summation over γ , follows from the resolution of the identity $\sum_{\gamma: |\alpha|+|\beta|} P_{\alpha,\beta}^\gamma = \text{id}_{V_\alpha^n \otimes V_\beta^n}$:

$$\varphi_\alpha(X) \varphi_\beta(Y) = \text{Tr}[\Phi_\alpha(X) \otimes \Phi_\beta(Y)] = \sum_{\gamma} \text{Tr}[P_{\alpha,\beta}^\gamma(\Phi_\alpha(X) \otimes \Phi_\beta(Y))] = \sum_{\gamma} \varphi_\gamma^{\alpha\beta}(X, Y), \quad (73)$$

where the last equality is just the definition of $\varphi_\gamma^{\alpha\beta}(X, Y)$. Once again, the condition that $|\gamma| = |\alpha| + |\beta|$ is implicitly present. □

Remark 28. One might consider what happens to the operator $\Phi_{\gamma}^{\alpha\beta}(X, Y)$ when one sums over γ instead of (α, β) . However, an analogous identity concerning a sum over γ of the operator $\Phi_{\gamma}^{\alpha\beta}(X, Y)$ does not immediately follow because the tensor product $\Phi_{\alpha}(X) \otimes \Phi_{\beta}(Y)$ is not necessarily block diagonalizable into γ -isotypic subspaces in any obvious way. One exception occurs when $X = Y$, in which case,

$$\pi_{\alpha}(X) \otimes \pi_{\beta}(X) = \sum_{\gamma} \iota_{\alpha\beta}^{\gamma} (\pi_{\gamma}(X) \otimes \text{id}_{C_{\alpha\beta}^{\gamma}}) \iota_{\alpha\beta}^{\gamma\dagger}, \quad (74)$$

which is the operator version of the character product formula $\chi_{\alpha}(X)\chi_{\beta}(X) = \sum_{\gamma} c_{\alpha\beta}^{\gamma} \chi_{\gamma}(X)$.

3.3 Inequalities for Horn's original problem

sec:new_ineq_horn_orig

In Section 3.2 we introduced to the quantity $\varphi_{\gamma}^{\alpha\beta}(X, Y)$ (Eq. (67)) and saw in Corollary 27 that it satisfied two summation identities. Under the assumption that the matrices X , and Y are positive semidefinite, we can conclude that $\varphi_{\gamma}^{\alpha\beta}(X, Y)$ is non-negative. This observation implies the following upper bounds on $\varphi_{\gamma}^{\alpha\beta}(X, Y)$.

lem:XY_marginal_upper_bounds

Lemma 29. *Let $X, Y \geq 0$ be positive semidefinite $n \times n$ matrices. For all Young diagrams (α, β, γ) , we have the inequality*

$$\varphi_{\gamma}^{\alpha\beta}(X, Y) \leq \min\{\varphi_{\alpha}(X)\varphi_{\beta}(Y), \varphi_{\gamma}(X + Y)\}. \quad (75)$$

Proof. The claim follows directly from Corollary 27 provided $\varphi_{\gamma}^{\alpha\beta}(X, Y) \geq 0$. However, this follows because $\varphi_{\gamma}^{\alpha\beta}(X, Y) = \text{Tr}[P_{\alpha\beta}^{\gamma}(\Phi_{\alpha}(X) \otimes \Phi_{\beta}(Y))]$ and $P_{\alpha\beta}^{\gamma} \geq 0$ (it is a projection operator) and $\Phi_{\alpha}(X), \Phi_{\beta}(Y) \geq 0$ because $X, Y \geq 0$. $\square$

cor:orig_horn_ineq

Corollary 30. *Let $X, Y \geq 0$ be positive semidefinite $n \times n$ matrices. Then the following inequalities hold:*

$$\forall \alpha, \beta: \quad \varphi_{\alpha}(X)\varphi_{\beta}(Y) \leq \sum_{\substack{\alpha, \beta \\ c_{\alpha\beta}^{\gamma} > 0}} \varphi_{\gamma}(X + Y), \quad (76)$$

$$\forall \gamma: \quad \varphi_{\gamma}(X + Y) \leq \sum_{\substack{\alpha, \beta \\ c_{\alpha\beta}^{\gamma} > 0}} \varphi_{\alpha}(X)\varphi_{\beta}(Y). \quad (77)$$

Proof. The claim follows by applying upper bounds from Lemma 29 to the summations in Corollary 27 being careful to note that if $c_{\alpha\beta}^{\gamma} = 0$, then $\varphi_{\gamma}^{\alpha\beta}(X, Y) = 0$ and one only needs to consider the cases where $c_{\alpha\beta}^{\gamma} \neq 0$. $\square$

3.4 The $6j$ symbol for $U(n)$

sec:sixj_tet

The purpose of this section is to introduce a representation theoretic object, namely the $6j$ symbol, and then in the subsequent sections show how it places constraints on whether a given sextuple of eigenvalues (a, b, c, d, e, f) is tetrahedral in a manner which generalizes how the Littlewood-Richardson coefficient encodes whether a given triple (a, b, c) is trigonal. For the case where $n = 2$, this correspondence is well-known [Rob99; PR68; Bac10], while for the case where $n \geq 3$, this correspondence was already established indirectly by Christandl, Şahinoğlu, and Walter [CSW18]. Here we establish this correspondence directly.

As we discussed in the introduction, the $6j$ symbol emerges naturally when considering successive decompositions of the triple tensor product $V_{\alpha}^n \otimes V_{\beta}^n \otimes V_{\delta}^n$.

defn:sixj

Definition 31. Let $(\alpha, \beta, \gamma, \delta, \epsilon, \phi)$ be a sextuple of Young diagrams with sizes satisfying

$$|\alpha| + |\beta| = |\gamma|, \quad |\beta| + |\delta| = |\phi|, \quad |\gamma| + |\delta| = |\epsilon|, \quad |\alpha| + |\phi| = |\epsilon|. \quad (78)$$

The $U(n)$ - $6j$ operator is the operator

$$\left\{ \begin{smallmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{smallmatrix} \right\} : C_{\alpha, \beta}^{\gamma} \otimes C_{\delta, \gamma}^{\epsilon} \rightarrow C_{\alpha, \phi}^{\epsilon} \otimes C_{\beta, \delta}^{\phi}, \quad (79)$$

defined by

$$\left\{ \begin{smallmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{smallmatrix} \right\} := \frac{1}{\dim(V_\epsilon)} \text{Tr}_{V_\epsilon} [\iota_{\alpha\phi}^{\epsilon;\dagger} (\text{id}_{V_\alpha^n} \otimes \iota_{\beta\delta}^{\phi;\dagger}) (\iota_{\alpha\beta}^\gamma \otimes \text{id}_{V_\delta^n}) \iota_{\gamma\delta}^\epsilon]. \quad \text{\texttt{\{eq:sixj\}}} \quad (80)$$

To illustrate the situation more concretely, consider the following three projection operators on the triple tensor product $V_\alpha^n \otimes V_\beta^n \otimes V_\delta^n$:

$$P_{\alpha\beta}^\gamma \otimes \text{id}_{V_\delta^n}, \quad P_{\alpha\beta\delta}^\epsilon, \quad \text{id}_{V_\alpha^n} \otimes P_{\beta\delta}^\phi. \quad (81)$$

In general, $P_{\alpha\beta}^\gamma$ and $P_{\beta\delta}^\phi$ do not necessarily commute with each other in the sense that for some values of $\alpha, \beta, \delta, \gamma$, and ϕ , one observes

$$[P_{\alpha\beta}^\gamma \otimes \text{id}_{V_\delta^n}, \text{id}_{V_\alpha^n} \otimes P_{\beta\delta}^\phi] \neq 0. \quad (82)$$

Nevertheless, the projection operators $P_{\alpha\beta}^\gamma$ and $P_{\beta\delta}^\phi$ each respectively commute with $P_{\alpha\beta\delta}^\epsilon$ in the sense that for all $\alpha, \beta, \gamma, \delta, \epsilon$, and ϕ ,

$$[P_{\alpha\beta\delta}^\epsilon, P_{\alpha\beta}^\gamma \otimes \text{id}_{V_\delta^n}] = 0, \quad [P_{\alpha\beta\delta}^\epsilon, \text{id}_{V_\alpha^n} \otimes P_{\beta\delta}^\phi] = 0. \quad (83)$$

Let $\Pi_{\alpha\beta\delta}^{\epsilon;\gamma}$ and $\Pi_{\alpha\beta\delta}^{\epsilon;\phi}$ be projection operators given by

$$\begin{aligned} \Pi_{\alpha\beta\delta}^{\epsilon;\gamma} &:= P_{\alpha\beta\delta}^\epsilon (P_{\alpha\beta}^\gamma \otimes \text{id}_{V_\delta^n}) = (\iota_{\alpha\beta}^\gamma \otimes \text{id}_{V_\delta^n}) (P_{\gamma\delta}^\epsilon \otimes \text{id}_{C_{\alpha\beta}^\gamma}) (\iota_{\alpha\beta}^\gamma \otimes \text{id}_{V_\delta^n})^\dagger, \\ \Pi_{\alpha\beta\delta}^{\epsilon;\phi} &:= P_{\alpha\beta\delta}^\epsilon (\text{id}_{V_\alpha^n} \otimes P_{\beta\delta}^\phi) = (\text{id}_{V_\alpha^n} \otimes \iota_{\beta\delta}^\phi) (P_{\alpha\phi}^\epsilon \otimes \text{id}_{C_{\beta\delta}^\phi}) (\text{id}_{V_\alpha^n} \otimes \iota_{\beta\delta}^\phi)^\dagger, \end{aligned} \quad \text{\texttt{\{eq:double_projs\}}} \quad (84)$$

\texttt{prop:6j_norm_double_proj}

Proposition 32. For any sextuple of Young diagrams $(\alpha, \beta, \gamma, \delta, \epsilon, \phi)$,

$$\left\| \left\{ \begin{smallmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{smallmatrix} \right\} \right\|_\infty = \left\| \Pi_{\alpha\beta\delta}^{\epsilon;\gamma} \Pi_{\alpha\beta\delta}^{\epsilon;\phi} \right\|_\infty, \quad \left\| \left\{ \begin{smallmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{smallmatrix} \right\} \right\|_2^2 = \frac{1}{\dim V_\epsilon^n} \text{Tr}[\Pi_{\alpha\beta\delta}^{\epsilon;\gamma} \Pi_{\alpha\beta\delta}^{\epsilon;\phi}]. \quad (85)$$

3.5 Inequalities for the tetrahedral Horn problem

\texttt{sec:new_ineq_horn_tet}

In Section 3.3, we saw how the quantity $\varphi_{\gamma\epsilon\phi}^{\alpha\beta}(X, Y)$ (Eq. (67)) could be used to produce inequalities applicable to the original Horn problem. In this section, we generalize this analysis to produce inequalities applicable to the tetrahedral Horn problem using a generalization of $\varphi_{\gamma\epsilon\phi}^{\alpha\beta}(X, Y)$ which depends on three matrices.

Definition 33. Let X, Y, Z be $n \times n$ matrices. For any sextuple $(\alpha, \beta, \gamma, \delta, \epsilon, \phi)$ of Young diagrams, let

$$\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) := \text{Tr}[(P_{\alpha\beta}^\gamma \otimes \text{id}_{V_\delta^n}) P_{\alpha\beta\delta}^\epsilon (\text{id}_{V_\alpha^n} \otimes P_{\beta\delta}^\phi) \Phi_\alpha(X) \otimes \Phi_\beta(Y) \otimes \Phi_\delta(Z)]. \quad (86)$$

Remark that in our definition of $\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)$, we permit the sizes of the involved Young diagrams to be arbitrary. However, there are numerous conditions on the sizes of the Young diagrams $(\alpha, \beta, \gamma, \delta, \epsilon, \phi)$ which are necessary (but not always sufficient) for $\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)$ to be non-zero; specifically,

$$|\alpha| + |\beta| = |\gamma|, \quad |\beta| + |\delta| = |\phi|, \quad |\alpha| + |\phi| = |\epsilon|, \quad |\gamma| + |\delta| = |\epsilon|. \quad \text{\texttt{\{eq:yd_sizes_tet\}}} \quad (87)$$

If we additionally let $\Omega_{\alpha\beta\delta}^{X,Y,Z} := \Phi_\alpha(X) \otimes \Phi_\beta(Y) \otimes \Phi_\delta(Z)$, then we can compactly express $\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)$ as

$$\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) = \text{Tr}[\Pi_{\alpha\beta\delta}^{\epsilon;\gamma} \Pi_{\alpha\beta\delta}^{\epsilon;\phi} \Omega_{\alpha\beta\delta}^{X,Y,Z}]. \quad \text{\texttt{\{eq:varphi_compact_tet\}}} \quad (88)$$

Unlike the quantity $\varphi_{\gamma\epsilon\phi}^{\alpha\beta}(X, Y) = \text{Tr}[P_{\alpha\beta}^\gamma \Phi_\alpha(X) \otimes \Phi_\beta(Y)]$ involving two matrices X, Y (see Section 3.2), the quantity defined here, $\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)$, is not necessarily non-negative or even real valued even under the assumption that $X, Y, Z \geq 0$ are positive semidefinite.

Nevertheless, summing over either γ or ϕ restores non-negativity because it eliminates one of the potentially non-commuting projection operators $\Pi_{\alpha\beta\delta}^{\epsilon;\gamma}$ or $\Pi_{\alpha\beta\delta}^{\epsilon;\phi}$. We denote these partial sums of $\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)$ over γ and ϕ respectively as

$$\varphi_{-\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) := \sum_{\gamma} \varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) = \text{Tr}[\Pi_{\alpha\beta\delta}^{\epsilon;\phi} \Omega_{\alpha\beta\delta}^{X,Y,Z}] = \text{Tr}[P_{\gamma\delta}^\epsilon (\Phi_\gamma^{\alpha\beta}(X, Y) \otimes \Phi_\delta(Z))], \quad (89)$$

$$\varphi_{\gamma\epsilon-}^{\alpha\beta\delta}(X, Y, Z) := \sum_{\phi} \varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) = \text{Tr}[\Pi_{\alpha\beta\delta}^{\epsilon;\gamma} \Omega_{\alpha\beta\delta}^{X,Y,Z}] = \text{Tr}[P_{\alpha\phi}^\epsilon (\Phi_\alpha(X) \otimes \Phi_\phi^{\beta\delta}(Y, Z))], \quad (90)$$

$$\begin{array}{ccccccc}
& & \xrightarrow{\sum_{\gamma}} & \varphi_{-\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) & \xrightarrow{\sum_{\epsilon}} & \varphi_{\alpha}(X) \varphi_{\phi}^{\beta\delta}(Y, Z) & \xrightarrow{\sum_{\beta\delta}} \varphi_{\alpha}(X) \varphi_{\phi}(Y + Z) \\
\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) & \xrightarrow{\sum_{\alpha\beta\delta\gamma\phi}} & \varphi_{\epsilon}(X + Y + Z) & \xleftarrow{\sum_{\alpha\beta\delta}} & \varphi_{\epsilon}^{\alpha\beta\delta}(X, Y, Z) & \xrightarrow{\sum_{\epsilon}} & \varphi_{\alpha}(X) \varphi_{\beta}(Y) \varphi_{\delta}(Z) \\
& & \xrightarrow{\sum_{\phi}} & \varphi_{\gamma\epsilon-}^{\alpha\beta\delta}(X, Y, Z) & \xrightarrow{\sum_{\epsilon}} & \varphi_{\gamma}^{\alpha\beta}(X, Y) \varphi_{\delta}(Z) & \xrightarrow{\sum_{\alpha\beta}} \varphi_{\gamma}(X + Y) \varphi_{\delta}(Z)
\end{array}$$

Figure 3: Various summation identities satisfied by $\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)$.

[fig:sum_ident](#)

where the sum over γ is implicitly such that $|\gamma| = |\alpha| + |\beta|$ and the sum over ϕ is implicitly such that $|\phi| = |\beta| + |\delta|$. Although $\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)$ is not necessarily real or positive, the above expression makes it evident that

$$\varphi_{-\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) \geq 0, \quad \varphi_{\gamma\epsilon-}^{\alpha\beta\delta}(X, Y, Z) \geq 0. \quad (91)$$

The various summation identities satisfied by Fig. 3 are summarized in Fig. 3.

The next series of results concerns upper bounds on the absolute value of $\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)$. These upper bounds follow from the following general lemma applied to the compact expression for $\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)$ appearing in Eq. (88).

[prop:cauchy_schwarz_ineq_double_proj](#)

Proposition 34. *Let $\Omega \geq 0$ be a positive semidefinite operator and let Π_1, Π_2 be Hermitian projection operators acting on a common vector space. Then*

$$|\text{Tr}(\Pi_1 \Pi_2 \Omega)| \leq \|\Pi_1 \Pi_2\|_{\infty} \text{Tr}(\Pi_1 \Omega)^{\frac{1}{2}} \text{Tr}(\Pi_2 \Omega)^{\frac{1}{2}}. \quad (92)$$

Proof. As $\Omega \geq 0$ is positive semidefinite, let $\Omega^{\frac{1}{2}} \geq 0$ denote its square root. Then let $L = \Pi_1 \Pi_2 \Omega^{\frac{1}{2}}$ and $R = \Pi_1 \Omega^{\frac{1}{2}}$ so that

$$\langle L, R \rangle = \text{Tr}(L^{\dagger} R) = \text{Tr}(\Pi_1 \Pi_2 \Omega \Pi_1) = \text{Tr}(\Pi_1 \Pi_2 \Omega), \quad (93)$$

where the last equality uses the assumption that $\Pi_1^2 = \Pi_1$. An application of the Cauchy-Schwarz inequality for the Hilbert-Schmidt inner product $\langle L, R \rangle$ then implies

$$|\text{Tr}(\Pi_1 \Pi_2 \Omega)|^2 = |\langle L, R \rangle|^2 \leq \text{Tr}[L^{\dagger} L] \text{Tr}[R^{\dagger} R] = \text{Tr}[\Pi_1 \Pi_2 \Omega \Pi_2 \Pi_1] \text{Tr}[\Pi_1 \Omega \Pi_1] = \text{Tr}[\Pi_1 \Pi_2 \Omega \Pi_2] \text{Tr}[\Pi_1 \Omega]. \quad (94)$$

Now let $\{v_j\}_j$ be an orthonormal eigenbasis for $\Pi_2 \Omega \Pi_2$ with eigenvalue $\omega_j \geq 0$. Then, because it was assumed that $\Pi_2^2 = \Pi_2$, we have

$$\text{Tr}[\Pi_1 \Pi_2 \Omega \Pi_2] = \text{Tr}[(\Pi_2 \Pi_1^2 \Pi_2)(\Pi_2 \Omega \Pi_2)] = \sum_j \omega_j \langle \Pi_1 \Pi_2 v_j, \Pi_1 \Pi_2 v_j \rangle \leq \sum_j \omega_j \|\Pi_1 \Pi_2\|_{\infty}^2 = \text{Tr}[\Pi_2 \Omega] \|\Pi_1 \Pi_2\|_{\infty}^2. \quad (95)$$

The claim then follows from combining these inequalities and taking the square root of both sides. $\square$

[lem:varphi_tet_main_ineq](#)

Lemma 35. *Let $X, Y, Z \geq 0$ be positive semidefinite $n \times n$ matrices. Then for all sextuples of Young diagrams $(\alpha, \beta, \gamma, \delta, \epsilon, \phi)$,*

$$\left| \varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) \right| \leq \left\| \begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix} \right\|_{\infty} \varphi_{-\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)^{\frac{1}{2}} \varphi_{\gamma\epsilon-}^{\alpha\beta\delta}(X, Y, Z)^{\frac{1}{2}}. \quad (96)$$

Proof. Under the assumption that $X, Y, Z \geq 0$, we have that $\Omega_{\alpha\beta\delta}^{X, Y, Z} = \Phi_{\alpha}(X) \otimes \Phi_{\beta}(Y) \otimes \Phi_{\delta}(Z) \geq 0$ is positive semidefinite also. Now applying Proposition 34 to the compact expression for $\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)$ in Eq. (88) yields

$$\left| \text{Tr}[\Pi_{\alpha\beta\delta}^{\epsilon; \gamma} \Pi_{\alpha\beta\delta}^{\epsilon; \phi} \Omega_{\alpha\beta\delta}^{X, Y, Z}] \right| \leq \left\| \Pi_{\alpha\beta\delta}^{\epsilon; \gamma} \Pi_{\alpha\beta\delta}^{\epsilon; \phi} \right\|_{\infty} \text{Tr}[\Pi_{\alpha\beta\delta}^{\epsilon; \gamma} \Omega_{\alpha\beta\delta}^{X, Y, Z}]^{\frac{1}{2}} \text{Tr}[\Pi_{\alpha\beta\delta}^{\epsilon; \phi} \Omega_{\alpha\beta\delta}^{X, Y, Z}]^{\frac{1}{2}} \quad (97)$$

which is equivalent to the claim due to Proposition 32. $\square$

Our next proposition is essential for places bounds on $\varphi_{-\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)$ and $\varphi_{\gamma\epsilon-}^{\alpha\beta\delta}(X, Y, Z)$ which depend only the eigenvalues of $X, Y, X + Y, Z, X + Y + Z$, and $Y + Z$. *prop:tet-partial-sum-ineq*

Proposition 36. *Let $X, Y, Z \geq 0$ be positive semidefinite $n \times n$ matrices. Then we have bounds on the partial sums,*

$$\begin{aligned}\varphi_{-\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) &\leq \varphi_\alpha(X)\varphi_\phi(Y + Z), \\ \varphi_{\gamma\epsilon-}^{\alpha\beta\delta}(X, Y, Z) &\leq \varphi_\gamma(X + Y)\varphi_\delta(Z),\end{aligned}\tag{98}$$

as well as bounds which are uniform in both γ and ϵ :

$$\max\{\varphi_{\gamma\epsilon-}^{\alpha\beta\delta}(X, Y, Z), \varphi_{-\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)\} \leq \varphi_\epsilon^{\alpha\beta\delta}(X, Y, Z) \leq \min\{\varphi_\epsilon(X + Y + Z), \varphi_\alpha(X)\varphi_\beta(Y)\varphi_\delta(Z)\}.\tag{99}$$

Proof. The bounds are consequences from summation identities and the fact that the partial sums $\varphi_{-\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)$ and $\varphi_{\gamma\epsilon-}^{\alpha\beta\delta}(X, Y, Z)$ are non-negative real-valued. The first pair of bounds follow from the summation identities

$$\begin{aligned}\varphi_\alpha(X)\varphi_\phi(Y + Z) &= \sum_{\epsilon\beta\delta} \varphi_{-\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z), \\ \varphi_\gamma(X + Y)\varphi_\delta(Z) &= \sum_{\epsilon\alpha\beta} \varphi_{\gamma\epsilon-}^{\alpha\beta\delta}(X, Y, Z).\end{aligned}\tag{100}$$

The last inequality comes from

$$\varphi_\epsilon^{\alpha\beta\delta}(X, Y, Z) = \sum_{\phi} \varphi_{-\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) = \sum_{\gamma} \varphi_{\gamma\epsilon-}^{\alpha\beta\delta}(X, Y, Z)\tag{101}$$

and the fact that

$$\begin{aligned}\varphi_\alpha(X)\varphi_\beta(Y)\varphi_\delta(Z) &= \sum_{\epsilon} \varphi_\epsilon^{\alpha\beta\delta}(X, Y, Z), \\ \varphi_\epsilon(X + Y + Z) &= \sum_{\alpha\beta\delta} \varphi_\epsilon^{\alpha\beta\delta}(X, Y, Z).\end{aligned}\tag{102}$$

□

We can now immediately prove the “only if” portion of the claim of Theorem 1. *cor:tet-eigenvalue-cubic*

Corollary 37. *Let $X, Y, Z \geq 0$ be positive semidefinite $n \times n$ matrices. Then for all triples of Young diagrams (α, β, δ) ,*

$$\varphi_\alpha(X)^{\frac{2}{3}}\varphi_\beta(Y)\varphi_\delta(Z)^{\frac{2}{3}} \leq \sum_{\gamma, \epsilon, \phi} \left\| \begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix} \right\|_\infty \varphi_\phi(Y + Z)^{\frac{1}{3}}\varphi_\gamma(X + Y)^{\frac{1}{3}}\varphi_\epsilon(X + Y + Z)^{\frac{1}{3}}.\tag{103}$$

Proof. The claim follows by summing the inequality in Lemma 35 over triples (γ, ϵ, ϕ) and relaxing both the sides of the resulting inequality. For the lower bound, we have $\varphi_\alpha(X)\varphi_\beta(Y)\varphi_\delta(Z) = \sum_{\gamma\epsilon\phi} \varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)$ which implies

$$\varphi_\alpha(X)\varphi_\beta(Y)\varphi_\delta(Z) \leq \sum_{\gamma\epsilon\phi} \left| \varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) \right|.\tag{104}$$

Then for the upper bound, Proposition 36 can be used to produce

$$\varphi_{-\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)\varphi_{\gamma\epsilon-}^{\alpha\beta\delta}(X, Y, Z) \leq [\varphi_\alpha(X)\varphi_\phi(Y + Z)\varphi_\gamma(X + Y)\varphi_\delta(Z)]^p \varphi_\epsilon(X + Y + Z)^{2(1-p)},\tag{105}$$

which holds for any $p \in [0, 1]$. The claim corresponds to the choice $p = \frac{2}{3}$ which makes the exponents match. Then the claimed result is obtained from dividing out both sides by a common factor of $(\varphi_\alpha(X)\varphi_\delta(Z))^{\frac{1}{3}}$ (note that if $\varphi_\alpha(X)\varphi_\delta(Z) = 0$, the inequality would hold trivially). □

The proof of the “if” portion of Theorem 1 can be found in Section 4.4 which proves the distance bound of Theorem 2 (this implies the “if” portion of Theorem 1 after sending $k \rightarrow \infty$).

3.6 The large n limit & an entropic inequality

sec:large_n

The purpose of this section is to show how to leverage the inverse polynomial bound on $\left\| \begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix} \right\|$ for tetrahedral (a, b, c, d, e, f) provided by Corollary 4 to derive inequalities on the entropies of the eigenvalues. Our strategy will be to prove an upper bound on the $6j$ symbol which depends only on the dimensions of the underlying irreducible representations and the multiplicity spaces between them. By combining this upper bound with the lower bound provided by Corollary 4 (specifically the n -independent bound in Eq. (170)), we will prove the following.

thm:entropic_ineq

Theorem 38. *Let (a, b, c, d, e, f) be tetrahedral eigenvalues. Then*

$$s(c) + s(f) - s(b) - s(e) \geq 0, \quad (106)$$

{eq:entropic_ineq}

where

$$s(x) := - \sum_{i=1}^n x_i \log x_i. \quad (107)$$

{eq:little_s_defn}

We begin with our upper bound on the $6j$ symbol.

lem:sixj_ssa_bound

Lemma 39. *Let $(\alpha, \beta, \gamma, \delta, \epsilon, \phi)$ be a sextuple of Young diagrams. Then*

$$\left\| \begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix} \right\|_2^2 \leq \left(c_{\alpha, \phi}^{\epsilon} c_{\beta, \delta}^{\phi} c_{\alpha, \beta}^{\gamma} c_{\gamma, \delta}^{\epsilon} \right)^{\frac{1}{2}} \frac{\dim(V_{\gamma}^n) \dim(V_{\phi}^n)}{\dim(V_{\beta}^n) \dim(V_{\epsilon}^n)}. \quad (108)$$

Proof. Here we provide a diagrammatic proof of the claimed inequality. For the sake of clarity, we abbreviate the irreducible representation labels, V_{λ}^n , and omit the multiplicity space labels on the intermediate steps.

$$\left(\begin{array}{c} \epsilon \\ \gamma \end{array} \right) \left\| \begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix} \right\|_2^2 = \left[\begin{array}{c} \gamma \\ \delta \\ \epsilon \end{array} \right] \leq \left[\begin{array}{c} \gamma \\ \delta \\ \epsilon \end{array} \right] = \left(\begin{array}{c} \epsilon \\ \gamma \end{array} \right) \left(\begin{array}{c} \beta \\ \alpha \end{array} \right) \left[\begin{array}{c} \epsilon \\ \gamma \\ \delta \\ \alpha \end{array} \right]^{\frac{1}{2}} \quad (109)$$

The first equality follows from Definition 31. The inequality follows from an application of the Cauchy-Schwarz inequality with respect to the annotated contraction. The final equality follows from applying Schur's lemma to components of the diagram mapping V_{β}^n to itself on the left and V_{ϵ}^n to itself on the right. Note that the factors of $\dim(V_{\epsilon}^n)^2$ appearing on left-most and right-most sides of the above inequality cancel producing the claimed inequality. $\square$

An immediate application of Lemma 39 is to combine it with Eq. (170). If (a, b, c, d, e, f) are tetrahedral eigenvalues, then by Corollary 4 there exists a converging sequence of Young diagrams $(\alpha_k, \beta_k, \gamma_k, \delta_k, \epsilon_k, \phi_k)$ such that

$$\frac{1}{q(k)} \leq \left\| \begin{Bmatrix} \alpha_k & \beta_k & \gamma_k \\ \delta_k & \epsilon_k & \phi_k \end{Bmatrix} \right\|_2^2 \leq \left(c_{\alpha_k, \phi_k}^{\epsilon_k} c_{\beta_k, \delta_k}^{\phi_k} c_{\alpha_k, \beta_k}^{\gamma_k} c_{\gamma_k, \delta_k}^{\epsilon_k} \right)^{\frac{1}{2}} \frac{\dim(V_{\gamma_k}^n) \dim(V_{\phi_k}^n)}{\dim(V_{\beta_k}^n) \dim(V_{\epsilon_k}^n)}. \quad (110)$$

{eq:combined_bound}

where $q(k)$ is some polynomial in k independent of the dimension n (see Eq. (170)). Remarkably, the Littlewood-Richardson coefficients $c_{\alpha_k, \phi_k}^{\epsilon_k}$, $c_{\beta_k, \delta_k}^{\phi_k}$, $c_{\alpha_k, \beta_k}^{\gamma_k}$, and $c_{\gamma_k, \delta_k}^{\epsilon_k}$ are also *independent* of the dimension parameter n defining the group $U(n)$ (assuming n is at least as large as the maximum depth of any of the involved Young diagrams. This observation provides us with an opportunity to optimize the above bound by varying the value of n .

In particular, it will be useful to better understand how the dimension, $\dim(V_{\lambda}^n)$, of the irreducible representation V_{λ}^n of $U(n)$ grows as n becomes much larger than the number of rows of λ . In general, for fixed λ , the function $n \mapsto \dim(V_{\lambda}^n)$ can grow at an exponential rate with increasing n . One can find estimates for

$\dim(V_\lambda^n)$ in the regime where n is large compared to $|\lambda|$ and also the last row $\lambda_{\min}$ is large compared to the depth of λ in paper by Itzykson and Nauenberg [IN66, Section III]. Here we use an elegant presentation of this scaling, which relates the asymptotics of $\dim(V_\lambda^n)$ to the value of $\dim(W_\lambda)$ (see [Har23, Lemma 2] or [CHW07, Equations (25, 26)].

prop:harrow_formula

Proposition 40. *Let λ be a Young diagram with depth at most n . Then*

$$\frac{\dim(V_\lambda^n)}{n^{|\lambda|}} = \frac{\dim(W_\lambda)}{|\lambda|!} \prod_{(i,j) \in \lambda} \left(1 + \frac{j-i}{n}\right). \quad (111)$$

Proof. The statement follows from the dimension formulas for $\dim(V_\lambda^n)$ and $\dim(W_\lambda)$ in Eq. (46) and Eq. (49). See also [Har23, Appendix A]. $\square$

From Proposition 40 we obtain the following corollary.

cor:harrow_ssa_lim

Corollary 41. *Let $\gamma, \phi, \beta, \epsilon$ be a quadruple of Young diagrams such that*

$$|\gamma| + |\phi| = |\beta| + |\epsilon|. \quad (112)$$

eq:balanced_sizes

Then

$$\lim_{n \rightarrow \infty} \frac{\dim(V_\gamma^n) \dim(V_\phi^n)}{\dim(V_\beta^n) \dim(V_\epsilon^n)} = \frac{|\beta|! |\epsilon|! \dim(W_\gamma) \dim(W_\phi)}{|\gamma|! |\phi|! \dim(W_\beta) \dim(W_\epsilon)} = \frac{H_\beta H_\epsilon}{H_\gamma H_\phi} \quad (113)$$

Proof. From Proposition 40 one evidently has for any Young diagram λ the limit

$$\lim_{n \rightarrow \infty} \frac{\dim(V_\lambda^n)}{n^{|\lambda|}} = \frac{\dim(W_\lambda)}{|\lambda|!} = \frac{1}{H_\lambda}. \quad (114)$$

The claim then follows because the missing prefactor, $\frac{n^{|\beta|} n^{|\epsilon|}}{n^{|\gamma|} n^{|\phi|}}$, is equal to one by Eq. (112). $\square$

We are now in a position to prove the main result of this section.

Proof of Theorem 38. As (a, b, c, d, e, f) are tetrahedral, the inverse polynomial bound $\left\| \begin{Bmatrix} \alpha_k & \beta_k & \gamma_k \\ \delta_k & \epsilon_k & \phi_k \end{Bmatrix} \right\|$ implies it is non-zero and thus Eq. (112) holds. Therefore, by Corollary 41 and the discussion surrounding the bound in Eq. (110) we conclude that along the convergent sequence of Young diagrams,

$$\lim_{k \rightarrow \infty} \frac{1}{k} \log \left(\frac{H_{\beta_k} H_{\epsilon_k}}{H_{\gamma_k} H_{\phi_k}} \right) \geq 0 \quad (115)$$

eq:hook_ratio_limit

provided the limit. To prove the limit exists, an application of Stirling's approximation, $\log n! \approx n \log n - n$ [Rob55], to the approximation formula $H_\lambda \approx \prod_{i=1}^n \lambda_i!$ produces for any constant η the formula

$$\frac{1}{\eta} \log H_\lambda \approx (\log \eta - 1) |\lambda| + \sum_{i=1}^n \frac{\lambda_i}{\eta} \log \frac{\lambda_i}{\eta} \quad (116)$$

Because $|\gamma| + |\phi| = |\beta| + |\epsilon|$ (Eq. (112)), the $(\log \eta - 1)$ terms will cancel in Eq. (115). Then using the fact that $(\beta_k, \gamma_k, \epsilon_k, \phi_k)$ converge in proportion to the eigenvalues (b, c, e, f) (with proportionality constant $\eta = \frac{k}{\text{Tr}[e]}$), we conclude that Eq. (115) has a limit and moreover it equals the left-hand side of Eq. (106). $\square$

4 Distance bounds via eigenvalue concentration

sec:distance_bounds

The purpose of this section is to provide a proof of the distance bound in Theorem 2. With Theorem 2 established, the “if” portion of Theorem 1 follows immediately. Note that “only if” portion of Theorem 1 is just a restatement of Corollary 37 proven in the previous sections.

These bounds make use of eigenvalue concentration techniques, which we review first in Section 4.1, as well as formulas from random matrix theory, which we review afterwards in Section 4.2. As a warm-up exercise, in Section 4.3 we demonstrate how these techniques can be applied to obtain a distance bound for the original Horn problem. Then we generalize the proof technique in Section 4.4 to prove Theorem 2.

4.1 Eigenvalue estimation & separation

sec:eig_est_and_sep

In this section we show how the representations $\pi_\lambda(X)$ of Hermitian matrices X encode useful information about the eigenvalues of X . These results apply to the case where $X \geq 0$ is a positive semidefinite matrix, in which case the representation $\pi_\lambda(X) \geq 0$ is positive semidefinite also. Thus whenever $X \geq 0$, one can define a probability distribution known as the *Schur-Weyl* distribution [OW16].

Definition 42. The Schur-Weyl distribution of order k for a positive semidefinite matrix $X \geq 0$ with eigenvalues $x = (x_1, \dots, x_n)$ is defined as the probability distribution over Young diagrams λ with k boxes with probability

$$p_x(\lambda) := \frac{\text{Tr}[P_\lambda X^{\otimes k}]}{\text{Tr}[X^{\otimes k}]} = \frac{\dim W_\lambda s_\lambda(x)}{\text{Tr}[x]^k} = \frac{|\lambda|!}{\text{Tr}[x]^k} \varphi_\lambda(x). \quad (117)$$

A key property of the Schur-Weyl distribution $p_x(\lambda)$ is that it behaves similarly to the multinomial distribution $m_x(\lambda)$ where

$$m_x(\lambda) := \frac{|\lambda|!}{\text{Tr}[x]^k} \frac{x_1^{\lambda_1}}{\lambda_1!} \frac{x_2^{\lambda_2}}{\lambda_2!} \cdots \frac{x_n^{\lambda_n}}{\lambda_n!}. \quad (118)$$

prop:multinomial_bound

Proposition 43. Given a Young diagram λ and non-negative eigenvalues x ,

$$p_x(\lambda) \leq \dim V_\lambda^n m_x(\lambda). \quad (119)$$

Proof. The proof follows from (i) an upper bound on the Schur-polynomial $s_\lambda(x)$,

$$s_\lambda(x) \leq \dim V_\lambda^n x_1^{\lambda_1} x_2^{\lambda_2} \cdots x_n^{\lambda_n}, \quad (120)$$

which follows because $s_\lambda(x)$ can be expressed as a sum over $\dim V_\lambda^n$ multinomials of the form $x_1^{\mu_1} x_2^{\mu_2} \cdots x_n^{\mu_n}$ with $x_1^{\lambda_1} x_2^{\lambda_2} \cdots x_n^{\lambda_n}$ the largest, and (ii) a lower bound on the Hook-product,

$$H_\lambda \geq \lambda_1! \cdots \lambda_n!, \quad (121)$$

which follows because $h_\lambda(b) \geq \lambda_i - j + 1$ holds for the box b in row i and column j . $\square$

Remark 44. One can eliminate the factor of $\dim V_\lambda^n$ using the dimension bound

$$\dim V_\lambda^n \leq (k+1)^{\frac{n(n-1)}{2}}. \quad (122)$$

The following is known result is known as eigenvalue or spectrum estimation [ARS88; KW01; HM02].

Proposition 45 (Eigenvalue estimation). *Let $X \geq 0$ be a positive semidefinite $n \times n$ matrix with eigenvalues $x = (x_1 \geq \cdots \geq x_n \geq 0)$. Then the Schur-Weyl distribution $p_x(\lambda)$ for $\lambda \vdash k$ satisfies*

$$p_x(\lambda) \leq (k+1)^{\frac{n(n-1)}{2}} \exp(-kD(\underline{\lambda} \parallel \underline{x})) \quad (123)$$

where $D(\underline{\lambda} \parallel \underline{x})$ denotes the Kullback-Leibler divergence between the normalized Young diagram $\underline{\lambda} = \lambda/|\lambda|$ and the normalized eigenvalues $\underline{x} = x/\text{Tr}[x]$.

This result captures the fact that, like the multinomial distribution $m_x(\lambda)$, the Schur-Weyl distribution $p_x(\lambda)$ is sharply concentrated around Young diagrams λ whose distribution, $\underline{\lambda}$, is similar to the eigenvalue distribution $\underline{x}$.

For our purposes, it will be useful to use a related result which captures the fact if two positive semidefinite matrices $X, Y \geq 0$ have dissimilar eigenvalue distributions then the fidelity between their Schur-Weyl distributions is exponentially small in k . Recall the fidelity $F(p, q) \in [0, 1]$ between distributions p, q on the set $\{1, \dots, n\}$ is defined by

$$F(p, q) = \sum_{i=1}^n \sqrt{p_i q_i}, \quad (124)$$

and is equal to $F(p, q) = 1$ if and only if $p = q$.

Lemma 46 (Eigenvalue separation). *Let (x, y) be eigenvalues of positive semidefinite $n \times n$ matrices $X, Y \geq 0$. Then the fidelity between Schur-Weyl distributions p_x and p_y of order k ,*

$$F_k(p_x, p_y) := \sum_{\lambda \vdash k} \sqrt{p_x(\lambda) p_y(\lambda)}, \quad (125)$$

is bounded by

$$F_k(p_x, p_y) \leq (k+1)^{n(n-1)/2} F(\underline{x}, \underline{y})^k. \quad (126)$$

where $F(\underline{x}, \underline{y})$ is the fidelity between the eigenvalue distributions of X and Y ,

$$F(\underline{x}, \underline{y}) = \sum_{i=1}^n \sqrt{\frac{x_i}{\text{Tr}[X]} \frac{y_i}{\text{Tr}[Y]}}. \quad (127)$$

Proof. The proof follows from Proposition 43 and the fact that fidelities are multiplicative with respect to multinomial distributions, i.e. $F_k(m_x, m_y) \leq F(\underline{x}, \underline{y})$.

$$F_k(p_x, p_y) \leq (k+1)^{n(n-1)/2} F_k(m_x, m_y) = (k+1)^{n(n-1)/2} F(\underline{x}, \underline{y})^k. \quad (128)$$

□

Remark 47. The eigenvalue separation lemma states that when two positive semidefinite matrices $X, Y \geq 0$ have the same trace $\tau = \text{Tr}(X) = \text{Tr}(Y)$ but distinct eigenvalues $x \neq y$, then we can conclude $F(\underline{x}, \underline{y}) < 1 - \varepsilon$ for some ε and therefore as $k \rightarrow \infty$, $F_k(p_x, p_y) \rightarrow 0$.

Remark 48. In order to quantify the separation between distributions more geometrically, we use the fact that the Bhattacharyya coefficient, defined as $B(p, q) = -\log F(p, q)$, satisfies the lower bound $B(p, q) \geq \frac{1}{4} \|p - q\|_1^2$ which implies

$$F(p, q)^k \leq \exp\left(-\frac{1}{4} \|p - q\|_1^2\right). \quad (129)$$

4.2 Results for random matrices

[sec:random_matrices](#)

The purpose of this section is record some results concerning sums of random Hermitian matrices with fixed eigenvalues sampled uniformly from their respective orbits. The first two propositions follow from standard applications of Schur's lemma, and are used to prove the others.

Proposition 49. *Let $x = (x_1 \geq \dots \geq x_n)$ be a set of eigenvalues and let $X \sim \mathcal{O}_x$ denote a random Hermitian matrix of the form $X = U[x]U^{-1}$ where $U \in \text{U}(n)$ is sampled uniformly with respect to the normalized Haar measure on $\text{U}(n)$. Then for any Young diagram λ ,*

$$\mathbb{E}_{X \sim \mathcal{O}_x} \pi_\lambda(X) = s_\lambda(x) \frac{\text{id}_{V_\lambda^n}}{\dim V_\lambda^n}. \quad (130)$$

Proposition 50. *Given a positive real number $\tau > 0$, let $X \sim \text{HS}_\tau$ denote a random Hermitian matrix with coefficients $X_{ij} = \frac{\tau}{n} \sum_{k,s,t=1}^n U_{ik;ss} U_{tt;jk}^{-1}$ where $U \in \text{U}(n^2)$ is a $n^2 \times n^2$ unitary matrix sampled uniformly with respect to the normalized Haar measure on $\text{U}(n^2)$. Then for any Young diagram λ ,*

$$\mathbb{E}_{X \sim \text{HS}_\tau} \pi_\lambda(X) = \tau^{|\lambda|} q_{n,\lambda} \frac{\text{id}_{V_\lambda^n}}{\dim V_\lambda^n}. \quad (131)$$

where $q_{n,\lambda}$ is the dimension ratio

$$q_{n,\lambda} := \frac{(\dim V_\lambda^n)^2}{\dim V_{(|\lambda|)}^{n^2}}. \quad (132)$$

Corollary 51. *Let (x, y) be a pair of eigenvalues. Then*

$$\mathbb{E}_{X \sim \mathcal{O}_x} \mathbb{E}_{Y \sim \mathcal{O}_y} \frac{\varphi_\gamma^{\alpha\beta}(X, Y)}{\dim V_\gamma^n} = c_{\alpha\beta}^\gamma \frac{\varphi_\alpha(x)}{\dim V_\alpha^n} \frac{\varphi_\beta(y)}{\dim V_\beta^n}. \quad (133)$$

$$\mathbb{E}_{X \sim \text{HS}_{\text{Tr}[x]}} \mathbb{E}_{Y \sim \text{HS}_{\text{Tr}[y]}} \varphi_\gamma^{\alpha\beta}(X, Y) = \frac{\text{Tr}[x]^{|\alpha|}}{|\alpha|!} \frac{\text{Tr}[y]^{|\beta|}}{|\beta|!} q_{n,\alpha} q_{n,\beta} \frac{c_{\alpha\beta}^\gamma \dim V_\gamma^n}{\dim(V_\alpha^n \otimes V_\beta^n)}. \quad (134)$$

cor:random_matrices_tet

Corollary 52. *Let (x, y, z) be a triple of eigenvalues. Then*

$$\mathbb{E}_{X \sim \mathcal{O}_x} \mathbb{E}_{Y \sim \mathcal{O}_y} \mathbb{E}_{Z \sim \mathcal{O}_z} \frac{\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)}{\dim V_\epsilon^n} = \left\| \begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix} \right\|_2^2 \frac{\varphi_\alpha(x)}{\dim V_\alpha^n} \frac{\varphi_\beta(y)}{\dim V_\beta^n} \frac{\varphi_\delta(z)}{\dim V_\delta^n}. \quad (135)$$

$$\mathbb{E}_{X \sim \text{HS}_{\text{Tr}[x]}} \mathbb{E}_{Y \sim \text{HS}_{\text{Tr}[y]}} \mathbb{E}_{Z \sim \text{HS}_{\text{Tr}[z]}} \varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) = \frac{\text{Tr}[x]^{|\alpha|}}{|\alpha|!} \frac{\text{Tr}[y]^{|\beta|}}{|\beta|!} \frac{\text{Tr}[z]^{|\delta|}}{|\delta|!} q_{n,\alpha} q_{n,\beta} q_{n,\delta} \frac{\dim V_\epsilon^n \left\| \begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix} \right\|_2^2}{\dim(V_\alpha^n \otimes V_\beta^n \otimes V_\delta^n)}. \quad (136)$$

4.3 Distance bound for the original Horn problem

sec:dist_bound_for_orig_horn

In this section, we prove the following proposition pertaining to the original Horn problem as a warm-up exercise before we prove its generalization for the tetrahedral Horn problem (Theorem 2) in Section 4.4.

prop:approx_tri

Proposition 53. *Let (a, b, c) be a triple of non-negative eigenvalues satisfying trace condition $\text{Tr}[a] + \text{Tr}[b] = \text{Tr}[c]$. If for all Young diagrams γ of size $|\gamma| = k$,*

$$\varphi_\gamma(c) \leq \sum_{\substack{\alpha, \beta \\ c_{\alpha\beta}^\gamma > 0}} \varphi_\alpha(a) \varphi_\beta(b), \quad (137)$$

then

$$D(a, b, c) \leq \text{Tr}[c] O \left(n \sqrt{\frac{\log k}{k}} \right). \quad (138)$$

where $D(a, b, c) := \min_{(A, B) \in \mathcal{O}_a \times \mathcal{O}_b} \|\text{eig}(A + B) - c\|_1$.

To begin we use the random matrix identities from Section 4.2 to prove that $\varphi_\gamma(A + B)$ does not differ too much when A, B are replaced with other matrices in their respective coadjoint orbits.

lem:random_tri_ineq

Lemma 54. *For a given γ , let $m_n(\gamma)$ be the following quantity*

$$m_n(\gamma) := \min_{\substack{\alpha, \beta \\ c_{\alpha\beta}^\gamma > 0}} \frac{c_{\alpha\beta}^\gamma \dim V_\gamma^n}{\dim V_\alpha^n \otimes V_\beta^n} \quad (139)$$

Then for all non-negative eigenvalues a and b ,

$$\sum_{\substack{\alpha, \beta \\ c_{\alpha\beta}^\gamma > 0}} \varphi_\alpha(A) \varphi_\beta(B) \leq \frac{1}{m_n(\gamma)} \max_{(X, Y) \sim \mathcal{O}_a \times \mathcal{O}_b} \varphi_\gamma(X + Y). \quad (140)$$

Proof. The result follows from Corollary 51 and the fact that the expectation value of $\varphi_\gamma(X + Y)$ over $(X, Y) \sim \mathcal{O}_a \times \mathcal{O}_b$ can be upper bounded by its maximum value. $\square$

Before continuing to the proof of Proposition 53, we remark that the prefactor $\frac{c_{\alpha\beta}^\gamma \dim V_\gamma^n}{\dim V_\alpha^n \otimes V_\beta^n}$ above can be viewed as a probability distribution over Young diagrams γ of size $|\gamma| = |\alpha| + |\beta|$ when conditioned on fixed α, β because it corresponds to the proportion of $V_\alpha^n \otimes V_\beta^n$ occupied by the γ -isotypic subspace. From this point of view, the quantity $m_n(\gamma)$ can be interpreted as the minimum likelihood for the outcome γ as one varies over all pairs of Young diagrams (α, β) of sizes fixed by γ .

Proof of Proposition 53. By Lemma 54, we conclude that for all $\gamma \vdash k$, (a, b, c) satisfies

$$\varphi_\gamma(c) \leq \frac{1}{m_n(\gamma)} \max_{(X,Y) \in \mathcal{O}_a \times \mathcal{O}_b} \varphi_\gamma(X+Y). \quad (141)$$

Since it was assumed that $\text{Tr}[c] = \text{Tr}[a] + \text{Tr}[b] = \text{Tr}(X)$, an application of the eigenvalue separation lemma produces:

$$\varphi_\gamma(c) = \sqrt{\varphi_\gamma(c)} \sqrt{\varphi_\gamma(c)}, \quad (142)$$

$$\leq \sqrt{\frac{1}{m_n(\gamma)} \max_{(X,Y) \in \mathcal{O}_a \times \mathcal{O}_b} \varphi_\gamma(c) \varphi_\gamma(X+Y)}, \quad (143)$$

$$\leq \frac{\text{Tr}[c]^k}{k!} \sqrt{\frac{(\dim V_\gamma^n)^2}{m_n(\gamma)} \exp\left(-\frac{k}{4\text{Tr}[c]^2} D(a, b, c)^2\right)}. \quad \text{\textcolor{blue}{eq:eig_sep_lr}} \quad (144)$$

Furthermore, note that $|\alpha|, |\beta|, |\gamma| \leq k$ so we also have the γ -independent bound

$$\frac{(\dim V_\gamma^n)^2}{m_n(\gamma)} = \max_{\substack{\alpha, \beta \\ c_{\alpha\beta}^\gamma > 0}} \frac{\dim V_\alpha^n \otimes V_\beta^n \otimes V_\gamma^n}{c_{\alpha\beta}^\gamma} \leq \left((k+1)^{n(n-1)/2}\right)^3. \quad (145)$$

After summing everything over $\gamma \vdash k$ and dividing both sides of Eq. (144) by $\frac{\text{Tr}[c]^k}{k!}$ (and noting there are at most $(k+1)^{n-1}$ summands) yields the claimed bound

$$\frac{1}{\text{Tr}[c]} D(a, b, c) \leq \sqrt{(3n^2 + n - 4) \frac{\log(k+1)}{k}}. \quad (146)$$

□

4.4 Distance for the tetrahedral Horn problem

\text{\textcolor{blue}{sec:dist_bound_for_tet_horn}}

The purpose of this section is to prove our main distance bound, Theorem 2. Note that Theorem 2 then immediately implies the “if” portion of the claim of Theorem 1 by taking $k \rightarrow \infty$. The “only if” portion of Theorem 1 was proven in Corollary 37.

Proof of Theorem 2. If a sextuple of eigenvalues (a, b, c, d, e, f) happens to satisfy Eq. (2) for a particular triple of Young diagrams (α, β, δ) , then multiplying both sides of Eq. (2) by $\varphi_\alpha(a)^{\frac{4}{3}} \varphi_\beta(b) \varphi_\delta(d)^{\frac{4}{3}}$ yields

$$(\varphi_\alpha(a) \varphi_\beta(b) \varphi_\delta(d))^2 \leq \sum_{\gamma, \epsilon, \phi} \left\| \left\{ \begin{smallmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{smallmatrix} \right\} \right\|_\infty \varphi_\alpha(a)^{\frac{4}{3}} \varphi_\beta(b) \varphi_\delta(d)^{\frac{4}{3}} \varphi_\phi(f)^{\frac{1}{3}} \varphi_\gamma(c)^{\frac{1}{3}} \varphi_\epsilon(e)^{\frac{1}{3}}. \quad \text{\textcolor{blue}{eq:abd_multiplied}} \quad (147)$$

Our objective is to sum this expression over all triples (α, β, δ) such that $|\alpha| + |\beta| + |\delta| = k$. Note that summing over $\varphi_\alpha(a) \varphi_\beta(b) \varphi_\delta(d)$ yields

$$\sum_{\substack{\alpha, \beta, \delta \\ |\alpha| + |\beta| + |\delta| = k}} \varphi_\alpha(a) \varphi_\beta(b) \varphi_\delta(d) = \frac{\text{Tr}[e]^k}{k!} \quad (148)$$

since we explicitly assumed the trace conditions held, i.e., $\text{Tr}[e] = \text{Tr}[a] + \text{Tr}[b] + \text{Tr}[d]$.

For convenience, let $W_k \left(\begin{smallmatrix} a & b & c \\ d & e & f \end{smallmatrix} \right)$ denote the corresponding sum of the right-hand side (appropriately normalized by $\frac{\text{Tr}[e]^k}{k!}$ so that W_k is invariant under rescaling):

$$W_k \left(\begin{smallmatrix} a & b & c \\ d & e & f \end{smallmatrix} \right) := \left(\frac{k!}{\text{Tr}[e]^k} \right)^2 \sum_{\substack{\alpha, \beta, \delta \\ \gamma, \epsilon, \phi \\ |\epsilon| = k}} \left\| \left\{ \begin{smallmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{smallmatrix} \right\} \right\|_\infty \varphi_\alpha(a)^{\frac{4}{3}} \varphi_\beta(b) \varphi_\delta(d)^{\frac{4}{3}} \varphi_\phi(f)^{\frac{1}{3}} \varphi_\gamma(c)^{\frac{1}{3}} \varphi_\epsilon(e)^{\frac{1}{3}}. \quad (149)$$

The bound in Eq. (147) then implies a moment inequality bound:

$$W_k \begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix} \geq \frac{1}{(k+1)^{3(n-1)}}, \quad \text{\texttt{eq:Wk_lower}} \quad (150)$$

where $(k+1)^{3(n-1)}$ is an upper bound on the number of triples (α, β, δ) such that $\varphi_\alpha(a)\varphi_\beta(b)\varphi_\delta(d) \neq 0$ as there are at most $(k+1)^{\ell-1}$ Young diagrams with at most k boxes and at most ℓ rows.

Our objective now is to derive a complementary upper bound on $W_k \begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix}$. To accomplish this, we note that an immediate corollary of Corollary 52 is the following bound which holds for any triple (x, y, z) of positive semidefinite eigenvalues:

$$\left\| \begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix} \right\|_2^2 \varphi_\alpha(x)\varphi_\beta(y)\varphi_\delta(z) \leq R_\epsilon^{\alpha\beta\delta} \max_{(X,Y,Z) \in \mathcal{O}_x \times \mathcal{O}_y \times \mathcal{O}_z} \left| \varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) \right|, \quad (151)$$

where $R_\epsilon^{\alpha\beta\delta} = \frac{\dim(V_\alpha^n \otimes V_\beta^n \otimes V_\delta^n)}{\dim(V_\epsilon^n)}$ and the where the maximum is taken over all triples (X, Y, Z) of positive semidefinite matrices with eigenvalues (x, y, z) . Applying Lemma 35 and Proposition 36 to $\left| \varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) \right|$ on the right-hand side above produces

$$\left| \varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z) \right| \leq \left\| \begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix} \right\|_\infty \varphi_\phi(Y+Z)^{\frac{1}{3}} \varphi_\gamma(X+Y)^{\frac{1}{3}} \varphi_\epsilon(X+Y+Z)^{\frac{1}{3}}. \quad \text{\texttt{eq:sup_bound}} \quad (152)$$

Then Eq. (152) can be applied to each summand in $W_k \begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix}$ (replacing (x, y, z) with (a, b, d)) to obtain

$$W_k \begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix} \leq \left(\frac{k!}{\text{Tr}[e]^k} \right)^2 \sum_{\substack{\alpha, \beta, \delta \\ \gamma, \epsilon, \phi \\ |\epsilon|=k}} R_\epsilon^{\alpha\beta\delta} \varphi_\alpha(a)^{\frac{2}{3}} \varphi_\delta(d)^{\frac{2}{3}} \max_{(X,Y,Z) \in \mathcal{O}_a \times \mathcal{O}_b \times \mathcal{O}_d} [\varphi_\phi(f)\varphi_\phi(Y+Z)\varphi_\gamma(c)\varphi_\gamma(X+Y)\varphi_\epsilon(e)\varphi_\epsilon(X+Y+Z)]^{\frac{1}{3}}. \quad \text{\texttt{eq:Wk_sum}} \quad (153)$$

Note that here we have made use of the inequality $\left\| \begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix} \right\|_2^2 \geq \left\| \begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix} \right\|_\infty^2$ to simplify the presentation slightly. Now the eigenvalue separation lemma implies a triple of bounds (because $|\alpha|+|\phi| = |\gamma|+|\delta| = |\epsilon| = k$)

$$\varphi_\alpha(a) \sqrt{\varphi_\phi(f)\varphi_\phi(Y+Z)} \leq \dim(V_\alpha^n \otimes V_\phi^n) \frac{\text{Tr}[e]^k}{k!} \exp \left(-\frac{k}{4 \text{Tr}[e]^2} \|f - \text{eig}(Y+Z)\|_1^2 \right), \quad (154)$$

$$\varphi_\delta(d) \sqrt{\varphi_\gamma(c)\varphi_\gamma(X+Y)} \leq \dim(V_\gamma^n \otimes V_\delta^n) \frac{\text{Tr}[e]^k}{k!} \exp \left(-\frac{k}{4 \text{Tr}[e]^2} \|c - \text{eig}(X+Y)\|_1^2 \right), \quad (155)$$

$$\sqrt{\varphi_\epsilon(e)\varphi_\epsilon(X+Y+Z)} \leq \dim(V_\epsilon^n) \frac{\text{Tr}[e]^k}{k!} \exp \left(-\frac{k}{4 \text{Tr}[e]^2} \|e - \text{eig}(X+Y+Z)\|_1^2 \right). \quad (156)$$

Taking the product of these three inequalities to the power of $2/3$, applying it to the upper bound on W_k and combining it with the lower bound on W_k then implies

$$\frac{1}{\text{Tr}[e]^2} D \begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix}^2 \leq \frac{6}{k} (3n(n-1) + 9(n-1)) \log(k+1). \quad (157)$$

where the deviation measure $D \begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix}$ was defined in Definition 9, where the $6(n-1)$ factor comes from the upper bound on the number of summands in Eq. (153), an additional factor of $3(n-1)$ comes from the lower bound on W_k in Eq. (150) and where the $3n(n-1)$ factor comes from an upper bound to product of dimensions:

$$\frac{\dim(V_\alpha^n \otimes V_\beta^n \otimes V_\delta^n)}{\dim(V_\epsilon^n)} \dim(V_\alpha^n \otimes V_\phi^n \otimes V_\gamma^n \otimes V_\delta^n \otimes V_\epsilon^n)^{\frac{2}{3}} \leq \left[(k+1)^{\frac{n(n-1)}{2}} \right]^{3+4\frac{2}{3}} \leq (k+1)^{3n(n-1)}. \quad (158)$$

Therefore, the deviation measure is bounded by

$$D \begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix} \leq \text{Tr}[e] \sqrt{(18n^2 + 36n - 54) \frac{\log(k+1)}{k}} \quad (159)$$

which proves the claim. $\square$

4.5 Asymptotics of the $6j$ symbols

In order to prove Corollary 4 it will be useful to have a bound on the quantity $|\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)|$ which directly relates the sextuple of Young diagrams $(\alpha, \beta, \gamma, \delta, \epsilon, \phi)$ to the eigenvalues of $(X, Y, X+Y, Z, X+Y+Z, Y+Z)$. [lem:asym_size_helper](#)

Lemma 55. *To begin let $X, Y, Z \geq 0$ be positive semidefinite $n \times n$ matrices and let (a, b, c, d, e, f) be the eigenvalues of the matrices $(X, Y, X+Y, Z, X+Y+Z, Y+Z)$. Then for all sextuples of Young diagrams, $(\alpha, \beta, \gamma, \delta, \epsilon, \phi)$, with sizes satisfying*

$$k = |\epsilon| = |\alpha| + |\beta| + |\delta| = |\alpha| + |\phi| = |\gamma| + |\delta|, \quad \text{\small \a{eq:yd_k_sizes}} \quad (160)$$

we have (assuming $\text{Tr}[e] \neq 0$)

$$\frac{k!}{\text{Tr}[e]^k} |\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)| \leq \min\left\{\left\|\left\{\frac{\alpha}{\delta} \frac{\beta}{\epsilon} \frac{\gamma}{\phi}\right\}\right\|_{\infty}, (k+1)^{n(n-1)} \exp\left(-\frac{k}{4} R_{abcdef}^{\alpha\beta\gamma\delta\epsilon\phi}\right)\right\}, \quad (161)$$

where

$$R_{abcdef}^{\alpha\beta\gamma\delta\epsilon\phi} = D\left(\frac{\alpha \oplus \beta \oplus \delta}{k} \left\| \frac{a \oplus b \oplus d}{\text{Tr}[e]}\right\|\right) + D\left(\frac{\gamma \oplus \delta}{k} \left\| \frac{c \oplus d}{\text{Tr}[e]}\right\|\right) + D\left(\frac{\alpha \oplus \phi}{k} \left\| \frac{a \oplus f}{\text{Tr}[e]}\right\|\right) + D\left(\frac{\epsilon}{k} \left\| \frac{e}{\text{Tr}[e]}\right\|\right). \quad (162)$$

Proof. A combination of Lemma 35 and Proposition 36 implies

$$|\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(X, Y, Z)| \leq \left\|\left\{\frac{\alpha}{\delta} \frac{\beta}{\epsilon} \frac{\gamma}{\phi}\right\}\right\|_{\infty} (\varphi_{\alpha}(a)\varphi_{\beta}(b)\varphi_{\delta}(d) \times \varphi_{\gamma}(c)\varphi_{\delta}(d) \times \varphi_{\epsilon}(e) \times \varphi_{\alpha}(a)\varphi_{\phi}(f))^{\frac{1}{4}}, \quad (163)$$

Additionally, the relative entropy bounds arising from the multinomial distribution places the following bounds on four probability distributions:

$$\frac{k!}{\text{Tr}[e]^k} \varphi_{\alpha}(a)\varphi_{\beta}(b)\varphi_{\delta}(d) \leq \dim(V_{\alpha}^n \otimes V_{\beta}^n \otimes V_{\delta}^n) \exp\left[-kD\left(\frac{\alpha \oplus \beta \oplus \delta}{k} \left\| \frac{a \oplus b \oplus d}{\text{Tr}[e]}\right\|\right)\right], \quad (164)$$

$$\frac{k!}{\text{Tr}[e]^k} \varphi_{\gamma}(c)\varphi_{\delta}(d) \leq \dim(V_{\gamma}^n \otimes V_{\delta}^n) \exp\left[-kD\left(\frac{\gamma \oplus \delta}{k} \left\| \frac{c \oplus d}{\text{Tr}[e]}\right\|\right)\right], \quad (165)$$

$$\frac{k!}{\text{Tr}[e]^k} \varphi_{\alpha}(a)\varphi_{\phi}(f) \leq \dim(V_{\alpha}^n \otimes V_{\phi}^n) \exp\left[-kD\left(\frac{\alpha \oplus \phi}{k} \left\| \frac{a \oplus f}{\text{Tr}[e]}\right\|\right)\right], \quad (166)$$

$$\frac{k!}{\text{Tr}[e]^k} \varphi_{\epsilon}(e) \leq \dim V_{\epsilon}^n \exp\left[-kD\left(\frac{\epsilon}{k} \left\| \frac{e}{\text{Tr}[e]}\right\|\right)\right]. \quad (167)$$

Since the product of the four distributions on the left-hand side above is at most the minimum of one or the product of relative entropy bounds on the right-hand side above (and the product of the dimensions to the fourth root is at most $[(k+1)^{\frac{n(n-1)}{2}}]^{\frac{8}{4}}$), obtain the claim. $\square$

Proof of Corollary 4. To begin we use Lemma 55 to prove the claimed inverse polynomial lower bound. Assume (a, b, c, d, e, f) are the eigenvalues of positive semidefinite matrices (A, B, C, D, E, F) satisfying the tetrahedral constraints. Summing $\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(A, B, D)$ over all sextuples of Young diagrams satisfying Eq. (160) produces

$$\sum_{\substack{\alpha, \beta, \delta \\ \gamma, \epsilon, \phi \\ |\epsilon|=k}} \varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(A, B, D) = \frac{\text{Tr}[e]^k}{k!}. \quad (168)$$

Therefore, the sextuple which maximizes the absolute value of $\varphi_{\gamma\epsilon\phi}^{\alpha\beta\delta}(A, B, D)$ over this k -dependent domain of summation, which we denote by $(\alpha_k, \beta_k, \gamma_k, \delta_k, \epsilon_k, \phi_k)$, must satisfy

$$\frac{k!}{\text{Tr}[e]^k} |\varphi_{\gamma_k \epsilon_k \phi_k}^{\alpha_k \beta_k \delta_k}(A, B, D)| \geq \frac{1}{(k+1)^{6(\text{rank}[e]-1)}} \quad (169)$$

where $(k+1)^{6(\text{rank}[e]-1)}$ is a crude upper bound on the number of terms in the above summation which are non-zero as there are at most $(k+1)^{r-1}$ Young diagrams of size at most k and depth at most r (note

that each of $\varphi_\alpha(a), \varphi_\beta(b), \varphi_\delta(d), \varphi_\phi(f), \varphi_\gamma(c)$ and $\varphi_\epsilon(e)$ are respectively only non-zero if the depth of the corresponding Young diagram is at most the rank of the corresponding set of eigenvalues and $\text{rank}[e]$ is always the maximum possible rank). Therefore, Lemma 55 immediately implies both

$$\left\| \begin{Bmatrix} \alpha_k & \beta_k & \gamma_k \\ \delta_k & \epsilon_k & \phi_k \end{Bmatrix} \right\|_\infty \geq \frac{1}{(k+1)^{6(\text{rank}[e]-1)}}, \quad \{\text{eq:n_independent_inverse_poly}\} \quad (170)$$

as well as

$$\exp\left(-\frac{k}{4} R_{abcdef}^{\alpha_k \beta_k \gamma_k \delta_k \epsilon_k \phi_k}\right) \geq \frac{1}{(k+1)^{n(n-1)}} \frac{1}{(k+1)^{6(\text{rank}[e]-1)}}. \quad (171)$$

Thus we have demonstrated the existence of a sequence of sextuples of Young diagrams for which (i) $\left\| \begin{Bmatrix} \alpha_k & \beta_k & \gamma_k \\ \delta_k & \epsilon_k & \phi_k \end{Bmatrix} \right\|_\infty$ satisfies the claimed inverse polynomial lower bound (which is actually independent of n), and (ii) $R_{abcdef}^{\alpha_k \beta_k \gamma_k \delta_k \epsilon_k \phi_k} \rightarrow 0$ which implies the convergence claim:

$$\frac{\alpha_k}{k} \rightarrow \frac{a}{\text{Tr}[e]}, \quad \frac{\beta_k}{k} \rightarrow \frac{b}{\text{Tr}[e]}, \quad \frac{\gamma_k}{k} \rightarrow \frac{c}{\text{Tr}[e]}, \quad \frac{\delta_k}{k} \rightarrow \frac{d}{\text{Tr}[e]}, \quad \frac{\epsilon_k}{k} \rightarrow \frac{e}{\text{Tr}[e]}, \quad \frac{\phi_k}{k} \rightarrow \frac{f}{\text{Tr}[e]}. \quad (172)$$

To prove the exponential decay on $\left\| \begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix} \right\|_\infty$ the case where (a, b, c, d, e, f) are not tetrahedral, it will be slightly easier to prove the exponential decay of $\left\| \begin{Bmatrix} \alpha & \beta & \gamma \\ \delta & \epsilon & \phi \end{Bmatrix} \right\|_2$ which is actually a bit stronger than necessary. Let (a, b, c, d, e, f) be the set of non-tetrahedral eigenvalues to which the sequence of Young diagrams $(\alpha_k, \beta_k, \gamma_k, \delta_k, \epsilon_k, \phi_k)$ is converging in proportion. We can henceforth assume the Young diagrams satisfy the size constraints Eq. (160) as before. Then from Corollary 52, we conclude

$$\left\| \begin{Bmatrix} \alpha_k & \beta_k & \gamma_k \\ \delta_k & \epsilon_k & \phi_k \end{Bmatrix} \right\|_2^2 \leq \max_{\substack{X, Y, Z \\ \text{Tr}(X)=\text{Tr}[a] \\ \text{Tr}(Y)=\text{Tr}[b] \\ \text{Tr}(Z)=\text{Tr}[d]}} (k+1)^{\frac{3n^2(n^2-1)}{2}} \frac{|\alpha_k|! |\beta_k|! |\delta_k|!}{\text{Tr}[a]^{|\alpha_k|} \text{Tr}[b]^{|\beta_k|} \text{Tr}[d]^{|\delta_k|}} \left| \varphi_{\gamma_k \epsilon_k \phi_k}^{\alpha_k \beta_k \delta_k}(X, Y, Z) \right|. \quad (173)$$

Now because $(\alpha_k, \beta_k, \gamma_k, \delta_k, \epsilon_k, \phi_k)$ are converging in proportion to the eigenvalues (a, b, c, d, e, f) we can assume that the sizes are such that the following multinomial distribution bound holds:

$$\frac{|\epsilon_k|!}{\text{Tr}[e]^{|\epsilon_k|}} \frac{\text{Tr}[a]^{|\alpha_k|} \text{Tr}[b]^{|\beta_k|} \text{Tr}[d]^{|\delta_k|}}{|\alpha_k|! |\beta_k|! |\delta_k|!} \geq \frac{1}{\text{poly}(k)}. \quad (174)$$

Using Lemma 55 then gives

$$\left\| \begin{Bmatrix} \alpha_k & \beta_k & \gamma_k \\ \delta_k & \epsilon_k & \phi_k \end{Bmatrix} \right\|_2^2 \leq \text{poly}(k) \exp\left(-\frac{k}{4} \min R_{a'b'c'd'e'f'}^{\alpha_k \beta_k \gamma_k \delta_k \epsilon_k \phi_k}\right). \quad (175)$$

where the minimum is taken over all tetrahedral eigenvalues (a', b', c', d', e', f') with matching traces. However, since $(\alpha_k, \beta_k, \gamma_k, \delta_k, \epsilon_k, \phi_k)$ are converging to non-tetrahedral eigenvalues (a, b, c, d, e, f) , there must exist a sufficiently large K such that for all $k \geq K$, $\min R_{a'b'c'd'e'f'}^{\alpha_k \beta_k \gamma_k \delta_k \epsilon_k \phi_k} \geq \delta$ for some positive constant $\delta > 0$ which proves the claim. $\square$

Acknowledgments

MC and TCF acknowledge financial support from the European Research Council (ERC Grant Agreement No. 818761), VILLUM FONDEN via the QMATH Centre of Excellence (Grant No. 10059) and the Novo Nordisk Foundation (grant NNF20OC0059939 ‘Quantum for Life’). They also thank the National Center for Competence in Research SwissMAP of the Swiss National Science Foundation and the Section of Mathematics at the University of Geneva for their hospitality. TCF would also like to thank Suvrit Sra, Harold Nieuwboer, Marco Fanizza and Jack Davis for helpful discussions.

A Review of previous work of eigenvalues of sums

sec:review

The purpose of this section is to describe various previously known inequalities which are necessarily satisfied by tetrahedral eigenvalues (a, b, c, d, e, f) . Readers exclusively interested in the techniques used to prove the results claimed in the introduction are encouraged to skip to Section 3. While our main theorems, Theorems 1 and 2, describe a countably infinite family of necessary inequalities which, when bundled together, are sufficient to characterize the set of tetrahedral eigenvalues for any fixed dimension n , the problem of providing a finite characterization, say in the form of a finite list of polynomial equalities and inequalities, remains an open problem for $n \geq 3$.

A.1 Horn's inequalities

Recall that (a, b, c, d, e, f) are tetrahedral only if each of (a, b, c) , (b, c, f) , (c, d, e) , and (a, f, e) are trigonal as in Eq. (14). Consequently, every inequality constraining trigonal eigenvalues can be lifted to produce a quadruplet of inequalities constraining the set of tetrahedral eigenvalues. These inequalities can be understood as generalizations of the triangle inequalities, Eq. (26), which hold for case where $n = 2$. Therefore, before proceeding, we briefly summarize the inequalities satisfied by trigonal eigenvalues (a, b, c) . Of particular interest are Horn's inequalities [Hor62] which are known to be necessary and sufficient [Kly98; KT99]. For a more complete historical account, the reader is directed to the survey articles [Bha01; Ful98; Knu00] or [NP06, Appendix D]. In addition to the aforementioned trace condition, $\sum_{i=1}^n a_i + \sum_{j=1}^n b_j = \sum_{k=1}^n c_k$, one always has the following pair of elementary inequalities,

$$a_1 + b_1 \geq c_1, \quad \text{and} \quad a_n + b_n \leq c_n, \quad (176)$$

which constrain the largest and smallest eigenvalues respectively (note that under inversion symmetry, these inequalities are equivalent). The first somewhat non-trivial inequality, derived by Weyl in 1912, is now known as *Weyl's inequality* and states for all i, j such that $i + j \leq n + 1$ [Wey12]⁵

$$a_i + b_j \geq c_{i+j-1}. \quad (177)$$

In 1949, Fan derived a majorization inequality constraining the first r largest eigenvalues [Fan49], and shortly thereafter Lidskii [Lid53] & Wielandt [Wie55] found proofs of another family of inequalities. Finally, in 1962, Horn determined a complete collection of inequalities for (a, b, c) are trigonal for the case where $n = 3$ [Hor62]. Moreover, also in [Hor62], Horn provided an inductively defined set of subsets $I, J, K \subset \{1, \dots, n\}$ of equal cardinality and conjectured that inequalities of the form

$$\sum_{i \in I} a_i + \sum_{j \in J} b_j \geq \sum_{k \in K} c_k, \quad \text{\texttt{eq:IJK_ineq}} \quad (178)$$

were both necessary and sufficient to ensure that (a, b, c) are trigonal. While proofs for some of these inequalities were found (see e.g., [TF71; TF70]), it took until the late 1990s for Horn's problem to be to be solved by Klyachko [Kly98] who proved that (a, b, c) if and only if in addition to the trace equality, $\text{Tr}[a] + \text{Tr}[b] = \text{Tr}[c]$, Eq. (178) holds for all (I, J, K) such that (i) $|I| = |J| = |K| = p$ for some p , and (ii) the Littlewood-Richardson coefficient $c_{\alpha, \beta}^\gamma$ is non-zero (here $\alpha = (i_p - p, \dots, i_1 - 1)$, $\beta = (j_p - p, \dots, j_1 - 1)$, and $\gamma = (k_p - p, \dots, k_1 - 1)$). Shortly thereafter, Knutson and Tao proved the saturation conjecture for Littlewood-Richardson coefficients [KT99], which when combined with Klyachko's solution, proved Horn's conjecture.

For the purposes of characterizing the set of tetrahedral eigenvalues further, we will henceforth be exclusively concerned with inequalities constraining the set of tetrahedral eigenvalues which are not already implied by Horn's inequalities.

⁵For 2×2 matrices one can see that these necessary inequalities are also sufficient to ensure that (a, b, c) are trigonal; upon shifting to traceless eigenvalues using Corollary 16, $a = (a, -a)$, $b = (b, -b)$ and $c = (c, -c)$, and these inequalities become equivalent to the triangle inequalities $c \leq a + b$ and $c \geq \max\{b - a, a - b\}$.

A.2 A simple quadratic inequality

In Section 2.4 we saw that the shifting symmetry implies, in a certain sense, one may assume without loss of generality that the candidate eigenvalues (a, b, c, d, e, f) are all non-negative or even strictly positive. While almost all of the following inequalities require this positivity assumption, we note that they can be extended to eigenvalues with negative entries in all cases.

Here we present an quadratic inequality which follows from a simple calculation. Let (A, B, C, D, E, F) be positive semidefinite $n \times n$ matrices satisfying $A + B = C$, $B + D = F$, and $A + F = C + D = E$. Then the following identity holds:

$$\begin{aligned}
& \text{Tr}[B^2] + \text{Tr}[E^2] - \text{Tr}[C^2] - \text{Tr}[F^2] \\
&= \text{Tr}[(B)^2] + \text{Tr}[(A + B + D)^2] - \text{Tr}[(A + B)^2] - \text{Tr}[(B + D)^2] \\
&= (\text{Tr}[B^2]) + (\text{Tr}[A^2] + \text{Tr}[B^2] + \text{Tr}[D^2] + 2 \text{Tr}[AB] + 2 \text{Tr}[BD] + 2 \text{Tr}[AD]) \\
&\quad - (\text{Tr}[A^2] + \text{Tr}[B^2] + 2 \text{Tr}[AB]) - (\text{Tr}[B^2] + \text{Tr}[D^2] + 2 \text{Tr}[BD]) \\
&= 2 \text{Tr}[AD].
\end{aligned} \tag{179}$$

Although the left-hand side of the above expression depends only on the set of eigenvalues (a, b, c, d, e, f) ,

$$\text{Tr}[B^2] + \text{Tr}[E^2] - \text{Tr}[C^2] - \text{Tr}[F^2] = \|b\|_2^2 + \|e\|_2^2 - \|c\|_2^2 - \|f\|_2^2, \tag{180}$$

the right-hand side does not. Nevertheless, one may note that for all $A, D \geq 0$ positive semidefinite, $\text{Tr}(AD) \geq 0$ and therefore,

$$\|b\|_2^2 + \|e\|_2^2 - \|c\|_2^2 - \|f\|_2^2 \geq 0. \tag{181}$$

This inequality is well-known and indeed follows as a special case of [Sag13, Proposition 2.3].

However, a minor improvement can be made here. While $\text{Tr}(AD)$ cannot, in general, be determined from the eigenvalues $a = \text{eig}(A)$ and $d = \text{eig}(D)$, one can apply the von Neumann trace inequalities ([MOA10, H.1.g & H.1.h] or [Bau+01, Proposition 4.4]) to bound the value of $\text{Tr}(AD)$ to obtain

$$\sum_j a_j d_{n-j+1} \leq \text{Tr}(AD) \leq \sum_j a_j d_j. \tag{182}$$

Therefore, we conclude

$$2 \sum_{j=1}^n a_j d_{n-j+1} \leq \|b\|_2^2 + \|e\|_2^2 - \|c\|_2^2 - \|f\|_2^2 \leq 2 \sum_{j=1}^n a_j d_j. \tag{183}$$

A.3 Ptolemy & Cayley-Menger inequalities

In this section, we prove inequalities constraining the 2-norms $\|\cdot\|_2$ of a tetrahedral sextuple of eigenvalues (a, b, c, d, e, f) associated to the Hermitian matrices (A, B, C, D, E, F) . These inequalities correspond to Ptolemy's inequality and the Cayley-Menger determinant inequality for the edge-lengths of a Euclidean tetrahedron in $\mathbb{R}^3$. These results rely on the observation that the real vector space of $n \times n$ Hermitian matrices equipped with the Hilbert-Schmidt inner product $\langle X, Y \rangle = \text{Tr}(X^\dagger Y)$ has norm $\|X\|_2 = \sqrt{\langle X, X \rangle}$ which is a function of the *eigenvalues* of X . Specifically, for any Hermitian matrix X with eigenvalues x , we have $\|X\|_2 = \|x\|_2 = \left(\sum_{j=1}^n x_j^2\right)^{\frac{1}{2}}$.

Recall that the four vertices of the tetrahedron of Hermitian matrices in Fig. 1 are

$$V_0 = 0, \quad V_1 = A, \quad V_2 = A + B = C, \quad V_3 = A + B + D = E. \tag{184}$$

Ptolemy's inequality is an inequality which holds for any four points V_0, V_1, V_2, V_3 in a normed space.⁶ It reads

$$\|V_3 - V_0\| \|V_1 - V_2\| \leq \|V_2 - V_0\| \|V_1 - V_3\| + \|V_1 - V_0\| \|V_2 - V_3\|. \tag{185}$$

⁶It is known that every real inner product space with norm $\|X\| = \sqrt{\langle X, X \rangle}$ is Ptolemaic in the sense that Ptolemy's inequality holds. A converse of this result also holds due to Schoenberg; namely, every real semi-normed space satisfying Ptolemy's inequality comes from a real inner product space [Sch52].

Applied to the vertices of the tetrahedron of Hermitian matrices with respect to the Hilbert-Schmidt norm $\|\cdot\|_2$ we obtain

$$\|E\|_2 \|B\|_2 \leq \|C\|_2 \|F\|_2 + \|A\|_2 \|D\|_2. \quad \text{\textcolor{blue}{(eq:abcdef_ptolemy)}} \quad (186)$$

More generally, note that the four Hermitian matrices $(V_0, V_1, V_2, V_3) = (0, A, C, E)$ necessarily lie inside a three-dimensional subspace of the set of $n \times n$ Hermitian matrices. Therefore, just as in the case of 2×2 matrices (Example 10) one can compute the square of its three-dimensional volume using the Cayley-Menger determinant to obtain the inequality:

$$\text{CM}(\|A\|, \|B\|, \|C\|, \|D\|, \|E\|, \|F\|) = \frac{1}{288} \det \begin{pmatrix} 0 & \|A\|^2 & \|C\|^2 & \|E\|^2 & 1 \\ \|A\|^2 & 0 & \|B\|^2 & \|F\|^2 & 1 \\ \|C\|^2 & \|B\|^2 & 0 & \|D\|^2 & 1 \\ \|E\|^2 & \|F\|^2 & \|D\|^2 & 0 & 1 \\ 1 & 1 & 1 & 1 & 0 \end{pmatrix} \geq 0 \quad (187)$$

A.4 Strongly subadditive/Submodular inequalities

Another strategy for understanding eigenvalues of positive semidefinite matrices and their sums is to consider the expansion of tensors powers of sums of matrices in relation to the tensor powers of their summands. For instance, given a positive integer k and positive semidefinite X and Y , using the non-commutative binomial expansion of $(X+Y)^{\otimes k}$ and dropping all summands involving both X and Y , one obtains [BS15, Proposition 1.1]:

$$(X+Y)^{\otimes k} \geq X^{\otimes k} + Y^{\otimes k}. \quad (188)$$

This technique, was generalized by [TCL11] [TCL11, Lemma 2.2] to obtain the following result.⁷

\text{\textcolor{blue}{prop:operator_ineq}}

Proposition 56. *Let $X, Y, Z \geq 0$ be positive semidefinite matrices and let k be a positive integer. Then*

$$(X+Y+Z)^{\otimes k} + Y^{\otimes k} \geq (X+Y)^{\otimes k} + (Y+Z)^{\otimes k}. \quad (189)$$

From Proposition 56 one can obtain an inequalities pertaining only to the eigenvalues of $Y, X+Y, Y+Z$, and $X+Y+Z$ by pairing the operator inequality with any $U(n)$ -equivariant positive semidefinite operator acting on $(\mathbb{C}^n)^{\otimes k}$.⁸

\text{\textcolor{blue}{cor:schur_ineq}}

Corollary 57. *Let $X, Y, Z \geq 0$ be positive semidefinite matrices and let s_λ be the Schur polynomial indexed by the Young diagram $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n)$. Then*

$$s_\lambda(\text{eig}(X+Y+Z)) + s_\lambda(\text{eig}(Y)) \geq s_\lambda(\text{eig}(X+Y)) + s_\lambda(\text{eig}(Y+Z)). \quad \text{\textcolor{blue}{(eq:schur_ineq)}} \quad (190)$$

Proof. Given a Young diagram λ , let P_λ be the Hermitian projection operator onto the isotypic subspace $V_\lambda^n \otimes W_\lambda \subseteq (\mathbb{C}^d)^{\otimes k}$ (see Section 3.1 for more details). Given positive semidefinite matrix $X \geq 0$, one can show that

$$\text{Tr}(P_\lambda X^{\otimes k}) = \text{Tr}(P_\lambda \text{diag}(\text{eig}(X))^{\otimes k}) = \dim(W_\lambda) s_\lambda(\text{eig}(X)) \geq 0. \quad (191)$$

Now we apply the positivity preserving map $\text{Tr}(P_\lambda \cdot)$ to Proposition 56 to obtain

$$\text{Tr}[P_x(X+Y+Z)^{\otimes k}] + \text{Tr}[P_x Y^{\otimes k}] \geq \text{Tr}[P_x(X+Y)^{\otimes k}] + \text{Tr}[P_x(Y+Z)^{\otimes k}] \quad (192)$$

and since λ has at most n rows, we conclude $\dim(W_\lambda) \neq 0$ and thus Eq. (190) holds. $\square$

⁷Further extensions of this technique include inequalities involving the sum $X+Z$ which are outside the scope of the tetrahedral Horn problem (see [Lin14; BS15; NS23])

⁸Paksoy, Turkmen, and Zhang has used a similar approach to prove analogous inequalities concerning so-called generalized matrix functions d_χ^H associated to irreducible characters χ of subgroups H of the symmetric group S_k [MM65; HM21]. Unfortunately, these generalized matrix functions and generally not invariant under unitary conjugation and thus depend on more just eigenvalues; the only exception is the determinant which is already a special case of Corollary 57.

In the special where where $\lambda = \underbrace{(1, \dots, 1)}_n$, one obtains the well-known determinant inequality [TCL11,

Lemma 2.2]:

$$\det(X + Y + Z) + \det(Y) \geq \det(X + Y) + \det(Y + Z). \quad \text{\texttt{eq:lin_det}} \quad (193)$$

Another well-known determinant inequality is the multiplicative variant of Eq. (193) which states that for all $X, Y, Z \geq 0$ positive semidefinite,

$$\det(X + Y + Z) \det(Y) \leq \det(X + Y) \det(Y + Z). \quad \text{\texttt{eq:multip_det}} \quad (194)$$

This inequality follows as a special case of a general result due to Sagnol [Sag13, Proposition 2.3]. \texttt{thm:sagnol_ineq}

Theorem 58. *Let $X, Y, Z \geq 0$ be positive semidefinite matrices. Let f be a real valued and differentiable function over the non-negative real numbers $\mathbb{R}_{\geq 0}$ whose derivative f' is an operator monotone function over the positive real numbers $\mathbb{R}_{> 0}$. Then the following inequality holds*

$$\text{Tr } f(X + Y + Z) + \text{Tr } f(Y) \geq \text{Tr } f(X + Y) + \text{Tr } f(Y + Z). \quad \text{\texttt{eq:sagnol_ineq}} \quad (195)$$

Proof. See [Sag13, Section 2.1]. □

There are numerous inequalities of the form Eq. (195) which immediately follow from Theorem 58. For example, letting $f(x) = \log x$ and noting $\text{Tr}(\log X) = \log \det(X)$, one recovers Eq. (194). Moreover, taking $f(x) = -x^p$ for $p \in [0, 1]$, one obtains [BGS08, Lemma 5.3]. In the limit where $p \rightarrow 0^+$, Sagnol also obtains at first order the well-known rank inequality [Sag13, Proposition A.1]

$$\text{rank}(X + Y) + \text{rank}(Y + Z) \geq \text{rank}(Y) + \text{rank}(X + Y + Z). \quad (196)$$

One can also let $f(x) = x^p$ for $p \in (1, 2]$. Of particular interest to us in Section 3.6 is the function $f(x) = x \log x$ which implies for all $X, Y, Z \geq 0$ the inequality

$$S(X + Y) + S(Y + Z) \geq S(Y) + S(X + Y + Z) \quad \text{\texttt{eq:ssa_like_ineq}} \quad (197)$$

where for a positive semidefinite matrix X with eigenvalues x , $S(X) := s(x) := -\sum_{i=1}^n x_i \log x_i$ is the *unnormalized* von Neumann entropy of X . Note that a similar inequality involving entropies of sums of positive semidefinite matrices (subject to additional eigenvalue constraints) has appeared in [CSW18]. Moreover, while it can be shown that Eq. (197) is strictly tighter than the inequality in [CSW18] for fixed matrices, it is also possible to obtain Eq. (197) from the result in [CSW18] by down-scaling the involved matrices in the sense of Proposition 11.

A.5 Seesaw reciprocity & eigenvalues of blocks

\texttt{sec:blocks_of_psd_matrices}

The purpose of this section is to reformulate both Horn's problem and tetrahedral Horn problem as a instance of a slightly more general kind of eigenvalue problem involving positive semidefinite matrices. Throughout this section, we will restrict our focus to the case of positive semidefinite matrices and their eigenvalues. As a reminder we note that this can be done without loss of generality due to Corollary 17.

Given a positive semidefinite matrix, X , with eigenvalues $\text{eig}(X) = \chi = (\chi_1 \geq \dots \geq \chi_n \geq 0)$ and eigen-decomposition $X = \sum_{i=1}^n \chi_i \Pi_i$, one can define $X^{\frac{1}{2}}$ by $X^{\frac{1}{2}} := \sum_{i=1}^n \chi_i^{\frac{1}{2}} \Pi_i$. Inspired by [Bha97, Theorem IV.2.13], we obtain the following fundamental lemma relating the eigenvalues c of a sum $C = A + B$ of $n \times n$ Hermitian matrices to the eigenvalues of a larger $2n \times 2n$ matrix M built from A and B . \texttt{lem:trigonal_block}

Lemma 59. *Let (A, B, C) be a triple of $n \times n$ Hermitian matrices with eigenvalues (a, b, c) such that $C = A + B$. Let C' be the $2n \times 2n$ Hermitian matrix*

$$C' = \begin{bmatrix} A & A^{\frac{1}{2}} B^{\frac{1}{2}} \\ B^{\frac{1}{2}} A^{\frac{1}{2}} & B \end{bmatrix}. \quad (198)$$

Then $C' \geq 0$ is positive semidefinite, has rank at most $2n$ and has eigenvalues given by c

$$\text{eig}(C') \simeq \text{eig}(A + B) = \text{eig}(C) = c. \quad (199)$$

Proof. That C' is positive semidefinite and has rank at most n follows because C' can be decomposed as $C' = T^*T$ where T is the $n \times 2n$ block matrix

$$T = \begin{bmatrix} A^{\frac{1}{2}} & B^{\frac{1}{2}} \end{bmatrix}. \quad (200)$$

That C' and $A+B$ share the same eigenvalues (up to padding by n zeros) follows because for any rectangular matrix T , the eigenvalues of T^*T and TT^* are equal (up to padding by zeros) and in this instance

$$T^*T = \begin{bmatrix} A^{\frac{1}{2}} \\ B^{\frac{1}{2}} \end{bmatrix} \begin{bmatrix} A^{\frac{1}{2}} & B^{\frac{1}{2}} \end{bmatrix} = \begin{bmatrix} A & A^{\frac{1}{2}}B^{\frac{1}{2}} \\ B^{\frac{1}{2}}A^{\frac{1}{2}} & B \end{bmatrix} = C' \quad TT^* = \begin{bmatrix} A^{\frac{1}{2}} & B^{\frac{1}{2}} \end{bmatrix} \begin{bmatrix} A^{\frac{1}{2}} \\ B^{\frac{1}{2}} \end{bmatrix} = A + B. \quad (201)$$

□

From the perspective of Lemma 59, one can view the Horn's problem involving the eigenvalues (a, b, c) of $n \times n$ Hermitian matrices (A, B, C) satisfying $A + B = C$ as a special case of a more general eigenvalue problem concerned with the relations between the eigenvalues of $2n \times 2n$ Hermitian matrices Σ of the form

$$C' = \begin{bmatrix} A' & Z \\ Z^* & B' \end{bmatrix}, \quad \text{\textcolor{blue}{eq:block_ABX}} \quad (202)$$

and the eigenvalues of its two $n \times n$ diagonal blocks Z and Z . It is known, by [HJ12, Theorem 7.7.9] or more generally [MKX19, Theorem 5.11] that if C' is a 2×2 block matrix such that $A', B' \geq 0$, then $C' \geq 0$ if and only if the off-diagonal block Z is of the form

$$Z = A'^{\frac{1}{2}} K B'^{\frac{1}{2}} \quad (203)$$

for some contraction matrix K (largest eigenvalue of $(K^*K)^{\frac{1}{2}}$ is at most unity). Additionally, when $Z = A'^{\frac{1}{2}} K B'^{\frac{1}{2}}$, one has the decomposition

$$C' = \begin{bmatrix} A' & Z \\ Z^* & B' \end{bmatrix} = \begin{bmatrix} A'^{\frac{1}{2}} & 0 \\ 0 & B'^{\frac{1}{2}} \end{bmatrix} \begin{bmatrix} I & K \\ K^* & I \end{bmatrix} \begin{bmatrix} A'^{\frac{1}{2}} & 0 \\ 0 & B'^{\frac{1}{2}} \end{bmatrix}, \quad (204)$$

noting that K is a contraction if and only if the middle block matrix above is positive semidefinite [MKX19, Lemma 5.10].

Therefore, the special case covered by Lemma 59 corresponds to the case where the corresponding contraction matrix K is the $n \times n$ identity matrix $K = I$ (or more generally any contractive matrix satisfying $A'^{\frac{1}{2}} K B'^{\frac{1}{2}} = A'^{\frac{1}{2}} B'^{\frac{1}{2}}$). A simple observation which distinguishes between eigenvalue problem concerning diagonal blocks and the eigenvalue problem concerning sums is simply that the rank of C' can in general be up to $2n$ whereas the rank of the sum $C = A + B$ is at most n . Despite this additional freedom, there remains non-trivial relations constraining the eigenvalues of C' and of the blocks A' and B' in terms of properties of Z (see [Gum+18; BL22]). Nevertheless, when the rank of C' is constrained to be at most n , we obtain the following theorem which provides an equivalent formulation of Horn's problem.

\text{\textcolor{blue}{thm:trigonal_block}}

Theorem 60. *Let (a, b, c) be a triple of positive semidefinite eigenvalues of length at most n . The following are equivalent:*

- (i) *there exists a triple of Hermitian matrices (A, B, C) with eigenvalues (a, b, c) satisfying $A + B = C$,*
- (ii) *there exists an $2n \times 2n$ Hermitian matrix,*

$$C' = \begin{bmatrix} A' & Z \\ Z^* & B' \end{bmatrix}, \quad (205)$$

such that $\text{eig}(A') = a$, $\text{eig}(B') = b$, and $\text{eig}(C') = c$.

Proof. Since we explicitly assume all of the eigenvalues in the triple (a, b, c) are non-negative, the Hermitian matrices A, B, C , and A', B', C' involved in the statements above are necessarily positive semidefinite. The

claim (i) $\implies$ (ii) follows from Lemma 59. That (ii) $\implies$ (i) follows by noticing that if C' has eigenvalues c , then it has rank at most n . Therefore, as $C' \geq 0$, there exists a $n \times 2n$ block matrix T of the form

$$T = \begin{bmatrix} T_A & T_B \end{bmatrix} \quad (206)$$

such that $C' = T^*T$. The diagonal blocks of C' are therefore $A' = T_A^*T_A$, and $B' = T_B^*T_B$. As in the proof of Lemma 59, we note that T^*T and TT^* have the same eigenvalues up to padding by zeros. Therefore, by letting $A = T_A T_A^*$ and $B = T_B T_B^*$ we see that

$$\begin{aligned} \text{eig}(A) &= \text{eig}(T_A T_A^*) \simeq \text{eig}(T_A^* T_A) = \text{eig}(A') = a, \\ \text{eig}(B) &= \text{eig}(T_B T_B^*) \simeq \text{eig}(T_B^* T_B) = \text{eig}(B') = b, \end{aligned} \quad (207)$$

while the eigenvalues of $C = A + B = T_A T_A^* + T_B T_B^*$ match the eigenvalues of C' because

$$\text{eig}(C) = \text{eig}(A + B) = \text{eig}(T_A T_A^* + T_B T_B^*) = \text{eig}(TT^*) \simeq \text{eig}(T^*T) = \text{eig}(C') = c. \quad (208)$$

□

Here we established a correspondence between (i) the existence of $n \times n$ positive semidefinite matrices (A, B, C) with eigenvalues (a, b, c) satisfying $A + B = C$ and (ii) the positivity of corresponding Littlewood-Richardson coefficients c_{ab}^c which govern the branching problem for the diagonal subgroup $\text{GL}(n) \hookrightarrow \text{GL}(n) \times \text{GL}(n)$ sending X to (X, X) . Recalling Theorem 60, there is a further correspondence to (iii) the existence of a $2n \times 2n$ positive semidefinite matrix Σ with eigenvalues c and $n \times n$ blocks A', B' with eigenvalues a, b . One may wonder how to show the correspondence between (ii) and (iii) *directly* without passing through (i). This can be accomplished by considering the branching multiplicities m_c^{ab} for the blockwise subgroup $\text{GL}(a) \times \text{GL}(b) \hookrightarrow \text{GL}(a+b)$ sending (A, B) to $A \oplus B$. In fact there exists a representation-theoretic analogue of Theorem 60, specifically [GW00, Theorem 9.2.3]. In particular, the values of m_c^{ab} and c_{ab}^c coincide provided the Young diagrams (a, b, c) involved have depths at most $(\min\{a, n\}, \min\{b, n\}, \min\{a+b, n\})$; this ensures that V_a^a, V_b^b, V_c^{a+b} and V_a^n, V_b^n, V_c^n are all non-zero representation spaces and therefore m_c^{ab} and c_{ab}^c are both meaningfully defined. The distinction between m_c^{ab} and c_{ab}^c lies with the fact that it is possible for c_{ab}^c to be zero while m_c^{ab} is non-zero (this can happen when the depth of c is greater than n but at most $a+b$ so that V_c^n is zero but V_c^{a+b} is non-zero). This parallels the fact that, in general, the rank of the $2n \times 2n$ matrix Σ in Theorem 60 could be greater than n while the rank of the $n \times n$ matrix C is at most n .

As a corollary of this correspondence between the Horn's problem and the eigenvalues of blocks of 2×2 matrices, we obtain an equivalent formulation of the tetrahedral Horn problem.

thm:tetrahedral_block

Theorem 61. *Let (a, b, c, d, e, f) be a sextuple of positive semidefinite eigenvalues of length at most n . Then (a, b, c, d, e, f) are tetrahedral if and only if there exists a $3n \times 3n$ matrix E' of the form*

$$E' = \begin{bmatrix} A' & Z & Y \\ Z^* & B' & X \\ Y^* & X^* & D' \end{bmatrix}, \quad \text{with blocks} \quad C' = \begin{bmatrix} A' & Z \\ Z^* & B' \end{bmatrix}, \quad F' = \begin{bmatrix} B' & X \\ X^* & D' \end{bmatrix} \quad \text{\textit{eq:block_3by3}} \quad (209)$$

such that

$$\text{eig}(A') = a, \quad \text{eig}(B') = b, \quad \text{eig}(C') = c, \quad \text{eig}(D') = d, \quad \text{eig}(E') = e, \quad \text{eig}(F') = f. \quad \text{\textit{eq:block_eigenvalue_tet}} \quad (210)$$

Proof. The proof is analogous to the proofs of Lemma 59 and Theorem 60. First we prove the “only if” portion of the claim which is a corollary of Lemma 59. Suppose (A, B, C, D, E, F) is a tetrahedral sextuple of Hermitian matrices with eigenvalues (a, b, c, d, e, c) . Let E' be given by

$$E' = \begin{bmatrix} A^{\frac{1}{2}} & B^{\frac{1}{2}} & D^{\frac{1}{2}} \end{bmatrix}^* \begin{bmatrix} A^{\frac{1}{2}} & B^{\frac{1}{2}} & D^{\frac{1}{2}} \end{bmatrix} = \begin{bmatrix} A & A^{\frac{1}{2}} B^{\frac{1}{2}} & A^{\frac{1}{2}} D^{\frac{1}{2}} \\ B^{\frac{1}{2}} A^{\frac{1}{2}} & B & B^{\frac{1}{2}} D^{\frac{1}{2}} \\ D^{\frac{1}{2}} A^{\frac{1}{2}} & D^{\frac{1}{2}} B^{\frac{1}{2}} & D \end{bmatrix} \quad (211)$$

Then the eigenvalues of the corresponding principal submatrices of E' satisfy Eq. (210) by Lemma 59. Now we prove the “if” portion of the claim which is a corollary of Theorem 60. If E' is a matrix of form given

by Eq. (209) with eigenvalues e with length at most n , then E' is positive semidefinite with rank at most n . Therefore, there exists an $n \times 3n$ block matrix

$$T = \begin{bmatrix} T_A & T_B & T_D \end{bmatrix} \quad (212)$$

such that $E' = T^*T$ and thus

$$\begin{aligned} A' &= T_A^*T_A, & B' &= T_B^*T_B, & D' &= T_D^*T_D, \\ X &= T_B^*T_D, & Y &= T_A^*T_D, & Z &= T_A^*T_B. \end{aligned} \quad (213)$$

Now let (A, B, C, D, E, F) be given by

$$\begin{aligned} A &= T_A T_A^*, & B &= T_B T_B^*, & C &= T_A T_A^* + T_B T_B^*, \\ D &= T_D T_D^*, & E &= T_A T_A^* + T_B T_B^* + T_D T_D^*, & F &= T_B T_B^* + T_D T_D^*. \end{aligned} \quad (214)$$

From here, one may directly verify that these $n \times n$ Hermitian matrices satisfy Eq. (12) and have eigenvalues (a, b, c, d, e, f) (using a similar argument presented in the proof of Theorem 60); i.e., one concludes that (a, b, c, d, e, f) are tetrahedral. $\square$

TODO: Describe the same correspondence for the blockwise $6j$ symbols.

Remark 62. One consequence of Theorem 61, which relates the eigenvalues of sums of positive semidefinite matrices to the eigenvalues of diagonal blocks of larger positive semidefinite matrices with bounded rank, is that many of the aforementioned inequalities can now be seen as consequences of more general inequalities which constrain the eigenvalues of blocks.

For instance, the Kotljanskii determinant inequality [JB93] (in the notation of Theorem 61) implies for any 3×3 block matrix, that

$$\det(Z') \det(F') \geq \det(E') \det(Y'). \quad (215)$$

Using Theorem 61, the Kotljanskii determinant inequality becomes Eq. (194). More generally, Audenaert, Hiai, and Petz were interested in functions f which are strongly superadditive in the sense that for any 3×3 block matrices E' (again in the notation of Theorem 61) the following inequality holds [AHP10]:

$$\mathrm{Tr} f(E') + \mathrm{Tr} f(Y') \geq \mathrm{Tr} f(Z') + \mathrm{Tr} f(F'). \quad (216) \quad \text{\textcolor{blue}{(eq:audenaert)}}$$

Their conclusion, also discovered by Friedland and Gaubert [FG13] was that it is *sufficient* for f' to be an operator monotone to conclude Eq. (216) holds. In light of Theorem 61, the results of Audenaert, Hiai, and Petz and [FG13] give an alternative proof of Theorem 58. In [LS14], Lewin and Sabin further proved that f' being an operator monotone is also *necessary* for Eq. (216) to hold for all 3×3 block matrices E' . Note that from the proof of Theorem 58 given in [Sag13, Section 2.1], it is not too difficult to see that f' being an operator monotone is also necessary for Eq. (195) to hold.

Remark 63. Much of the interest in characterizing the eigenvalues of principal submatrices of positive semidefinite matrices is closely related to the correspondence between positive semidefinite matrices and covariance matrices of Gaussian multivariate random variables [Lam+17]. From this perspective, principal submatrices become covariance matrices of Gaussian marginal distributions, and thus one may view the principal submatrix eigenvalue problem as a kind of “spectral” classical Gaussian marginal problem. Note that from this perspective, the differential entropies of the underlying Gaussian distributions are equal to logarithms of determinants up to an additive constants [Lam+17]. The Gaussian entropy region has been partially characterized by Shadbakht and Hassibi [SH11; HS08].

A.6 The Courtade-Sra determinant inequality

Many of the previously discussed inequalities constrain the eigenvalues of positive semidefinite matrices $Y, X + Y, Y + Z$ and $X + Y + Z$. Here we show how to improve the determinant inequality $\det(X + Y) \det(Y + Z) \geq \det(X + Y + Z) \det(Y)$ in a manner which also accounts for the determinants $\det(X)$ and

$\det(Z)$. This inequality exploits Minkowski's determinant inequality for positive semidefinite $n \times n$ matrices X and Y which reads (see [HJ85, Theorem 7.8.21] or [Bha97, Equation II.44]):

$$\det(X + Y)^{\frac{1}{n}} \geq \det(X)^{\frac{1}{n}} + \det(Y)^{\frac{1}{n}}. \quad (217)$$

The following inequality can be attributed to an information-theoretic result obtained by Courtade [Cou16]⁹, but the elementary proof we present here is do actually due to Sra [Sra]. This inequality also appears in the recent review paper by Niculescu and Sra [NS23, Section 6.2].

Proposition 64 (Courtade-Sra determinant inequality). *Let $X, Y, Z \geq 0$ be positive semidefinite $n \times n$ matrices. Then*

$$\det(X + Y)^{\frac{1}{n}} \det(Y + Z)^{\frac{1}{n}} \geq \det(X + Y + Z)^{\frac{1}{n}} \det(Y)^{\frac{1}{n}} + \det(X)^{\frac{1}{n}} \det(Z)^{\frac{1}{n}}. \quad (218)$$

Proof. By standard continuity arguments, it suffices to prove the inequality for $Y > 0$ positive definite (if it holds for positive definite $Y + \varepsilon I$, then continuity permits taking the limit $\varepsilon \rightarrow 0$). As Y is positive definite, it is also invertible and we may define

$$\tilde{X} = Y^{-\frac{1}{2}} X Y^{-\frac{1}{2}}, \quad \tilde{Z} = Y^{-\frac{1}{2}} Z Y^{-\frac{1}{2}}. \quad (219)$$

By dividing Eq. (218) by $\det(Y)^{\frac{2}{n}} \neq 0$, it then suffices to prove

$$\det(I + \tilde{X})^{\frac{1}{n}} \det(I + \tilde{Z})^{\frac{1}{n}} \geq \det(I + \tilde{X} + \tilde{Z})^{\frac{1}{n}} + \det(\tilde{X})^{\frac{1}{n}} \det(\tilde{Z})^{\frac{1}{n}}. \quad (220)$$

Let $x = \text{eig}(\tilde{X})$ and $z = \text{eig}(\tilde{Z})$. Then for any permutation $\pi \in S_n$ let

$$L_\pi = \text{diag}\{1 + x_i + z_{\pi(i)}\}_{i=1}^n, \quad R_\pi = \text{diag}\{x_i z_{\pi(i)}\}_{i=1}^n. \quad (221)$$

Then we have the identity

$$\det(I + \tilde{X}) \det(I + \tilde{Z}) = \prod_{i=1}^n (1 + x_i) \prod_{i=1}^n (1 + z_i), \quad (222)$$

$$= \prod_{i=1}^n (1 + x_i + y_{\pi(i)} + x_i z_{\pi(i)}), \quad (223)$$

$$= \det(L_\pi + R_\pi) \quad (224)$$

Here we can apply Minkowski's theorem to obtain

$$\det(L_\pi + R_\pi)^{\frac{1}{n}} \geq \det(L_\pi)^{\frac{1}{n}} + \det(R_\pi)^{\frac{1}{n}}. \quad (225)$$

By definition, we have $\det(R_\pi) = \det(\tilde{X}) \det(\tilde{Z})$ for any permutation π , while a theorem do to Fiedler [Fie71] states (see also [Bha97, Exercise VI.7.2, Theorem VI.7.1])

$$\det(I + \tilde{X} + \tilde{Y}) \leq \max_{\pi} \prod_{i=1}^n (1 + x_i + z_{\pi(i)}) = \max_{\pi} \det(L_\pi). \quad (226)$$

Combining these bounds together yields Eq. (220) and thus the claim holds. $\square$

A.7 Some conjectures

TC: On [mathoverflow](https://mathoverflow.net/questions/257562/a-determinant-inequality) <https://mathoverflow.net/questions/257562/a-determinant-inequality> Sra also proves

$$E_k(Y) E_k(X + Y + Z) \leq E_k(X + Y) E_k(Y + Z) \quad (227)$$

where $E_k(X) = e_k(\text{eig}(X))$ where e_k is the k -th elementary symmetric polynomial. TC: I conjecture that Eq. (218) also holds for more general hyperbolic polynomials, although I do not have a proof yet:

⁹Note in [Cou17], Courtade conjectures a generalization of the Brunn-Minkowski inequality for bounded convex sets X, Y, Z in $\mathbb{R}^n$ of the form $V(X + Y)^{\frac{1}{n}} V(Y + Z)^{\frac{1}{n}} \geq V(X + Y + Z)^{\frac{1}{n}} V(Y)^{\frac{1}{n}} + V(X)^{\frac{1}{n}} V(Z)^{\frac{1}{n}}$. See also [CC84].

Conjecture 65. *Let $X, Y, Z \geq 0$ be positive semidefinite $n \times n$ matrices. Then*

$$E_k^{\frac{1}{k}}(X+Y)E_k^{\frac{1}{k}}(Y+Z) \geq E_k^{\frac{1}{k}}(X+Y+Z)E_k^{\frac{1}{k}}(Y) + E_k^{\frac{1}{k}}(X)E_k^{\frac{1}{k}}(Z), \quad \{\text{eq:cs_ineq_conjecture}\} \quad (228)$$

where $E_k(X) = e_k(\text{eig}(X))$ and where e_k is the k -th elementary symmetric polynomial.

Proving this conjecture has proven quite challenging.

What I believe I'm managed to prove is the following.

Let $\phi : \text{PSD}_n \rightarrow \mathbb{R}$ be a function on the cone of positive semidefinite matrices. If ϕ is such that $\log \phi$ is 2-monotone, i.e.,

$$-(\partial_X \log \phi)(Y) \geq 0, \quad +(\partial_Z \partial_X \log \phi)(Y) \geq 0, \quad (229)$$

then it is fairly clear that

$$\log \phi(X+Y) + \log \phi(Y+Z) \geq \log \phi(Y) + \log \phi(X+Y+Z), \quad (230)$$

or equivalently,

$$\phi(X+Y)\phi(Y+Z) \geq \phi(Y)\phi(X+Y+Z). \quad (231)$$

If in addition to $\log \phi$ being 2-monotone, ϕ satisfies a kind of 3-montonicity, namely

$$-\partial_W(\phi^2 \partial_Z \partial_X \log \phi)(Y) \geq 0, \quad (232)$$

then,

$$[\phi(X+Y+W)\phi(Y+Z+W) - \phi(X+Y+Z+W)\phi(Y+W)] - [\phi(X+Y)\phi(Y+Z) - \phi(X+Y+Z)\phi(Y)] \geq 0 \quad (233)$$

which means, assuming $\phi(0) = 0$ and for $Y \mapsto 0$ and $W \mapsto Y$, we have desired inequality:

$$\phi(X+Y)\phi(Y+Z) - \phi(X+Y+Z)\phi(Y) - \phi(X)\phi(Z) \geq 0. \quad (234)$$

Then what remains is to prove that $\phi = E_k^{\frac{1}{k}}$ has this property.

TC: References that might be useful: The strongly subadditive inequality relies on the log concavity result for hyperbolic polynomials due to Bauschke et al. [Bau+01]. An excellent crash course into Gårding's theory of hyperbolic polynomials (as well as some important results) is also [Bau+01]. For the particular case of the real vector space of $n \times n$ Hermitian matrices of dimension $n(n+1)/2$, the polynomial $X \mapsto \det(X)$ is hyperbolic in the direction of the identity matrix id_n (i.e., Hermitian matrices only have real eigenvalues). Moreover, the homogeneous polynomial $p_m(X) = \text{Tr}(P_{(1,m,1)} X^{\otimes m})$ is the sum of the principal minors of X of size m and also equals the elementary symmetric polynomial of the eigenvalues of X , i.e.,

$$p_m(X) = E_m(\text{eig}(X)). \quad (235)$$

The homogeneous polynomial $p_m(X)$, like the determinant $\det(X) = p_n(X)$ itself, is also hyperbolic in the direction of the identity matrix id_n . In this context, together with the positivity $p_m(\text{id}_n) > 0$, Gårding's inequality generalizes the Minkowski determinant inequality and yields superlinearity of $p_m(X)^{\frac{1}{m}}$:

$$p_m(X+Y)^{\frac{1}{m}} \geq p_m(X)^{\frac{1}{m}} + p_m(Y)^{\frac{1}{m}} \quad (236)$$

for all X, Y positive semidefinite (the hyperbolicity cone for p_m). This inequality is due to Marcus and Lopes [ML57, Theorem 2], but it can also be proven using joint concavity of $(X_1, \dots, X_m) \mapsto X_1^{\frac{1}{m}} \otimes \dots \otimes X_m^{\frac{1}{m}}$ and thus concavity of $(X^{\frac{1}{m}})^{\otimes m}$ and thus by homogeneity the operator inequality $((X+Y)^{\frac{1}{m}})^{\otimes m} \geq (X^{\frac{1}{m}})^{\otimes m} + (Y^{\frac{1}{m}})^{\otimes m}$ ([Bha09, Exercise 4.2.9]).

See [ML57; McL59; AMT84] for using inequalities and proof techniques which might be relevant. See also [AMT84] or [Bha09, Chapter 4] for many properties of matrix means and their ability to prove Minkowski-like inequalities. See also Khovanskii [Kho84] for inequalities mirroring the Aleksandrov-Fenchel type inequalities for hyperbolic polynomials (i.e., the mixed discriminant inequalities of Aleksandrov... see Shenfeld and Van Handel for proofs [SV19].) See [KMS19] for representation theorems and other results concerning complete monotonicity.

References

- [AH11] Daiki Akimoto and Masahito Hayashi. “Discrimination of the change point in a quantum setting”. In: *Physical Review A* 83.5 (2011), p. 052328.
- [AHP10] Koenraad Audenaert, Fumio Hiai, and Dénes Petz. “Strongly subadditive functions”. In: *Acta Mathematica Hungarica* 128.4 (2010), pp. 386–394.
- [AI19] Arseniy Akopyan and Ivan Izmestiev. “The Regge symmetry, confocal conics, and the Schläfli formula”. In: *Bulletin of the London Mathematical Society* 51.5 (2019), pp. 765–775.
- [Alc+23] Judith Alcock-Zeilinger et al. “Wigner 6j symbols for SU (N): Symbols with at least two quark-lines”. In: *Journal of Mathematical Physics* 64.2 (2023).
- [Ale+25] Anton Alekseev et al. “Multiple Horn problems for planar networks and invertible matrices”. In: *arXiv preprint arXiv:2503.05277* (2025).
- [AMS20] Victor Alekseev, Andrey Morozov, and Alexey Sleptsov. “Multiplicity-free $U_q(\mathfrak{sl}N)$ 6-j symbols: relations, asymptotics, symmetries”. In: *Nuclear Physics B* 960 (2020), p. 115164.
- [AMT84] WN Anderson Jr, TD Morley, and GE Trapp. “Symmetric function means of positive operators”. In: *Linear algebra and its applications* 60 (1984), pp. 129–143.
- [AMW01] Anton Alekseev, Eckhard Meinrenken, and Chris Woodward. “Linearization of Poisson actions and singular values of matrix products”. In: *Annales de l’institut Fourier*. Vol. 51. 6. 2001, pp. 1691–1717.
- [Aqu+12] Vincenzo Aquilanti et al. “Semiclassical mechanics of the Wigner 6j-symbol”. In: *Journal of Physics A: Mathematical and Theoretical* 45.6 (2012), p. 065209.
- [ARS88] Robert Alicki, Slawomir Rudnicki, and Slawomir Sadowski. “Symmetry properties of product states for the system of N n-level atoms”. In: *Journal of mathematical physics* 29.5 (1988), pp. 1158–1162.
- [Art23] DV Artamonov. “Classical 6j-symbols of finite-dimensional representations of the algebra $\mathfrak{gl}_3$ ”. In: *Theoretical and Mathematical Physics* 216.1 (2023), pp. 909–923.
- [Art24] Dmitry Artamonov. “Calculation of 6j-symbols for the Lie algebra $\mathfrak{gl}_n$ ”. In: *arXiv preprint arXiv:2405.05628* (2024).
- [AW97] Sharad Agnihotri and Chris Woodward. “Eigenvalues of products of unitary matrices and quantum Schubert calculus”. In: *arXiv preprint alg-geom/9712013* (1997).
- [Bac10] Miriam Backens. “The 6j symbols and an extended version of Horn’s problem”. In: (2010).
- [Bae96] John C Baez. “Spin networks in gauge theory”. In: *Advances in Mathematics* 117.2 (1996), pp. 253–272.
- [Bae99] John C Baez. “An introduction to spin foam models of quantum gravity and BF theory”. In: (1999). arXiv: [gr-qc/9905087](https://arxiv.org/abs/gr-qc/9905087).
- [Bar03] John W Barrett. “Geometrical measurements in three-dimensional quantum gravity”. In: *International Journal of Modern Physics A* 18.suppl02 (2003), pp. 97–113.
- [Bau+01] Heinz H Bauschke et al. “Hyperbolic polynomials and convex analysis”. In: *Canadian Journal of Mathematics* 53.3 (2001), pp. 470–488.
- [BC98] John W Barrett and Louis Crane. “Relativistic spin networks and quantum gravity”. In: *Journal of Mathematical Physics* 39.6 (1998), pp. 3296–3302.
- [Bel02] Prakash Belkale. “Geometric proofs of Horn and saturation conjectures”. In: *arXiv preprint math/0208107* (2002).
- [Bel05] Prakash Belkale. “Geometric proof of a conjecture of Fulton”. In: *arXiv preprint math/0511664* (2005).
- [Bel08] Prakash Belkale. “Quantum generalization of the Horn conjecture”. In: *Journal of the American Mathematical Society* 21.2 (2008), pp. 365–408.

- [BGS08] Mustapha Bouhtou, Stéphane Gaubert, and Guillaume Sagnol. “Optimization of network traffic measurement: a semidefinite programming approach”. In: *Proceedings of the International*. Vol. 230. 2008.
- [Bha01] Rajendra Bhatia. “[Linear algebra to quantum cohomology: the story of Alfred Horn’s inequalities](#)”. In: *The American Mathematical Monthly* 108.4 (2001), pp. 289–318.
- [Bha09] Rajendra Bhatia. *Positive definite matrices*. Princeton university press, 2009.
- [Bha97] Rajendra Bhatia. *Matrix analysis*. New York: Springer, 1997.
- [BK16] Prakash Belkale and Shrawan Kumar. “The multiplicative eigenvalue problem and deformed quantum cohomology”. In: *Advances in Mathematics* 288 (2016), pp. 1309–1359.
- [BL22] Jean-Christophe Bourin and Eun-Young Lee. “Eigenvalue inequalities for positive block matrices with the inradius of the numerical range”. In: *International Journal of Mathematics* 33.01 (2022), p. 2250009.
- [BN23] Bruce Bartlett and Nzaganya Nzaganya. “Coherent loop states and angular momentum”. In: *arXiv preprint arXiv:2306.17293* (2023).
- [Boa07] Philip P Boalch. “[Regge and Okamoto symmetries](#)”. In: *Communications in mathematical physics* 276.1 (2007), pp. 117–130.
- [BS10] John Baez and Mike Stay. “[Physics, topology, logic and computation: a Rosetta Stone](#)”. In: *New structures for physics*. Springer, 2010, pp. 95–172. arXiv: 0903.0340v3.
- [BS15] Wolfgang Berndt and Suvrit Sra. “Hlawka–Popoviciu inequalities on positive definite tensors”. In: *Linear Algebra and its Applications* 486 (2015), pp. 317–327.
- [BVW18] Nicole Berline, Michèle Vergne, and Michael Walter. “The Horn inequalities from a geometric point of view”. In: *L’Enseignement Mathématique* 63.3 (2018), pp. 403–470.
- [CC84] Max Costa and Thomas Cover. “On the similarity of the entropy power inequality and the Brunn–Minkowski inequality (corresp.)”. In: *IEEE Transactions on Information Theory* 30.6 (1984), pp. 837–839.
- [Chr08] Matthias Christandl. “A Quantum Information-Theoretic Proof of the Relation between Horn’s Problem and the Littlewood–Richardson Coefficients”. In: *Lecture Notes in Computer Science*. Springer Berlin Heidelberg, 2008, pp. 120–128.
- [CHW07] Andrew M Childs, Aram W Harrow, and Paweł Wocjan. “Weak Fourier–Schur sampling, the hidden subgroup problem, and the quantum collision problem”. In: *Annual Symposium on Theoretical Aspects of Computer Science*. Springer. 2007, pp. 598–609.
- [CIS99] Li-Chen Chen, Mourad EH Ismail, and Plamen Simeonov. “Asymptotics of Racah coefficients and polynomials”. In: *Journal of Physics A: Mathematical and General* 32.3 (1999), p. 537.
- [CMZ19] Robert Coquereaux, Colin McSwiggen, and Jean-Bernard Zuber. “Revisiting Horn’s problem”. In: *Journal of Statistical Mechanics: Theory and Experiment* 2019.9 (2019), p. 094018.
- [Coq23] Robert Coquereaux. “About integer-valued variants of the theta and 6j symbols”. In: *Journal of Mathematical Physics* 64.3 (2023).
- [Cou16] Thomas A Courtade. “Strengthening the entropy power inequality”. In: *2016 IEEE International Symposium on Information Theory (ISIT)*. IEEE. 2016, pp. 2294–2298.
- [Cou17] Thomas A Courtade. “Concavity of entropy power: Equivalent formulations and generalizations”. In: *2017 IEEE International Symposium on Information Theory (ISIT)*. IEEE. 2017, pp. 56–60.
- [CŞW18] Matthias Christandl, M Burak Şahinoğlu, and Michael Walter. “Recoupling coefficients and quantum entropies”. In: *Annales Henri Poincaré*. Vol. 19. 2. Springer. 2018, pp. 385–410.
- [CZ18] Robert Coquereaux and Jean-Bernard Zuber. “From orbital measures to Littlewood–Richardson coefficients and hive polytopes”. In: *Annales de l’Institut Henri Poincaré D* 5.3 (2018), pp. 339–386.

- [DH05] Sumit Daftuar and Patrick Hayden. “Quantum state transformations and the Schubert calculus”. In: *Annals of Physics* 315.1 (2005), pp. 80–122.
- [DJN93] Bergfinnur Durhuus, Hans Plesner Jakobsen, and Ryszard Nest. “Topological quantum field theories from generalized 6j-symbols”. In: *Reviews in Mathematical Physics* 5.01 (1993), pp. 1–67.
- [Edm96] Alan Robert Edmonds. *Angular momentum in quantum mechanics*. Vol. 4. Princeton university press, 1996.
- [Fan49] Ky Fan. “On a theorem of Weyl concerning eigenvalues of linear transformations I”. In: *Proceedings of the National Academy of Sciences* 35.11 (1949), pp. 652–655.
- [FG13] S Friedland and S Gaubert. “Submodular spectral functions of principal submatrices of a hermitian matrix, extensions and applications”. In: *Linear Algebra and its Applications* 438.10 (2013), pp. 3872–3884.
- [Fie71] Miroslav Fiedler. “Bounds for the determinant of the sum of hermitian matrices”. In: *Proceedings of the American Mathematical Society* (1971), pp. 27–31.
- [FK00] Laurent Freidel and Kirill Krasnov. “Simple spin networks as Feynman graphs”. In: *Journal of Mathematical Physics* 41.4 (2000), pp. 1681–1690.
- [FKL10] Laurent Freidel, Kirill Krasnov, and Etera R Livine. “Holomorphic factorization for a quantum tetrahedron”. In: *Communications in Mathematical Physics* 297.1 (2010), pp. 45–93.
- [FL03] Laurent Freidel and David Louapre. “Asymptotics of 6j and 10j symbols”. In: *Classical and Quantum Gravity* 20.7 (2003), p. 1267.
- [Fra22] Thomas C Fraser. *A sufficient family of necessary inequalities for the compatibility of quantum marginals*. 2022. arXiv: [2211.00685v1](https://arxiv.org/abs/2211.00685v1).
- [Ful00] William Fulton. “Eigenvalues, invariant factors, highest weights, and Schubert calculus”. In: *Bulletin of the American Mathematical Society* 37.3 (2000), pp. 209–249.
- [Ful98] William Fulton. “Eigenvalues of sums of Hermitian matrices”. In: *Séminaire Bourbaki* 40.255-269 (1998), p. 5.
- [GS82] Victor Guillemin and Shlomo Sternberg. “Geometric quantization and multiplicities of group representations”. In: *Inventiones mathematicae* 67.3 (1982), pp. 515–538.
- [Gum+18] Mehmet Gumus et al. “Positive semi-definite 2×2 block matrices and norm inequalities”. In: *Linear Algebra and Its Applications* 551 (2018), pp. 83–91.
- [Gur08] Razvan Gurau. “The Ponzano–Regge Asymptotic of the 6 j Symbol: An Elementary Proof”. In: *Annales Henri Poincaré*. Vol. 9. Springer. 2008, pp. 1413–1424.
- [GW00] Roe Goodman and Nolan R Wallach. *Representations and invariants of the classical groups*. Cambridge University Press, 2000.
- [Hal08] Matthew John Hall. “Representations of the Symmetric Groups and Combinatorics of the Frobenius-Young Correspondence”. PhD thesis. Citeseer, 2008.
- [Har23] Aram W Harrow. “Approximate orthogonality of permutation operators, with application to quantum information”. In: *Letters in Mathematical Physics* 114.1 (2023), p. 1.
- [Hay17] Masahito Hayashi. *A Group Theoretic Approach to Quantum Information*. Springer, 2017.
- [HJ12] Roger A Horn and Charles R Johnson. *Matrix analysis*. Cambridge university press, 2012.
- [HJ85] RA Horn and CR Johnson. “Matrix Analysis, Cambridge Univ”. In: *Press. MR0832183* (1985).
- [HK96] Jean-Claude Hausmann and Allen Knutson. “Polygon spaces and Grassmannians”. In: *arXiv preprint dg-ga/9602012* (1996).
- [HM02] Masahito Hayashi and Keiji Matsumoto. “Quantum universal variable-length source coding”. In: *Physical Review A* 66.2 (2002), p. 022311.
- [HM21] Felix Huber and Hans Maassen. “Matrix forms of immanant inequalities”. In: *arXiv preprint arXiv:2103.04317* (2021).

- [Hor62] Alfred Horn. “Eigenvalues of sums of Hermitian matrices.” In: (1962).
- [HS08] Babak Hassibi and Sormeh Shadbakht. “The entropy region for three gaussian random variables”. In: *2008 IEEE International Symposium on Information Theory*. IEEE. 2008, pp. 2634–2638.
- [IN66] Claude Itzykson and Michael Nauenberg. “Unitary groups: Representations and decompositions”. In: *Reviews of Modern Physics* 38.1 (1966), p. 95.
- [IP23] Christian Ikenmeyer and Greta Panova. “All Kronecker coefficients are reduced Kronecker coefficients”. In: *arXiv preprint arXiv:2305.03003* (2023).
- [JB93] Charles R Johnson and Wayne W Barrett. “Determinantal inequalities for positive definite matrices”. In: *Discrete Mathematics* 119.1-3 (1993), pp. 97–106.
- [Kho84] Askold Georgievich Khovanskii. “Analogues of Aleksandrov–Fenchel inequalities for hyperbolic forms”. In: *Doklady Akademii Nauk*. Vol. 276. 6. Russian Academy of Sciences. 1984, pp. 1332–1334.
- [Kir04] Aleksandr Aleksandrovich Kirillov. *Lectures on the orbit method*. Vol. 64. American Mathematical Soc., 2004.
- [Kly00] Alexander A Klyachko. “Random walks on symmetric spaces and inequalities for matrix spectra”. In: *Linear Algebra and its Applications* 319.1-3 (2000), pp. 37–59.
- [Kly98] Alexander A Klyachko. “Stable bundles, representation theory and Hermitian operators”. In: *Selecta Mathematica* 4 (1998), pp. 419–445.
- [KMS19] Khazhgali Kozhasov, Mateusz Michałek, and Bernd Sturmfels. “Positivity certificates via integral representations”. In: *Facets of algebraic geometry* 2 (2019), pp. 84–114.
- [Knu00] Allen Knutson. “[The symplectic and algebraic geometry of Horn’s problem](#)”. In: *Linear Algebra and its Applications* 319.1-3 (2000), pp. 61–81.
- [KPS24] Stefan Keppeler, Simon Plätzer, and Malin Sjö Dahl. “Wigner 6j symbols with gluon lines: completing the set of 6j symbols required for color decomposition”. In: *Journal of High Energy Physics* 2024.5 (2024), pp. 1–25.
- [KT99] Allen Knutson and Terence Tao. “The honeycomb model of $GL_n(\mathbb{C})$ tensor products I: Proof of the saturation conjecture”. In: *Journal of the American Mathematical Society* 12.4 (1999), pp. 1055–1090.
- [KW01] Michael Keyl and Reinhard F Werner. “[Estimating the spectrum of a density operator](#)”. In: *Physical Review A* 64.5 (2001), p. 052311.
- [Lam+17] Ludovico Lami et al. “From log-determinant inequalities to Gaussian entanglement via recoverability theory”. In: *IEEE Transactions on Information Theory* 63.11 (2017), pp. 7553–7568.
- [Lid53] V B Lidskiĭ. “The proper values of the sum and product of symmetric matrices”. Trans. by Curtis D. Benster. In: (1953). Ed. by George E. Forsythe Wallace Givens.
- [Lin14] Minghua Lin. “A determinantal inequality for positive semidefinite matrices”. In: *The Electronic Journal of Linear Algebra* 27 (2014), pp. 821–826.
- [Lit58] Dudley E Littlewood. “Products and plethysms of characters with orthogonal, symplectic and symmetric groups”. In: *Canadian Journal of Mathematics* 10 (1958), pp. 17–32.
- [LS14] Mathieu Lewin and Julien Sabin. “[A family of monotone quantum relative entropies](#)”. In: *Letters in Mathematical Physics* 104.6 (2014), pp. 691–705. arXiv: [1309.4046v1](#).
- [LS23] Elena Lanina and Alexey Sleptsov. “Tug-the-hook symmetry for quantum 6j-symbols”. In: *Physics Letters B* 845 (2023), p. 138138.
- [McL59] John Bryce McLeod. “On four inequalities in symmetric functions”. In: *Proceedings of the Edinburgh Mathematical Society* 11.4 (1959), pp. 211–219.
- [MKX19] Mohammad Sal Moslehian, Mohsen Kian, and Qingxiang Xu. “Positivity of 2×2 block matrices of operators”. In: (2019).

- [ML57] M Marcus and L Lopes. “Inequalities for symmetric functions and Hermitian matrices”. In: *Canadian Journal of Mathematics* 9 (1957), pp. 305–312.
- [MM65] Marvin Marcus and Henryk Minc. “Generalized matrix functions”. In: *Transactions of the American Mathematical Society* 116 (1965), pp. 316–329.
- [MOA10] A.W. Marshall, I. Olkin, and B.C. Arnold. *Inequalities: Theory of Majorization and Its Applications*. Springer Series in Statistics. Springer New York, 2010.
- [Mou84] J Moussouris. “Quantum models of space-time based on recoupling theory”. PhD thesis. University of Oxford, 1984.
- [MR05] Annalisa Marzuoli and Mario Rasetti. “Computing spin networks”. In: *Annals of Physics* 318.2 (2005), pp. 345–407.
- [MS18] Andrey Morozov and Alexey Sleptsov. “New Symmetries for the $U_q(\mathfrak{sl}_N)$ $6j$ Symbols from the Eigenvalue Conjecture”. In: *JETP Letters* 108 (2018), pp. 697–704.
- [Mur55] Francis D Murnaghan. “On the analysis of the Kronecker product of irreducible representations of S_n ”. In: *Proceedings of the National Academy of Sciences* 41.7 (1955), pp. 515–518.
- [Nis00] Kyo Nishiyama. “Restriction of the irreducible representations of GL_n to the symmetric group S_n ”. In: *preprint* (2000).
- [NP06] Constantin Niculescu and Lars-Erik Persson. *Convex functions and their applications*. Vol. 23. Springer, 2006.
- [NPZ13] Satoshi Nawata, Ramadevi Pichai, and Zodinmawia. “Multiplicity-free Quantum $6j$ -Symbols for $U_q(\mathfrak{sl}_N)$ ”. In: *Letters in Mathematical Physics* 103 (2013), pp. 1389–1398.
- [NS23] Constantin P Niculescu and Suvrit Sra. “Functions with Positive Differences on Convex Cones”. In: *Results in Mathematics* 78.6 (2023), p. 217.
- [OW16] Ryan O’Donnell and John Wright. “Efficient quantum tomography”. In: *Proceedings of the forty-eighth annual ACM symposium on Theory of Computing*. 2016, pp. 899–912. arXiv: [1508.01907v2](#).
- [Pen71] Roger Penrose. “Applications of negative dimensional tensors”. In: *Combinatorial mathematics and its applications* 1 (1971), pp. 221–244.
- [PR68] Giorgio Ponzano and Tullio Eugenio Regge. “Semiclassical limit of Racah coefficients”. In: *Spectroscopic and group theoretical methods in physics* (1968).
- [PTZ14] Vehbi Paksoy, Ramazan Turkmen, and Fuzhen Zhang. “Inequalities of generalized matrix functions via tensor products”. In: *The Electronic Journal of Linear Algebra* 27 (2014), pp. 332–341.
- [Rac42] Giulio Racah. “Theory of complex spectra. II”. In: *Physical Review* 62.9-10 (1942), p. 438.
- [Reg59] Tullio Regge. “Symmetry properties of Racah’s coefficients”. In: *Il Nuovo Cimento (1955-1965)* 11.1 (1959), pp. 116–117.
- [Res87] N Yu Reshetikhin. *Quantized universal enveloping algebras, the Yang-Baxter equation and invariants of links, 1*. Tech. rep. 1987.
- [Ric02] G Richardson. “The trigonometry of the tetrahedron”. In: *The Mathematical Gazette* 2.32 (1902), pp. 149–158.
- [Rob02] Justin Roberts. “Asymptotics and $6j$ -symbols”. In: *Geom. Topol. Monogr* 4 (2002), pp. 245–261.
- [Rob55] Herbert Robbins. “[A remark on Stirling’s formula](#)”. In: *The American mathematical monthly* 62.1 (1955), pp. 26–29.
- [Rob99] Justin Roberts. “[Classical \$6j\$ -symbols and the tetrahedron](#)”. In: *Geometry & Topology* 3.1 (1999), pp. 21–66.
- [Rov04] Carlo Rovelli. *Quantum gravity*. Cambridge university press, 2004.

- [Sag13] Guillaume Sagnol. “Approximation of a maximum-submodular-coverage problem involving spectral functions, with application to experimental designs”. In: *Discrete Applied Mathematics* 161.1-2 (2013), pp. 258–276.
- [Sch52] Isaac J Schoenberg. “A remark on MM Day’s characterization of inner-product spaces and a conjecture of LM Blumenthal”. In: *Proceedings of the American Mathematical Society* 3.6 (1952), pp. 961–964.
- [SH11] Sormeh Shadbakht and Babak Hassibi. “On the entropy region of Gaussian random variables”. In: *arXiv preprint arXiv:1112.0061* (2011).
- [She15] Cass Sherman. “Geometric proof of a conjecture of King, Tollu, and Toumazet”. In: *arXiv preprint arXiv:1505.06551* (2015).
- [SJS18] Meera Sitharam, Audrey John, and Jessica Sidman, eds. *Handbook of geometric constraint systems principles*. Discrete Mathematics and Its Applications. New York, NY: Productivity Press, July 2018.
- [Spe05] David E Speyer. “Horn’s problem, Vinnikov curves, and the hive cone”. In: (2005).
- [Sra] Suvrit Sra. “[Reverse Minkowski \(and related\) determinant inequalities](#)”. In: *MathOverflow*. (version: 10/10/2016) ().
- [SV19] Yair Shenfeld and Ramon Van Handel. “Mixed volumes and the Bochner method”. In: *Proceedings of the American Mathematical Society* 147.12 (2019), pp. 5385–5402.
- [TCL11] Lin Tie, Kai-Yuan Cai, and Yan Lin. “Rearrangement inequalities for Hermitian matrices”. In: *Linear algebra and its applications* 434.2 (2011), pp. 443–456.
- [TF70] Robert C. Thompson and Linda J. Freede. “[On the Eigenvalues of Sums of Hermitian Matrices. II.](#)” In: *Aequationes mathematicae* 5 (1970), pp. 103–115.
- [TF71] Robert C Thompson and Linda J Freede. “On the eigenvalues of sums of Hermitian matrices”. In: *Linear Algebra and Its Applications* 4.4 (1971), pp. 369–376.
- [TT71] RC Thompson and S Therianos. “The eigenvalues and singular values of matrix sums and product, viii: displacement of indices”. In: *aequationes mathematicae* 7 (1971), pp. 219–242.
- [TV92] Vladimir G Turaev and Oleg Ya Viro. “State sum invariants of 3-manifolds and quantum 6j-symbols”. In: *Topology* 31.4 (1992), pp. 865–902.
- [TW06] Yuka U Taylor and Christopher T Woodward. “6 j symbols for and non-Euclidean tetrahedra”. In: *Selecta Mathematica* 11.3-4 (2006), p. 539.
- [VMK88] Dmitriĭ Aleksandrovich Varshalovich, Anatolij Nikolaevič Moskalev, and Valerii Kel’manovich Khersonskii. [Quantum theory of angular momentum](#). World Scientific, 1988.
- [Wat99] James KG Watson. “Asymptotic approximations for certain 6-j and 9-j symbols”. In: *Journal of Physics A: Mathematical and General* 32.39 (1999), p. 6901.
- [WD09] Karl Wirth and André S Dreiding. “Edge lengths determining tetrahedrons”. In: *Elemente der Mathematik* 64.4 (2009), pp. 160–170.
- [Wey12] Hermann Weyl. “Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen (mit einer Anwendung auf die Theorie der Hohlraumstrahlung)”. In: *Mathematische Annalen* 71.4 (1912), pp. 441–479.
- [Wie55] Helmut Wielandt. “An extremum property of sums of eigenvalues”. In: *Proceedings of the American Mathematical Society* 6.1 (1955), pp. 106–110.
- [Wig60] Eugene Paul Wigner. “Group theory and its application to the quantum mechanics of atomic spectra”. In: *American journal of physics* 28.4 (1960), pp. 408–409.
- [Wig93] Eugene P Wigner. “On the matrices which reduce the Kronecker products of representations of SR groups”. In: *The Collected Works of Eugene Paul Wigner: Part A: The Scientific Papers*. Springer, 1993, pp. 608–654.
- [Zub18] Jean-Bernard Zuber. “Horn’s problem and Harish-Chandra’s integrals. Probability density functions”. In: *Annales de l’Institut Henri Poincaré D* 5.3 (2018), pp. 309–338.