

OLD DRAFT

[WORKING DOCUMENT]

On the spectra of partial products of matrices and the asymptotics of recoupling symbols

Thomas C. Fraser, Matthias Christandl

May 30, 2024

1 Introduction

1.1 Thompson's conjecture for 2×2 matrices

[sec:2by2case_intuitively](#)

The purpose of this section is to sketch how to prove Thompson's conjecture for the case of 2×2 matrices. Given $\lambda \geq 0$, let $\mathcal{O}_\lambda$ denotes the set of 2×2 traceless Hermitian operators with eigenvalues $\pm\lambda$ and $\mathcal{C}_{\exp(\lambda)}$ denote the set of 2×2 unit determinant Hermitian operators with singular-values $\exp(\pm\lambda)$. Note that the matrix exponential $\exp : \mathcal{O}_\lambda \rightarrow \mathcal{C}_{\exp(\lambda)}$ here is a bijective map with inverse $\log : \mathcal{C}_{\exp(\lambda)} \rightarrow \mathcal{O}_\lambda$.

The central idea is to show that exists a diffeomorphism on $\mathcal{O}_\alpha \times \mathcal{O}_\beta$, sending each pair of matrices (A, B) with eigenvalues $(\pm\alpha, \pm\beta)$ to another pair of matrices

$$(A, B) \mapsto (\tilde{A}, \tilde{B}) \in \mathcal{O}_\alpha \times \mathcal{O}_\beta \quad (1)$$

with the same eigenvalues as (A, B) but with the additional property:

$$A + B \in \mathcal{O}_\gamma \iff \exp(\tilde{A}) \exp(\tilde{B}) \in \mathcal{C}_{\exp(\gamma)}. \quad (2)$$

From the existence of this map one can therefore prove Thompson's conjecture: there exists $(A, B) \in \mathcal{O}_\alpha \times \mathcal{O}_\beta$ satisfying $A + B \in \mathcal{O}_\gamma$ if and only if there exists $(X, Y) \in \mathcal{C}_{\exp(\alpha)} \times \mathcal{C}_{\exp(\beta)}$ satisfying $XY \in \mathcal{C}_{\exp(\gamma)}$. For the only if portion of the proof, simply take (X, Y) to be $(\exp(\tilde{A}), \exp(\tilde{B}))$. For the if portion of the proof, define $(\tilde{A}, \tilde{B})$ to equal $(\log X, \log Y)$ and then apply the inverse of the above diffeomorphism to obtain the desired (A, B) from $(\tilde{A}, \tilde{B})$.

1.1.1 Euclidean triangles and eigenvalues

Letting σ_1, σ_2 and σ_3 be the 2×2 Pauli matrices,

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (3)$$

it becomes possible to identify the space of traceless 2×2 Hermitian matrices, $i\mathfrak{su}(2)$, with $\mathbb{R}^3$ via the standard parameterization associating $(x_1, x_2, x_3) \in \mathbb{R}^3$ to

$$X = x \cdot \sigma = x_1\sigma_1 + x_2\sigma_2 + x_3\sigma_3 = \begin{pmatrix} x_3 & x_1 - ix_2 \\ x_1 + ix_2 & -x_3 \end{pmatrix}. \quad (4)$$

The spectrum of X obtained in this way is simply $(+\chi, -\chi)$ where $\chi = \|x\|$ and $\|x\|$ the Euclidean norm of the vector $x \in \mathbb{R}^3$

$$\|x\| = \sqrt{x_1^2 + x_2^2 + x_3^2}. \quad (5)$$

From this point of view, it is possible to characterize the set of possible spectra, $(\alpha, \beta, \gamma) \in \mathbb{R}^3$, for the set of two Hermitian operators,

$$A = a_1\sigma_1 + a_2\sigma_2 + a_3\sigma_3, \quad B = b_1\sigma_1 + b_2\sigma_2 + b_3\sigma_3 \quad (6)$$

with spectra $(\alpha, \beta) = (\|a\|, \|b\|)$ and there sum $A + B$

$$A + B = (a_1 + b_1)\sigma_1 + (a_2 + b_2)\sigma_2 + (a_3 + b_3)\sigma_3, \quad (7)$$

with spectrum $\gamma = \|a + b\|$ where


$$\|a + b\|^2 = \|a\|^2 + \|b\|^2 + 2\|a\|\|b\|\cos\theta, \quad \cos\theta = \frac{a \cdot b}{\|a\|\|b\|}. \quad (8)$$

As the only parameter not fixed by spectra is the angle θ , it is easy to see that $(\alpha, \beta, \gamma) \in \mathbb{R}^3$ is precisely determined by the triangle inequalities for the Euclidean norm in $\mathbb{R}^3$:

$$|\|a\| - \|b\|| \leq \|a + b\| \leq \|a\| + \|b\|, \quad (9)$$

$$|\alpha - \beta| \leq \gamma \leq \alpha + \beta, \quad (10)$$

where the extreme cases are characterized by anti-alignment $\cos\theta = -1$ and alignment $\cos\theta = 1$ respectively.



In summary, given a pair $(A, B) = (a \cdot \sigma, b \cdot \sigma)$ of traceless 2×2 Hermitian operators with spectra $(\pm\alpha, \pm\beta)$ we saw that the spectrum $\pm\gamma$ of their sum, $A + B$, is given by the Euclidean cosine law with $\cos\theta = \frac{a \cdot b}{\alpha\beta}$ ¹

$$\gamma^2 = \alpha^2 + \beta^2 + 2\alpha\beta\cos\theta. \quad (12)$$

1.1.2 Hyperbolic triangles and singular values

An analogous calculation can be performed to character the spectra of certain invertible matrices in $\text{SL}(2, \mathbb{C})$ (obtained by exponentiation of the traceless Hermitian matrices previously considered) and their products. Given the traceless Hermitian matrix $X = x \cdot \sigma$, the identity $X^2 = \|x\|^2 I$ yields a well-known pair of results (see [FT14]):

$$\exp(iX) = \cos\|x\| I + i \frac{\sin\|x\|}{\|x\|} X, \quad \exp(X) = \cosh\|x\| I + \frac{\sinh\|x\|}{\|x\|} X. \quad (13)$$

¹Note that the angle θ used here is not the interior angle typically used in the statement of cosine law, hence the difference in minus sign.

As before, consider traceless Hermitian matrices, $A = a \cdot \sigma$ and $B = b \cdot \sigma$ with $\alpha = \|a\|$ and $\beta = \|b\|$. Now consider their exponentials $\exp(A)$ and $\exp(B)$ and their product $\exp(A) \exp(B)$:

$$\exp(A) \exp(B) = \left(\cosh \alpha I + \frac{\sinh \alpha}{\alpha} A \right) \left(\cosh \beta I + \frac{\sinh \beta}{\beta} B \right) \quad (14)$$

Notice the singular values of $\exp(A)$ and $\exp(B)$ are given by $\exp(\pm\alpha)$ and $\exp(\pm\beta)$. The singular values of their product, $\exp(A) \exp(B)$, are however equal to the positive square-root of the eigenvalues of

$$\exp(A) \exp(B) (\exp(A) \exp(B))^\dagger = \exp(A) \exp(B)^2 \exp(A). \quad (15)$$

Using the identity $\{A, B\} = AB + BA = 2(a \cdot b)I$ yields

$$\exp(A) \exp(B)^2 \exp(A) \quad (16)$$

$$= \cosh(2\alpha) \cosh(2\beta) I + \frac{\sinh(2\alpha) \cosh(2\beta)}{\alpha} A + \frac{\sinh(2\beta)}{\beta} B \dots \quad (17)$$

$$\dots + \sinh(2\alpha) \sinh(2\beta) \cos \theta I + \frac{2 \sinh^2 \alpha \sinh(2\beta) \cos \theta}{\alpha} A \quad (18)$$

While there is an asymmetry in $\exp(A) \exp(B)^2 \exp(A)$ with respect to $A \leftrightarrow B$, this asymmetry disappears when calculating the eigenvalues; the eigenvalues are given by $\eta \pm \delta$ where $\eta^2 - \delta^2 = 1$ and

$$\eta = \cosh(2\alpha) \cosh(2\beta) + \sinh(2\alpha) \sinh(2\beta) \cos \theta, \quad (19)$$

$$\delta = \frac{1}{4} \left(8 \sinh^2(2\alpha) \sinh^2(2\beta) \cos(2\theta) + 8 \sinh(4\alpha) \sinh(4\beta) \cos(\theta) + \dots \right. \quad (20)$$

$$\left. \dots + 3 \cosh(4(\alpha - \beta)) + 3 \cosh(4(\alpha + \beta)) + 2 \cosh(4\alpha) + 2 \cosh(4\beta) - 10 \right)^{\frac{1}{2}}. \quad (21)$$

Alternatively, defining $\gamma' \geq 0$ such that

$$\cosh(2\gamma') = \eta, \quad \sinh(2\gamma') = \delta, \quad (22)$$

One has that the singular values of $\exp(A) \exp(B)$ are equal to $\exp(\pm\gamma')$. Once again, there exists only one parameter that is not fixed by specifying the singular values ($\exp(\pm\alpha)$, $\exp(\pm\beta)$) of $\exp(A)$ and $\exp(B)$; specifically the angle $\theta \in [0, 2\pi)$ fixes the singular values of $\exp(A) \exp(B)$ by

$$\cosh(2\gamma') = \cosh(2\alpha) \cosh(2\beta) + \sinh(2\alpha) \sinh(2\beta) \cos \theta, \quad (23)$$

which is the *hyperbolic* cosine law with Gaussian curvature -4 (again note that the angle θ used is not interior to the hyperbolic triangle, hence the difference in minus sign).

Also note the singular values $\exp(\pm\gamma')$ of $\exp(A) \exp(B)$ at the values of $\theta \in \{0, \pi\}$ are

$$\theta = 0 \implies \exp(\pm(\alpha + \beta)), \quad \theta = \pi \implies \exp(\pm(\alpha - \beta)). \quad (24)$$

1.1.3 Thompson's conjecture via angle modifications

Thompson's conjecture in the context of 2×2 matrices corresponds to the following conjecture: there exists a pair (A, B) of traceless 2×2 Hermitian matrices with spectra $(\pm\alpha, \pm\beta)$

such that $A + B$ has spectra $\pm\gamma$ if and only if there exists a pair (X, Y) of unit determinant positive-definite Hermitian matrices with singular values $(\exp(\pm\alpha), \exp(\pm\beta))$ such that XY has singular values $\exp(\pm\gamma)$.

By the above analysis, letting $(X, Y) = (\exp A, \exp B)$ does *not* prove the conjecture; while the singular values of $(\exp A, \exp B)$ are $(\exp(\pm\alpha), \exp(\pm\beta))$ as desired, the singular values of their product $\exp A \exp B$ are equal to $\exp(\pm\gamma')$ which, in general, are not equal to $\exp(\pm\gamma)$ (the singular values of $\exp(A + B)$). This difference is to be expected because in general, one has

$$\exp(A) \exp(B) = \exp(A + B + \frac{1}{2}[A, B] + \dots) \neq \exp(A + B), \quad (25)$$

or alternatively

$$\exp(A) \exp(B)^2 \exp(A) \neq \exp(2(A + B)). \quad (26)$$


Nevertheless, there is a relationship between γ and γ' provided by the Golden-Thompson inequality $\text{Tr}(\exp(A' + B')) \leq \text{Tr}(\exp(A') \exp(B'))$. Letting $A' = 2A$ and $B' = 2B$ implies

$$\text{Tr}(\exp(2(A + B))) \leq \text{Tr}(\exp(2A) \exp(2B)) = \text{Tr}(\exp(A) \exp(B)^2 \exp(A)). \quad (27)$$

Or by dividing by 2, one obtains

$$\cosh(2\gamma) \leq \cosh(2\gamma') \implies \gamma \leq \gamma', \quad (28)$$

which essentially states that the side-lengths, γ' , of hyperbolic triangles are generally longer than their Euclidean counter-parts, γ :



Therefore, in order to prove Thompson's conjecture in the context of 2×2 matrices, one needs to pick the pair (X, Y) to be something other than $(\exp(A), \exp(B))$ but with the *same* singular values. The solution comes from choosing a different pair $(\tilde{A}, \tilde{B}) = (\tilde{a} \cdot \sigma, \tilde{b} \cdot b)$ of Hermitian matrices with the *same* spectra $(\pm\alpha, \pm\beta)$ as (A, B) such that $\exp(\tilde{A}) \exp(\tilde{B})$ has as singular values $\exp(\pm\gamma)$ where $(\pm\gamma)$ are the eigenvalues of $A + B$.

As we have shown, the only freedom in this choice for $(\tilde{A}, \tilde{B})$ is the angle $\tilde{\theta}$ between $\tilde{a}$ and $\tilde{b}$:

$$\cos \tilde{\theta} = \frac{\tilde{a} \cdot \tilde{b}}{\alpha \beta}. \quad (30)$$

The value $\tilde{\theta}$ determines the value $\tilde{\gamma}'$, and thus the singular values of $\exp(\tilde{A}) \exp(\tilde{B})$ by the previously derived formula:

$$\cosh(2\tilde{\gamma}') = \cosh(2\alpha) \cosh(2\beta) + \sinh(2\alpha) \sinh(2\beta) \cos \tilde{\theta}. \quad (31)$$





Setting $\tilde{\gamma}' = \gamma$ and solving for $\tilde{\theta}$ yields

$$\cos \tilde{\theta} = \cosh(2\gamma) \text{csch}(2\alpha) \text{csch}(2\beta) - \tanh(2\alpha) \tanh(2\beta). \quad (32)$$

Recalling that γ itself is fixed by the angle θ between the original (A, B) , one obtains a formula for $\tilde{\theta}$ in terms of (θ, α, β) :

$$\cos \tilde{\theta} = \cosh(2\sqrt{\alpha^2 + \beta^2 + 2\alpha\beta \cos \theta}) \operatorname{csch}(2\alpha) \operatorname{csch}(2\beta) - \tanh(2\alpha) \tanh(2\beta). \quad (33)$$

In summary, when the spectra α, β , and γ are fixed and there exists a pair of Hermitian matrices (A, B) with spectra $(\pm\alpha, \pm\beta)$ whose sum $A + B$ has spectra $\pm\gamma$ (thus there exists a value for θ), the above formula describes precisely how to reduce the angle θ between A and B to the value $\tilde{\theta}$ obtaining another pair of Hermitian matrices $(\tilde{A}, \tilde{B})$ such that $(\exp(\tilde{A}), \exp(\tilde{B}))$ have singular values $(\exp(\pm\alpha), \exp(\pm\beta))$ and their product $\exp(\tilde{A})\exp(\tilde{B})$ has singular values $\exp(\pm\gamma)$.

 $\lambda(A) = \pm\alpha$ $\lambda(B) = \pm\beta$ $\lambda(A + B) = \pm\gamma$	 $\sigma(\exp(A)) = \exp(\pm\alpha)$ $\sigma(\exp(B)) = \exp(\pm\beta)$ $\sigma(\exp(A)\exp(B)) = \exp(\pm\gamma')$	$\gamma \leq \gamma'$ $\tilde{\gamma} \leq \tilde{\gamma}'$
(34)		
 $\lambda(\tilde{A}) = \pm\alpha$ $\lambda(\tilde{B}) = \pm\beta$ $\lambda(\tilde{A} + \tilde{B}) = \pm\tilde{\gamma}$	 $\sigma(\exp(\tilde{A})) = \exp(\pm\alpha)$ $\sigma(\exp(\tilde{B})) = \exp(\pm\beta)$ $\sigma(\exp(\tilde{A})\exp(\tilde{B})) = \exp(\pm\gamma)$	$\tilde{\gamma} \leq \gamma$ $\tilde{\gamma}' = \gamma$

Of course, this only proves one half of Thompson's conjecture in this setting, one also needs to prove the reverse: given positive definite (X, Y, XY) with singular values $\exp(\pm\alpha), \exp(\pm\beta)$ and $\exp(\pm\gamma)$, find the corresponding Hermitian matrices (A, B) . This can be done simply by defining $(\tilde{A}, \tilde{B}) = (\log X, \log Y)$ with separated by angle $\tilde{\theta}$ and then modifying this angle to obtain (A, B) .

1.1.4 Scaling hyperbolic curvature

In the previous section, it was noted that in general $\exp(A)\exp(B) \neq \exp(A + B)$ because of the Baker-Campbell-Hausdorff term:

$$\exp(A)\exp(B) = \exp(A + B + \frac{1}{2}[A, B] + \dots) \neq \exp(A + B). \quad (35)$$

If $(A, B) \in \mathcal{O}_\alpha \times \mathcal{O}_\beta$ is replaced with $(sA, sB) \in \mathcal{O}_{s\alpha} \times \mathcal{O}_{s\beta}$ above for some $s \geq 0$, then one obtains

$$\exp(sA)\exp(sB) = \exp(sA + sB + \frac{s^2}{2}[A, B] + \dots) \neq \exp(s(A + B)). \quad (36)$$

Consequently, one still should expect the singular values of $\exp(sA)\exp(sB)$ to be different from the singular values, $\exp(\pm s\gamma)$, of $\exp(s(A+B))$. Nevertheless, for small $s \approx 0$, one might expect an approximation of the form

$$\exp(sA)\exp(sB) \approx \exp(s(A+B)). \quad (37)$$

and thus the singular values of $\exp(sA)\exp(sB)$ might approximate $\exp(\pm s\gamma)$ in a suitable sense. This intuition is also encoded in the hyperbolic geometry; rescaling $(\alpha, \beta) \mapsto (s\alpha, s\beta)$ amounts to changing the singular values $\exp(\pm \gamma')$ of $\exp(A)\exp(B)$ to $\exp(\pm s\gamma'_s)$ where γ'_s is given by the formula

$$\cosh(2s\gamma'_s) = \cosh(2s\alpha)\cosh(2s\beta) + \sinh(2s\alpha)\sinh(2s\beta)\cos\theta. \quad (38)$$

This formula is the hyperbolic cosine law for triangles in hyperbolic space whose Gaussian curvature is $-4s^2$, meaning γ'_s is the side length of a hyperbolic triangle whose opposite sides are α, β with opposite angle $\pi - \theta$ in a space with Gaussian curvature $-4s^2$. For small $s \approx 0$, space is approximately Euclidean: $\cosh(2sx) = 1 + 2s^2x^2 + O(s^4)$ and $\sinh(2sx) = 2sx + O(s^3)$ together yield

$$1 + 2s^2(\gamma'_s)^2 = 1 + 2s^2\alpha^2 + 2s^2\beta^2 + 4s^2\alpha\beta\cos\theta + O(s^3). \quad (39)$$

Simplifying yields the expected result that $\gamma'_s \approx \gamma$ when $s \approx 0$:

$$(\gamma'_s)^2 = \alpha^2 + \beta^2 + 2\alpha\beta\cos\theta + O(s) = \gamma^2 + O(s). \quad (40)$$

1.2 General theory

[sec:general_theory](#)

1.2.1 The additive case & symplectic geometry

Horn's problem is concerned with a fundamental question about how the eigenvalues of a sum of Hermitian matrices relates to the eigenvalues of the individual summands. More concretely, let $\mathcal{O}_\lambda$ denote the set of Hermitian matrices with fixed eigenvalues λ , Horn's problem is to determine for a given triple of eigenvalues (α, β, γ) if there exists a pair of Hermitian matrices $(A, B) \in \mathcal{O}_\alpha \times \mathcal{O}_\beta$ such that $A + B \in \mathcal{O}_\gamma$.

Modern approaches to Horn's problem use tools and results from *symplectic geometry* and, in particular, the notion of *Hamiltonian K-manifold*. Hamiltonian K -manifolds are symplectic manifolds (M, ω) together with an action of a Lie group K on M which is both symplectomorphic and for which there exists a K -equivariant map $\Phi : M \rightarrow \mathfrak{k}^*$ called the *moment map* that satisfies the condition

$$d\langle \Phi, \xi \rangle = \iota_{\xi_M} \omega \quad (41)$$

for each element $\xi \in \mathfrak{k}$ of the Lie algebra of K where ξ_M denotes the vector field on M induced by the K -action.

This theory is applicable to Horn's problem when $K = \mathrm{U}(n)$ because it is possible to identify the set of $n \times n$ Hermitian matrices with the dual $\mathfrak{k}^* = \mathfrak{u}(n)^*$ of the Lie algebra $\mathfrak{k} = \mathfrak{u}(n)$ of $n \times n$ skew-Hermitian matrices such that the orbits $\mathcal{O}_\lambda$ of the coadjoint action of $K = \mathrm{U}(n)$ on $\mathfrak{u}(n)^*$ labelled by λ are identified with sets of Hermitian matrices with fixed eigenvalues λ .

In general, each coadjoint orbit $M = \mathcal{O}_\lambda$ of K on $\mathfrak{k}^*$ labelled by λ can be equipped with a symplectic form (the Kirillov-Kostant-Souriau symplectic form) such that the inclusion map

$\mathcal{O}_\lambda \hookrightarrow \mathfrak{k}^*$ is the corresponding moment map. The property of Hamiltonian K -manifolds, (M, ω, Φ) which is crucial for Horn's problem is the following: given a pair (M_1, ω_1, Φ_1) and (M_2, ω_2, Φ_2) of Hamiltonian K -manifolds the diagonal action of K on $M_1 \times M_2$ given by

$$k \cdot (m_1, m_2) = (k \cdot m_1, k \cdot m_2) \quad (42)$$

turns the product symplectic manifold $(M_1 \times M_2, \omega_1 + \omega_2)$ into a Hamiltonian K -manifold with moment map $\Phi : M_1 \times M_2 \rightarrow \mathfrak{k}^*$ given by

$$\Phi(m_1, m_2) = \Phi_1(m_1) + \Phi_2(m_2). \quad (43)$$

In other words, moment maps behave additivity with respect to the Cartesian product of symplectic manifolds.

In the context of Horn's problem, when $K = \mathrm{U}(n)$ acts on $M = \mathcal{O}_\alpha \times \mathcal{O}_\beta$ via the diagonal coadjoint action, the corresponding $\mathrm{U}(n)$ -equivariant moment map Φ sends the pair $(A, B) \in \mathcal{O}_\alpha \times \mathcal{O}_\beta$ to their sum $\Phi(A + B) = A + B$.

1.2.2 The multiplicative case & Poisson geometry

As it turns out, there is a generalization of the theory of symplectic manifolds and Hamiltonian K -manifolds with $\mathfrak{k}^*$ -valued moment maps to the setting of Poisson manifolds and Hamiltonian K -manifolds with K^* -valued moment maps where K^* is a group dual to K in a sense which will be defined shortly. As we will see, while $\mathfrak{k}^*$ -valued moment maps, $\Phi : M \rightarrow \mathfrak{k}^*$, can be added together in a commutative way, K^* -valued moment maps, $\Psi : M \rightarrow K^*$, can be multiplied together in a non-commutative way. It is in this sense that Alekseev, Meinrenken, and Woodward generalize the additive Horn problem to a multiplicative one [AMW01].

In this generalized setting, the manifold M upon which K acts need not be a symplectic manifold, but can be a Poisson manifold equipped with a Poisson bracket $\{\cdot, \cdot\}_M : C^\infty(M) \times C^\infty(M) \rightarrow C^\infty(M)$ or equivalently a Poisson bivector field $\pi_M \in \Gamma(\wedge^2 TM)$. This is a generalization of a symplectic manifold in the sense that every symplectic form $\omega_M \in \Gamma(\wedge^2 T^*M)$ on M gives rise to Poisson bivector $\pi_M \in \Gamma(\wedge^2 TM)$ on M but not all Poisson bivectors arise in this way.

In addition to the generalized structure on the manifold M , the Lie group K itself is equipped with its own Poisson structure $\pi_K \in \Gamma(\wedge^2 TK)$ which is compatible with group multiplication of K turning (K, π_K) into a Poisson Lie group. It turns out that Lu and Weinstein constructively showed how every connected, compact Lie group, like $K = \mathrm{U}(n)$, admits of a non-trivial Poisson bivector π_K , called the *Lu-Weinstein Poisson-Lie* structure, making (K, π_K) into a Poisson-Lie group [LW90] (see also [LS91]). In general, Poisson Lie groups (K, π_K) have very rich structure. For instance, viewing the Poisson bivector field $\pi_K \in \Gamma(\wedge^2 TK)$ as a map $K \rightarrow \mathfrak{k} \wedge \mathfrak{k}$, its linearization is a map $\delta : \mathfrak{k} \rightarrow \mathfrak{k} \wedge \mathfrak{k}$ which is a 1-cocycle on $\mathfrak{k}$ and for which the dual map $\delta^* : \mathfrak{k}^* \wedge \mathfrak{k}^* \rightarrow \mathfrak{k}^*$ defines a Lie bracket on $\mathfrak{k}^*$ turning $(\mathfrak{k}, \mathfrak{k}^*)$ into the structure of a *Lie bialgebra*. The connected, simply connected Lie group K^* with whose Lie algebra is $\mathfrak{k}^*$ with bracket δ^* is called the *Poisson dual* of K .

Moreover, there is a way to equip the vector space $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{k}^*$ with a Lie algebra structure such that it is the Lie algebra of a Lie group G (called the *double* for K) with closed subgroups K and K^* such that the map $K \times K^* \rightarrow G$ sending $(k, b) \in K \times K^*$ to G is a diffeomorphism. Typically, the diffeomorphism $K \times K^* \rightarrow G$ does not respect the group structure of G in the sense that $(k_1 b_1)(k_2 b_2) \neq (k_1 k_2)(b_1 b_2)$. Nevertheless, the diffeomorphism defines so-called

dressing actions of K on K^ (where $k \in K$ sends $b \in K^*$ to some element denoted by $b^k \in K^*$), and K^* on K (where $b \in K^*$ sends $k \in K$ to some element denoted by $k^b \in K$) such that for all $k \in K$ and $b \in K^*$,*

$$bk = k^b b^k. \quad (44)$$

With respect to the Lu-Weinstein Poisson structure on K , the double G is equivalent to the *complexification* of K with Iwasawa decomposition $G = K^{\mathbb{C}} = KAN$ and the dual K^* is $K^* = AN$.

Now, when a Poisson manifold (M, π_M) is acted upon by a Poisson-Lie group (K, π_K) there are additional compatibility conditions which generalize the moment map conditions above. The moment map itself if generalized to a map $\Psi : M \rightarrow K^*$, called the a *K^* -valued moment*. At the very least, $\Psi : M \rightarrow K^*$ must be a Poisson map (in that it respects the Poisson structures on π_M and π_{K^*}). Using the Poisson map, it is possible to define a Lie algebra action of $\mathfrak{k}$ on M sending every $\xi \in \mathfrak{k}$ to the vector field ξ_M defined in the following way. First from $\xi \in \mathfrak{k} \simeq T_e K^*$ construct a 1-form on K^* , denoted by $\tilde{\xi} \in \Gamma(T^* K^*)$ which, for all vector fields $\zeta \in \Gamma(TK^*)$, satisfies at $b \in K^*$

$$\langle \tilde{\xi}_b, \zeta_b \rangle = \langle (R_{b^{-1}})_* \zeta, \xi \rangle \quad (45)$$

where $(R_{b'})_{*,b} : T_b K^* \rightarrow T_{bb'} K^* \simeq \mathfrak{k}^*$ is the pushforward of the right-translate $R_{b'} : K^* \rightarrow K^*$ (sending b to bb').² Second, pullback this 1-form on K^* through the Poisson map $\Psi : M \rightarrow K^*$ to obtain the 1-form on M denoted by $\Psi^* \tilde{\xi} \in \Gamma(T^* M)$. Finally, using the Poisson bivector $\pi_M \in \Gamma(\wedge^2 TM)$ on M , identify $\Psi^* \tilde{\xi}$ with the unique vector field on M , denoted by ξ_M , satisfying for all 1-forms $\alpha \in \Gamma(T^* M)$ the equality

$$\langle \alpha, \xi_M \rangle = \langle \alpha \wedge \Psi^* \tilde{\xi}, \pi_M \rangle. \quad (46)$$

It can be shown that the map sending $\xi \in \mathfrak{k}$ to $\xi_M \in \Gamma(TM)$ just defined is a Lie algebra action of $\mathfrak{k}$ on M in the sense that $[\xi, \eta]_M = [\xi_M, \eta_M]$. If this action integrates to a full K -action on M , then the triple (M, π_M, Ψ) is referred to as a *Hamiltonian K -manifold with K^* -valued moment map*.³

As previously mentioned, K^* -valued moment maps can be multiplied in the following way. If (M_1, π_1, Ψ_1) and (M_2, π_2, Ψ_2) are a pair of Hamiltonian K -manifolds with K^* -valued moment maps, the Poisson map $\Psi : M_1 \times M_2 \rightarrow K^*$ defined by

$$\Psi(m_1, m_2) = \Psi_1(m_1) \Psi_2(m_2) \quad (47)$$

is indeed a K^* -valued moment map for the Poisson manifold $(M_1 \times M_2, \pi_1 + \pi_2)$ but with respect to *twisted (diagonal) K -action*

$$k \cdot (m_1, m_2) = (k \cdot m_1, k^{\Psi_1(m_1)} \cdot m_2) \quad (48)$$

Just as coadjoint action of K on $\mathfrak{k}^*$ produces coadjoint orbits $\mathcal{O}_\lambda \subset \mathfrak{k}^*$ with the structure of a symplectic manifold such that the inclusion map is the corresponding $\mathfrak{k}^*$ -valued moment map, the dressing action of K on K^* produces dressing orbits $\mathcal{D}_{\lambda'} \subset K^*$ with Poisson structure such that the inclusion map is the corresponding K^* -valued moment map. This gives rise to a multiplicative generalization of the “additive” Horn-type problem: given a triple $(\mathcal{D}_\alpha, \mathcal{D}_\beta, \mathcal{D}_\gamma)$ of dressing orbits, does there exist a pair $(X, Y) \in \mathcal{D}_\alpha \times \mathcal{D}_\beta$ such that $XY \in \mathcal{D}_\gamma$?

²Note that $\tilde{\xi} = \theta^R(\cdot \otimes \xi)$ where θ^R is the right Maurer-Cartan form.

³Note that Lamb [Lam05], in summarizing Alekseev, Meinrenken, and Woodward’s paper [Lam05], uses the more manageable terminology of a *Poisson K -space*. I am not sure if this terminology is standard.

1.2.3 The linearization theorem

Under suitable conditions, it was shown by [Ale97] that there exists an equivalence between the category of Hamiltonian K -manifolds (M, ω, Φ) with $\mathfrak{k}^*$ -valued moment maps $\Phi : M \rightarrow \mathfrak{k}^*$ and the category of Hamiltonian K -manifolds (M, π, Ψ) with K^* -valued moment maps $\Psi : M \rightarrow K^*$ [Ale97]. Generally speaking, this process is referred to as *linearization*.

The conditions require K to be a compact connected Lie group with Lu-Weinstein Poisson-Lie structure and for the Poisson structure π on M to be symplectic in the sense that it comes from symplectic form Ω on M . Associated to K are its dual $K^* = AN$ and its double $G = K^{\mathbb{C}} = KAN$. Furthermore, in [AMW01], Alekseev, Meinrenken, and Woodward define an invertible K -equivariant map

$$E : \mathfrak{k}^* \rightarrow K^* \quad (49)$$

together with a 1-form $\beta \in \Omega(\mathfrak{k}^*)$. The two objects E and β depend implicitly upon a choice of a K -invariant inner product $B : \mathfrak{k} \times \mathfrak{k} \rightarrow \mathbb{R}$ on $\mathfrak{k}$ (think negative scalar multiples of the Killing form) used to identify $\mathfrak{k}^*$ with $\mathfrak{a} \oplus \mathfrak{n}$ in the Iwasawa decomposition of the Lie algebra $\mathfrak{g}$ of the double G . Using these maps, the following theorem holds.

Theorem 1 (Linearization). *Let M be a K -manifold and let $\pi \in \Gamma(\wedge^2 TM)$ be the Poisson bivector induced by a symplectic form $\Omega \in \Gamma(\wedge^2 T^*M)$ on M . Let $\Psi : M \rightarrow K^*$ be a map and define*

$$\Phi : M \rightarrow \mathfrak{k}^*, \quad \omega \in \Gamma(\wedge^2 T^*M) \quad (50)$$

in terms of E, β (depending on K), and Ψ, Ω (depending on (M, Ω)) by

$$\Phi = E^{-1} \circ \Psi, \quad \omega = \Omega - d\Phi^*\beta. \quad (51)$$

Then (M, π, Ψ) is a Hamiltonian K -manifold with K^ -valued moment map Ψ if and only if (M, ω, Φ) is a Hamiltonian K -manifold with $\mathfrak{k}^*$ -valued moment map.*

The linearization theorem plays a crucial role in Alekseev, Meinrenken, and Woodward's proving a kind of equivalence between the additive-type Horn problems to the multiplicative-type Horn problems. Suppose that (M_1, π_1, Ψ_1) and (M_2, π_2, Ψ_2) are a pair of Hamiltonian K -manifolds with K^* -valued moment maps (where the Poisson bivector π_1 comes from a symplectic form Ω_1 on M_1 and likewise π_2 comes from Ω_2 on M_2). As we have seen before, $(M_1 \times M_2, \pi_1 + \pi_2, \Psi_1 \Psi_2)$ is a Hamiltonian K -manifold with K^* -valued moment with respect to the twisted diagonal action of K on $M_1 \times M_2$. Similarly, one could consider $(M_2 \times M_1, \pi_1 + \pi_2, \Psi_2 \Psi_1)$ where the order of Ψ_1 and Ψ_2 is swapped. By applying the linearization theorem to the triplet of K^* -valued moment maps, one obtains three symplectic manifolds with $\mathfrak{k}^*$ -valued moment maps:

$$(M_1, \omega_1, \Phi_1) \Leftarrow (M_1, \pi_1, \Psi_1) \quad (52)$$

$$(M_2, \omega_2, \Phi_2) \Leftarrow (M_2, \pi_2, \Psi_2) \quad (53)$$

$$(M_1 \times M_2, \omega_{1,2}, \Phi_{1,2}) \Leftarrow (M_1 \times M_2, \pi_1 + \pi_2, \Psi_1 \Psi_2) \quad (54)$$

What is the relationship between $\Phi_{1,2}$ and $\Phi_1 + \Phi_2$? In general, they are distinct maps, meaning

$$\Phi_{2,1}(m_2, m_1) \neq \Phi_1(m_1) + \Phi_2(m_2) \neq \Phi_{1,2}(m_1, m_2), \quad (55)$$

but they are both moment maps, albeit associated to two different symplectic forms (namely $\omega_{1,2}$ and $\omega_1 + \omega_2$). Amazingly, by appropriately rescaling the E and β maps by a parameter

$s \geq 0$, Alekseev, Meinrenken, and Woodward [AMW01] derive a smooth family $\{\omega_{1,2}^s\}_{s \in [0,1]}$ of symplectic forms of $M_1 \times M_2$ together with a smooth family of $\mathfrak{k}^*$ -valued moment maps $\{\Phi_{1,2}^s\}_{s \in [0,1]}$ such that the $s = 0$ case recovers the sum of linearizations

$$(M_1 \times M_2, \omega_{1,2}^0, \Phi_{1,2}^0) = (M_1 \times M_2, \omega_1 + \omega_2, \Phi_1 + \Phi_2) \quad (56)$$

whereas the $s = 1$ case recovers the linearization of the product, i.e.

$$(M_1 \times M_2, \omega_{1,2}^1, \Phi_{1,2}^1) = (M_1 \times M_2, \omega_{1,2}, \Phi_{1,2}). \quad (57)$$

Furthermore, using a Moser isotropy argument, they prove that there exists a K -equivariant family of symplectomorphisms $\{\phi_s\}_{s \in [0,1]}$ (symplectomorphic from $(M_1 \times M_2, \omega_1 + \omega_2)$ to $(M_1 \times M_2, \omega_{1,2}^s)$, and K -equivariant from diagonal action to an s -scaled twisted diagonal action) such that

$$\omega_{1,2}^s = (\phi_s)^*(\omega_1 + \omega_2), \quad \Phi_{1,2}^s = (\phi_s)^*(\Phi_1 + \Phi_2). \quad (58)$$

As a corollary, one can show, in a certain sense, that the space of solutions to the additive Horn-type problem is symplectomorphic to the space of solutions to the space of solutions of the multiplicative Horn-type problem. This proves Thompson's conjecture when one takes $K = U(n)$.

Reference notes

- A popular textbook on quantum groups with an emphasis on the connection to Poisson manifolds, Lie bialgebras and their quantizations by Chari and Pressley [CP95]
- The general subject of multivariate trace inequalities involving exponentials of sums of Hermitian and products of their exponentials is, of course, intimately related. See for instance the paper by Sutter, Berta, and Tomamichel [SBT17].
- Also the article [FT14] by Forrester and Thompson which describes the connection to hyperbolic geometry quite explicitly.
- Course notes on Poisson geometry with an eye toward quantization by Ciccoli and Witkowski [CW06]
- An excellent expository overview of the big picture by Knutson [Knu00]
- The masters essay/thesis of Backens (TODO: check if the pdf Matthias sent to me is the final version) [Bac10]
- An excellent introduction to these ideas is the recent masters thesis on various aspects of additive and multiplicative Horn problems in connection to Vinnikov curves and the Lax conjecture due to Hess and supervised by S. Kuhlmann and M. Infusino [Hes20]
- The PhD thesis of Esposito (from the maths department here in Copenhagen) serves as a introduction to the world of Poisson momentum maps and the notion of a “quantum” momentum map as a deformation of the ordinary “classical” momentum map (where deformation is in the sense of deformation quantization) [Esp12].
- Alekseev [Ale97]
- Papers contained results pertaining to the generalization of Duistermaat-Heckman measures for Hamiltonian group actions with group-valued moment maps by Alekseev, Meinrenken, and Woodward [AMW99; AMW00]
- The proof of Thompson’s conjecture in the style of Alekseev, Meinrenken, and Woodward is the “linearization paper” due to [AMW01].
- See also the related article due to Lamb which summarizes the ideas of [AMW01] while also providing more technical background [Lam05]
- Lu and Evens extend the proof technique of [AMW01] to a larger class of Lie groups [LE03]
- Alekseev, Podkopaeva, and Szenes [APS17a]
- Alekseev, Podkopaeva, and Szenes [APS17b]
- Alekseev and Meinrenken [AM07]
- Deforming the symplectic volume by the s parameter Alekseev et al. [Ale+18]
- Klyachko [Kly00]
- Christandl [Chr08]

- Christandl, Şahinoğlu, and Walter [CŞW18]
- Kirillov and Reshetikhin [KR90]
- Usage of Wigner’s $6j$ symbol to compute asymptotic errors rates for the promise problem of distinguishing between two possible “change points” for a sequence of unknown pure states to transition from one unknown type to another [AH11]
- A great summary by Roberts [Rob02]; also the original result for the asymptotics of the $6j$ symbols due to Roberts [Rob99]
- Coquereaux and Zuber explicitly outline the link between the orbital measures (i.e. the induced distributions in the sense of Klyachko’s random walk approach) and the magnitude of the stretched Littlewood-Richardson coefficients [CZ18]. See also the papers [CMZ20; CMZ19; Zub18].
- A recent article by McSwiggen is an excellent introduction to the Harish-Chandra integral [McS21]. The proof of the Harish-Chandra formula provide by McSwiggen using the heat equation methods from Fegan’s 1978 paper [Feg78] seems very promising for deriving a connection to the character formula for the norm of the recoupling operator!
- See also Coquereaux’s later paper which promotes the alternative, albeit equivalent, usage of the integer-valued gon and tet functions in place of the rational-valued theta (triangular) and irrational-valued $6j$ symbols [Coq23]. Most relevant for this project is [Coq23, Section IV D] which discusses the apparent lack of substantial generalization from $SU(2)$ to $SU(N)$ (“*apart from very few special cases (low dimension of irreps and no multiplicity) worked out by physicists at the time of the eightfold way, we are not aware of any published general result*”).
- The recent paper by Alekseev, Morozov, and Sleptsov (different Alekseev!) covers the case of some multiplicity free formulas for the quantum $6j$ symbols for $U_q(\mathfrak{sl}_N)$ [AMS20]
- Boalch [Boa07]
- Taylor and Woodward [TW06]
- Henriques and Kamnitzer [HK05]
- Octahedron Recurrence/Rule Robbins and Rumsey Jr [RR86] applied to the “Hive” ring Knutson, Tao, and Woodward [KTW03], and also Henriques and Kamnitzer [HK06]
- The integral formula for the square of the $SU(2)$ $6j$ symbol can be performed by an appropriate change of variables [FL03]. See also other papers by Laurent Freidel and collaborators that are relevant [FK00; FNR07; FKL10].
- Speyer [Spe05]
- Generalizations of Horn’s problem to the product of unitary matrices [AW97; Bel08; BK16]

- Bhatia [Bha01]
- History account offered by Daftuar and Hayden [DH05], and technical review article by Fulton [Ful00].
- Connecting group-valued moment maps to spherical, flat, and hyperbolic tetrahedron via Minkowski's theorem from the reconstruction of flat tetrahedron Haggard, Han, and Riello. See also the comments in [HHR16, Section 11] on the utility of *quasi-poisson* (controlled failure of the Jacobi identity) manifolds and group-valued moment maps.
- Foundational results concerning the relationships between the edge lengths, face areas, dihedral and solid angles of a tetrahedron by Richardson [Ric02]
- Potential application to inverse kinematics of 3-link arms in three dimensions? (Thanks Harold!)
- **TODO:** track down pdfs for the original triple of papers by Thompson and Therianos titled “On the singular values of matrix products” I, II, and III from 1973
- **TODO:** track down pdfs for the original triple of papers by Thompson and Freede titled “On the eigenvalues of sums of Hermitian matrices” I, II, and III from 1973
- **TODO:** track down a pdf for Thompson's 1988 “Matrix Spectral Inequalities” referenced by Klyachko
- Thompson and Therianos [TT73] present a number of interesting counter-examples to the question of whether the inequalities pertaining to the spectra of general matrices can be deduced from looking at the spectra of diagonal matrices (both in the additive and multiplicative cases).
- [Bar03]

2 Asymptotic recoupling from spectral estimation

2.1 Converting Hermitian matrices to density matrices

The approach taken in this section and the next is based on extending the application of Keyl and Werner's spectral estimation theorem [KW01] to Horn's problem (i.e. relations between the spectra of A , B and $A + B$) provided by Christandl [Chr08] to the generalised setting concerning relations between the spectra of A , B , C , $A + B$, $B + C$, $A + C$, and $A + B + C$. Our approach will parallel the techniques developed by Christandl, Şahinoğlu, and Walter [CSW18] for the purposes of tackling the spectral marginal problem.

Consider the following decision problems.

[prob:thp](#)

Problem 1 (Trivariate Horn's problem). Let $d \in \mathbb{N}$ and let

$$(s_A, s_B, s_C, s_{AB}, s_{BC}, s_{AC}, s_{ABC}) \in (\mathbb{R}^d)^{\times 7} \quad (59)$$

be given as input. Does there exist a triple of $d \times d$ Hermitian matrices, A, B and C , such that

$$\begin{aligned} \text{spec}(A) &= s_A, & \text{spec}(A + B) &= s_{AB}, \\ \text{spec}(B) &= s_B, & \text{spec}(B + C) &= s_{BC}, & \text{spec}(A + B + C) &= s_{ABC}, \\ \text{spec}(C) &= s_C, & \text{spec}(A + C) &= s_{AC}. \end{aligned} \quad (60)$$

There are two properties of the spectral map $\text{spec} : \text{Herm}_d \rightarrow \mathbb{R}^d$ worth emphasizing, namely for all $X \in \text{Herm}_d$ and $\lambda \in \mathbb{R}$,

$$\text{spec}(\lambda X) = \lambda \text{spec}(X), \quad \text{spec}(X + \lambda I_d) = \text{spec}(X) + \lambda \text{spec}(I_d) \quad (61)$$

where $u := \text{spec}(I_d) = (1, \dots, 1)$.

Consequently, the collection of tuples of spectra $(s_A, \dots, s_{ABC})$ for which there exists Hermitian matrices A, B and C satisfying the above equations is closed under scalar multiplication and affine shifts by scalar multiples of $(u, \dots, u)$.

[lem:shifting_scaling](#)

Lemma 2. *Given a $d \times d$ Hermitian operator, X , then the operator*

$$T_\lambda(X) := \frac{X + \lambda I_d}{\text{Tr}(X) + \lambda d}, \quad (62)$$

satisfies $\text{Tr}(T_\lambda(X)) = 1$ and, for sufficiently large $\lambda \in \mathbb{R}$, $T_\lambda(X) > 0$ is positive definite.

Therefore, a solution to Problem 1 can always be extracted from a solution to the following related problem.

[prob:dmhp](#)

Problem 2 (Trivariate density matrix Horn's problem). Let $d \in \mathbb{N}$, let $p = (p_A, p_B, p_C) \in \Delta_3$ have strictly positive entries and let

$$(r_A, r_B, r_C, r_{AB}, r_{BC}, r_{AC}, r_{ABC}) \in (\Delta_d^{\geq})^{\times 7} \quad (63)$$

be given as input. Does there exist a triple of $d \times d$ density matrices, ρ_A, ρ_B and ρ_C such

that

$$\begin{aligned}
\text{spec}(\rho_A) &= r_A, & \text{spec}\left(\frac{p_A \rho_A + p_B \rho_B}{p_A + p_B}\right) &= r_{AB}, \\
\text{spec}(\rho_B) &= r_B, & \text{spec}\left(\frac{p_B \rho_B + p_C \rho_C}{p_B + p_C}\right) &= r_{BC}, \\
\text{spec}(\rho_C) &= r_C, & \text{spec}\left(\frac{p_A \rho_A + p_C \rho_C}{p_A + p_C}\right) &= r_{AC}, \\
\text{spec}(p_A \rho_A + p_B \rho_B + p_C \rho_C) &= r_{ABC}.
\end{aligned} \tag{64}$$

Remark 1. The conversion from Problem 1 to Problem 2 lies with picking $\lambda \in \mathbb{R}$ sufficiently large (see Lemma 2) and then letting

$$(\rho_A, \rho_B, \rho_C) := (T_\lambda(A), T_\lambda(B), T_\lambda(C)), \tag{65}$$

and furthermore letting

$$(p_A, p_B, p_C) := \frac{(\text{Tr}(A) + \lambda d, \text{Tr}(B) + \lambda d, \text{Tr}(C) + \lambda d)}{\text{Tr}(A + B + C) + 3\lambda d} \tag{66}$$

such that, for example,

$$\frac{p_A \rho_A + p_B \rho_B}{p_A + p_B} = \frac{A + B + 2\lambda d}{\text{Tr}(A + B) + 2\lambda d}, \tag{67}$$

and furthermore if r_{AB} is related to $s_{AB} = \text{spec}(A + B)$ by the formula

$$r_{AB} = \text{spec}\left(\frac{p_A \rho_A + p_B \rho_B}{p_A + p_B}\right) = \frac{1}{\text{Tr}(A + B) + 2\lambda d} (s_{AB} + 2\lambda \text{spec}(I_d)) \tag{68}$$

2.2 Spectral Estimation

The following lemma [Chr08] places an upper bound on the value of $\text{Tr}(P_\lambda \rho^{\otimes n})$ for given λ and ρ .

Lemma 3. Let $\mathcal{H} \cong \mathbb{C}^d$ and let $\mathcal{H}^{\otimes n} \cong \bigoplus_{\lambda \vdash_{dn}} V_\lambda \otimes U_\lambda^d$ be the Schur-Weyl decomposition and P_λ the Hermitian projection operator acting on $\mathcal{H}^{\otimes n}$ which projects into the isotypic component $V_\lambda \otimes U_\lambda^d$. Then for any density operator ρ with spectrum r ,

$$\text{Tr}(P_\lambda \rho^{\otimes n}) \leq (n+1)^{d(d-1)/2} \exp(-nD(\bar{\lambda} \| r)), \tag{69}$$

where $D(\cdot \| \cdot)$ is the Kullback-Leibler divergence and $\bar{\lambda}_j := \lambda_j/n$.

Corollary 4. Let $\epsilon > 0$, let $n \in \mathbb{N}$ and let $\Lambda_{<\epsilon}^{nr}$ denote the set of integer partitions, $\lambda \vdash k$, satisfying $D(\bar{\lambda} \| r) < \epsilon$. Define

$$E_{<\epsilon}^{nr} = \sum_{\lambda \in \Lambda_{<\epsilon}^{nr}} P_\lambda. \tag{70}$$

For any ρ with spectrum $\text{spec}(\rho) = r$,

$$\text{Tr}(E_{<\epsilon}^{nr} \rho^{\otimes n}) \geq 1 - (n+1)^{(d+2)(d-1)/2} \exp(-n\epsilon). \tag{71}$$

2.3 Recoupling operators

Definition 1 (F-Symbols). Let $U_\alpha^d, U_\beta^d, U_\gamma^d, U_\mu^d, U_\nu^d$ and U_λ^d be irreducible highest-weight representations of $SU(d)$. Define the *recoupling operator* or *F-symbol* as the linear operator

$$\left\{ \begin{array}{ccc} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{array} \right\} : C_\mu^{\alpha,\beta} \otimes C_\lambda^{\mu,\gamma} \rightarrow C_\lambda^{\alpha,\nu} \otimes C_\nu^{\beta,\gamma}, \quad (72)$$

between the multiplicity spaces of irreps of $SU(d)$ defined as

$$\left\{ \begin{array}{ccc} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{array} \right\} := \frac{1}{\text{Tr}(U_\lambda^d)} \begin{array}{c} \begin{array}{ccc} & C_\lambda^{\alpha,\nu} & \\ & \nearrow U_\nu^d & \\ C_\mu^{\alpha,\beta} & & C_\nu^{\beta,\gamma} \\ & \nwarrow U_\beta^d & \\ & C_\lambda^{\mu,\gamma} & \end{array} \\ \begin{array}{ccc} & U_\alpha^d & \\ & \nearrow U_\lambda^d & \\ & U_\mu^d & \\ & \nwarrow U_\gamma^d & \\ & C_\mu^{\alpha,\beta} & \end{array} \end{array} . \quad (73)$$

2.4 Ensemble Approximation Lemma

TODO: this section needs lots of work

Lemma 5. Let $(p_A, p_B, p_C) \in \Delta_3$ be a probability distribution, let $n \in \mathbb{N}$ and let X be any S_n -equivariant and positive semi-definite operator acting on $\mathcal{H}^{\otimes n}$. Then there exists an explicit triple of sequences $(k_A^{(n)}, k_B^{(n)}, k_C^{(n)})_{n \in \mathbb{N}}$ such that for all $n \in \mathbb{N}$, $k_A^{(n)} + k_B^{(n)} + k_C^{(n)} = n$ and

$$\lim_{n \rightarrow \infty} \left(\frac{k_A^{(n)}}{n}, \frac{k_B^{(n)}}{n}, \frac{k_C^{(n)}}{n} \right) = (p_A, p_B, p_C), \quad (74)$$

and furthermore for all $n \in \mathbb{N}$,

$$\text{Tr} [X(p_A \rho_A + p_B \rho_B + p_C \rho_C)^{\otimes n}] \geq (n+1)^{-2} \text{Tr} \left[X(\rho_A^{\otimes k_A^{(n)}} \otimes \rho_B^{\otimes k_B^{(n)}} \otimes \rho_C^{\otimes k_C^{(n)}}) \right] \quad (75)$$

Proof. The S_n -equivariance of X simplifies the non-commutative multinomial expansion of $(p_A \rho_A + p_B \rho_B + p_C \rho_C)^{\otimes n}$ to obtain an ordinary multinomial expansion of the form:

$$\begin{aligned} & \text{Tr} [X(p_A \rho_A + p_B \rho_B + p_C \rho_C)^{\otimes n}] \\ &= \sum_{\substack{k_1, k_2, k_3=0 \\ k_1+k_2+k_3=n}}^n \binom{n}{k_1, k_2, k_3} p_A^{k_1} p_B^{k_2} p_C^{k_3} \text{Tr} [X(\rho_A^{\otimes k_1} \otimes \rho_B^{\otimes k_2} \otimes \rho_C^{\otimes k_3})]. \end{aligned} \quad (76)$$

Since the summand is always non-negative (because $X \geq 0$ by assumption) the overall sum can be lower-bounded by the value of any particular summand. Consider the term, indexed by (k_1, k_2, k_3) , which (approximately) maximizes the probability

$$m(k_1, k_2, k_3) := \binom{n}{k_1, k_2, k_3} p_A^{k_1} p_B^{k_2} p_C^{k_3}. \quad (77)$$

Specifically, take (k'_1, k'_2, k'_3) to be a rational approximation to the distribution (p_A, p_B, p_C) satisfying

$$\|(k'_1, k'_2, k'_3) - n(p_A, p_B, p_C)\|_1 \leq 3. \quad (78)$$

Then “it can be shown” that $m(k'_1, k'_2, k'_3)$, viewed as a function of n , cannot decay to zero faster than one over the number of summands up to a constant independent of n **TODO**. A crude approximation of $m(k'_1, k'_2, k'_3) \geq (n+1)^{-2}$ is then obtained from ignoring the condition $k_1 + k_2 + k_3 = n$. $\square$

$$\sum_{i=1}^k p_i \operatorname{Tr}[E\rho_i] = \operatorname{Tr}[E\rho] \quad (79)$$

$$\sum_{i,j=1}^k p_i p_j \operatorname{Tr}[E(\rho_i \otimes \rho_j)] = \operatorname{Tr}[E(\rho \otimes \rho)] \quad (80)$$

Let $\tilde{E}$ be S_n -equivariant, then

$$\sum_{n_1, \dots, n_k} m_p(n_1, \dots, n_k) \operatorname{Tr}[\tilde{E}(\rho_1^{\otimes n_1} \otimes \dots \otimes \rho_k^{\otimes n_k})] = \operatorname{Tr}[\tilde{E}\rho^{\otimes n}] \quad (81)$$

{eq:symmetric-multinomial-expansion}

where m_p is the multinomial distribution

$$m_p(n_1, \dots, n_k) = \binom{n}{n_1, \dots, n_k} p_1^{n_1} \dots p_k^{n_k} \quad (82)$$

If $q = (n_1/n, \dots, n_k/n)$ is the empirical distribution, then a classical result due to [San61; DZ10] places the following bounds on m_p in terms of the relative entropy or Kullback-Liebler divergence [Kul97] of q from p :

$$(n+1)^{-k} \exp(-nD(q\|p)) \leq m_p(n_1, \dots, n_k) \leq \exp(-nD(q\|p)). \quad (83)$$

Since $\operatorname{Tr}[\tilde{E}(\rho_1^{\otimes n_1} \otimes \dots \otimes \rho_k^{\otimes n_k})] \in [0, 1]$, one obtains from Eq. (81) the following lower bound where $(n_1, \dots, n_k)$ is chosen such that $D(q\|p) \leq C_p n^{-2}$ where C_p is a constant independent of n but dependent upon p (TODO: Prove that this choice can be made using reverse Pinsker’s inequality or by constructing q explicitly):

$$\operatorname{Tr}[\tilde{E}\rho^{\otimes n}] \geq (n+1)^{-k} \exp(-C_p/n) \operatorname{Tr}[\tilde{E}(\rho_1^{\otimes nq_1} \otimes \dots \otimes \rho_k^{\otimes nq_k})] \quad (84)$$

In fact, one should be able to do much better by noting that the number of partitions $(n_1, \dots, n_k)$ of n (with at most k rows) is upper bounded by the number of compositions (partitions without order) which admits the following upper bound

$$\binom{n+k-1}{n} = \frac{n+k-1}{k-1} \frac{n+k-2}{k-2} \dots \frac{n+1}{1}, \quad (85)$$

$$= \left(1 + \frac{n}{k-1}\right) \left(1 + \frac{n}{k-2}\right) \dots \left(1 + \frac{n}{1}\right), \quad (86)$$

$$\leq (1+n)^{k-1}. \quad (87)$$

The maximum likelihood q (taken with respect to m_p and also dependent on n) must be greater than the average likelihood which can be lower-bounded by the reciprocal of the above upper bound and therefore:

$$\mathrm{Tr}[\tilde{E}\rho^{\otimes n}] \geq (n+1)^{-(k-1)} \mathrm{Tr}[\tilde{E}(\rho_1^{\otimes nq_1} \otimes \cdots \otimes \rho_k^{\otimes nq_k})] \quad (88)$$

Purifying all the states involved separately ($\rho_i = \mathrm{Tr}_{\mathcal{Z}}(\psi_i)$ with ψ_i rank one projection) yields

$$\mathrm{Tr}[(\tilde{E} \otimes I^{\otimes n})\psi^{\otimes n}] \geq (n+1)^{-(k-1)} \mathrm{Tr}[(\tilde{E} \otimes I^{\otimes n})(\psi_1^{\otimes nq_1} \otimes \cdots \otimes \psi_k^{\otimes nq_k})] \quad (89)$$


2.5 From states to recoupling

In this section we repeat the proof technique used in [Chr08] to prove the existence of a sextuple of convergent sequences of Young diagrams satisfying the recoupling operator lower bound.


Let $(p_1, p_2, p_3) \in \Delta_3$, let $\rho_1, \rho_2, \rho_3, \rho_4, \rho_5, \rho_6$ be a sextuple of $d \times d$ density operators (with spectra $r_i = \mathrm{spec}(\rho_i)$), and suppose that the density matrices $\{\rho_1, \dots, \rho_6\}$ are subject to the constraints

$$\begin{aligned} (p_1 + p_2)\rho_4 &= p_1\rho_1 + p_2\rho_2, \\ (p_2 + p_3)\rho_5 &= p_2\rho_2 + p_3\rho_3, \\ (p_1 + p_2 + p_3)\rho_6 &= p_1\rho_1 + p_2\rho_2 + p_3\rho_3. \end{aligned} \quad \begin{array}{l} \text{\texttt{\textcolor{blue}{eq:sextuple_config_constraints}}} \\ (90) \end{array}$$

In the case of $d = 2$, this configuration can be depicted faithfully as a configuration of points in $\mathbb{R}^3$:



$$(91)$$



$$(92)$$

Using spectra estimation theorem, we have for each $i \in \{1, \dots, 6\}$, $k_i \in \mathbb{N}$, and $\lambda \vdash k_i$,

$$\mathrm{Tr}(P_\lambda \rho_i^{\otimes k_i}) \leq (k_i + 1)^{d(d-1)/2} \exp(-k_i D(\bar{\lambda} \| r_i)). \quad (93)$$

In particular, if $D(\bar{\lambda} \| r_i) < \epsilon$, then

$$\text{Tr}(P_\lambda \rho_i^{\otimes k_i}) \leq (k_i + 1)^{d(d-1)/2} \exp(-k_i \epsilon). \quad (94)$$

Furthermore, the ensemble approximation lemma applied to $\rho_4 = x\rho_1 + (1-x)\rho_2$ with $x = \frac{p_1}{p_1+p_2}$ and $\mu \vdash k_1 + k_2 = k_4$ (satisfying $k_4 x \approx k_1$ and $k_4(1-x) \approx k_2$)

$$\text{Tr}(P_\mu \rho_4^{\otimes k_4}) = \text{Tr}(P_\mu (x\rho_1 + (1-x)\rho_2)^{\otimes k_4}) \geq \frac{1}{k_4 + 1} \text{Tr}(P_\mu \rho_1^{\otimes k_1} \otimes \rho_2^{\otimes k_2}) \quad (95)$$

Combining this with spectral estimation yields the following bound

$$\text{Tr}(P_\mu \rho_1^{\otimes k_1} \otimes \rho_2^{\otimes k_2}) \leq (k_4 + 1) \text{Tr}(P_\mu \rho_4^{\otimes k_4}) \leq (k_4 + 1)^{d(d-1)/2+1} \exp(-k_4 D(\bar{\mu} \| r_4)). \quad (96)$$

Let $k_6 = k_1 + k_2 + k_4$ and note that $k_i \leq k_6$ for all i . Then one obtains the following collection of upper bounds

$$\text{Tr}(P_\alpha \rho_1^{\otimes k_1}) \leq (k_1 + 1)^{d(d-1)/2} \exp(-k_1 D(\bar{\alpha} \| r_1)), \quad (97)$$

$$\text{Tr}(P_\beta \rho_2^{\otimes k_2}) \leq (k_2 + 1)^{d(d-1)/2} \exp(-k_2 D(\bar{\beta} \| r_2)), \quad (98)$$

$$\text{Tr}(P_\gamma \rho_3^{\otimes k_3}) \leq (k_3 + 1)^{d(d-1)/2} \exp(-k_3 D(\bar{\gamma} \| r_3)), \quad (99)$$

$$\text{Tr}(P_\mu \rho_1^{\otimes k_1} \otimes \rho_2^{\otimes k_2}) \leq (k_4 + 1)^{d(d-1)/2+1} \exp(-k_4 D(\bar{\mu} \| r_4)), \quad (100)$$

$$\text{Tr}(P_\nu \rho_2^{\otimes k_2} \otimes \rho_3^{\otimes k_3}) \leq (k_5 + 1)^{d(d-1)/2+1} \exp(-k_5 D(\bar{\nu} \| r_5)), \quad (101)$$

$$\text{Tr}(P_\lambda \rho_1^{\otimes k_1} \otimes \rho_2^{\otimes k_2} \otimes \rho_3^{\otimes k_3}) \leq (k_6 + 1)^{d(d-1)/2+2} \exp(-k_6 D(\bar{\lambda} \| r_6)). \quad (102)$$

At the moment, there is an important subtlety which has NOT been addressed. In order to apply the ensemble approximation lemma it must be the case that the vales of k_1, k_2 , and k_3 are reasonably good approximations of p_1, p_2 , and p_3 in the sense that the multinomial probability $m_p(k_1, k_2, k_3)$ is taken larger than the crude, albeit polynomial, lower bound of $(k_1 + k_2 + k_3 + 1)^{-2}$. Additionally, in order to apply the ensemble approximation lemma in parallel three times as we have done here, one must show that these reasonably good approximations can be made for (p_1, p_2, p_3) , $(p_1, p_2)/(p_1 + p_2)$ and $(p_2, p_3)/(p_2 + p_3)$ for the *same* tuple of integers (k_1, k_2, k_3) . While this is not too difficult to demonstrate, we omit a proof of this claim here.

At this stage, consider the following sextuple of projection operators actings on $\mathcal{H}^{\otimes k_6}$,

$$\Pi_\alpha := P_\alpha \otimes I^{\otimes k_2+k_3}, \quad (103)$$

$$\Pi_\beta := I^{\otimes k_1} \otimes P_\beta \otimes I^{\otimes k_3}, \quad (104)$$

$$\Pi_\gamma := I^{\otimes k_1+k_2} \otimes P_\gamma, \quad (105)$$

$$\Pi_\mu := P_\mu \otimes I^{\otimes k_3}, \quad (106)$$

$$\Pi_\nu := I^{\otimes k_1} \otimes P_\nu, \quad (107)$$

$$\Pi_\lambda := P_\lambda. \quad (108)$$

It can be shown (using applications of Schur's lemma for $U(d)$ for example) that almost all pairs of projection operators in $\{\Pi_\alpha, \Pi_\beta, \dots, \Pi_\lambda\}$ are commuting; the only exception is the

pair $\{\Pi_\mu, \Pi_\nu\}$. Define the following projection operators,

$$\Pi_{\alpha\beta\gamma} = \Pi_\alpha \Pi_\beta \Pi_\gamma, \quad (109)$$

$$\Pi_{\alpha\beta\gamma\lambda} = \Pi_\alpha \Pi_\beta \Pi_\gamma \Pi_\lambda, \quad (110)$$

$$\Pi_{\alpha\beta\gamma\lambda}^\mu = \Pi_\mu \Pi_\alpha \Pi_\beta \Pi_\gamma \Pi_\lambda, \quad (111)$$

$$\Pi_{\alpha\beta\gamma\lambda}^\nu = \Pi_\nu \Pi_\alpha \Pi_\beta \Pi_\gamma \Pi_\lambda. \quad (112)$$

Additionally, define

$$\Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3} := \rho_1^{\otimes k_1} \otimes \rho_2^{\otimes k_2} \otimes \rho_3^{\otimes k_3}. \quad (113)$$

In any case, the upper bounds above are equivalent to the following upper bounds:

$$\text{Tr}(\Pi_\alpha \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \leq (k_1 + 1)^{d(d-1)/2} \exp(-k_1 D(\bar{\alpha} \| r_1)), \quad (114)$$

$$\text{Tr}(\Pi_\beta \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \leq (k_2 + 1)^{d(d-1)/2} \exp(-k_2 D(\bar{\beta} \| r_2)), \quad (115)$$

$$\text{Tr}(\Pi_\gamma \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \leq (k_3 + 1)^{d(d-1)/2} \exp(-k_3 D(\bar{\gamma} \| r_3)), \quad (116)$$

$$\text{Tr}(\Pi_\mu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \leq (k_4 + 1)^{d(d-1)/2+1} \exp(-k_4 D(\bar{\mu} \| r_4)), \quad (117)$$

$$\text{Tr}(\Pi_\nu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \leq (k_5 + 1)^{d(d-1)/2+1} \exp(-k_5 D(\bar{\nu} \| r_5)), \quad (118)$$

$$\text{Tr}(\Pi_\lambda \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \leq (k_6 + 1)^{d(d-1)/2+2} \exp(-k_6 D(\bar{\lambda} \| r_6)). \quad (119)$$

Henceforth let the polynomial prefactors be accounted for using the expression

$$L_c^d(k) := \left(\frac{d(d-1)}{2} + c \right) \log(k+1). \quad (120)$$

Using $L_c^d(k)$, the above bounds can be re-expressed as

$$\text{Tr}(\Pi_\alpha \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \leq \exp(-k_1 D(\bar{\alpha} \| r_1) + L_0^d(k_1)), \quad (121)$$

$$\text{Tr}(\Pi_\beta \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \leq \exp(-k_2 D(\bar{\beta} \| r_2) + L_0^d(k_2)), \quad (122)$$

$$\text{Tr}(\Pi_\gamma \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \leq \exp(-k_3 D(\bar{\gamma} \| r_3) + L_0^d(k_3)), \quad (123)$$

$$\text{Tr}(\Pi_\mu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \leq \exp(-k_4 D(\bar{\mu} \| r_4) + L_1^d(k_4)), \quad (124)$$

$$\text{Tr}(\Pi_\nu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \leq \exp(-k_5 D(\bar{\nu} \| r_5) + L_1^d(k_5)), \quad (125)$$

$$\text{Tr}(\Pi_\lambda \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \leq \exp(-k_6 D(\bar{\lambda} \| r_6) + L_2^d(k_6)). \quad (126)$$

An application of the Cauchy-Schwarz inequality to the Hilbert-Schmidt inner product $\langle X, Y \rangle = \text{Tr}(X^\dagger Y)$ with $X = \Omega^{\frac{1}{2}} \Pi_1$ and $Y = \Omega^{\frac{1}{2}} \Pi_2$ (where Π_i Hermitian projector and Ω positive-semi definite Hermitian) yields

$$|\text{Tr}(\Pi_1 \Omega \Pi_2)|^2 = \left| \text{Tr}((\Omega^{\frac{1}{2}} \Pi_1)^\dagger (\Omega^{\frac{1}{2}} \Pi_2)) \right|^2 \quad (127)$$

$$= \left| \langle \Omega^{\frac{1}{2}} \Pi_1, \Omega^{\frac{1}{2}} \Pi_2 \rangle \right|^2 \quad (128)$$

$$\leq \langle \Omega^{\frac{1}{2}} \Pi_1, \Omega^{\frac{1}{2}} \Pi_1 \rangle \langle \Omega^{\frac{1}{2}} \Pi_2, \Omega^{\frac{1}{2}} \Pi_2 \rangle \quad (129)$$

$$= \text{Tr}(\Pi_1 \Omega) \text{Tr}(\Pi_2 \Omega) \quad (130)$$

Additionally, if $[\Pi_1, \Pi_2] = 0$ then $|\text{Tr}(\Pi_1 \Omega \Pi_2)| = |\text{Tr}(\Pi_2 \Pi_1 \Pi_2 \Omega)| = \text{Tr}(\Pi_2 \Pi_1 \Pi_2 \Omega) \geq 0$. Similarly, taking $X = (\Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3})^{\frac{1}{2}} \Pi_{\alpha\beta\gamma\lambda}^\mu$ and $Y = (\Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3})^{\frac{1}{2}} \Pi_{\alpha\beta\gamma\lambda}^\nu$ yields

$$\left| \text{Tr}(\Pi_{\alpha\beta\gamma\lambda}^\mu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3} \Pi_{\alpha\beta\gamma\lambda}^\nu) \right|^2 \leq \text{Tr}(\Pi_{\alpha\beta\gamma\lambda}^\mu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \text{Tr}(\Pi_{\alpha\beta\gamma\lambda}^\nu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}). \quad (131)$$

Chaining these bounds together produces a (perhaps sub-optimal but nevertheless sufficient) upper bound

$$\left| \text{Tr}(\Pi_{\alpha\beta\gamma\lambda}^\mu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3} \Pi_{\alpha\beta\gamma\lambda}^\nu) \right|^4 \quad (132)$$

$$\leq \text{Tr}(\Pi_{\alpha\beta\gamma\lambda}^\mu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3})^2 \text{Tr}(\Pi_{\alpha\beta\gamma\lambda}^\nu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3})^2, \quad (133)$$

$$\leq \text{Tr}(\Pi_\mu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \text{Tr}(\Pi_\nu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \text{Tr}(\Pi_{\alpha\beta\gamma\lambda} \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3})^2, \quad (134)$$

$$\leq \text{Tr}(\Pi_\mu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \text{Tr}(\Pi_\nu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) \text{Tr}(\Pi_\alpha \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3})^{\frac{1}{2}} \times \dots \quad (135)$$

$$\times \text{Tr}(\Pi_\beta \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3})^{\frac{1}{2}} \text{Tr}(\Pi_\gamma \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3})^{\frac{1}{2}} \text{Tr}(\Pi_\lambda \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3})^{\frac{1}{2}} \quad (136)$$

Therefore,

$$\left| \text{Tr}(\Pi_{\alpha\beta\gamma\lambda}^\mu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3} \Pi_{\alpha\beta\gamma\lambda}^\nu) \right| \leq \exp(-k_6 I_{r_1, r_2, r_3, r_4, r_5, r_6}^{\alpha, \beta, \gamma, \mu, \nu, \lambda} + O(\log(k_6))) \quad (\text{eq:sextuple_total_upper_bound}) \quad (137)$$

where (under the hand-wavy approximations $p_i \approx k_i/k_6$) we define

$$\begin{aligned} I_{r_1, r_2, r_3, r_4, r_5, r_6}^{\alpha, \beta, \gamma, \mu, \nu, \lambda} &= \frac{p_1}{2} D(\bar{\alpha} \| r_1) + \frac{p_2}{2} D(\bar{\beta} \| r_2) + \frac{p_3}{2} D(\bar{\gamma} \| r_3) + \dots \\ &\dots + \frac{p_1 + p_2}{4} D(\bar{\mu} \| r_4) + \frac{p_2 + p_3}{4} D(\bar{\nu} \| r_5) + \frac{1}{2} D(\bar{\lambda} \| r_6) \end{aligned} \quad (138)$$

The crucial property of the rate function here is that $I_{r_1, r_2, r_3, r_4, r_5, r_6}^{\alpha, \beta, \gamma, \mu, \nu, \lambda} \geq 0$ always while $I_{r_1, r_2, r_3, r_4, r_5, r_6}^{\alpha, \beta, \gamma, \mu, \nu, \lambda} = 0$ only if $\bar{\alpha} = r_1, \bar{\beta} = r_2, \dots, \bar{\lambda} = r_6$.

At this stage, one can use the completeness property $\sum_\lambda \Pi_\lambda = I$ to prove

$$1 = \text{Tr}(\Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}) = \sum_{\alpha, \beta, \gamma, \mu, \nu, \lambda} \text{Tr}(\Pi_{\alpha\beta\gamma\lambda}^\mu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3} \Pi_{\alpha\beta\gamma\lambda}^\nu). \quad (139)$$

Taking absolute values (and using subadditivity) and applying Eq. (137) produces

$$1 \leq \sum_{\alpha, \beta, \gamma, \mu, \nu, \lambda} \left| \text{Tr}(\Pi_{\alpha\beta\gamma\lambda}^\mu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3} \Pi_{\alpha\beta\gamma\lambda}^\nu) \right|, \quad (140)$$

$$\leq \sum_{\alpha, \beta, \gamma, \mu, \nu, \lambda} \exp(-k_6 I_{r_1, r_2, r_3, r_4, r_5, r_6}^{\alpha, \beta, \gamma, \mu, \nu, \lambda} + O(\log(k_6))). \quad (141)$$

As there are at most

$$U(k_1, k_2, k_3) := (k_1+1)^{d-1} (k_2+1)^{d-1} (k_3+1)^{d-1} (k_1+k_2+1)^{d-1} (k_2+k_3+1)^{d-1} (k_1+k_2+k_3+1)^{d-1} \quad (142)$$

possible summands, the above equation implies there exists a (possibly non-unique) sextuple $\alpha', \beta', \gamma', \mu', \nu', \lambda'$ (implicitly dependent upon the k_i 's) such that

$$\exp(-k_6 I_{r_1, r_2, r_3, r_4, r_5, r_6}^{\alpha', \beta', \gamma', \mu', \nu', \lambda'} + O(\log(k_6))) \geq \left| \text{Tr}(\Pi_{\alpha'\beta'\gamma'\lambda'}^{\mu'} \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3} \Pi_{\alpha'\beta'\gamma'\lambda'}^{\nu'}) \right| \geq U(k_1, k_2, k_3)^{-1} \quad (\text{eq:occasionality_lower_bound}) \quad (143)$$

which implies

$$I_{r_1, r_2, r_3, r_4, r_5, r_6}^{\alpha', \beta', \gamma', \mu', \nu', \lambda'} \leq \frac{1}{k_6} \log U(k_1, k_2, k_3) + \frac{1}{k_6} O(\log(k_6)) = O(\log(k_6)/k_6) \quad (144)$$

which tends to zero as k_i 's tend to infinity. Therefore, this sequence of sextuples of integer partitions converge to the given spectra in the sense that

$$\bar{\alpha}' \rightarrow r_1, \quad \bar{\beta}' \rightarrow r_2, \quad \bar{\gamma}' \rightarrow r_3, \quad \bar{\mu}' \rightarrow r_4, \quad \bar{\nu}' \rightarrow r_5, \quad \bar{\lambda}' \rightarrow r_6. \quad (145)$$

In addition, by considering any purification of $\Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}$ one additionally obtains a bound on the *operator norm*, $\|\cdot\|_\infty$, of the product of all Π 's (noting that the operator norm also satisfies $\|X \otimes I\|_\infty = \|X\|_\infty$). This bound takes the form

$$\|\Pi_\mu \Pi_\nu \Pi_\alpha \Pi_\beta \Pi_\gamma \Pi_\lambda\|_\infty = \left\| \Pi_{\alpha\beta\gamma\lambda}^\mu \Pi_{\alpha\beta\gamma\lambda}^\nu \right\|_\infty \geq \left| \text{Tr}(\Pi_{\alpha\beta\gamma\lambda}^\mu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3} \Pi_{\alpha\beta\gamma\lambda}^\nu) \right|. \quad (146)$$

(The inequality comes from $\|XY\|_\infty^2 \geq \langle XYv, XYv \rangle \geq |\langle v, XYv \rangle|^2$ for all v with unit norm.) Furthermore, one can show by a detailed diagrammatic calculation that there exists appropriate isometries ι_I and ι_{II} such that

$$\iota_{II} \left(I_{U_\lambda^d} \otimes I_{V_\alpha} \otimes I_{V_\beta} \otimes I_{V_\gamma} \otimes \begin{Bmatrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{Bmatrix} \right) \iota_I^\dagger = \Pi_\mu \Pi_\nu \Pi_\alpha \Pi_\beta \Pi_\gamma \Pi_\lambda \quad (147)$$

which in turn implies an equality of operator norms

$$\left\| \begin{Bmatrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{Bmatrix} \right\|_\infty = \|\Pi_\mu \Pi_\nu \Pi_\alpha \Pi_\beta \Pi_\gamma \Pi_\lambda\|_\infty \quad (148)$$

By combining Eqs. (143), (146) and (148) we obtain a inverse polynomial lower bound for the operator norm of the recoupling operator of the form

$$\left\| \begin{Bmatrix} \alpha' & \beta' & \mu' \\ \gamma' & \lambda' & \nu' \end{Bmatrix} \right\|_\infty \geq U(k_1, k_2, k_3)^{-1} \geq (k_1 + k_2 + k_3 + 1)^{-6(d-1)} \quad (149)$$

In summary, we have shown that the existence of a certain configuration of density matrices implies a polynomial lower bound on the operator norm of the recoupling operator for $U(d)$ representations (modulo the $p_i \approx k_i/k_6$ approximation which needs to be justified),

Theorem 6. *Let $0 < p_1, p_2, p_3 < 1$ satisfying $p_1 + p_2 + p_3 = 1$ be given and let $(r_1, r_2, \dots, r_6) \in \Delta_d^{\downarrow; \times 6}$ be a sextuple of candidate spectra. If there exists a sextuple of density operators $\rho_1, \rho_2, \rho_3, \rho_4, \rho_5$ and ρ_6 with spectra $r_i = \text{spec}(\rho_i)$ satisfying Eq. (90), then there exists a sequence of sextuples $(\alpha^{(n)}, \beta^{(n)}, \gamma^{(n)}, \mu^{(n)}, \nu^{(n)}, \lambda^{(n)})$ of integer partitions satisfying both*

$$\left\| \begin{Bmatrix} \alpha^{(n)} & \beta^{(n)} & \mu^{(n)} \\ \gamma^{(n)} & \lambda^{(n)} & \nu^{(n)} \end{Bmatrix} \right\|_\infty \geq (n+1)^{-6(d-1)} \quad (150)$$

(for sufficiently large n) and the following convergences

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\alpha^{(n)}}{|\alpha^{(n)}|} &= r_1, & \lim_{n \rightarrow \infty} \frac{\beta^{(n)}}{|\beta^{(n)}|} &= r_2, & \lim_{n \rightarrow \infty} \frac{\gamma^{(n)}}{|\gamma^{(n)}|} &= r_3, \\ \lim_{n \rightarrow \infty} \frac{\mu^{(n)}}{|\mu^{(n)}|} &= r_4, & \lim_{n \rightarrow \infty} \frac{\nu^{(n)}}{|\nu^{(n)}|} &= r_5, & \lim_{n \rightarrow \infty} \frac{\lambda^{(n)}}{|\lambda^{(n)}|} &= r_6, \\ \lim_{n \rightarrow \infty} \frac{|\alpha^{(n)}|}{|\lambda^{(n)}|} &= p_1, & \lim_{n \rightarrow \infty} \frac{|\beta^{(n)}|}{|\lambda^{(n)}|} &= p_2, & \lim_{n \rightarrow \infty} \frac{|\gamma^{(n)}|}{|\lambda^{(n)}|} &= p_3. \end{aligned} \quad (151)$$

Remark 2. Note that the satisfaction of Eq. (150) for some $n \in \mathbb{N}$ automatically entails that $\begin{Bmatrix} \alpha^{(n)} & \beta^{(n)} & \mu^{(n)} \\ \gamma^{(n)} & \lambda^{(n)} & \nu^{(n)} \end{Bmatrix} \neq 0$ and thus $c_{\alpha,\beta}^\mu, c_{\beta,\gamma}^\nu, c_{\mu,\gamma}^\lambda, c_{\alpha,\nu}^\lambda \neq 0$ and thus the sequences of integer partitions also satisfy equality the following equality constraints

$$|\alpha^{(n)}| + |\beta^{(n)}| = |\mu^{(n)}|, \quad (152)$$

$$|\beta^{(n)}| + |\gamma^{(n)}| = |\nu^{(n)}|, \quad (153)$$

$$|\alpha^{(n)}| + |\beta^{(n)}| + |\gamma^{(n)}| = |\lambda^{(n)}|. \quad (154)$$

2.6 From recoupling to states

In this section, our objective is to prove the converse of the above theorem. Specifically, if the norm of a sequence of recoupling operators with components satisfying Eq. (151) admits of a polynomial lower bound, then there necessarily exists of a configuration of density matrices satisfying Eq. (90). In fact, this converse statement follows from a stronger claim that when there does not exist a configuration of density matrices satisfying the spectral constraints, every tuple of sequences of integer partitions satisfying Eq. (90) has recoupling operator norm eventually decaying at an exponential rate.

Consider the following operator defined by an integral over the normalized Hilbert-Schmidt measure $d\rho$ on density matrices and a given $k \in \mathbb{N}$:

$$\mathbb{E} [\sigma^{\otimes k}] = \int d\sigma \sigma^{\otimes k} \geq \binom{k+d^2-1}{k}^{-1} \rho^{\otimes k} \geq (k+1)^{-(d^2-1)} \rho^{\otimes k}. \quad (155)$$

The three-fold tensor product of the above operator is an integral over $\Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}$ and has the form

$$\mathbb{E} [\Omega_{\sigma_1, \sigma_2, \sigma_3}^{k_1, k_2, k_3}] := \int d\sigma_1 \int d\sigma_2 \int d\sigma_3 \Omega_{\sigma_1, \sigma_2, \sigma_3}^{k_1, k_2, k_3}. \quad (156)$$

The operator $\mathbb{E} [\Omega_{\sigma_1, \sigma_2, \sigma_3}^{k_1, k_2, k_3}]$ is equivariant with respect to the natural action of

$$S_{k_1} \times S_{k_2} \times S_{k_3} \times U(d) \times U(d) \times U(d). \quad (157)$$

Schur's lemma therefore implies that the operator $\mathbb{E} [\Omega_{\sigma_1, \sigma_2, \sigma_3}^{k_1, k_2, k_3}]$ is equal to a positive sum of projection operators of the form $\Pi_{\alpha\beta\gamma}$ (where $\alpha \vdash k_1, \beta \vdash k_2$, and $\gamma \vdash k_3$) with the following coefficients:

$$\mathbb{E} [\Omega_{\sigma_1, \sigma_2, \sigma_3}^{k_1, k_2, k_3}] = \sum_{\alpha, \beta, \gamma} R_{\alpha\beta\gamma} \Pi_{\alpha\beta\gamma} \quad \text{\textcolor{blue}{\{eq:mean_tensor_power_decomp\}}} \quad (158)$$

with coefficient $R_{\alpha\beta\gamma}$ given by a ratio of dimensions

$$R_{\alpha\beta\gamma} = \frac{1}{\dim(U_{(k_1)}^{d^2}) \dim(U_{(k_2)}^{d^2}) \dim(U_{(k_3)}^{d^2})} \frac{\dim(U_\alpha^d) \dim(U_\beta^d) \dim(U_\gamma^d)}{\dim(V_\alpha) \dim(V_\beta) \dim(V_\gamma)}. \quad (159)$$

Consequently, $\mathbb{E} [\Omega_{\sigma_1, \sigma_2, \sigma_3}^{k_1, k_2, k_3}]$ commutes with the projection operators Π_μ, Π_ν and Π_λ and thus also $\Pi_{\alpha\beta\gamma\lambda}^\mu$ and $\Pi_{\alpha\beta\gamma\lambda}^\nu$. In fact the following holds

$$\Pi_{\alpha\beta\gamma\lambda}^\mu \mathbb{E} [\Omega_{\sigma_1, \sigma_2, \sigma_3}^{k_1, k_2, k_3}] \Pi_{\alpha\beta\gamma\lambda}^\nu = R_{\alpha\beta\gamma} \Pi_{\alpha\beta\gamma\lambda}^\mu \Pi_{\alpha\beta\gamma\lambda}^\nu \quad (160)$$

Also since operator X acting on a D -dimensional vector space satisfies $\|X\|_\infty \leq \|X\|_{\text{HS}} \leq \sqrt{D} \|X\|_\infty$, one has a polynomial equivalence between the norms of the recoupling symbol

$$\left\| \begin{Bmatrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{Bmatrix} \right\|_\infty \leq \left\| \begin{Bmatrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{Bmatrix} \right\|_{\text{HS}} \leq \left(c_{\alpha,\beta}^\mu c_{\beta,\gamma}^\nu c_{\mu,\gamma}^\lambda c_{\alpha,\nu}^\lambda \right)^{\frac{1}{2}} \left\| \begin{Bmatrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{Bmatrix} \right\|_\infty \quad (161)$$

For the product of projectors, $\Pi_{\alpha\beta\gamma\lambda}^\mu \Pi_{\alpha\beta\gamma\lambda}^\nu$, the only (polynomial or better) bound one has is simply

$$\left\| \Pi_{\alpha\beta\gamma\lambda}^\mu \Pi_{\alpha\beta\gamma\lambda}^\nu \right\|_\infty \leq \left\| \Pi_{\alpha\beta\gamma\lambda}^\mu \Pi_{\alpha\beta\gamma\lambda}^\nu \right\|_{\text{HS}} = \sqrt{\text{Tr}(\Pi_{\alpha\beta\gamma\lambda}^\mu \Pi_{\alpha\beta\gamma\lambda}^\nu)} \quad (162)$$

Using the decomposition provided by Eq. (147), together with Eq. (137) and Eq. (158) yields

$$\left\| \begin{Bmatrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{Bmatrix} \right\|_{\text{HS}}^2 = \text{Tr} \left(\begin{Bmatrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{Bmatrix}^\dagger \begin{Bmatrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{Bmatrix} \right) \quad (163)$$

$$= \frac{\text{Tr} \left(\Pi_{\alpha\beta\gamma\lambda}^\mu \Pi_{\alpha\beta\gamma\lambda}^\nu \right)}{\dim(U_\lambda^d) \dim(V_\alpha) \dim(V_\beta) \dim(V_\gamma)} \quad (164)$$

$$= Q_{\alpha\beta\gamma\lambda} \text{Tr} \left(\Pi_{\alpha\beta\gamma\lambda}^\mu \mathbb{E} [\Omega_{\sigma_1, \sigma_2, \sigma_3}^{k_1, k_2, k_3}] \Pi_{\alpha\beta\gamma\lambda}^\nu \right) \quad (165)$$

$$\leq \sup_{\rho_1 \rho_2 \rho_3} Q_{\alpha\beta\gamma\lambda} \left| \text{Tr} \left(\Pi_{\alpha\beta\gamma\lambda}^\mu \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3} \Pi_{\alpha\beta\gamma\lambda}^\nu \right) \right| \quad (166)$$

$$\leq Q_{\alpha\beta\gamma\lambda} \exp(-k_6 \inf_{\rho_1 \rho_2 \rho_3} I_{r_1, r_2, r_3, r_4, r_5, r_6}^{\alpha, \beta, \gamma, \mu, \nu, \lambda} + O(\log(k_6))) \quad (167)$$

where the infimum is taken over all density matrices ρ_1, ρ_2, ρ_3 with $r_i = \text{spec}(\rho_i)$ for $i \in \{1, 2, 3\}$ and r_4, r_5, r_6 the spectra of the appropriate convex combinations of ρ_1, ρ_2, ρ_3 with weights determined by the relative sizes of the integer partitions involved (again here we are making the rather dubious substitution $p_i \approx k_i/k_6$ when substituting the function $I_{r_1, r_2, r_3, r_4, r_5, r_6}^{\alpha, \beta, \gamma, \mu, \nu, \lambda}$ **TODO: resolve this issue formally**). Also the scalar $Q_{\alpha\beta\gamma\lambda}$ is

$$Q_{\alpha\beta\gamma\lambda} := \frac{1}{R_{\alpha\beta\gamma} \dim(U_\lambda^d) \dim(V_\alpha) \dim(V_\beta) \dim(V_\gamma)} \quad (168)$$

$$= \frac{\dim(U_{(k_1)}^{d^2}) \dim(U_{(k_2)}^{d^2}) \dim(U_{(k_3)}^{d^2})}{\dim(U_\alpha^d) \dim(U_\beta^d) \dim(U_\gamma^d) \dim(U_\lambda^d)} \quad (169)$$

As we have seen prior, the rate function $I_{r_1, r_2, r_3, r_4, r_5, r_6}^{\alpha, \beta, \gamma, \mu, \nu, \lambda}$ only vanishes if all of the spectra match exactly. Furthermore, it can be shown that the infimum over ρ_1, ρ_2, ρ_3 of $I_{r_1, r_2, r_3, r_4, r_5, r_6}^{\alpha, \beta, \gamma, \mu, \nu, \lambda}$ is always bounded below by a constant $I^* > 0$ if there does not exist a configuration of density matrices satisfying the requirements. Consequently, we conclude the following theorem.

Theorem 7. Suppose $(\alpha^{(n)}, \beta^{(n)}, \gamma^{(n)}, \mu^{(n)}, \nu^{(n)}, \lambda^{(n)})$ is a sequence of integer partitions satisfying Eq. (151) for some $0 < p_1, p_2, p_3 < 1$ satisfying $p_1 + p_2 + p_3 = 1$ and $(r_1, r_2, \dots, r_6) \in \Delta_d^{\downarrow; \times 6}$. If there does not exist a collection of density matrices $\rho_1, \rho_2, \rho_3, \rho_4, \rho_5$ and ρ_6 with spectra $r_i = \text{spec}(\rho_i)$ satisfying Eq. (90), then there exists a constant $I^* > 0$ such that

$$\left\| \begin{Bmatrix} \alpha^{(n)} & \beta^{(n)} & \mu^{(n)} \\ \gamma^{(n)} & \lambda^{(n)} & \nu^{(n)} \end{Bmatrix} \right\|_\infty \leq \exp(-nI^* + O(\log n)) \quad (170) \quad \text{\textcolor{blue}{\small{eq:sequential_recoupling_norm_upper_bound}}}$$

Remark 3. It appears that I did not make explicit use of the following operator inequalities when proving the above theorem.

$$\mathbb{E} [\sigma^{\otimes k}] \geq (k+1)^{-(d^2-1)} \rho^{\otimes k}. \quad (171)$$

$$\mathbb{E} [\Omega_{\sigma_1, \sigma_2, \sigma_3}^{k_1, k_2, k_3}] \geq ((k_1+1)(k_2+1)(k_3+1))^{-d^2+1} \Omega_{\rho_1, \rho_2, \rho_3}^{k_1, k_2, k_3}. \quad (172)$$

2.7 From the perspective of strong duality

It would be very desirable to articulate the above result as an application of the strong duality and shifted/deformed strong duality results of non-commutative optimization theory [Bür+19], specifically the presentation offered by Franks and Walter [FW20].

Given a representation $\Phi : G \rightarrow \text{GL}(V)$ of complex reductive group $G = K_{\mathbb{C}}$ on a finite dimensional complex vector space V equipped with K -invariant inner product $\langle \cdot, \cdot \rangle$, the *moment map* associated to viewing $\mathbb{P}V$ as a symplectic, Hamiltonian G -manifold is (perhaps up to some constants) equal to

$$\mu(\psi)(X) = \text{Tr}(P_\psi \phi(X)) = \frac{\langle v, \phi(X)v \rangle}{\langle v, v \rangle}. \quad (173)$$

In the special case where $G = \text{SL}(d)$, $\mu(\psi) \in \mathfrak{isu}(d)^*$ can be identified with a $d \times d$ density operator ρ_ψ in a canonical way such that $\text{Tr}(\rho_\psi X) = \mu(\psi)(X)$ for all Hermitian operators $X \in \mathfrak{isu}(d)$. The (projective) capacity of $\psi \in \mathbb{P}V$ is

$$\text{cap}(\psi) = \inf_{g \in G} \text{Tr}(P_\psi \Phi(g^*g)) = \inf_{g \in G} \frac{\langle \Phi(g)v, \Phi(g)v \rangle}{\langle v, v \rangle} \quad (174)$$

In particular, it can be seen that $0 \leq \text{cap}(\psi) \leq 1$ and it can be shown that $\text{cap}(\psi) = 1$ if and only if $\mu(\psi) = 0$. Furthermore, define $\gamma_n(\psi)$ for $n \in \mathbb{N}$ to be equal to

$$\gamma_n(\psi) = \text{Tr}(P_\psi^{\otimes n} \Gamma_n)^{\frac{1}{n}} \quad (175)$$

where $\Gamma_n = \int_K dk \Phi(k)^{\otimes n}$ is the orthogonal projection operator onto the subspace of invariant vectors $(V^{\otimes n})^G$. The strong duality result in [FW20] simply states the following

$$\sup_{n \in \mathbb{N}} \gamma_n(\psi) = \text{cap}(\psi) \quad (176)$$

In particular, $\text{cap}(\psi) = 1$ if and only if $\mu(\psi) = 0$ if and only if $\limsup_{n \rightarrow \infty} \gamma_n(\psi) = 1$.

TODO

2.8 Comparison with S_k recoupling

A similar problem concerning sextuples of spectra is to characterize the admissible A, B, C , AB , BC , and ABC subsystem spectra of a tripartite mixed state ρ_{ABC} (or equivalently quadripartite pure state ψ_{ABCD}). This spectral problem was considered by Christandl, Şahinoğlu, and Walter [CSW18]. The approach adopted in [CSW18] was to use the spectral estimation method developed by Keyl and Werner [KW01] which essentially establishes an approximate correspondence between (i) density operators ρ with spectra r and (ii) representations of the symmetric group S_k labelled by Young diagrams λ such that $\lambda \approx kr$. This approximate

correspondence becomes exact in the limit of large symmetric groups S_k (specifically as $k \rightarrow \infty$).

The second half of the paper [CŞW18] considers how to reduce the partial Horn problem (concerning the spectra of Hermitian matrices $H_A, H_B, H_C, H_A + H_B, H_B + H_C$ and $H_A + H_B + H_C$) to a special case of the spectral marginal problem stated above. Their demonstration of this reduction consists of two complementary results: their first result shows how to embed the triple of Hermitian matrices H_A, H_B , and H_C into a tripartite quantum state in a manner which *preserves* the spectral information of the Hermitian matrices and their partial sums, while their second result demonstrates how to recover certain combinatorial data for the representation theory of the unitary group $U(k)$ as a special case of the analogous combinatorial data for the representation theory of the symmetric group S_k .

2.9 The representation-theoretic connection

The second result follows from a theorem due to [Nis00], which describes the restriction of the irrep of $U(k)$ labelled by $\lambda \vdash k$ to the symmetric subgroup $S_k \subseteq U(k)$, namely [Nis00, Theorem 2.3].

Theorem 8. *Let $k \in \mathbb{N}$ and let $\lambda \vdash k$. Let $\mathcal{U}_\lambda^{(k)}$ denote the irrep of $U(k)$ and V_λ the irrep of S_k . Given a partition μ with $\ell(\mu) \leq k$, let Υ_μ denote the Young subgroup $\Upsilon_\mu = S_{\mu_1} \times \cdots \times S_{\mu_k} \subseteq S_k$ and let $\tilde{\mu}$ denote the (not necessarily sorted) partition of $\ell(\mu)$ corresponding to the multiplicities of μ . Alternatively, $\tilde{\mu}$ can be characterised in terms of the normaliser of Υ_μ inside S_k via the formula*

$$N_{S_k}(\Upsilon_\mu) = \Upsilon_{\tilde{\mu}} \ltimes \Upsilon_\mu \quad (177)$$

Then the restriction of $\mathcal{U}_\lambda^{(k)}$ to S_k decomposes as follows:

$$\text{Res}_{S_k}^{U(k)} \mathcal{U}_\lambda^{(k)} \simeq \bigoplus_{\mu \vdash k, |\lambda|} \text{Ind}_{S_{k-\ell(\mu)} \times \Upsilon_{\tilde{\mu}}}^{S_k} (1 \otimes \mathcal{V}_\lambda^{\Upsilon_\mu}) \quad \text{\textcolor{blue}{(eq:nishiyama-decomp)}} \quad (178)$$

where 1 is the trivial representation of $S_{k-\ell(\mu)}$ and $\mathcal{V}_\lambda^{\Upsilon_\mu}$ is the Υ_μ -invariant subspace of V_λ supporting a representation of $\Upsilon_{\tilde{\mu}}$.

While the summands appearing on the right-hand side of Eq. (178) are quite complicated in general, it was shown in [CŞW18] that the irreducible representations of S_k , labelled by $\nu \vdash k$, which appear in the decomposition of $\text{Res}_{S_k}^{U(k)} \mathcal{U}_\lambda^{(k)}$ and constrained in a particular manner. Specifically, it can be shown that the size of first row ν_1 of ν can be lowered bounded by $k - |\lambda|$ when k is sufficiently large. Here is [CŞW18, Lemma 13]:

Theorem 9. *Consider a sorted partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_{\ell(\lambda)})$ of size $|\lambda|$. If $k \geq \lambda_1 + |\lambda|$, then it is possible to view λ as a Young sub-diagram of the Young diagram λ^{+k} defined by*

$$\lambda^{+k} := (k - |\lambda|, \lambda_1, \lambda_2, \dots, \lambda_{\ell(\lambda)}) \quad (179)$$

with length $\ell(\lambda^{+k}) = \ell(\lambda) + 1$. In this case, $k \geq \lambda_1 + |\lambda|$ and λ^{+k} is well-defined, and Eq. (178) implies

$$\text{Res}_{S_k}^{U(k)} \mathcal{U}_\lambda^{(k)} \simeq V_{\lambda^{+k}} \oplus \bigoplus_{\substack{\mu \vdash k \\ \mu_1 > k - |\lambda|}} V_\mu \otimes \text{Hom}(V_\mu, \text{Res}_{S_k}^{U(k)} \mathcal{U}_\lambda^{(k)}). \quad (180)$$

In other words the multiplicity of $V_{\lambda+k}$ appearing in $\text{Res}_{S_k}^{U(k)} \mathcal{U}_{\lambda}^{(k)}$ is one and all other irreps of S_k have first rows larger than $\lambda_1^{+k} = k - |\lambda|$.

On the other hand, if $k < \lambda_1 + |\lambda|$ is too small, then

$$\text{Hom}(V_{\mu}, \text{Res}_{S_k}^{U(k)} \mathcal{U}_{\lambda}^{(k)}) > 0 \implies \mu_1 > k - |\lambda|. \quad (181)$$

From this result, one can consider k sufficiently large to obtain a relationship between the Littlewood-Richardson coefficients,

$$c_{\gamma}^{\alpha, \beta} = \dim \text{Hom}(\mathcal{U}_{\gamma}^{(k)}, \mathcal{U}_{\alpha}^{(k)} \otimes \mathcal{U}_{\beta}^{(k)}) \quad (182)$$

(which are independent of k for $k \geq \max\{\ell(\alpha), \ell(\beta), \ell(\gamma)\}$) and the Kronecker coefficients

$$g_{\lambda}^{\mu, \nu} = \dim \text{Hom}(\mathcal{V}_{\lambda}, \mathcal{V}_{\mu} \otimes \mathcal{V}_{\nu}) \quad (183)$$

(which are implicitly dependent on k because $k = |\mu| = |\nu| = |\lambda|$).

Specifically, it can be shown that the Littlewood-Richardson coefficients can be obtained from the Kronecker coefficients by appropriately augmenting the Young diagrams involved:

Corollary 10. *Let α, β, γ be Young diagrams and let k be sufficiently large such that α^{+k}, β^{+k} and γ^{+k} are all well-defined. Then $c_{\gamma^{+k}}^{\alpha^{+k}, \beta^{+k}} = g_{\gamma^{+k}}^{\alpha^{+k}, \beta^{+k}}$.*

Remark 4. Note that this result indirectly implies a stability result for the Kronecker coefficients, namely that the value of $g_{\lambda^{+k}}^{\mu^{+k}, \nu^{+k}}$ is eventually independent of k and reaches the value $c_{\lambda_{-1}}^{\mu_{-1}, \nu_{-1}}$ where λ_{-1} is obtained from λ by removal of its first row.

2.10 The quantum-information connection

The purpose of this section is to describe the spectrum-preserving embedding of a triple of Hermitian operators into the space of tripartite density matrices used in [CŞW18] to reduce the partial-sum Horn problem to the quantum marginal problem.

2.10.1 Bipartite case

Before describing this embedding, it is worth considering an analogue of this embedding in the bivariate case; in particular, given a pair of $d \times d$ Hermitian operators σ_X and σ_Y (on $\mathbb{C}^d \simeq \text{span}\{|1\rangle, \dots, |d\rangle\}$) (which I will assume are density operators), is there a way to construct a pure tripartite quantum state $|\psi\rangle_{XYZ}$ which has marginals ρ_X, ρ_Y and ρ_Z satisfying some relationship to σ_X and σ_Y ?

Let σ_X and σ_Y have purifications $|\phi\rangle_{XZ}$ and $|\phi\rangle_{YZ}$ on a shared purifying space Z . Furthermore, let $p \in (0, 1)$ be given. Then define a control register C which is a two dimensional $\{|0\rangle_C, |1\rangle_C\}$ and define the pure state

$$|\psi\rangle_{CXYZ} = \sqrt{p} |0\rangle_C |0\rangle_Y |\phi\rangle_{XZ} + \sqrt{1-p} |1\rangle_C |0\rangle_X |\phi\rangle_{YZ} \quad (184)$$

The reduced state on XYZ is simply

$$\rho_{XYZ} = p |0\rangle \langle 0|_Y |\phi\rangle \langle \phi|_{XZ} + (1-p) |0\rangle \langle 0|_X |\phi\rangle \langle \phi|_{YZ} \quad (185)$$

In this way, one can interpret this state as a description of a controlled preparation wherein a Bernoulli trial (with probability p) is performed to prepare either $|\phi\rangle_{XZ}$ or $|\phi\rangle_{YZ}$. While

the outcome of the trial can be determined directly by a measurement of system C , it can also be determined by measuring either subsystem X or Y ; if the outcome on system X is zero $|0\rangle_X$, then it is known that C obtained outcome $|1\rangle_C$, if the outcome on system X is non-zero, then it is known that C obtained the complementary outcome $|0\rangle_C$. As this information is accessible from subsystem X , it is perhaps not surprising that the reduced state on subsystem XZ and Y has the form

$$\rho_{XZ} = p|\phi\rangle\langle\phi|_{XZ} + (1-p)|0\rangle\langle 0|_X \otimes (\sigma_Y)_Z, \quad (186)$$

$$\rho_X = p\sigma_X + (1-p)|0\rangle\langle 0|_X. \quad (187)$$

In particular, the information encoded in σ_Y is not accessible in subsystem X . The somewhat interesting case here is the reduced state on σ_Z :

$$\rho_Z = p(\sigma_X)_Z + (1-p)(\sigma_Y)_Z. \quad (188)$$

In otherwords, performing local measurements of system Z alone is indistinguishable from performing measurements on the convex combination of ρ_X and ρ_Y . Where this becomes interesting is when one considers the moment map of the ray defined by $|\psi\rangle_{CXYZ}$ with respect to the representation of $K = U(1) \times \text{SU}(d) \times \text{SU}(d) \times \text{SU}(d)$ on $\mathbb{C}^2 \otimes \mathbb{C}^{1+d} \otimes \mathbb{C}^{1+d} \otimes \mathbb{C}^d$ by the formula,

$$(e^{i\theta}, U_1, U_2, U_3) \mapsto \begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & U_1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & U_2 \end{pmatrix} \otimes U_3 \quad (189)$$

It is not too difficult to show that the moment map can be identified in a suitable sense with the data

$$\mu(\psi_{CXYZ}) \simeq (p, p\sigma_X, (1-p)\sigma_Y, p\sigma_X + (1-p)\sigma_Y). \quad (190)$$

Note that another equivalent option is to consider a representation without the control register C of the form

$$(e^{i\theta}, U_1, U_2, U_3) \mapsto \begin{pmatrix} e^{-i\theta} & 0 \\ 0 & U_1 \end{pmatrix} \otimes \begin{pmatrix} e^{i\theta} & 0 \\ 0 & U_2 \end{pmatrix} \otimes U_3. \quad (191)$$

Furthermore, the K -orbit of ψ_{CXYZ} under the moment map μ takes the form

$$\begin{aligned} \mu(K \cdot \psi_{CXYZ}) \simeq \{ & (p, pU_1\sigma_X U_1^{-1}, (1-p)U_2\sigma_Y U_2^{-1}, \\ & pU_3\sigma_X U_3^{-1} + (1-p)U_3\sigma_Y U_3^{-1}) \mid (U_1, U_2, U_3) \in \text{SU}(d)^{\times 3} \}. \end{aligned} \quad (192)$$

The unique intersection of this orbit under μ with the dominant Weyl chamber for K yields the spectral data (p, r_X, r_Y, r_{XY}) where r_{XY} is the spectra of $p\sigma_X + (1-p)\sigma_Y$.

The existence or non-existence of a pair σ_X and σ_Y satisfying this spectral constraint is therefore, by spectral estimation theory, encoded in the asymptotics of the appropriate representation theoretic data (in this case, the data is the Littlewood-Richardson coefficient $c_{\alpha, \beta}^\gamma$ where $|\alpha| \approx p|\gamma|$, $\bar{\alpha} \approx r_X$, $\bar{\beta} \approx r_Y$, and $\bar{\gamma} \approx r_{XY}$ [Chr08]).

2.10.2 Tripartite case

Given any collection of three $d \times d$ Hermitian matrices H_A , H_B , and H_C , one can assume without loss of generality that $H_A, H_B, H_C \geq 0$ are all positive semidefinite and normalized such that the eigenvalues of $A + B + C$ are small in the sense that

$$\text{Tr}(H_A + H_B + H_C) + \|H_A + H_B + H_C\|_\infty \leq 1. \quad (193)$$

If H'_A , H'_B and H'_C did not satisfy this condition, then one can define

$$H_A = \mu(H'_A + \nu I), \quad H_B = \mu(H'_B + \nu I), \quad H_C = \mu(H'_C + \nu I). \quad (194)$$

for $\mu > 0$ and $\nu \geq 0$ such that H_A, H_B and H_C satisfy the desired conditions. Specifically, let $\nu \geq |\min(0, \lambda_d(H_A), \lambda_d(H_B), \lambda_d(H_C))|$ (the magnitude of the largest negative eigenvalue of all three matrices), and let $\mu > 0$ is choosen sufficiently small so that the following holds

$$\text{Tr}(H'_A + H'_B + H'_C) + \|H'_A + H'_B + H'_C\|_\infty + 3(d+1)\nu \leq \frac{1}{\mu}. \quad (195)$$

For each positive semidefinite Hermitian matrix, H_X , let $|\phi_X\rangle$ be the vector in $\mathbb{C}^d \times \mathbb{C}^d$ defined by

$$|\phi\rangle_{XZ} = \sum_{i=1}^d H_X^{\frac{1}{2}} |i\rangle_X \otimes |i\rangle_Z. \quad (196)$$

Then the (non-normalized) reduced state on the first subsystem satisfies

$$\rho_X := \text{Tr}_Z(|\phi\rangle_{XZ} \langle\phi|_{XZ}) = H_X. \quad (197)$$

Therefore, one can define a quadripartite pure state $|\psi\rangle_{AB CZ}$ on $(\mathbb{C}^{d+1})^{\otimes 4}$ by

$$|\psi\rangle_{AB CZ} = \kappa_{ABC}^{\frac{1}{2}} |0000\rangle_{AB CZ} + |\phi\rangle_{AZ} |00\rangle_{BC} + |\phi\rangle_{BZ} |00\rangle_{AC} + |\phi\rangle_{CZ} |00\rangle_{AB}, \quad (198)$$

$$= \kappa_{ABC}^{\frac{1}{2}} |0000\rangle_{AB CZ} + \sum_{i=1}^d |\omega_i\rangle_{ABC} |i\rangle_Z, \quad (199)$$

where $\kappa_{ABC} = 1 - \text{Tr}(H_A + H_B + H_C)$ and

$$|\omega_i\rangle_{ABC} := H_A^{\frac{1}{2}} |i\rangle_A |00\rangle_{BC} + H_B^{\frac{1}{2}} |i\rangle_B |00\rangle_{AC} + H_C^{\frac{1}{2}} |i\rangle_C |00\rangle_{AB}. \quad (200)$$

The coefficients of $|\omega_i\rangle_{ABC}$ in the computational basis $|j_1 j_2 j_3\rangle_{ABC}$ are therefore

$$\langle j_1 j_2 j_3 | \omega_i \rangle_{ABC} = \delta_{j_1,0} \delta_{j_2,0} \langle j_3 | H_C^{\frac{1}{2}} | i \rangle + \delta_{j_1,0} \delta_{j_3,0} \langle j_2 | H_B^{\frac{1}{2}} | i \rangle + \delta_{j_2,0} \delta_{j_3,0} \langle j_1 | H_A^{\frac{1}{2}} | i \rangle \quad (201)$$

which implies

$$\sum_{j_1, j_2, j_3=0}^d \langle j_1 j_2 j_3 | \omega_{i_1} \rangle \langle \omega_{i_2} | j_1 j_2 j_3 \rangle = \langle i_2 | H_A | i_1 \rangle + \langle i_2 | H_B | i_1 \rangle + \langle i_2 | H_C | i_1 \rangle \quad (202)$$

The marginals of $|\psi\rangle_{AB CZ}$ encode faithfully the spectra of H_A , H_B , H_C and their partial sums in the following manner:

$$\rho_Z = \kappa_{ABC} |0\rangle \langle 0|_Z + \sum_{i_1, i_2=1}^d \langle i_2 | H_A + H_B + H_C | i_1 \rangle |i_1\rangle \langle i_2|_Z \quad (203)$$

$$= \kappa_{ABC} |0\rangle \langle 0|_Z + \overline{H_A + H_B + H_C}. \quad (204)$$

Similarly, letting $|\eta\rangle := |\phi\rangle_{CZ} + \kappa_{ABC}^{\frac{1}{2}} |00\rangle_{CZ}$, one obtains

$$\rho_{CZ} = |\eta\rangle \langle \eta|_{CZ} + |0\rangle \langle 0|_C \otimes \overline{H_A + H_B} \quad (205)$$

2.10.3 On the entropics sub-ensembles

TC: While at dinner on May 29th, 2024, Matthias suggested the possibility for an additive, Horn-like notion of an entropy cone which would serve as an analogue to the ordinary quantum entropy cone in the sense of Pippenger [Pip03]. The purpose of this section is to articulate the proposal more carefully.

A *ensemble of states* $\mathcal{E}$ is simply a finite collection of *weight-state* pairs:

$$\mathcal{E} = \{(p_1, \rho_1), \dots, (p_k, \rho_k)\}, \quad (206)$$

where the weights, p_i for $i \in [k]$, are *strictly positive* real numbers summing to one, while the states, ρ_i for $i \in [k]$, are $d \times d$ density matrices (Hermitian matrices, positive semidefinite, trace one).

Given an ensemble $\mathcal{E}$, there are two derived states of interest, $\sigma(\mathcal{E})$ and $\tau(\mathcal{E})$, defined as follows; $\sigma(\mathcal{E})$ is the bipartite *quantum-classical* (QC) state $\sigma(\mathcal{E}) \in \mathcal{S}(\mathbb{C}^k \otimes \mathbb{C}^d)$ defined by

$$\sigma(\mathcal{E}) := p_1 \rho_1 \otimes |1\rangle\langle 1| + \dots + p_k \rho_k \otimes |k\rangle\langle k| \quad (207)$$

whereas $\tau(\mathcal{E}) \in \mathcal{S}(\mathbb{C}^d)$ is

$$\tau(\mathcal{E}) := p_1 \rho_1 + \dots + p_k \rho_k. \quad (208)$$

Note that $\tau(\mathcal{E})$ is simply the partial trace of $\sigma(\mathcal{E})$ over the subsystem encoding the index j . Moreover, we remark that given a QC state of the above form, it is possible to extract the corresponding ensemble $\mathcal{E}$.

Now given a non-empty subset $S \subseteq [k]$, the subensemble associated to S , called the S -ensemble, is

$$\mathcal{E}_S = \{(q_j^S, \rho_j) \mid j \in S\}, \quad (209)$$

where the j th weight of the S -ensemble, denoted by q_j^S , is defined by renormalization:

$$q_j^S := \frac{p_j}{\sum_{i \in S} p_i}. \quad (210)$$

The state $\sigma(\mathcal{E}_S)$ associated to the S -ensemble can be interpreted as the post-measurement state obtained from performing the binary S vs. not S measurement on the classical register of $\sigma(\mathcal{E}_S)$.

Ultimately, we are interested in the *entropics* of ensembles and subensembles and the relations between them. Specifically, if $\mathcal{E}$ is an ensemble, then we are interested in the von Neumann entropies of the states $\tau(\mathcal{E}_S)$ for various subsets $S \subseteq [k]$. For example, we say that a given collection $(p_1, p_2, h_1, h_2, h_{12})$ of non-negative real numbers is *admissible* (realizable/compatible/consistent/etc.) if there exists density matrices ρ_1 and ρ_2 such that the ensemble $\mathcal{E} = \{(p_1, \rho_1), (p_2, \rho_2)\}$ simultaneously satisfies

$$h_{12} = H(\tau(\mathcal{E}_{12})) = H(p_1 \rho_1 + p_2 \rho_2), \quad (211)$$

$$h_1 = H(\tau(\mathcal{E}_1)) = H(\rho_1), \quad (212)$$

$$h_2 = H(\tau(\mathcal{E}_2)) = H(\rho_2). \quad (213)$$

For example, the convexity of the logarithm implies a concavity of von Neumann entropy:

$$\chi(\mathcal{E}_{12}) := H(p_1 \rho_1 + p_2 \rho_2) - p_1 H(\rho_1) - p_2 H(\rho_2) \geq 0 \quad (214)$$

where $\chi(\mathcal{E})$ is called the *Holevo information* of the ensemble. This implies that the collection of numbers $(p_1, p_2, h_1, h_2, h_{12})$ are admissible *only if* they satisfy the linear constraint

$$h_{12} - p_1 h_1 - p_2 h_2 \geq 0. \quad (215)$$

Similarly, $(p_1, p_2, p_3, h_1, h_2, h_3, h_{12}, h_{23}, h_{13})$, are said to be admissible if there exists (ρ_1, ρ_2, ρ_3) forming the ensemble

$$\mathcal{E}_{123} := \{(p_1, \rho_1), (p_2, \rho_2), (p_3, \rho_3)\}, \quad (216)$$




such that

$$\begin{aligned} h_1 &= H(\tau(\mathcal{E}_1)), & h_{12} &= H(\tau(\mathcal{E}_{12})), \\ h_2 &= H(\tau(\mathcal{E}_2)), & h_{23} &= H(\tau(\mathcal{E}_{23})), & h_{123} &= H(\tau(\mathcal{E}_{123})), \\ h_3 &= H(\tau(\mathcal{E}_3)), & h_{13} &= H(\tau(\mathcal{E}_{13})). \end{aligned} \quad (217)$$

Just as in the bipartite case, this allocation of entropies is admissible *only if* if all instances of concavity hold: **TODO**

2.11 Miscellaneous Diagrammatics

$$\begin{aligned}
& \left(U_{\lambda}^d \right)^2 C_{\lambda}^{\alpha, \nu} C_{\mu}^{\alpha, \beta} C_{\lambda}^{\mu, \gamma} C_{\nu}^{\beta, \gamma} = \\
& = \prod_i \int du_i \text{ (complex network of nodes and arcs) } \\
& = \prod_i \int du_i \text{ (simplified network of nodes } u_1, u_2, u_3, u_4 \text{ and arcs } \alpha, \mu, \gamma, \beta, \lambda, \nu \text{)} \\
& \quad (218)
\end{aligned}$$



(220)

2.12 Integral formula for the squared recoupling operator

By the above diagrammatic reasoning, it can be shown that the *square* of the $6j$ symbols can be written in terms of an integral over a quadruple $(k_1, k_2, k_3, k_4) \in \text{SU}(2)^{\times 4}$ according to the formula

$$S[\{l_{xy}\}] := \begin{Bmatrix} l_{12} & l_{13} & l_{14} \\ l_{34} & l_{24} & l_{23} \end{Bmatrix}^2 = \int_{\text{SU}(2)^4} [dk_x] \prod_{x < y} \chi_{l_{xy}}(k_x k_y^{-1}) \quad (221)$$

where $x, y \in [4] = \{1, 2, 3, 4\}$ and l_{xy} labels a spin $j_{xy} = l_{xy}/2$ representation $V_{l_{xy}}$ of $\text{SU}(2)$ of dimension $l_{xy} + 1$ and $\chi_{l_{xy}}$ its character (see also [FL03, Equation (1)]). The proof essentially

follows by noting that each k_x for $x \in [4]$ appears exactly three times and moreover, for each such x the following integral over k_x ,

$$\int_{\text{SU}(2)} dk_x \bigotimes_{y \neq x} \pi_{l_{xy}}(k_x) \quad (222)$$

is a projection operator onto the subspace of $\text{SU}(2)$ -invariant vectors $\text{Inv}(\bigotimes_{y \neq x} V_{l_{xy}})$ in $\bigotimes_{y \neq x} V_{l_{xy}}$ which is either zero or one dimensional.

3 Random walks on symmetric spaces

The purpose of this section is to summarize the conceptual and technical contributions of Klyachko's paper "Random walks on symmetric spaces and inequalities for matrix spectra" [Kly00].

3.1 Symmetric spaces

3.1.1 Definitions

As formulated in [Kly00, Sec. 2.1], *symmetric spaces* are parametrized by so-called *Cartan pairs* (G, σ) consisting of a Lie group G and an involution $\sigma : G \rightarrow G$ with compact centralizer $K = \{g \in G \mid g\sigma = \sigma g\}$. Given a Cartan pair (G, σ) the space of cosets $X := G/K$ can be equipped with a G -invariant metric (unique up to proportionality) which makes $X = G/K$ a homogeneous space (upon which G acts transitively by left multiplication). The space $X = G/K$ is a *symmetric space* in the sense that the geodesic symmetry, σ , with center at $x_0 = g_0K \in X = G/K$ is an isometry, sending points x lying on geodesics through x_0 to their symmetric point x' lying on the same geodesic at the same distance from x_0 . Specifically, given $x = gK \in X = G/K$, the isometry with center at $x_0 = g_0K$ is

$$gK \mapsto g_0 g_0^{-\sigma} g^\sigma K. \quad (223)$$

Following [Kly00, Sec. 2.2.2], a compact Lie group K can be used to define the following three symmetric spaces:

- (Positive curvature) The compact group K itself can be considered as a symmetric space with isometry group $K \times K$, acting by left and right multiplication sending $x \in K$ to $(k_1, k_2) \cdot x \mapsto k_1 x k_2^{-1} \in K$. The Cartan involution here, $\sigma : K \times K \rightarrow K \times K$ swaps the factors. The isotropy group is K itself embedded diagonally in $K \times K$.
- (Zero curvature) The Lie algebra $\mathfrak{k} = \text{Lie}(K)$ of K is a symmetric space with noncompact isometry group generated by translations and the adjoint action of K .
- (Negative curvature) The complexification of K , denoted $G = K_{\mathbb{C}}$, is a complex reductive group with Lie algebra $\mathfrak{g} = \mathfrak{k} \otimes \mathbb{C}$. The coset space $H = G/K$ is a symmetric space, called the *dual symmetric space*, with complex conjugation in G as a Cartan involution.

Notice that in all three cases, the compact centralizer of the involution σ is (isomorphic to) the compact group K itself; what distinguishes the three cases is the enveloping group G and the involution itself: positive curvature $G = K \times K$, negative curvature $G = K_{\mathbb{C}}$ and zero curvature $G = \mathfrak{k}$.

3.1.2 Examples

The example of a compact group which will receive the most attention here is the group $K = \text{SU}(d)$ of $d \times d$ complex unitary matrices with unit determinant. For this special case, the three symmetric spaces are as follows.

In the *zero curvature* case, the associated symmetric space is the real Lie algebra $X = \mathfrak{su}(d)$ of skew-Hermitian traceless matrices. The presentation of $\mathfrak{su}(d)$ as a quotient takes the isometry group to be equal to the group generated by both the translations in

$\mathfrak{su}(d)$ as a vector space together with the adjoint action of $SU(d)$ on $\mathfrak{su}(d)$. The geodesic involution $\sigma : \mathfrak{su}(d) \rightarrow \mathfrak{su}(d)$ with center taken at the zero element in $\mathfrak{su}(d)$ sends every skew-Hermitian matrix $A \in \mathfrak{su}(d)$ to $\sigma(A) = -A$. The compact isometry subgroup (subgroup of translations and adjoint actions on $\mathfrak{su}(d)$) which fixes the center of involution or equivalently is the centralizer of the geodesic involution by be identified with $SU(d)$ itself through the aforementioned coadjoint action on $\mathfrak{su}(d)$. In the subcase where $d = 2$, as a real vector space $\mathfrak{su}(2)$ is isomorphic to $\mathbb{R}^3$ and as a Riemannian manifold is isomorphic to Euclidean space $\mathbb{E}^3$.

In the *positive curvature* case, the associated symmetric space is $SU(d)$ itself. The group $SU(d) \times SU(d)$ acts on $SU(d)$ by left and right actions $(g_1, g_2) \cdot x = g_1 x g_2^{-1}$ and the involution on $SU(d) \times SU(d)$ simply acts by swapping $\sigma(g_1, g_2) = (g_2, g_1)$. The compact centralizer of σ in this case is the diagonal subgroup $SU(d) \simeq \{(g, g) \mid g \in SU(d)\} \subset SU(d) \times SU(d)$; one can see that this subgroup under the left/right action fixes the element $x_0 = e \in SU(d)$.

In the subcase where $d = 2$, the manifold $SU(2)$ may be identified with the 3-sphere $\mathbb{S}^3$ in the sense that $\begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix} \in SU(2)$ for $a, b \in \mathbb{C}$ satisfies $|a|^2 + |b|^2 = 1$.

In the *negative curvature* case, the symmetric space (called the *dual* symmetric space) is a quotient of the complexification of $SU(d)$. The complexification of $SU(d)$ is the group $SL(d, \mathbb{C})$ of $d \times d$ complex matrices with unitary determinant. The involution $\sigma : SL(d, \mathbb{C}) \rightarrow SL(d, \mathbb{C})$ for this example is the complex conjugation of $SL(d, \mathbb{C})$ (as a complexification). Specifically, the local involution here leaves skew-Hermitian matrices $A \in \mathfrak{su}(d)$ unchanged but sends Hermitian matrices $H \in i\mathfrak{su}(d)$ to $-H \in i\mathfrak{su}(d)$, and thus the global involution leaves unitary matrices $U = \exp(A)$ unchanged but sends positive definite matrices $P = \exp(H)$ to their inverses $P^{-1} = \exp(-H)$. The centralizer of this involution is therefore the subgroup $SU(d) \subset SL(d, \mathbb{C})$ of unitary matrices and thus the symmetric space is the quotient $SL(d, \mathbb{C})/SU(d)$. By the (left-) polar-decomposition (also called the Cartan decomposition) every $M \in SL(d, \mathbb{C})$ can be written uniquely as $M = P \cdot U$ for some positive definite matrix P and unitary matrix U . Through this decomposition, the symmetric space $SU(d) \subset SL(d, \mathbb{C})$ may be identified with the space of positive definite matrices. Equivalently, every $M \in SL(d, \mathbb{C})$ can be used to calculate this P by the formula $M\sigma(M)^{-1} = (PU)(P^{-1}U)^{-1} = P^2$ and thus $P = \sqrt{M\sigma(M)^{-1}}$ where $M \mapsto \sigma(M)^{-1}$ is equivalent to complex conjugate transpose of matrices. The positive numbers forming the eigenvalues of P are called the *singular values* of M . It is important to note that there exists another way to think about this symmetric space. Since every positive-definite matrix P admits of a unique *Cholesky decomposition* $P = T^*T$ where T is an upper triangular matrix with real positive diagonal entries, one can also identify this dual symmetric space with the Borel subgroup $B(d) \subseteq SL(d, \mathbb{C})$ of such upper triangular matrices. This further decomposition is of course consistent with the Iwasawa decomposition $SL(d, \mathbb{C}) \simeq SU(d) \times A(d) \times N(d)$ where $B(d) \simeq A(d) \times N(d)$.

In the subcase where $d = 2$, the symmetric space of positive Hermitian 2×2 matrices is the hyperbolic three-space or Lobachevskii space $\mathbb{L}^3$. In particular, since $\det(P) = 1$ the most general form has P diagonalize to $\text{diag}(e^{r/2}, e^{-r/2})$ for some $r \geq 0$. Given $P_1, P_2 \in SL(2, \mathbb{C})/SU(2)$, the metric is simply $d(P_1, P_2) = r$ where r is obtained by calculating the P such that $PU = P_2^{-1}P_1$ for some $U \in SU(2)$ or equivalently just calculating

$$P = \sqrt{(P_2^{-1}P_1)(\sigma(P_2^{-1}P_1)^{-1})} = \sqrt{P_2^{-1}P_1^2P_2^{-1}} \quad (224)$$

and diagonalizing to find r . To get a sense, consider the extremely special case wherein $P_i = \text{diag}(e^{r_i/2}, e^{-r_i/2})$ are both diagonal in the same basis. Then the distance between P_1

and P_2 has a particularly simply form because

$$\sqrt{P_2^{-1}P_1^2P_2^{-1}} = \text{diag}(e^{(r_1-r_2)/2}, e^{(r_2-r_1)/2}) \quad (225)$$

and therefore $d(P_1, P_2) = |r_1 - r_2|$.

3.2 Random walks

Given a symmetric space $X = G/K$, the Haar measure μ on K induces a well-defined uniform measure, μ_i , over each double coset $X_i \in K \backslash G/K$. A *random walk* in the symmetric space $X = G/K$ depends upon on a fixed sequence of (double) cosets $(X_1, X_2, X_3, \dots)$ where each X_i is a left-coset of $X = G/K$ or equivalently a double coset of G given by $X_i \in K \backslash G/K$. Treating this sequence as input, one can sample a sequence $(g_i \in X_i)$ where each $g_i \in X_i$ is chosen uniformly and independently from all other g_j by sampling a Haar random $k_i \in K$ and letting $g_i = k_i r_i$ where $r_i \in X = G/K$ is any representative element of the coset $X_i = K r_i \in K \backslash G/K$. While the “step sizes” of the random walk on $X = G/K$ are the given cosets $X_i \in K \backslash X$, the “locations” of the random walk after a fixed number of “steps” are interpreted as follows:

- The starting location is the element $x_0 := eK \in G/K = X$.
- The first step is the element $g_1 \in X_1 = K r_1 \in K \backslash X$.
- The first location is the element $x_1 := g_1 K = (k_1 r_1) K \in G/K = X$.
- The second step is the element $g_2 \in X_2 = K r_2 \in K \backslash X$.
- The second location is the element $x_2 := g_1 g_2 K = (k_1 r_1)(k_2 r_2) K \in G/K = X$.
- ...
- The n th step is the element $g_n \in X_n = K r_n \in K \backslash X$.
- The n th location is the element $x_n := g_1 \cdots g_n K = (k_1 r_1) \cdots (k_n r_n) K \in G/K = X$.

The result quantity of interest is the induced probability measure/density for the final point $x := x_n$ after n steps. The probability density, $P_n(x) = P(x \mid X_1, \dots, X_n)$, is density for the pushforward of the product Haar measure $\mu_K^{\otimes n}$ on $K^{\times n}$ through the map $\psi_n : K^{\times n} \rightarrow G/K$ defined by

$$\psi_n(k_1, \dots, k_n) := (k_1 r_1) \cdots (k_n r_n) K \quad \text{\textcolor{blue}{\{eq:n_step_process\}}} \quad (226)$$

relative to the Haar measure $dx \cong d\mu_X(x)$ on $X = G/K$:

$$(\psi_n)_*(\mu^{\otimes n})(dx) = \mu^{\otimes n}(\psi_n^{-1}(dx)) = P_n(x)dx. \quad (227)$$

3.3 Spherical Functions

Below is a preliminary summary of the utility of spherical functions in this context. The textbooks by Helgason are a fantastic references [Hel84; Hel94].

Given some continuous measurable function $f : G/K \rightarrow \mathbb{C}$, the expected value of $f(x)$ when $x \in X = G/K$ is distributed according to $P_n(x)dx$ is therefore expressible as an integral over $K^{\times n}$.

$$\mathbb{E}[f] = \int_{x \in X} f(x) P_n(x) dx = \int_{K^{\times n}} dk_1 \cdots dk_n f(k_1 r_1 \cdots k_n r_n K) \quad (228)$$

The technique used by Klyachko to derive an expression for $P_n(x)$ is to consider a special family of functions, denoted by $\varphi_\lambda \in L_2(G/K)$ and indexed by $\lambda \in \Lambda$ for some set Λ , called *spherical functions*. A non-zero function $\varphi \in L_2(G)$ is spherical if it is normalized $\varphi(e) = 1$ and satisfies the following decomposition for all $x, y \in G$:

$$\varphi(x)\varphi(y) = \int_{k \in K} dk \varphi(xky). \quad (229)$$

This property implies that φ is also bi-invariant in that $\varphi(k_1 x k_2) = \varphi(x)$ for all $k_1, k_2 \in K$ because

$$\int_{k \in K} dk \varphi(kx) = \varphi(e)\varphi(x) = \varphi(x) = \varphi(x)\varphi(e) = \int_{k \in K} dk \varphi(xk) \quad (230)$$

As a consequence, we can identify spherical functions $\varphi \in L_2(G)$ on G with their pushforward through the coset projections $G \mapsto G/K \mapsto K \backslash G/K$.

An elementary example of a spherical function occurs when π_λ is an irreducible unitary representation of G on a Hilbert space $\mathcal{H}_\lambda$ (with inner product $\langle \cdot, \cdot \rangle$) such that the subspace of vector $v \in \mathcal{H}_\lambda$ which are fixed by K , i.e.,

$$\mathcal{H}_\lambda^K := \{v \in \mathcal{H} \mid \forall k \in K : \pi_\lambda(k)v = v\}, \quad (231)$$

is exactly one dimensional, i.e., $\dim(\mathcal{H}_\lambda^K) = 1$. In this setting, the map $\varphi_\lambda : G \rightarrow \mathbb{C}$, defined for all $g \in G$ by

$$\varphi_\lambda(g) = \frac{\langle v, \pi_\lambda(g)v \rangle}{\langle v, v \rangle} \quad (232)$$

is an elementary spherical function.

Whenever $f = \varphi$, as considered above, is a spherical function (or rather the pullback of f through $G \mapsto G/K$ is the spherical function φ), the expectation value $\mathbb{E}[\varphi]$ can be computed directly:

$$\mathbb{E}[\varphi] = \int_{x \in X} \varphi(x) P_n(x) dx, \quad (233)$$

$$= \int_{K^{\times n}} dk_1 \cdots dk_n \varphi(k_1 r_1 k_2 r_2 \cdots k_n r_n K), \quad (234)$$

$$= \int_{K^{\times n-1}} dk_2 \cdots dk_n \varphi(e) \varphi(r_1 k_2 r_2 \cdots k_n r_n K), \quad (235)$$

$$= \int_{K^{\times n-2}} dk_3 \cdots dk_n \varphi(e) \varphi(r_1) \varphi(r_2 \cdots k_n r_n K), \quad (236)$$

$$= \varphi(r_1) \varphi(r_2) \cdots \varphi(r_n), \quad (237)$$

$$= \varphi(X_1) \varphi(X_2) \cdots \varphi(X_n). \quad (238)$$

The space of bi-invariant functions on G , identified with $f \in L_2(K \backslash G / K)$, there exists a *spherical transform* sending $f \in L_2(K \backslash G / K)$ to the function $\hat{f} \in L_2(\Lambda)$ where $\lambda \in \Lambda$ indexes the spherical functions φ_λ such that

$$\int_{g \in G} |f(g)|^2 dg = \int_{\lambda \in \Lambda} |\hat{f}(\lambda)|^2 d\lambda \quad (239)$$

where dg is the Haar measure on G and $d\lambda$ is known as the *Plancherel measure* on Λ . Explicitly, the spherical transform of $f \in L_2(K \backslash G / K)$ is

$$\hat{f}(\lambda) = \int_{g \in G} f(g) \varphi_\lambda^*(g) dg. \quad (240)$$

The inversion formula for the spherical transform is

$$f(g) = \int_{\lambda \in \Lambda} \hat{f}(\lambda) \varphi_\lambda(g) d\lambda. \quad (241)$$

As mentioned in [Kly00, Remark 3.4.3], the spherical transform can be described as a homomorphism from convolution of bi-invariant functions $p, q \in L_2(K \backslash G / K)$,

$$(p * q)(x) = \int_{g \in G} p(xg) q(g^{-1}x) dg, \quad (242)$$

to pointwise multiplication of transformed functions $\hat{p}, \hat{q} \in L_2(\Lambda)$ such that $\widehat{p * q} = \hat{p} \cdot \hat{q}$.

Furthermore, as the product of spherical functions is a bi-invariant, it is possible to perform a spherical transform of a product of spherical functions. Specifically, letting

$$(\varphi_\alpha \cdot \varphi_\beta)(g) := \varphi_\alpha(g) \varphi_\beta(g), \quad (243)$$

and letting $\Phi_{\alpha, \beta}^\gamma$ be the coefficient

$$\Phi_{\alpha, \beta}^\gamma := \widehat{\varphi_\alpha \cdot \varphi_\beta}(\gamma) = \int_{g \in G} \varphi_\alpha(g) \varphi_\beta(g) \varphi_\gamma^*(g) dg, \quad (244)$$

the product of spherical functions can be written as a linear combination of spherical functions as

$$(\varphi_\alpha \cdot \varphi_\beta)(g) = \int_{\gamma \in \Lambda} \Phi_{\alpha, \beta}^\gamma \varphi_\gamma(g) d\gamma \quad (245)$$

The probability density $P_n(x)$ above is bi-invariant, therefore its spherical transform gives rise to a formula for $P_n(x)$ in terms of an integration over $\lambda \in \Lambda$:

$$P_n(x) = \int_{\lambda \in \Lambda} \hat{P}_n(\lambda) \varphi_\lambda(x) d\lambda \quad (246)$$

$$= \int_{\lambda \in \Lambda} \left(\int_{g \in G} P_n(g) \varphi_\lambda^*(g) dg \right) \varphi_\lambda(x) d\lambda \quad (247)$$

$$= \int_{\lambda \in \Lambda} \left(\int_{y \in X} P_n(y) \varphi_\lambda^*(y) dy \right) \varphi_\lambda(x) d\lambda \quad (248)$$

$$= \int_{\lambda \in \Lambda} \varphi_\lambda^*(X_1) \cdots \varphi_\lambda^*(X_n) \varphi_\lambda(x) d\lambda. \quad (249)$$

At this stage, two steps remain to compute the density $P_n(x)$. The first is to identify the space of spherical functions, $\{\varphi_\lambda\}_{\lambda \in \Lambda}$, (together with the Plancherel measure on Λ). The second is then to perform the above integral over Λ . As the spherical functions and the Plancherel measures are known in great detail for many symmetric spaces (for arbitrary complex semisimple groups, the formula is due to Harish-Chandra [Hel84, Theorem 5.7]), all that remains is to perform the integral which Klyachko performs/sketches in [Kly00].

3.4 Intermediate random walks

The purpose of this subsection is to turn our attention to a situation where one is not merely interested in the distribution of the *terminal* point x_n but also potentially a subsequence $x_{i_1}, \dots, x_{i_\ell}$ of length $\ell > 1$ or additionally intervals describing intermediate random walks. Our primary focus for now will be a short process of $n = 3$ steps: let (k_1, k_2, k_3) are uniformly sampled from the Haar measure $\mu_K^{\otimes 3}$ on $K^{\times 3}$ and let $r_1, r_2, r_3 \in G$ be fixed double coset representatives. Then let $\Psi : K^{\times 3} \rightarrow (G/K)^{\times 3}$ be the map defined

$$\Psi(k_1, k_2, k_3) := (\Psi_{12}(k_1, k_2), \Psi_{23}(k_2, k_3), \Psi_{123}(k_1, k_2, k_3)) \quad (250)$$

where the component maps $\Psi_{12} : K^{\times 2} \rightarrow G/K$, $\Psi_{23} : K^{\times 2} \rightarrow G/K$ and $\Psi_{123} : K^{\times 3} \rightarrow G/K$ are defined as follows:

$$\Psi_{12}(k_1, k_2) := k_1 r_1 k_2 r_2 K, \quad (251)$$

$$\Psi_{23}(k_2, k_3) := k_2 r_2 k_3 r_3 K, \quad (252)$$

$$\Psi_{123}(k_1, k_2, k_3) := k_1 r_1 k_2 r_2 k_3 r_3 K. \quad (253)$$

Note that $\Psi_{12} = \psi_2$ and $\Psi_{123} = \psi_3$ where ψ_n is defined by Eq. (226). The induced probability measure of interest is now the joint measure over $X^3 = (G/K)^{\times 3}$ obtained from the pushforward of $\mu_K^{\otimes 3}$ through Ψ :

$$P(x_1, x_2, x_3) dx_1 dx_2 dx_3 = (\Psi)_*(\mu^{\otimes 3})(dx_1 dx_2 dx_3) = \mu^{\otimes 3}(\Psi^{-1}(dx_1 dx_2 dx_3)). \quad (254)$$

Given a function $f \in L_2((G/K)^3)$ its expected value is therefore

$$\mathbb{E}[f] = \int_{K^{\times 3}} dk_1 dk_2 dk_3 f(k_1 r_1 k_2 r_2 K, k_2 r_2 k_3 r_3 K, k_1 r_1 k_2 r_2 k_3 r_3 K). \quad (255)$$

TODO: Figure out how to perform these types of integrals and relate them back to recoupling symbols.


In the special case where $f(x_1, x_2, x_3) = \varphi_\mu(x_1)\varphi_\nu(x_2)\varphi_\lambda(x_3)$ happens to be a product of spherical functions, only a moderate amount of simplification occurs

$$\mathbb{E}[\varphi_\mu \varphi_\nu \varphi_\lambda] = \int_{K^{\times 3}} dk_1 dk_2 dk_3 \varphi_\mu(k_1 r_1 k_2 r_2 K) \varphi_\nu(k_2 r_2 k_3 r_3 K) \varphi_\lambda(k_1 r_1 k_2 r_2 k_3 r_3 K) \quad (256)$$

$$= \int_{K^{\times 2}} dk_2 dk_3 \varphi_\mu(r_1 k_2 r_2) \varphi_\nu(r_2 k_3 r_3) \varphi_\lambda(r_1 k_2 r_2 k_3 r_3). \quad (257)$$

3.5 Joint statistics for partial sums of traceless 2×2 matrices

In this section we consider a three-step random walk in the symmetric space $\mathfrak{su}(2) \simeq \mathbb{E}^3$ starting at the origin $O = (0, 0, 0)$ and terminating at C with intermediate steps A, B .



Supposing the step sizes $(a, b, c) \in \mathbb{R}_{\geq 0}^3$ are fixed, we are interested in the induced correlations between the interval sizes $(x, y, z) \in \mathbb{R}_{\geq 0}^3$ when the angles planar angles $(\phi, \psi) \in [0, \pi]$ and dihedral angle $\eta \in [0, \pi]$ are sampled independently according to the measures $\frac{1}{2} \sin \phi d\phi$, $\frac{1}{2} \sin \psi d\psi$ and $\pi^{-1} d\eta$ respectively (note it is a non-trivial observation that sampling the vectors A, B , and C uniformly from spheres of radii a, b , and c induces the same correlations between x, y , and z as the correlations induced by the random angles ϕ, ψ and η).

Begin by noting that x and y are fixed by ϕ and ψ respectively by the law of cosines:


$$x^2 = a^2 + b^2 - 2ab \cos \phi, \quad (259)$$

$$y^2 = b^2 + c^2 - 2bc \cos \psi. \quad (260)$$

As x is functionally (and thus statistically) independent of ϕ and η and y is independent of ψ and η , one can expect x and y to be independent of each other and thus the support of the marginal distribution over x and y is a rectangle:

$$(x, y) \in [|a - b|, a + b] \times [|b - c|, b + c] \quad (261)$$

Comparatively less obvious is the relationship between z and the variables a, b, c, x, y . To process, we begin by introducing two more planar angles, ω and ζ , at the vertex A :



The cosine laws for the planar angles incident to A are

$$\cos \phi = \frac{a^2 + b^2 - z^2}{2ab}, \quad \cos \omega = \frac{b^2 + y^2 - c^2}{2by}, \quad \cos \zeta = \frac{a^2 + y^2 - z^2}{2ay}. \quad (263)$$

To make progress, one needs to use a relationship holding between the dihedral angle η and the other specified variables. For instance [Ric02],

$$\cos \eta = \frac{\cos \zeta - \cos \phi \cos \omega}{\sin \phi \sin \omega}. \quad (264)$$

Combining the above identities, one obtains

$$z^2 = a^2 + y^2 - 2ay \cos \zeta, \quad (265)$$

$$\cos \zeta = \cos \eta \sin \phi \sin \omega + \cos \phi \cos \omega, \quad (266)$$

$$\cos \omega = \frac{b - c \cos \psi}{\sqrt{b^2 + c^2 - 2bc \cos \psi}}, \quad (267)$$

$$\sin \omega = \frac{c \sin \psi}{\sqrt{b^2 + c^2 - 2bc \cos \psi}}. \quad (268)$$

Thus, one can determine z as a function of the side lengths a, b, c and the angles η, ϕ, ψ :

$$z^2 = a^2 + y^2 - 2ay \cos \zeta, \quad (269)$$

$$= a^2 + y^2 - 2ay \cos \eta \sin \phi \sin \omega - 2ay \cos \phi \cos \omega, \quad (270)$$

$$= a^2 + b^2 + c^2 - 2bc \cos \psi - 2a \cos \eta \sin \phi (c \sin \psi) - 2a \cos \phi (b - c \cos \psi), \quad (271)$$

$$= a^2 + b^2 + c^2 - 2bc \cos \psi - 2ac \cos \eta \sin \phi \sin \psi + 2ac \cos \phi \cos \psi - 2ab \cos \phi, \quad (272)$$

$$= a^2 + b^2 + c^2 - 2bc \cos \psi - 2ab \cos \phi - 2ac(\cos \eta \sin \phi \sin \psi - \cos \phi \cos \psi), \quad (273)$$

or alternatively as function of a, b, c and x, y, η :

$$z^2 = x^2 + y^2 - b^2 + \frac{(a^2 + b^2 - x^2)(b^2 + c^2 - y^2)}{2b^2} + \dots \quad (274)$$

$$\dots - \cos \eta \frac{\sqrt{4a^2b^2 - (a^2 + b^2 - x^2)^2} \sqrt{4b^2c^2 - (b^2 + c^2 - y^2)^2}}{2b^2}$$

Consequently, the admissible values of z (found by varying $\cos \eta \in [-1, +1]$) depend upon x, y according to $Z_-(x, y) \leq z \leq Z_+(x, y)$ where

$$Z_{\pm}(x, y) = \left(x^2 + y^2 - b^2 + \frac{(a^2 + b^2 - x^2)(b^2 + c^2 - y^2)}{2b^2} + \dots \right. \quad (275)$$

$$\left. \dots \pm \frac{\sqrt{4a^2b^2 - (a^2 + b^2 - x^2)^2} \sqrt{4b^2c^2 - (b^2 + c^2 - y^2)^2}}{2b^2} \right)^{\frac{1}{2}}$$

The change of variables from (ϕ, ψ, η) to (x, y, z) (for constant a, b, c) is

$$\begin{pmatrix} x^2 \\ y^2 \\ z^2 \end{pmatrix} = \begin{pmatrix} a^2 + b^2 - 2ab \\ b^2 + c^2 - 2bc \\ a^2 + b^2 + c^2 - 2bc \cos \psi - 2ab \cos \phi - 2ac(\cos \eta \sin \phi \sin \psi - \cos \phi \cos \psi) \end{pmatrix}. \quad (276)$$

After taking square-roots, it has a Jacobian determinant equal to

$$\det J(\phi, \psi, \eta) = \frac{a^2 b^2 c^2 \sin \eta \sin^2 \phi \sin^2 \psi}{xyz} \quad (277)$$

If (ϕ, ψ, η) is sampled according to the measure

$$\frac{1}{4\pi} \sin \phi \sin \psi d\phi d\psi d\eta, \quad (278)$$

then the joint probability density for (x, y, z) can be shown to be equal to

$$p(x, y, z|a, b, c) = \left(\frac{1}{4\pi} \sin \phi \sin \psi \right) (\det J(\phi, \psi, \eta))^{-1}, \quad (279)$$

$$= \frac{1}{4\pi} \frac{xyz}{a^2 b^2 c^2 \sin \eta \sin \phi \sin \psi}, \quad (280)$$

$$= \frac{xyz}{2\pi abc} \frac{1}{\sqrt{Q(a, b, c, x, y, z)}} \quad (281)$$

where $Q(a, b, c, x, y, z)$ is the following homogeneous degree 6 polynomial:

$$Q(a, b, c, x, y, z) := -a^4 c^2 + a^2 b^2 c^2 - a^2 b^2 x^2 + a^2 b^2 z^2 - a^2 c^4 + a^2 c^2 x^2 \quad (282)$$

$$+ a^2 c^2 y^2 + a^2 c^2 z^2 + a^2 x^2 y^2 - a^2 y^2 z^2 - b^4 z^2 - b^2 c^2 y^2 \quad (283)$$

$$+ b^2 c^2 z^2 + b^2 x^2 y^2 + b^2 x^2 z^2 + b^2 y^2 z^2 - b^2 z^4 \quad (284)$$

$$+ c^2 x^2 y^2 - c^2 x^2 z^2 - x^4 y^2 - x^2 y^4 + x^2 y^2 z^2 \quad (285)$$

Further analysis [Ric02] (being careful to note the notational convention in [Ric02] is incompatible with the one used here $(a, b, c, x, y, z) \leftrightarrow (a, x, z, c, y, b)$) reveals that $Q(a, b, c, x, y, z)$ is equal to 114 times the volume squared of the tetrahedron with side lengths a, b, c, x, y, z , i.e.

$$Q(a, b, c, x, y, z) = 144 \text{Vol}(a, b, c, x, y, z)^2, \quad (286)$$

$$= \frac{1}{2} \det \begin{pmatrix} 2a^2 & a^2 + b^2 - x^2 & a^2 + y^2 - z^2 \\ a^2 + b^2 - x^2 & 2b^2 & b^2 - c^2 + y^2 \\ a^2 + y^2 - z^2 & b^2 - c^2 + y^2 & 2y^2 \end{pmatrix} \quad (287)$$

Therefore, the resulting probability density is simply

$$p(x, y, z|a, b, c) = \frac{xyz}{24\pi abc \text{Vol}(a, b, c, x, y, z)} \quad (288)$$




Figure 1: Admissible eigenvalues of partial sums $(A+B, B+C$ and $A+B+C)$ of traceless 2×2 matrices (A, B, C) with spectra $(\alpha, \beta, \gamma) = (a, b, c) = (0.5, 0.6, 0.7)$. Left is a density plot of the analytic formula $p(x, y, z|a, b, c)$ restricted to the admissible region. Right is a numerical sample with admissible region boundary plotted in green.

3.6 Admissibility of Hyperbolic tetrahedron

TODO, TC: Type up the proof from my tablet notes “Positive definite products of 2 by 2 matrices” The space of admissible (a, b, c, x, y, z) is given by triangle inequalities on x, y in terms of a, b, c as well as the following constraint on z in terms of x, y, a, b, c :

$$\begin{aligned}
& + 2 \cosh(2s(a - b - c - z)) + 2 \cosh(2s(a + b - c - z)) + 2 \cosh(2s(a - b + c - z)) \\
& + 2 \cosh(2s(a + b + c - z)) + 2 \cosh(2s(a - b - c + z)) + 2 \cosh(2s(a + b - c + z)) \\
& + 2 \cosh(2s(a - b + c + z)) + 2 \cosh(2s(a + b + c + z)) + 2 \cosh(2s(a - c - x - y)) \\
& + 2 \cosh(2s(a + c - x - y)) + 2 \cosh(2s(a - c + x - y)) + 2 \cosh(2s(a + c + x - y)) \\
& + 2 \cosh(2s(a - c - x + y)) + 2 \cosh(2s(a + c - x + y)) + 2 \cosh(2s(a - c + x + y)) \\
& + 2 \cosh(2s(a + c + x + y)) + 2 \cosh(2s(b - x - y - z)) + 2 \cosh(2s(b + x - y - z)) \\
& + 2 \cosh(2s(b - x + y - z)) + 2 \cosh(2s(b + x + y - z)) + 2 \cosh(2s(b - x - y + z)) \\
& + 2 \cosh(2s(b + x - y + z)) + 2 \cosh(2s(b - x + y + z)) + 2 \cosh(2s(b + x + y + z)) \\
& + 2 \cosh(4as) + 2 \cosh(4bs) + 2 \cosh(4cs) + 2 \cosh(4sx) + 2 \cosh(4sy) + 2 \cosh(4sz) + 10 \\
& \geq \\
& + 4 \cosh(2s(a - b - x)) + 4 \cosh(2s(a + b - x)) + 4 \cosh(2s(a - b + x)) \\
& + 4 \cosh(2s(a + b + x)) + 4 \cosh(2s(a - y - z)) + 4 \cosh(2s(a + y - z)) \\
& + 4 \cosh(2s(a - y + z)) + 4 \cosh(2s(a + y + z)) + 4 \cosh(2s(b - c - y)) \\
& + 4 \cosh(2s(b + c - y)) + 4 \cosh(2s(b - c + y)) + 4 \cosh(2s(b + c + y)) \\
& + 4 \cosh(2s(c - x - z)) + 4 \cosh(2s(c + x - z)) + 4 \cosh(2s(c - x + z)) \\
& + 4 \cosh(2s(c + x + z)) \\
& + \cosh(4s(x - y)) + \cosh(4s(x + y)) + \cosh(4s(a - c)) \\
& + \cosh(4s(a + c)) + \cosh(4s(b - z)) + \cosh(4s(b + z))
\end{aligned} \tag{289}$$

4 Multiplicative and additive Horn problems

4.1 On the validity of Thompson's conjecture

To begin, we use the notation of Speyer [Spe05].

Given a triple of n -tuples of real numbers, $(\alpha, \beta, \gamma) \in (\mathbb{R}^n)^{\times 3}$, Horn's problem is concerned with whether there exists a triple (A, B, C) of $n \times n$ Hermitian matrices with eigenvalues (α, β, γ) that also satisfies the constraint $A + B + C = 0$. Without loss of generality, one can always assume the eigenvalues are sorted such that one can restrict their attention to the subset of non-increasing lists of real numbers, $\mathbb{R}_\downarrow^n \subset \mathbb{R}^n$. In addition, if the triple (A, B, C) of Hermitian operators satisfies the conditions required, then so will the triple of matrices

$$U \cdot (A, B, C) := (UAU^{-1}, UBU^{-1}, UCU^{-1}) \quad (290)$$

where $U \in \mathrm{U}(n)$ is an $n \times n$ unitary matrix and U^{-1} its inverse.

There is a variation of this problem also concerned with a triple of n -tuples of real numbers, (α, β, γ) , that is concerned with singular values instead of eigenvalues. The singular values of an $n \times n$ matrix A are simply the positive square-roots of the eigenvalues of the matrix positive semidefinite matrix A^*A (or equivalently AA^*). Just as the eigenvalues of a matrix A characterize the orbit under the action $U \cdot A = UAU^{-1}$ by $U \in \mathrm{U}(d)$, the singular values of a matrix A characterize the orbit under the action $(U, V) \cdot A = UAV^{-1}$ by $(U, V) \in \mathrm{U}(d) \times \mathrm{U}(d)$. In a similar fashion, without loss of generality, one can always assume the singular values are non-increasing but also non-negative. Furthermore, if one restricts their attention to matrices (A, B, C) which are additionally *invertible*, then their singular values are necessarily strictly positive. The multiplicative variation of Horn's problem is concerned with whether there exists a triple (A, B, C) of $n \times n$ matrices with singular values (α, β, γ) that also satisfy the constraint $ABC = I_n$ where I_n is the $n \times n$ identity matrix. While the additive problem possesses a $\mathrm{U}(d)$ symmetry, the multiplicative problem possesses a $\mathrm{U}(d)^{\times 3}$ symmetry given by the action

$$(U, V, W) \cdot (A, B, C) := (UAV^{-1}, VBW^{-1}, WCU^{-1}). \quad (291)$$

The following result, originally due to Klyachko [Kly00], demonstrates a correspondence between the solutions to the multiplicative and additive variants of this problem.

Theorem 11 (Klyachko's Theorem). *There exists a triple of Hermitian matrices (A, B, C) with eigenvalues $(\alpha, \beta, \gamma) \in (\mathbb{R}_\downarrow^n)^{\times 3}$ satisfying $A + B + C = 0$ if and only if there exists a triple of invertible matrices (A', B', C') with singular values $(e^\alpha, e^\beta, e^\gamma)$ satisfying $ABC = I_n$.*

The method used by Klyachko [Kly00] to prove this statement is rather indirect when compared to the method used by Alekseev, Meinrenken, and Woodward [AMW01]. In [AMW01], Alekseev, Meinrenken, and Woodward construct an implicit homeomorphism, $\alpha : \mathcal{O}^{\mathrm{add}} \rightarrow \mathcal{O}^{\mathrm{mult}}$, between the set $\mathcal{O}^{\mathrm{add}}$ of triples of Hermitian matrices (A, B, C) (modulo $U \in \mathrm{U}(d)$) and the set $\mathcal{O}^{\mathrm{mult}}$ of triples of invertible matrices (A', B', C') (modulo $(U, V, W) \in \mathrm{U}(d)^{\times 3}$).

This homeomorphism preserves the spectral information in the sense that it maps the fiber $\mathcal{O}_{\alpha, \beta, \gamma}^{\mathrm{add}}$ of triples of Hermitian matrices (A, B, C) (modulo $U \in \mathrm{U}(d)$) with eigenvalues (α, β, γ) to the set $\mathcal{O}_{e^\alpha, e^\beta, e^\gamma}^{\mathrm{mult}}$ of triples of invertible matrices (A', B', C') (modulo $(U, V, W) \in \mathrm{U}(d)^{\times 3}$) with singular values $(e^\alpha, e^\beta, e^\gamma)$. In summary, one has $\mathrm{sing} \circ \alpha = \exp \circ \mathrm{eigen}$.

4.2 The implicit homeomorphism

The article by Alekseev, Meinrenken, and Woodward [AMW01] proves a large generalization of the above result, from which the above result (concerning the relation between $\mathcal{O}_{\alpha,\beta,\gamma}^{\text{add}}$ and $\mathcal{O}_{e^\alpha, e^\beta, e^\gamma}^{\text{mult}}$) follows as a special case. The purpose of this section is to sketch how the techniques of the general method specialize to this case (see [AMW01, Section 4] for comparison).

4.2.1 A Naive Guess

Given a triple of *mutually commuting* Hermitian matrices (A, B, C) satisfying $A+B+C=0$ it is not too difficult to see that

$$(A', B', C') = (\exp(A), \exp(B), \exp(C)) \quad (292)$$

is a triple of positive semidefinite matrices satisfying $A'B'C' = I_n$ such that if (α, β, γ) were the eigenvalues of (A, B, C) , then $(e^\alpha, e^\beta, e^\gamma)$ would be the eigenvalues of (A', B', C') . If the triple (A, B, C) are *not* mutually commuting, then it becomes possible that $A'B'C' \neq I_n$ (see [Hes20, Example 3, 31] for explicit examples and counterexamples) because

$$A'B'C' = \exp(A)\exp(B)\exp(C) \neq \exp(A+B+C) = \exp(0_n) = I_n. \quad (293)$$

In general, one has that

$$\exp(A)\exp(B)\exp(C) = \exp(A+B+C + \text{BCH}_3^\delta(A, B, C)) \quad (294)$$

where BCH_k for $k \geq 2$ is the Baker-Campbell-Hausdorff formula where for $k=2$

$$\text{BCH}_2(X, Y) = X + Y + \frac{1}{2}[X, Y] + \frac{1}{12}[X, [X, Y]] - \frac{1}{12}[Y, [X, Y]] + \dots \quad (295)$$

and, because BCH_2 is associative, BCH_3 is defined by

$$\text{BCH}_3(A, B, C) = \text{BCH}_2(\text{BCH}_2(A, B), C) = \text{BCH}_2(A, \text{BCH}_2(B, C)). \quad (296)$$

Moreover, we let

$$\text{BCH}_2^\delta(X, Y) := \text{BCH}_2(X, Y) - X - Y, \quad (297)$$

$$\text{BCH}_3^\delta(X, Y, Z) := \text{BCH}_3(X, Y, Z) - X - Y - Z. \quad (298)$$

In any case, we note that applying the matrix exponential is insufficient, in general, for transporting solutions from the additive problem to the multiplicative problem.⁴

Nevertheless, this naive attempt points in the right direction. If (A, B) are small in some suitable sense, say $(A, B) = (\varepsilon X, \varepsilon Y)$, then $\text{BCH}_2^\delta(A, B)$ will be even smaller (of order ε^2) because

$$\exp(\varepsilon X)\exp(\varepsilon Y) = \exp(\varepsilon X + \varepsilon Y + \frac{\varepsilon^2}{2}[X, Y] + O(\varepsilon^3)). \quad (299)$$

⁴It is worth noting that the matrix exponential still preserves some spectral information; for example, the well-known Golden-Thompson inequality states that $\text{Tr}(\exp(A+B)) \leq \text{Tr}(\exp(A)\exp(B))$ holds for all Hermitian matrices A, B . Also note that there exists generalizations of this inequality to three or more matrices [SBT17].

4.2.2 The Alekseev-Meinrenken-Woodward E -map

Given a Hermitian operator A with eigenvalues α , let $E(A)$ denote the unique upper triangular matrix with real diagonal entries satisfying⁵

$$E(A)E(A)^* = \exp(2A). \quad (300)$$

Under this definition, the singular values of $E(A)$ are equal to e^α .⁶ This definition places $E(A)$ in the canonical Borel subgroup B of $\mathrm{GL}(n, \mathbb{C})$ (which constitutes the Poisson-Lie group $K^* = \mathrm{U}(n)^*$ which is dual to $K = \mathrm{U}(n)$). Given an upper triangular matrix invertible matrix with real diagonals T , define

$$E^{-1}(T) = \log(\sqrt{TT^*}). \quad (301)$$

If A is a positive *definite* Hermitian matrix, $E(A)$ is invertible and thus $E^{-1}(E(A)) = A$.

The E -map is extended to a family of associated maps E_s parametrized by $s \in [0, \infty]$ (with $E \cong E_1$ and $E_s = E(s \cdot)$) as follows:

$$E_s(A)E_s(A)^* = \exp(2sA). \quad (302)$$

The singular values of $E_s(A)$ are therefore equal to $e^{s\alpha}$. Similarly, define

$$E_s^{-1}(T) = s^{-1} \log \sqrt{TT^*}. \quad (303)$$

Given a pair of Hermitian operators (A, B) with eigenvalues (α, β) , the singular values of $(E(A), E(B))$ will be (e^α, e^β) . It is not too surprising to notice that in general

$$A + B \neq E^{-1}(E(A)E(B)) \quad (304)$$

Moreover, the singular values of $E(A)E(B)$ (eigenvalues of $\sqrt{E(A)E(B)E(B)^*E(A)^*}$) will not necessarily be equal to the eigenvalues of $\exp(A + B)$. Consider instead the pair $(E_s(A), E_s(B))$; their singular values will be $(e^{s\alpha}, e^{s\beta})$ and the singular values of $E_s(A)E_s(B)$ will also be distinct from the eigenvalues of $\exp(s(A + B))$. However, when $s \approx 0$ is small, one might expect to observe approximation of the following type:

$$A + B \approx E_s^{-1}(E_s(A)E_s(B)). \quad (305)$$

Similarly, one might expect for small $s \approx 0$,

$$A + B + C \approx E_s^{-1}(E_s(A)E_s(B)E_s(C)). \quad (306)$$

⁵Note that $E(A)$ is related to the Cholesky decomposition of the positive semidefinite $\exp(2A)$ as follows. The Cholesky decomposition of $\exp(2A)$ is an upper triangular C such that $C^*C = \exp(2A)$. Given the unitary involutive permutation P which reverses the rows and columns, the Cholesky decomposition of $P\exp(2A)P$ is therefore an upper triangular $\tilde{C}$ such that $\tilde{C}^*\tilde{C} = P\exp(2A)P$. From this, one notes that $\exp(2A) = (P\tilde{C}^*P)(P\tilde{C}^*P)^*$ and therefore concludes that $E(A) = P\tilde{C}^*P$ where $\tilde{C}$ is the upper triangular matrix in the Cholesky decomposition of $P\exp(2A)P$.

⁶Note that [AMW01, Lemma 4.1] allows us, without loss of generality, to restrict our attention to the multiplicative problem for dressing orbits in $\mathrm{U}(n)^*$ instead of the double coset orbits $\mathrm{U}(n) \backslash \mathrm{GL}(n, \mathbb{C}) / \mathrm{U}(n)$.
TODO: Elaborate on the triple (quadruplet with unitary case) of problems earlier in this working document.

4.3 Moser's isotropy lemma

Here we summarize parts of the wikipedia page https://en.wikipedia.org/wiki/Moser%27s_trick.

Moser's isotropy lemma provides sufficient conditions to conclude when a parameterized family $\{\omega_t\}_{t \in [0,1]} \subset \Omega^k(M)$ of differential forms on a manifold M are related by a corresponding family of diffeomorphisms $\{\psi_t\}_{t \in [0,1]} \subset \text{Diff}(M)$ such that ω_0 is the pullback of ω_t by ψ_t , i.e. $\psi_t^* \omega_t = \omega_0$ where $\psi_0 = \text{id}_M$.

The sufficient condition is the existence of a smooth family $\{X_t\}_{t \in [0,1]} \subset \mathfrak{X}(M)$ of vector fields satisfying a certain differential equation involving $\{\omega_t\}$ and $\{X_t\}$:

$$\frac{d}{dt} \omega_t + \mathcal{L}_{X_t} \omega_t = 0. \quad \text{\textcolor{blue}{\code{eq:mosers_condition}}} \quad (307)$$

Given such a family of vector fields, with initial condition $\psi_0 = \text{id}_M$, one can define ψ_t by the solution to the equation $\frac{d}{dt} \psi_t = X_t \circ \psi_t$ for every $t \in [0, 1]$. Intuitively speaking, the diffeomorphism ψ_t sends the point $p_0 \in M$ to $p_t = \psi_t(p_0) \in M$ for all $t \in [0, 1]$ in a manner such that p_{t+dt} is obtained from p_t by flowing along the vector field X_t for a duration of dt . If X_t is a solution to Eq. (307), then the family $\{\psi_t^* \omega_t\}_{t \in [0,1]}$ of differential forms is actually independent of $t \in [0, 1]$ in the sense that

$$\frac{d}{dt} (\psi_t^* \omega_t) = \psi_t^* \left(\frac{d}{dt} \omega_t + \mathcal{L}_{X_t} \omega_t \right) = 0. \quad (308)$$

In the special case where the family of differential forms $\{\omega_t\}_{t \in [0,1]}$ under consideration is a family of symplectic forms, the conditions of Moser's lemma simplify both because ω_t is closed ($d\omega_t = 0$) and because ω_t is non-degenerate meaning that the family of vector fields $\{X_t\}_{t \in [0,1]}$ may be related to a corresponding family of 1-forms $\{\alpha_t\}_{t \in [0,1]}$ defined by

$$\alpha_t = -\iota_{X_t} \omega_t. \quad (309)$$

In this setting, the existence of vector fields $\{X_t\}_{t \in [0,1]}$ satisfying Eq. (307) is supplanted by the existence 1-forms satisfying Eq. (310):

$$\frac{d}{dt} \omega_t = d\alpha_t. \quad \text{\textcolor{blue}{\code{eq:symplectic_mosers_condition}}} \quad (310)$$

The satisfaction of Eq. (307) then follows from Cartan's formula because

$$\mathcal{L}_{X_t}(\omega_t) = \underbrace{\iota_{X_t}(d\omega_t)}_{=0} + d(\underbrace{\iota_{X_t} \omega_t}_{=-\alpha_t}) = -d\alpha_t. \quad (311)$$

4.4 The β form

An important entity in the “linearization” paper due to Alekseev, Meinrenken, and Woodward [AMW01] is the 1-form on $\mathfrak{k}^*$ denoted by β . It is defined according to [AMW01, Equation (8)] by

$$\beta = \frac{1}{2i} \mathcal{H} (E^* B^C(\theta^L, \theta^{\dagger;L})) \in \Omega^1(\mathfrak{k}^*) \quad (312)$$

In order to understand this 1-form better we analyze each of its pieces separately.

4.4.1 The Homotopy Operator

The operator $\mathcal{H} : \Omega^2(\mathfrak{k}^*) \rightarrow \Omega^1(\mathfrak{k}^*)$ is the homotopy operator sending 2-forms to 1-forms on $\mathfrak{k}^*$ (it also sends k -forms to $k-1$ forms) which has the exterior derivative $d : \Omega^1(\mathfrak{k}^*) \rightarrow \Omega^2(\mathfrak{k}^*)$ as its left-inverse when restricted to closed 2-forms in the sense that $d\omega = 0$ implies $\omega = d(\mathcal{H}\omega)$ and thus ω is exact. This operator appears in the context of the Poincaré lemma which states that closed k -forms on a open ball in $\mathbb{R}^n$ are also exact for all $1 \leq k \leq n$.

As we are only interested in the application of $\mathcal{H}$ to 2-forms on $\mathfrak{k}^* \simeq \mathbb{R}^n$, it is relatively easy to state $\mathcal{H}$ as follows. Let $h : \mathfrak{k}^* \times [0, 1] \rightarrow \mathfrak{k}^*$ be defined by the scaling $h_t : \mathfrak{k}^* \rightarrow \mathfrak{k}^*$ by $t \in [0, 1]$, i.e.

$$h(x_1, \dots, x_n, t) := h_t(x_1, \dots, x_n) = (tx_1, \dots, tx_n). \quad (313)$$

Note that h_t is a diffeomorphism of $\mathfrak{k}^*$ for all $t \in (0, 1]$.

Consider a generic 2-form, denoted by ω , on $\mathfrak{k}^* \simeq \mathbb{R}^n$ in the coordinates $(x_1, \dots, x_n)$:

$$\omega = \sum_{i < j} \omega_{i,j} dx^i \wedge dx^j \quad (314)$$

The pullback of ω along h_t is therefore simply

$$h_t^* \omega = \sum_{i < j} (\omega_{i,j} \circ h_t) d(x^i \circ h_t) \wedge d(x^j \circ h_t), \quad (315)$$

$$= t^2 \sum_{i < j} (\omega_{i,j} \circ h_t) dx^i \wedge dx^j \quad (316)$$

which means ω can be re-expressed as an integral:

$$\omega = \underbrace{h_1^* \omega}_{=\omega} - \underbrace{h_0^* \omega}_{=0} = \int_0^1 \frac{d}{dt} (h_t^* \omega) dt = \int_0^1 h_t^* (\mathcal{L}_{\xi_t} \omega) dt = \int_0^1 h_t^* (d\iota_{\xi_t} \omega + \iota_{\xi_t} d\omega) dt \quad (317)$$

where ξ_t is the vector field on $\mathfrak{k}^*$ such that $(\xi_t)_x$ is the tangent vector to the curve $t \mapsto h_t(x)$, **TODO: Figure out what this vector field is.** If ω is additionally closed ($d\omega = 0$), then

$$\omega = \int_0^1 (h_t^* d\iota_{\xi_t} \omega) dt = d \left(\int_0^1 (h_t^* \iota_{\xi_t} \omega) dt \right) \quad (318)$$

where the exterior derivative commutes with the pullback by h_t^* and with the integration over $t \in [0, 1]$.

Given a closed $\omega \in \Omega^2(\mathfrak{k}^*)$, the 1-form $\mathcal{H}(\omega)$ is therefore defined as

$$\mathcal{H}(\omega) := \int_0^1 dt h_t^* \iota_{\xi_t} \omega. \quad (319)$$

Given a vector field $Y \in \mathfrak{X}(\mathfrak{k}^*)$, **TODO: Figure out how to calculate this contraction in local coordinates explicitly.**

4.4.2 The Maurer-Cartan Forms

TODO:

4.4.3 Bringing it altogether

TODO:

4.5 An application of the Moser isotropy lemma

In this section, we borrow some of the notation of Lu and Evens [LE03]. Let $U(n)^*$ ($= B = AN$) denote the upper triangular Borel subgroup of $GL(n, \mathbb{C})$ and define two maps

$$\mathbf{m} : (U(n)^*)^{\times 3} \rightarrow U(n)^* :: (A', B', C') \mapsto A'B'C', \quad (320)$$

$$\mathbf{a} : (iu(n))^{\times 3} \rightarrow iu(n) :: (A, B, C) \mapsto A + B + C. \quad (321)$$

In the previous section, we defined a pair of maps

$$E_s : iu(n) \rightarrow U(n)^*. \quad (322)$$

$$E_s^{-1} : U(n)^* \rightarrow iu(n). \quad (323)$$

The proof of Thompson's conjecture follows if one can prove there exists a $U(n)$ -equivariant diffeomorphism

$$L : (U(n)^*)^{\times 3} \rightarrow (iu(n))^{\times 3} \quad (324)$$

such that $\mathbf{m} = E_1 \circ \mathbf{s} \circ L$, i.e. for all $(A', B', C') \in (U(n)^*)^{\times 3}$ one has a corresponding triple

$$(A, B, C) := L(A', B', C') \quad (325)$$

(with components possibly depending on each of (A', B', C') non-trivially) such that

$$E_1(A + B + C) = E_1(\mathbf{s}(A, B, C)) = \mathbf{m}(A', B', C') = A'B'C' \quad (326)$$

In this manner, if $A'B'C' = I_n$, then one automatically has a triple (A, B, C) such that $A + B + C = 0_n$. The equivariance of L ensures that the appropriate spectral data is preserved. In fact, the desired map will be of the form $L = (E_1^{-1})^{\times 3} \circ \psi_1$ where ψ_1 is a diffeomorphism of $(U(n)^*)^{\times 3}$, meaning there exists

$$(A'', B'', C'') = \psi_1(A', B', C') \in (U(n)^*)^{\times 3} \quad (327)$$

such that the following holds

$$E_1(A + B + C) = E_1(E_1^{-1}(A'') + E_1^{-1}(B'') + E_1^{-1}(C'')) = A'B'C' \quad (328)$$

4.6 Linearization commutes with products

In an earlier article by Alekseev, Meinrenken, and Woodward [AMW01], a map sending $\mathfrak{k}^*$ to the dual Poisson-Lie group K^* , denoted by $E : \mathfrak{k}^* \rightarrow K^*$, is defined as follows. Given the (local) Cartan involution $\kappa : \mathfrak{g} \rightarrow \mathfrak{g}$ (complex conjugation) (denoted by θ in my thesis), the map $\dagger = -\kappa$ becomes an anti-involution (called the $*$ -involution in my thesis, “anti-” refers to anti-linear).

Now let $B : \mathfrak{k} \times \mathfrak{k} \rightarrow \mathbb{R}$ be an invariant inner product on $\mathfrak{k}$ (for $\mathfrak{k} = \mathfrak{su}(d)$ think the negative of the Killing form $K(a, b) = \text{Tr}(\text{ad}_a \text{ad}_b)$ (which is $-2d \text{Tr}(ab)$ for $\mathfrak{su}(d)$) and let $B^\sharp : \mathfrak{k}^* \rightarrow \mathfrak{k}$ denote the isomorphism identifying linear functionals $\mathfrak{k}^*$ on $\mathfrak{k}$ with elements of $\mathfrak{k}$ induced by B (i.e., $B(B^\sharp(F), X) = F(X)$). Then $iB^\sharp(F) \in i\mathfrak{k}$ and $\exp(iB^\sharp(F)) \in P \subseteq G$. Furthermore, every element in P can be written uniquely as $ll^\dagger$ for some $l \in K^* = AN$. Therefore, there exists a map $E : \mathfrak{k}^* \rightarrow K^*$ sending $F \in \mathfrak{k}^*$ to $E(F) \in K^*$ uniquely determined by the non-linear equation

$$E(F)E(F)^\dagger = \exp(iB^\sharp(F)). \quad \text{\texttt{\{eq:E-map\}}} \quad (329)$$

This map is equivariant in the sense that the coadjoint action of K on $F \in \mathfrak{k}^*$ coincides with the so-called *dressing* action on K^* .

For the group $K = U(d)$ of unitary matrices, its Lie algebra $\mathfrak{k} = \mathfrak{u}(d)$ consists of skew-Hermitian matrices and the invariant inner product B on $\mathfrak{u}(d)$ can be taken to be $B(A_1, A_2) = -\text{Tr}(A_1 A_2)$ (or any constant multiple of this). Then $B^\sharp$ identifies linear functional over skew-Hermitian matrices $F \in \mathfrak{u}(d)^*$ with the skew-Hermitian matrix $A_F := B^\sharp(F)$ defined by $B(A_F, \cdot) = -\text{Tr}(A_F \cdot) = F(\cdot)$. Then iA_F is a Hermitian matrix and $\exp(iA_F)$ is positive semidefinite matrix with Cholesky decomposition $\exp(iA_F) = L_F^\dagger L_F$ where L_F is upper-triangular.

In the context of additive and multiplicative Horn problems, the map E (acting on Hermitian matrices now) satisfies the equation

$$\text{Sing} \circ E = \exp \circ \text{Eigen} \quad (330)$$

Given the eigenvalues $\text{Eigen}(H) = \lambda(H)$ of a Hermitian matrix H , the values appearing in $\exp(\lambda(H))$ (taken component-wise) are equivalent to the singular values of $E(H)$, i.e. the square roots of the eigenvalues of $E(H)^\dagger E(H)$. Note that this statement is not totally consistent with Eq. (329); the square-root in the definition of singular values must be related to the choice of inner product B ?

Now given an invariant inner product $B_s : \mathfrak{k} \times \mathfrak{k} \rightarrow \mathbb{R}$ on $\mathfrak{k}$ (possibly depending on some data s), one can consider the corresponding equivariant map $E_s : \mathfrak{k}^* \rightarrow K^*$. Given a moment map $\Phi : M \rightarrow \mathfrak{k}^*$ for a Hamiltonian K -manifold M , the map $\Psi_s : M \rightarrow K^*$ formed by composing Φ with E_s yields the K^* -valued moment map Ψ_s .

The family of invariant inner products eventually considered are those which are obtained by rescalings by a positive $s > 0$, i.e. $B_s = s^{-1}B$. In this setting, the map E_s satisfies

$$E_s(F)E_s(F)^\dagger = \exp(iB_s^\sharp(F)), \quad \text{\textcolor{blue}{(eq:Es-map)}} \quad (331)$$

but $s^{-1}B(B_s^\sharp(F), \cdot) = B_s(B_s^\sharp(F), \cdot) = F(X)$ should be independent of s so clearly $B_s^\sharp(F) = sB^\sharp(F)$.

In any case, if $\Phi_1 : M_1 \rightarrow \mathfrak{k}^*$ and $\Phi_2 : M_2 \rightarrow \mathfrak{k}^*$ are two moment maps, then the moment map on the product $M_1 \times M_2$ is given by a sum

$$\Phi_{1,2}(x_1, x_2) = \Phi_1(x_1) + \Phi_2(x_2). \quad (332)$$

Now for each s , there exists a moment map $\Phi_{1,2}^s : M_1 \times M_2 \rightarrow \mathfrak{k}^*$ (with modified symplectic form ω_s on $M_1 \times M_2$) such that $\Phi_{1,2}^s$ satisfies

$$\Phi_{1,2}^s = E_s^{-1}[(E_s \circ \Phi_1)(E_s \circ \Phi_2)] \quad (333)$$

and a linearization property $\lim_{s \rightarrow 0^+} \Phi_{1,2}^s = \Phi_1 + \Phi_2$.

In this setting, $K = U(n)$, its double is its complexification $K^\mathbb{C} = \text{GL}(n, \mathbb{C})$, and the Lie algebra $\mathfrak{k} = \mathfrak{u}(n)$ consists of anti-Hermitian $n \times n$ matrices. One can identify the dual $\mathfrak{u}(n)^*$ with the space of $n \times n$ Hermitian matrices using the invertible linear map

$$H :: \xi \mapsto H_\xi \quad (334)$$

such that each $\xi \in \mathfrak{u}(n)^*$ acting on the skew-Hermitian matrix $x \in \mathfrak{u}(n)$ is identified with the Hermitian matrix H_ξ satisfying

$$\xi(x) = \langle \xi, x \rangle = \text{Im Tr}(H_\xi x) = -i \text{Tr}(H_\xi x). \quad (335)$$

In the notation of [AMW01], we have $H = iB^\sharp$.

Note there exists a unique $\mathfrak{u}(n)^*$ with the Poisson bracket $\{\cdot, \cdot\}$ which is the unique bracket satisfying $\{f_x, f_y\} = f_{[x, y]}$ where f_x is the linear function on $\mathfrak{u}(n)^*$ defined by $f_x(\xi) = \langle \xi, x \rangle = \xi(x)$. Through the identification $\xi \mapsto H_\xi$ made above, the space of Hermitian $n \times n$ matrices inherits this Poisson structure in a manner which is dependent upon the identification made.

4.7 Coordinates for $\mathrm{SL}_2(\mathbb{C})$

In the multiplicative variant of Horn's problem, one is interested in matrices $A \in \mathrm{SL}(n, \mathbb{C})$ and their *singular values* $\sigma(A) := \lambda(\sqrt{AA^*})$. For example, when $n = 2$, we have coordinates $(a, b, c, d) \in \mathbb{C}^4$ subject to the constraint $ad - bc = 1$ specifying $A \in \mathrm{SL}(2, \mathbb{C})$ via

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}. \quad (336)$$

Alternatively, using the polar decomposition we have coordinates $(x, y, z) \in \mathbb{R}^2 \times \mathbb{C}$ subject to $x^2 - y^2 - |z|^2 = 1$ and $(u, v) \in \mathbb{C}^2$ subject to $|u|^2 + |v|^2 = 1$

$$A = \exp \left[\begin{pmatrix} x+y & z \\ z^* & x-y \end{pmatrix} \right] \begin{pmatrix} u & v \\ -v^* & u^* \end{pmatrix}. \quad (337)$$

Using these coordinates, AA^* depends only on $(x, y, z) \in \mathbb{R}^2 \times \mathbb{C}$,

$$AA^* = \exp \left[2 \begin{pmatrix} x+y & z \\ z^* & x-y \end{pmatrix} \right], \quad (338)$$

and then $\sqrt{AA^*}$ is simply

$$\sqrt{AA^*} = \exp \left[\begin{pmatrix} x+y & z \\ z^* & x-y \end{pmatrix} \right]. \quad (339)$$

The characteristic polynomial of the exponent is

$$\det \begin{pmatrix} x+y-\lambda & z \\ z^* & x-y-\lambda \end{pmatrix} = \lambda^2 - 2x\lambda + 1. \quad (340)$$

At this stage, it is useful to let $w = \sqrt{x^2 - 1} = \sqrt{y^2 + |z|^2}$. Therefore, the eigenvalues of the exponent are

$$x \pm \sqrt{x^2 - 1} = x \pm \sqrt{y^2 + |z|^2} = x \pm w. \quad (341)$$

Therefore the eigenvalues of $\sqrt{AA^*}$ (singular values of A) are given by

$$\lambda(\sqrt{AA^*}) = (e^{x+w}, e^{x-w}) \quad (342)$$

4.8 The Ginzburg-Weinstein diffeomorphism

At the core of the relationships holding between the multiplicative and additive perspectives seems to be this mysterious *Ginzburg-Weinstein* diffeomorphism from $\mathfrak{u}(n)^*$ (the dual of the Lie algebra of $U(n)$) equipped with the KKS Poisson bracket and the Poisson Lie group

$U(n)^*$ with some unique Poisson Lie bracket which is compatible with the KKS Poisson bracket in some sense. The details of this construction are the content of Alekseev and Meinrenken [AM07].

The Lie algebra $\mathfrak{u}(n)$ consists of $n \times n$ complex matrices X which are skew-Hermitian ($X^* = -X$). The dual of this algebra can be identified with the space of $n \times n$ complex matrices H which are Hermitian ($H^* = H$). Specifically, one can identify H with the linear map sending $X \in \mathfrak{u}(n)$ to $\langle X, H \rangle := \text{Im Tr}(HX) \in \mathbb{R}$. Moreover, the Poisson Lie group $U(n)^*$ can be identified with the space of $n \times n$ complex upper triangular matrices with positive real numbers along the diagonal.

In the special case where $n = 2$, the aforementioned isomorphism $\varphi : \mathfrak{u}(2)^* \rightarrow U(2)^*$ admits an of a simple expression. Let $H \in \mathfrak{u}(2)^*$ be written in Pauli coordinates $(x, y, z) \in \mathbb{R}^3$ (without loss of generality, H is taken to be traceless because $\varphi(H + tI) = e^t \varphi(H)$),

$$H = \begin{pmatrix} z & x + iy \\ x - iy & -z \end{pmatrix}, \quad (343)$$

and furthermore define $(\rho, r, \phi) \in \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0} \times [0, 2\pi)$ such that

$$r = (x^2 + y^2 + z^2)^{\frac{1}{2}}, \quad x + iy = \rho e^{i\phi}. \quad (344)$$

Then the Ginzburg-Weinstein map φ sends H to (taken from [Anton's ICM2014 talk on youtube](#)):

$$\varphi \begin{pmatrix} z & x + iy \\ x - iy & -z \end{pmatrix} = \begin{pmatrix} e^z & e^{i\phi}(e^{+2r} + e^{-2r} - e^{+2z} - e^{-2z})^{\frac{1}{2}} \\ 0 & e^{-z} \end{pmatrix}. \quad (345)$$

This map is interesting because it is a Poisson map in the sense that for all $H \in \mathfrak{u}(n)^*$ and all $F, G \in C^\infty(U(2)^*)$,

$$\{F, G\}_{U(2)^*}(\varphi(H)) = \{F \circ \varphi, G \circ \varphi\}_{\mathfrak{u}(2)^*}(H) \quad (346)$$

where $\{\cdot, \cdot\}_{U(2)^*}$ is the unique Poisson bracket on $U(2)^*$ satisfying certain conditions (according to Drinfeld, Semenov-Tian-Shansky, Lu-Weinstein (see the youtube talk)) where $\{\cdot, \cdot\}_{\mathfrak{u}(2)^*}$ is the Kirillov-Kostant-Souriau (KKS) Poisson bracket on $\mathfrak{u}(n)^*$ defined to be

$$\{f_{A_1}, f_{A_2}\}_{\mathfrak{u}(2)^*} = f_{[A_1, A_2]} \quad (347)$$

where $f_A : H \rightarrow \langle H, A \rangle = \text{Im Tr}(HA)$ is a linear function on $\mathfrak{u}(n)^*$.

Furthermore, this Poisson diffeomorphism belongs to a family of Poisson diffeomorphisms obtained by scaling all of the exponential quantities except ϕ :

$$\varphi_\tau \begin{pmatrix} z & x + iy \\ x - iy & -z \end{pmatrix} = \begin{pmatrix} e^{\tau z} & e^{i\phi}(e^{+2\tau r} + e^{-2\tau r} - e^{+2\tau z} - e^{-2\tau z})^{\frac{1}{2}} \\ 0 & e^{-\tau z} \end{pmatrix}. \quad (348)$$

If $\tau = 1$, then $\varphi_\tau = \varphi$. Also if $\tau \rightarrow \infty$, then

$$\varphi_{\tau \rightarrow \infty} \begin{pmatrix} z & x + iy \\ x - iy & -z \end{pmatrix} \approx \begin{pmatrix} e^{\tau z} & e^{i\phi + \tau r} \\ 0 & e^{-\tau z} \end{pmatrix}, \quad (349)$$

because $r \geq z$ always.

4.9 Leading coefficient of L-R polynomials

The probability measure associated to the pushforward of the G -invariant probability measure over the product of orbits, $\mathcal{O}_\alpha \times \mathcal{O}_\beta$, through the addition map $(A, B) \mapsto A + B$ and then through the orbital projection $V \mapsto V/G$ is termed the *Horn probability measure*. In [CZ18], Coquereaux and Zuber consider two special cases of the Horn probability measure: the coadjoint case wherein $V = \mathfrak{g}^*$ and the orbits under consideration are coadjoint orbits and the “self-adjoint” case where the groups $\mathrm{SO}(n)$, $\mathrm{SU}(n)$, and $\mathrm{USp}(n)$ are acting on real-symmetric, complex-Hermitian, and quaternionic self-dual matrices respectively. Focusing on the coadjoint case, their derivation of the integral form of the Horn probability measure is essentially identical (modulo notation and normalization conventions) to the form considered by Klyachko [Kly00]. In particular, the Horn probability density for the “random” orbit $\mathcal{O}_\gamma$ is

$$p(\gamma \mid \alpha, \beta) = \frac{1}{(2\pi)^r |W|} \left(\frac{\Delta_{\mathfrak{g}}(\gamma)}{\Delta_{\mathfrak{g}}(\rho)} \right)^2 \int_{\mathfrak{t}} dt \Delta_{\mathfrak{g}}(t) \mathcal{H}(\alpha, it) \mathcal{H}(\beta, it) \mathcal{H}(\gamma, it)^* \quad (350)$$

Here the expression $\Delta_{\mathfrak{g}}(x) = \prod_{\alpha > 0} \langle \alpha, x \rangle$ is a product over the positive roots of $\mathfrak{g}$, $|W|$ is the order of the Weyl group and r is the rank of K . The expression $\mathcal{H}(\alpha, it)$ for $\alpha, t \in C_+$ lying in the dominant Weyl chamber is given by the Harish-Chandra orbital function:

$$\mathcal{H}(\alpha, it) = \int_G dg \exp(i \langle g \cdot \alpha, x \rangle) \quad (351)$$

where $\langle \cdot, \cdot \rangle$ is -1 times the Killing form on $\mathfrak{g}$.

The so-called *volume function* considered in [CZ18] is the expression

$$J(\alpha, \beta; \gamma) = \frac{\Delta_{\mathfrak{g}}(\alpha) \Delta_{\mathfrak{g}}(\beta)}{\Delta_{\mathfrak{g}}(\gamma) \Delta_{\mathfrak{g}}(\rho)} p(\gamma \mid \alpha, \beta). \quad (352)$$

In the case where $G = \mathrm{SU}(n)$, the explicit connection to the Littlewood-Richardson coefficients made by [CZ18] is to related the integral form of $p(\gamma \mid \alpha, \beta)$ over Harish-Chandra orbital functions in terms of an integral expression reminiscent of the integral expression for the Littlewood-Richardson coefficient $c_{\lambda, \mu}^\nu$ (note that $N_{\lambda, \mu}^\nu$ is used in [CZ18]) in terms of characters:

$$c_{\lambda, \mu}^\nu = \int_{\mathbb{T}_n} dT \chi_\lambda(T) \chi_\mu(T) \chi_\nu^*(T). \quad (353)$$

where dT is the normalized Haar measure on $\mathbb{T}_n$ given by [CZ18, Equation (17)]. Their method for establishing the connection is through Weyl’s character formula which yields [CZ18, Section 2.2.1]:

$$\frac{\chi_\lambda(e^{it})}{\chi_\lambda(1)} = \frac{\Delta_{\mathfrak{g}}(it)}{\Delta_{\mathfrak{g}}(e^{it})} \mathcal{H}(\lambda', it) \quad (354)$$

where the prime $\lambda' = \lambda + \rho$ is λ shifted by the Weyl vector $\rho = \sum_{\alpha > 0} \alpha/2$ (the half sum of positive roots). Altogether, a sketch of their asymptotic correspondence is provided by taking $k \in \mathbb{N}$ very large and then [CZ18, Equation (44)] reads:

$$c_{k\lambda, k\mu}^{k\nu} \approx \mathcal{J}(k\lambda + \rho, k\mu + \rho; k\nu + \rho) \approx \mathcal{J}(k\lambda, k\beta; k\gamma) = s^{(n-1)(n-2)/2} \mathcal{J}(\alpha, \beta; \gamma). \quad (355)$$

4.10 Singular values of partial products

One objective of this research project is to solve the following generalization of the multiplicative version of Horn’s problem.

Problem 3 (Trivariate Multiplicative Horn’s problem). Let $d \in \mathbb{N}$ and let

$$(s_A, s_B, s_C, s_{AB}, s_{BC}, s_{ABC}) \in (\mathbb{R}^d)^{\times 6} \quad (356)$$

be given as input. Decide if there exists matrices $M_A, M_B, M_C \in \mathrm{SL}(d, \mathbb{C})$ satisfying

$$\begin{aligned} \mathrm{sing}(M_A) &= s_A, & \mathrm{sing}(M_A M_B) &= s_{AB}, \\ \mathrm{sing}(M_B) &= s_B, & \mathrm{sing}(M_B M_C) &= s_{BC}, \\ \mathrm{sing}(M_C) &= s_C, & \mathrm{sing}(M_A M_B M_C) &= s_{ABC}, \end{aligned} \quad (357)$$

or equivalently,

$$\begin{aligned} \mathrm{spec}(\sqrt{M_A M_A^*}) &= s_A, & \mathrm{spec}(\sqrt{M_A M_B M_B^* M_A^*}) &= s_{AB}, \\ \mathrm{spec}(\sqrt{M_B M_B^*}) &= s_B, & \mathrm{spec}(\sqrt{M_A M_B M_B^* M_A^*}) &= s_{BC}, \\ \mathrm{spec}(\sqrt{M_C M_C^*}) &= s_C, & \mathrm{spec}(\sqrt{M_A M_B M_C M_C^* M_B^* M_A^*}) &= s_{ABC}, \end{aligned} \quad (358)$$

References

- [AH11] Daiki Akimoto and Masahito Hayashi. “Discrimination of the change point in a quantum setting”. In: *Physical Review A* 83.5 (2011), p. 052328.
- [Ale+18] Anton Alekseev et al. “Concentration of symplectic volumes on Poisson homogeneous spaces”. In: *arXiv preprint arXiv:1808.06975* (2018).
- [Ale97] A Yu Alekseev. “On Poisson actions of compact Lie groups on symplectic manifolds”. In: *Journal of Differential Geometry* 45.2 (1997), pp. 241–256.
- [AM07] Anton Alekseev and Eckhard Meinrenken. “Ginzburg-Weinstein via Gelfand-Zeitlin”. In: *Journal of Differential Geometry* 76.1 (2007), pp. 1–34.
- [AMS20] Victor Alekseev, Andrey Morozov, and Alexey Sleptsov. “Multiplicity-free $\mathrm{Uq}(\mathfrak{slN})$ 6-j symbols: relations, asymptotics, symmetries”. In: *Nuclear Physics B* 960 (2020), p. 115164.
- [AMW00] Anton Alekseev, Eckhard Meinrenken, and Christopher Woodward. “Group-valued equivariant localization”. In: *Inventiones mathematicae* 140.2 (2000), pp. 327–350.
- [AMW01] Anton Alekseev, Eckhard Meinrenken, and Chris Woodward. “Linearization of Poisson actions and singular values of matrix products”. In: *Annales de l’institut Fourier*. Vol. 51. 6. 2001, pp. 1691–1717.
- [AMW99] Anton Alekseev, Eckhard Meinrenken, and Chris Woodward. “Duistermaat-Heckman distributions for group valued moment maps”. In: *arXiv preprint math/9903087* (1999).
- [APS17a] Anton Alekseev, Masha Podkopaeva, and Andras Szenes. “A symplectic proof of the Horn inequalities”. In: *Advances in Mathematics* 318 (2017), pp. 711–736.

- [APS17b] Anton Alekseev, Masha Podkopaeva, and Andras Szenes. “The Horn problem and planar networks”. In: *Advances in Mathematics* 318 (2017), pp. 678–710.
- [AW97] Sharad Agnihotri and Chris Woodward. “Eigenvalues of products of unitary matrices and quantum Schubert calculus”. In: *arXiv preprint alg-geom/9712013* (1997).
- [Bac10] Miriam Backens. “The 6j symbols and an extended version of Horn’s problem”. In: (2010).
- [Bar03] John W Barrett. “Geometrical measurements in three-dimensional quantum gravity”. In: *International Journal of Modern Physics A* 18.suppl02 (2003), pp. 97–113.
- [Bel08] Prakash Belkale. “Quantum generalization of the Horn conjecture”. In: *Journal of the American Mathematical Society* 21.2 (2008), pp. 365–408.
- [Bha01] Rajendra Bhatia. “[Linear algebra to quantum cohomology: the story of Alfred Horn’s inequalities](#)”. In: *The American Mathematical Monthly* 108.4 (2001), pp. 289–318.
- [BK16] Prakash Belkale and Shrawan Kumar. “The multiplicative eigenvalue problem and deformed quantum cohomology”. In: *Advances in Mathematics* 288 (2016), pp. 1309–1359.
- [Boa07] Philip P Boalch. “[Regge and Okamoto symmetries](#)”. In: *Communications in mathematical physics* 276.1 (2007), pp. 117–130.
- [Bür+19] Peter Bürgisser et al. “Towards a theory of non-commutative optimization: Geodesic 1st and 2nd order methods for moment maps and polytopes”. In: *2019 IEEE 60th Annual Symposium on Foundations of Computer Science (FOCS)*. IEEE. 2019, pp. 845–861.
- [Chr08] Matthias Christandl. “A Quantum Information-Theoretic Proof of the Relation between Horn’s Problem and the Littlewood-Richardson Coefficients”. In: *Lecture Notes in Computer Science*. Springer Berlin Heidelberg, 2008, pp. 120–128.
- [CMZ19] Robert Coquereaux, Colin McSwiggen, and Jean-Bernard Zuber. “Revisiting Horn’s problem”. In: *Journal of Statistical Mechanics: Theory and Experiment* 2019.9 (2019), p. 094018.
- [CMZ20] Robert Coquereaux, Colin McSwiggen, and Jean-Bernard Zuber. “On Horn’s problem and its volume function”. In: *Communications in Mathematical Physics* 376.3 (2020), pp. 2409–2439.
- [Coq23] Robert Coquereaux. “About integer-valued variants of the theta and 6j symbols”. In: *Journal of Mathematical Physics* 64.3 (2023).
- [CP95] Vyjayanthi Chari and Andrew N Pressley. *A guide to quantum groups*. Cambridge university press, 1995.
- [CŞW18] Matthias Christandl, M Burak Şahinoğlu, and Michael Walter. “Recoupling coefficients and quantum entropies”. In: *Annales Henri Poincaré*. Vol. 19. 2. Springer. 2018, pp. 385–410.
- [CW06] Nicola Ciccoli and Pawel Witkowski. “From Poisson to quantum geometry”. In: *Lecture Notes, available at <http://toknotes.mimuw.edu.pl>* (2006).

- [CZ18] Robert Coquereaux and Jean-Bernard Zuber. “From orbital measures to Littlewood–Richardson coefficients and hive polytopes”. In: *Annales de l’Institut Henri Poincaré D* 5.3 (2018), pp. 339–386.
- [DH05] Sumit Daftuar and Patrick Hayden. “Quantum state transformations and the Schubert calculus”. In: *Annals of Physics* 315.1 (2005), pp. 80–122.
- [DZ10] Amir Dembo and Ofer Zeitouni. *Large deviations techniques and applications*. Berlin: Springer, 2010.
- [Esp12] Chiara Esposito. “On the classical and quantum momentum map”. In: (2012). 1203.4112: [arxiv](#).
- [Feg78] HD Fegan. “The heat equation on a compact Lie group”. In: *Transactions of the American Mathematical Society* 246 (1978), pp. 339–357.
- [FK00] Laurent Freidel and Kirill Krasnov. “Simple spin networks as Feynman graphs”. In: *Journal of Mathematical Physics* 41.4 (2000), pp. 1681–1690.
- [FKL10] Laurent Freidel, Kirill Krasnov, and Etera R Livine. “Holomorphic factorization for a quantum tetrahedron”. In: *Communications in Mathematical Physics* 297.1 (2010), pp. 45–93.
- [FL03] Laurent Freidel and David Louapre. “Asymptotics of 6j and 10j symbols”. In: *Classical and Quantum Gravity* 20.7 (2003), p. 1267.
- [FNR07] L Freidel, K Noui, and Ph Roche. “6J symbols duality relations”. In: *Journal of mathematical physics* 48.11 (2007).
- [FT14] Peter J Forrester and Colin J Thompson. “The Golden-Thompson inequality: Historical aspects and random matrix applications”. In: *Journal of Mathematical Physics* 55.2 (2014).
- [Ful00] William Fulton. “Eigenvalues, invariant factors, highest weights, and Schubert calculus”. In: *Bulletin of the American Mathematical Society* 37.3 (2000), pp. 209–249.
- [FW20] Cole Franks and Michael Walter. “Minimal length in an orbit closure as a semi-classical limit”. In: (2020). arXiv: [2004.14872v2](#).
- [Hel84] Sigurdur Helgason. *Groups and geometric analysis: integral geometry, invariant differential operators, and spherical functions*. Pure and applied mathematics, 1984.
- [Hel94] Sigurdur Helgason. *Geometric Analysis on Symmetric Spaces*. Vol. 39. American Mathematical Soc., 1994.
- [Hes20] Sarah Tanja Hess. “Examination of the connection between the Horn Problem and the Lax Conjecture”. In: (2020).
- [HHR16] Hal M Haggard, Muxin Han, and Aldo Riello. “Encoding curved tetrahedra in face holonomies: phase space of shapes from group-valued moment maps”. In: *Annales Henri Poincaré*. Vol. 17. Springer. 2016, pp. 2001–2048.
- [HK05] Andre Henriques and Joel Kamnitzer. *Crystals and coboundary categories*. 2005. arXiv: [math/0406478](#) [[math.QA](#)].
- [HK06] André Henriques and Joel Kamnitzer. “The octahedron recurrence and gln crystals”. In: *Advances in Mathematics* 206.1 (2006), pp. 211–249.

- [Kly00] Alexander A Klyachko. “Random walks on symmetric spaces and inequalities for matrix spectra”. In: *Linear Algebra and its Applications* 319.1-3 (2000), pp. 37–59.
- [Knu00] Allen Knutson. “[The symplectic and algebraic geometry of Horn’s problem](#)”. In: *Linear Algebra and its Applications* 319.1-3 (2000), pp. 61–81.
- [KR90] AN Kirillov and N Yu Reshetikhin. “Representations of the Algebra $U_q(\mathfrak{sl}(2))$, q -Orthogonal Polynomials and Invariants of Links”. In: *New developments in the theory of knots* 11 (1990), p. 202.
- [KTW03] Allen Knutson, Terence Tao, and Christopher T Woodward. “A positive proof of the Littlewood-Richardson rule using the octahedron recurrence”. In: *arXiv preprint math/0306274* (2003).
- [Kul97] Solomon Kullback. *Information theory and statistics*. Courier Corporation, 1997.
- [KW01] Michael Keyl and Reinhard F Werner. “[Estimating the spectrum of a density operator](#)”. In: *Physical Review A* 64.5 (2001), p. 052311.
- [Lam05] McKenzie Lamb. “A proof of the Thompson conjecture using Poisson geometry”. In: (2005).
- [LE03] Jiang-Hua Lu and Sam Evens. “Thompson’s conjecture for real semi-simple Lie groups”. In: (2003). arXiv: [math/0310098](#).
- [LS91] Serge Levendorskii and Yan Soibelman. “Algebras of functions on compact quantum groups, Schubert cells and quantum tori”. In: *Communications in Mathematical Physics* 139.1 (1991), pp. 141–170.
- [LW90] Jiang-Hua Lu and Alan Weinstein. “Poisson Lie groups, dressing transformations, and Bruhat decompositions”. In: *Journal of Differential geometry* 31.2 (1990), pp. 501–526.
- [McS21] Colin McSwiggen. “The Harish-Chandra integral: an introduction with examples”. In: *L’Enseignement Mathématique* 67.3 (2021), pp. 229–299.
- [Nis00] Kyo Nishiyama. “Restriction of the irreducible representations of GL_n to the symmetric group S_n ”. In: *preprint* (2000).
- [Pip03] Nicholas Pippenger. “[The inequalities of quantum information theory](#)”. In: *IEEE Transactions on Information Theory* 49.4 (2003), pp. 773–789.
- [Ric02] G Richardson. “The trigonometry of the tetrahedron”. In: *The Mathematical Gazette* 2.32 (1902), pp. 149–158.
- [Rob02] Justin Roberts. “Asymptotics and 6j-symbols”. In: *Geom. Topol. Monogr* 4 (2002), pp. 245–261.
- [Rob99] Justin Roberts. “[Classical 6j-symbols and the tetrahedron](#)”. In: *Geometry & Topology* 3.1 (1999), pp. 21–66.
- [RR86] David P Robbins and Howard Rumsey Jr. “Determinants and alternating sign matrices”. In: *Advances in Mathematics* 62.2 (1986), pp. 169–184.
- [San61] Ivan Nicolaevich Sanov. “On the probability of large deviations of random variables”. In: *Selected Translations in Mathematical Statistics and Probability* 1 (1961), pp. 213–244.

- [SBT17] David Sutter, Mario Berta, and Marco Tomamichel. “[Multivariate trace inequalities](#)”. In: *Communications in Mathematical Physics* 352.1 (2017), pp. 37–58. arXiv: [1604.03023v3](#).
- [Spe05] David E Speyer. “Horn’s problem, Vinnikov curves, and the hive cone”. In: (2005).
- [TT73] RC Thompson and S Therianos. “The Eigenvalues and Singular Values of Matrix Sums and Products. VII (1)”. In: *Canadian Mathematical Bulletin* 16.4 (1973), pp. 561–569.
- [TW06] Yuka U Taylor and Christopher T Woodward. “6 j symbols for and non-Euclidean tetrahedra”. In: *Selecta Mathematica* 11.3-4 (2006), p. 539.
- [Zub18] Jean-Bernard Zuber. “Horn’s problem and Harish-Chandra’s integrals. Probability density functions”. In: *Annales de l’Institut Henri Poincaré D* 5.3 (2018), pp. 309–338.