

# Eigenvalues of tetrahedral sums of Hermitian matrices

Thomas C. Fraser<sup>1</sup>, Matthias Christandl<sup>1</sup>, and Anton Alekseev<sup>2</sup>

<sup>1</sup>Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, 2100  
Denmark

<sup>2</sup>Anton's affiliations

June 20, 2024

## Abstract

This is the abstract.

## Contents

|          |                                                                                                      |           |
|----------|------------------------------------------------------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                                                                  | <b>2</b>  |
| 1.1      | Triangles and tetrahedron . . . . .                                                                  | 2         |
| 1.2      | Notational conventions . . . . .                                                                     | 2         |
| <b>2</b> | <b>Generalizations of Horn's problem</b>                                                             | <b>3</b>  |
| 2.1      | Graphical Horn problems . . . . .                                                                    | 3         |
| <b>3</b> | <b>A representation-theoretic characterization</b>                                                   | <b>8</b>  |
| 3.1      | Preliminaries . . . . .                                                                              | 8         |
| 3.2      | Reformulating the cycle condition . . . . .                                                          | 9         |
| 3.3      | Trigonal spectra & Littlewood-Richardson coefficients . . . . .                                      | 10        |
| 3.4      | Tetrahedral spectra & recoupling operators . . . . .                                                 | 11        |
| 3.5      | The $\mathbf{U}(\mathbf{d})$ -recoupling operator as a $\mathbf{S}_d$ -recoupling operator . . . . . | 11        |
| 3.6      | An entropic inequality for tetrahedral spectra . . . . .                                             | 13        |
| <b>4</b> | <b>The spectral quantum marginal problem</b>                                                         | <b>14</b> |
| 4.1      | A spectral embedding . . . . .                                                                       | 14        |
| <b>5</b> | <b>Discussion</b>                                                                                    | <b>17</b> |
| <b>A</b> | <b>Proof of symmetric group recoupling operator upper bound</b>                                      | <b>21</b> |
| <b>B</b> | <b>Proof of recoupling operator norm equivalences</b>                                                | <b>23</b> |

# Outline Draft May 30 2024

Below is the proposed outline of the paper as of May 30, 2024. Our proposed journal is *Communications in Mathematical Physics*. It will be ideal to illustrate with figures.

- introduction (include 2x2 case)
- general definition
- relationship to Horn's problem, (how does it generalize the problem)
- representation theoretic/quantization analogue (Ponzano-Regge and connection to state-sum models? etc.)
- comment about deviation from Thompson's conjecture (correspondence between eigenvalues of sums and logs of singular values of products)
- spectral embedding into the quantum marginal problem
- derivation of entropic inequalities (both from rep theory directly, and from embedding into QMP, things like concavity and data processing inequality for Renyi-entropies, etc), then prove they go beyond face constraints

## 1 Introduction

### 1.1 Triangles and tetrahedron

TODO: Describe the moduli space of Euclidean triangles (up to rigid motions) and Euclidean tetrahedron (again up to rigid motions) and describe how applying the triangle inequalities to each face of the tetrahedron, while necessary, are ultimately insufficient for the existence of a tetrahedron with specified side lengths.

### 1.2 Notational conventions

Throughout this paper,  $d \in \mathbb{N}_{>0}$  will denote the size of the matrices under consideration. We also exclusively work with matrices with coefficients in the field of complex numbers  $\mathbb{C}$ , so for notational brevity we write  $\text{GL}(d)$  as shorthand for  $\text{GL}(d, \mathbb{C})$ . Weights for  $\text{GL}(d)$  will be denoted using Greek letters and will be identified with  $d$ -tuples of integers, e.g.  $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{Z}^d$ . The weight lattice for  $\text{GL}(d)$  is denoted by  $\Lambda^{\mathbb{Z}}(d) \simeq \mathbb{Z}^d$  in the usual way; a weight  $\alpha \in \Lambda^{\mathbb{Z}}(d)$  is dominant if  $\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_d$  and the set of all dominant weights for  $\text{GL}(d)$  is denoted by  $\Lambda_{\geq}^{\mathbb{Z}}(d)$ .

Irreducible highest weight representations of  $\text{GL}(d)$  with weight  $\lambda \in \Lambda^{\mathbb{Z}}(d)$  are denoted by  $L_{\lambda}$  and irreducible representations of  $S_n$  of highest weight  $\lambda \in \Upsilon_n$  are denoted by  $V_{\lambda}$ .

The set of all  $d \times d$  complex Hermitian matrices will be denoted by  $\mathcal{H}(d)$  while the set of all possible eigenvalues of a  $d \times d$  complex Hermitian matrix is denoted by  $\Lambda_{\geq}^{\mathbb{R}}(d) \subset \mathbb{R}^d$ . The set of  $d \times d$  Hermitian matrices  $X$  with eigenvalues  $\text{eig}(X) = (x_1, \dots, x_d) \in \Lambda_{\geq}^{\mathbb{R}}(d)$  will be denoted by  $\mathcal{O}_x$ .




Figure 1: Examples of graphical Horn problems.

## 2 Generalizations of Horn's problem

This section will include a formal statement of the problem under investigation, its relation to Horn's original problem and other generalizations of Horn's problem. This section will hopefully briefly describe what is known about the eigenvalues of triangular sums of matrices. Regarding tetrahedral spectra, we will describe many of the symmetries of the problem statement and various “gauge” choices that one could make. Additional generalizations of the problem statement, like to more matrices, linear combinations, etc. will be mentioned here. It will be important to mention the manner in which the analogue of the “Thompson-Klyachko” theorem does not hold, i.e. singular values of tetrahedral products are not the exponentials of eigenvalues of tetrahedral sums in general.

### 2.1 Graphical Horn problems

**Definition 1.** Let  $\mathcal{G} = (\mathcal{V}, \mathcal{E})$  be a finite directed graph with vertex set  $\mathcal{V}$  and edge set  $\mathcal{E}$  together with source and target maps  $s, t : \mathcal{E} \rightarrow \mathcal{V}$ . Let  $d$  be a positive integer. A *spectral map* (for  $\mathcal{G}$  of dimension  $d$ ) is a function  $\lambda : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  assigning to each edge  $e \in \mathcal{E}$  in the directed graph a  $d$ -tuple of real numbers  $\lambda(e) = (\lambda_1(e), \dots, \lambda_d(e))$ . The map  $\lambda : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  is said to be *realizable (by Hermitian matrices)* if there exists a map  $H : \mathcal{E} \rightarrow \mathcal{H}(d)$  assigning to each edge  $e \in \mathcal{E}$  a  $d \times d$  Hermitian matrix  $H(e)$  such that

- for each edge  $e \in \mathcal{E}$ ,  $H(e)$  has eigenvalues  $\lambda(e)$ ,

$$\text{eig}(H(e)) = \lambda(e), \tag{1} \quad \text{\texttt{\{eq:spec_cond\}}}$$

- for each cycle  $\ell \subseteq \mathcal{E}$  with orientation map  $\gamma_{\ell} : \ell \rightarrow \{\pm 1\}$ ,

$$\sum_{e \in \ell} \gamma_{\ell}(e) H(e) = 0. \tag{2} \quad \text{\texttt{\{eq:cycle_cond\}}}$$

Alternatively, one can equivalently encode the constraints imposed by Eq. (2) in terms of a  $\{-1, 0, +1\}$ -valued  $L \times |E|$  matrix (where  $L$  is the number of (generating) cycles in  $\mathcal{G}$ ), denoted by  $\Gamma(\mathcal{G})$ , where

$$\Gamma(\mathcal{G})_{\ell,e} = \begin{cases} 0 & e \notin \ell, \\ \gamma_\ell(e) & e \in \ell. \end{cases} \quad (3)$$

In terms of  $\Gamma(\mathcal{G})$ , Eq. (2) reads for all cycles  $\ell$ :

$$\sum_{e \in \mathcal{E}} \Gamma(\mathcal{G})_{\ell,e} H(e) = 0 \quad (4)$$

prob:ghp

**Problem 1** (Graphical Horn problem). Let  $\mathcal{G} = (\mathcal{V}, \mathcal{E})$  be a finite directed graph, let  $d$  be a positive integer. The *graphical Horn problem* is to characterize the set of realizable spectral maps  $\lambda : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  for  $\mathcal{G}$  of dimension  $d$ .

There might be an important connection to the theory of quivers and representations of quivers. In particular, the notion of a morphism between a representation of a quiver in terms of operators on vertices forming commuting squares models precisely the kinds of continuous symmetries which may be present in a given graphical Horn problem. Indeed, the spectral data we are ultimately interested in is equivalent to the orbit under such “local” isomorphisms (i.e. conjugation by isomorphisms).

TC: It turns out that the only relevant data for statement of a graphical Horn problem is the matrix  $\Gamma(\mathcal{G})$ ; perhaps it's best to drop reference to  $\mathcal{G}$  altogether and just have the problem statement depend on a matrix  $\Gamma$ ? Moreover, one could in principle take the matrix to be  $\mathbb{R}$ -valued instead of  $\{-1, 0, +1\}$ -valued, although that would make the connection to the representation theory even weaker. Generalizing to  $\mathbb{Z}$ -valued  $\Gamma$  is also essentially the same I think... One downside of dropping the graph  $\mathcal{G}$  is that the matrix  $\Gamma$  loses the non-abelian information about the ordering of the elements in a cycle; this is problematic when discussing the “quantum” generalizations to multiplicative Horn-type problems. One upside of dropping the graph  $\mathcal{G}$ , is that it becomes possible to ignore certain cycles of the graph (by dropping rows of  $\Gamma$ ) and/or additionally including constraints non-easily associated to cycles of a common graph (like  $A + C = Q, B + Q = L$  constraints for the tetrahedral problem).

There are a few operations which preserve the set of realizable spectral maps.

**Lemma 1.** *Consider a graphical Horn problem of dimension  $d$ .*

- *The zero spectral map  $0_d : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  assigning to  $e \in \mathcal{E}$ , the spectrum  $0_d(e) = (0, \dots, 0)$  is realizable.*
- *If  $\alpha : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(a)$  and  $\beta : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(b)$  are realizable, then their direct sum  $\alpha \oplus \beta : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(a+b)$  and tensor product  $\alpha \otimes \beta : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(ab)$  (defined pointwise in the obvious manner), are both realizable.*
- *If  $\lambda : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  is a realizable and  $p \in \mathbb{R}_{>0}$  a positive real number, then  $p\lambda : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$ , defined by  $(p\lambda)(e) = (p\lambda_1(e), \dots, p\lambda_d(e))$  is also a realizable.*
- *If  $\lambda : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  is a realizable, then  $\lambda^* : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$ , defined by a pointwise application of the negation-reversal involution, is also realizable.*

**Example 1** (The trigonal Horn problem). Perhaps the smallest non-trivial example of a graphical Horn problem is the one for which Horn originally conjectured a solution [Hor62], namely, what are the possible eigenvalues,  $(\alpha, \beta, \gamma)$ , of a triple of  $d \times d$  Hermitian matrices  $(A, B, C)$  satisfying the condition  $A + B = C$ ? As a graphical Horn problem,

$$\Gamma = \begin{pmatrix} 1 & 1 & -1 \end{pmatrix} \quad (5)$$

From the perspective of Problem 1, we refer to this question as the *trigonal Horn problem*.

**Example 2** (Polygonal Horn problems). If the directed graph  $\mathcal{G}$  consists of a single closed cycle, then the corresponding Horn problem will be referred to as a *polygonal Horn problem*; the polygonal Horn problem associated to an *oriented* cycle  $(e_1, e_2, \dots, e_r, e_1)$  consisting of  $r$  edges asks what are the possible eigenvalues  $(\lambda_1, \dots, \lambda_r)$  of  $r$  Hermitian matrices  $(H_1, \dots, H_r)$  with zero-sum  $H_1 + \dots + H_r = 0$ .

$$\Gamma = \begin{pmatrix} 1 & \dots & 1 \end{pmatrix} \quad (6)$$

**Remark 1.** Evidently, there is a bijection between the set of realizable spectra  $(\alpha, \beta, \gamma)$  and the set of realizable spectra  $(\lambda_1, \lambda_2, \lambda_3)$  for the polygonal Horn problem where  $r = 3$ : namely

$$(\lambda_1, \lambda_2, \lambda_3) = (\alpha, \beta, \gamma^*) \quad (7)$$

where  $\gamma \mapsto \gamma^*$  is the involution on  $\Lambda_{\geq}^{\mathbb{R}}(d)$  defined by negation of entries and then reversing to preserve the ordering:

$$\gamma^* = (-\gamma_d, \dots, -\lambda_1). \quad (8)$$

**Example 3** (Tetrahedral Horn problem). In this paper, a considerable focus will be placed on the *tetrahedral Horn problem* which is the graphical Horn problem associated with the directed graph above forming a tetrahedron. Specifically, a sextuple of candidate spectra,

$$(\alpha, \beta, \gamma, \mu, \nu, \lambda) \in (\Lambda_{\geq}^{\mathbb{R}}(d))^{\times 6} \quad (9)$$

are said to be *tetrahedral* if they are the respective eigenvalues of Hermitian matrices

$$(A, B, C, M, N, L) \in \mathcal{H}(d)^{\times 6} \quad (10)$$

satisfying the following four conditions:

$$A + B = M, \quad M + C = L, \quad B + C = N, \quad A + N = L. \quad (11)$$

As a graphical Horn problem,

$$\Gamma = \begin{pmatrix} 1 & 1 & 0 & -1 & 0 & 0 \\ 0 & 1 & 1 & 0 & -1 & 0 \\ 0 & 0 & 1 & 1 & 0 & -1 \\ 1 & 0 & 0 & 0 & 1 & -1 \end{pmatrix} \quad (12)$$

**Example 4** (One-dimensional graphical Horn problems). An important special case of graphical Horn problem for a digraph  $\mathcal{G}$  is the case where  $d = 1$ . If  $d = 1$ , then vector space of  $1 \times 1$  Hermitian matrices becomes isomorphic to the real numbers,  $\mathcal{H}(1) \simeq \mathbb{R}$ , and similarly the space of candidate eigenvalues becomes  $\Lambda_{\geq}^{\mathbb{R}}(1) = \mathbb{R}$ . Moreover, the spectral map  $\lambda : \mathcal{E} \rightarrow \mathbb{R}$  becomes an assignment of real numbers of the edges of  $\mathcal{G}$ , the map  $H : \mathcal{E} \rightarrow \mathbb{R}$  must necessarily equal spectral map  $\lambda$  by the spectral condition Eq. (1), and the cycle condition characterizing realizable spectral maps simplifies to the condition that  $\lambda$ , viewed as a vector  $(\lambda(e))_{e \in \mathcal{E}} \in \mathbb{R}^{|\mathcal{E}|}$ , must lie in the kernel of  $\Gamma(\mathcal{G})$ ,

$$\lambda \in \ker(\Gamma(\mathcal{G})). \quad \text{\textcolor{blue}{\small{eq:ker_cond_d_1}}} \quad (13)$$

In fact, this simple condition provides a complete solution to all graphical Horn problems where  $d = 1$ .

The solution to all one-dimensional graphical Horn problems provided by Eq. (13) also gives rise to a simple necessary condition for  $d > 1$  graphical Horn problems. \textcolor{blue}{\small{lem:trace\_cond}}

**Lemma 2.** *Consider the graphical Horn problem of dimension  $d$  for the digraph  $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ . Then the spectral map  $\lambda : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  is realizable only if*

$$\text{Tr} \circ \lambda \in \ker(\Gamma(\mathcal{G})), \quad (14)$$

where, by abuse of notation,  $\text{Tr} : \Lambda_{\geq}^{\mathbb{R}}(d) \rightarrow \mathbb{R}$  denotes the sum

$$\text{Tr}((\lambda_1, \dots, \lambda_d)) := \lambda_1 + \dots + \lambda_d. \quad (15)$$

*Proof.* Suppose that  $\lambda : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  is a spectral map realized by the assignment  $H : \mathcal{E} \rightarrow \mathcal{H}(d)$ , meaning  $\text{eig}(H(e)) = \lambda(e)$  for all edges  $e \in \mathcal{E}$ , and  $\sum_{e \in E} \Gamma(\mathcal{G})_{\ell, e} H(e) = 0$  for all cycles  $\ell$ . Taking the trace of cycle condition Eq. (2) yields

$$0 = \sum_{e \in E} \Gamma(\mathcal{G})_{\ell, e} \text{Tr}(H(e)) = \sum_{e \in E} \Gamma(\mathcal{G})_{\ell, e} \text{Tr}(\lambda(e)). \quad (16)$$

In other words,  $\text{Tr} \circ \lambda$  lies in the kernel of  $\Gamma(\mathcal{G})$  which implies  $\text{Tr} \circ \lambda$  is a realizable spectral map for the  $d = 1$  case. \textcolor{blue}{\small{lem:ker\_shift}}  $\square$

**Lemma 3.** *Consider the graphical Horn problem of dimension  $d$  for the digraph  $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ . Suppose  $\mu : \mathcal{E} \rightarrow \mathbb{R}$  lies in the kernel of  $\Gamma(\mathcal{G})$  and define  $\mu \cdot 1_d : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  to be the spectral map defined by*

$$(\mu \cdot 1_d)(e) := (\mu(e), \dots, \mu(e)). \quad (17)$$

*Then  $\mu \cdot 1_d : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  is a realizable spectral map. Moreover,  $\lambda : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  is a realizable spectral map if and only if  $\lambda + \mu \cdot 1_d$  is a realizable.*

*Proof.* The spectral map  $\mu \cdot 1_d$  is realized by the assignment  $H_{\mu} : \mathcal{E} \rightarrow \mathcal{H}(d)$  defined by

$$H_{\mu}(e) := \mu(e)I_d \quad (18)$$

where  $I_d$  is the  $d \times d$  identity matrix. This is because it was assumed that  $\mu \in \ker(\Gamma(\mathcal{G}))$  and therefore

$$\sum_{e \in E} \Gamma(\mathcal{G})_{\ell, e} H_\mu(e) = \left( \sum_{e \in E} \Gamma(\mathcal{G})_{\ell, e} \mu(e) \right) I_d = 0. \quad (19)$$

Now suppose  $\lambda : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  is realized by the assignment  $H_\lambda : \mathcal{E} \rightarrow \mathcal{H}(d)$ , then it follows that  $H_\lambda + H_\mu$ , defined by

$$(H_\lambda + H_\mu)(e) = H_\lambda(e) + \mu(e)I_d \quad (20)$$

realizes the spectral map  $\lambda + \mu \cdot 1_d$  because the spectrum of  $H_\lambda(e) + \mu(e)I_d$  is simply  $\lambda(e) + \mu \cdot 1_d(e)$  (the eigenvalues add because  $[H_\lambda(e), I_d] = 0$ ).  $\square$

A direct corollary of Lemma 3 is that the set of realizable spectral maps,  $\lambda : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$ , is entirely determined by the set of realizable and *traceless* spectral maps in the sense that  $\text{Tr} \circ \lambda = 0_d$ .

**Corollary 4.** *Let  $\lambda : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  be a spectral map such that  $\text{Tr} \circ \lambda \in \ker(\Gamma(\mathcal{G}))$  and let  $\lambda^0 : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  be the traceless spectral map defined by*

$$\lambda^0(e) := (\lambda_1(e) - \text{Tr}(\lambda(e)), \dots, \lambda_d(e) - \text{Tr}(\lambda(e))). \quad (21)$$

*Then  $\lambda$  is realizable if and only if  $\lambda^0$  is realizable.*

*Proof.* The spectral map  $\lambda^0$  is obtained from  $\lambda$  by subtracting the spectral map  $\text{Tr} \circ \lambda$ . By assumption,  $\text{Tr} \circ \lambda$  lies in the kernel of  $\Gamma(\mathcal{G})$  by Lemma 2, and thus Lemma 3 (when  $\mu \mapsto \pm \text{Tr} \circ \lambda$ ) implies  $\lambda^0$  is realizable if and only if  $\lambda$  is realizable.  $\square$

**Corollary 5.** *Suppose  $\mathcal{G}$  is a graphical Horn problem such that  $\ker(\Gamma(\mathcal{G}))$  contains a one-dimensional spectral map  $\rho : \mathcal{E} \rightarrow \mathbb{R}$  which is strictly positive, meaning*

$$\forall e \in \mathcal{E} : \rho(e) > 0. \quad (22)$$

*Let  $\lambda : \mathcal{E} \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  be a spectral map. Then there exists a sufficiently large positive  $p \in \mathbb{R}_{>0}$  such that  $\lambda^+ := \lambda + p\rho \cdot 1_d$  is a spectral map consisting of strictly positive spectra meaning*

$$\forall e \in \mathcal{E} : \lambda_d^+(e) > 0. \quad (23)$$

*Proof.* The value of  $\lambda_d^+(e)$  for  $e \in \mathcal{E}$  is determined by

$$\lambda_d^+(e) = \lambda_d(e) + p\rho(e). \quad (24)$$

In order to satisfy Eq. (23) it suffices to let  $p$  satisfy

$$p > \max_{e \in \mathcal{E}} \frac{-\lambda_d(e)}{\rho(e)}. \quad (25)$$

$\square$

**Example 5.** The tetrahedral Horn problem defined above admits of strictly positive one-dimensional spectral maps. Specifically,  $\rho = (a, b, c, a + b, b + c, a + b + c)$  lies in the kernel of  $\Gamma$  and is strictly positive whenever  $a, b, c > 0$ .

### 3 A representation-theoretic characterization

The purpose of this section is to widen the bridge between representation theory and Horn-type problems. We will begin with a treatment of the connection between triangular spectra and the Littlewood-Richardson coefficients and discuss its connection to symplectic geometry and moment maps. Then we will articulate the manner in which tetrahedral spectra are related to the norm of the recoupling operator for  $SU(d)$  and discuss the manner in which convexity results for moment maps cannot be applied and thus one must not expect the space of tetrahedral spectra to be convex (indeed it is not).

There is a well-established and fundamental connection between the eigenvalues of sums of Hermitian matrices and the representation theory of the general linear group [Kly98; KT99; Knu00; Ful00; Bha01; Chr08; BVW18]. The purpose of this section is to briefly summarize the nature of this connection (Section 3.3) and then describe the manner in which this fundamental connection generalizes to graphical Horn problems.

#### 3.1 Preliminaries

sec:prelim

There are a number of approaches that can be taken to begin building the bridge which connects eigenvalues of sums of  $d \times d$  Hermitian matrices to the representation theory of  $GL(d)$ ; the method taken here is through notions of stability in the sense of geometric invariant theory. We begin by recalling prerequisite notions from representation theory, symplectic geometry and geometric invariant theory, establishing our notation along the way.

As always, we identify the set of  $d \times d$  Hermitian matrices,  $\mathcal{H}(d)$  with the set  $\mathfrak{u}(d)^*$  of linear functionals on the Lie algebra  $\mathfrak{u}(d)$  of  $U(d)$  using the non-degenerate bilinear pairing  $(H, X) = i \operatorname{Tr}(XH)$  for  $H \in \mathcal{H}(d)$  Hermitian and  $X \in \mathfrak{u}(d)$  skew-Hermitian. Given  $\lambda \in \Lambda_{\geq}^{\mathbb{R}}(d)$ , let  $D_{\lambda} = \operatorname{diag}(\lambda_1, \dots, \lambda_d)$  be the diagonal  $d \times d$  Hermitian matrix. The set  $\mathcal{O}_{\lambda}$  of  $d \times d$  Hermitian matrices with eigenvalues  $\lambda$  can be identified with the  $U(d)$ -orbit of  $D_{\lambda}$  under the adjoint action,  $U \cdot D_{\lambda} = UD_{\lambda}U^{-1}$ , i.e.

$$\mathcal{O}_{\lambda} = U(d) \cdot D_{\lambda} = \{U \cdot D_{\lambda} \mid U \in U(d)\}, \quad (26)$$

or equivalently identified with a coadjoint orbit in  $\mathfrak{u}(d)^*$ . When  $\lambda \in \Lambda_{\geq}^{\mathbb{Z}}(d)$  is integer-valued, it may additionally be considered as a dominant weight for  $GL(d)$ . In such cases, let  $\pi_{\lambda} : GL(d) \rightarrow \operatorname{Aut}(L_{\lambda})$ <sup>1</sup> denote the unique (up to isomorphism) irreducible representation of  $GL(d)$  with highest weight  $\lambda$  and let  $v_{\lambda} \in L_{\lambda}$  be a highest weight vector and let  $[v_{\lambda}] \in \mathbb{P}(L_{\lambda})$  be the associated highest weight ray. One important consequence of  $v_{\lambda}$  being a highest

<sup>1</sup>The representation space  $L_{\lambda}$  can be realized concretely in the sense of Borel-Weil as follows. Let  $B(d)$  be the Borel subgroup of upper triangular matrices in  $GL(d)$  and  $F(d) = GL(d)/B(d)$  the corresponding flag variety/manifold with  $GL(d) \rightarrow F(d)$  the principle  $B(d)$ -bundle over  $F(d)$ . Given a weight  $\mu \in \Lambda^{\mathbb{Z}}(d)$ , let  $\chi_{\mu}$  be the one-dimensional complex representation of  $B(d)$  sending the invertible diagonal  $D_t = \operatorname{diag}(t_1, \dots, t_d)$  to  $\chi_{\mu}(D_t) = t_1^{\mu_1} \cdots t_d^{\mu_d} \in \mathbb{C}$ . The representation  $\chi_{\mu}$  of  $B(d)$  gives rise to the associated line bundle,  $GL(d) \times_{B(d)} \mathbb{C} \rightarrow F(d)$ . With respect to this construction, the representation space  $L_{\lambda}$  can be identified with the vector space of holomorphic sections of the associated line bundle given by  $\chi_{-\lambda^*}$ , i.e. holomorphic functions  $s : GL(d) \rightarrow \mathbb{C}$  satisfying  $s(gb) = s(g)\chi_{\lambda^*}(b)$  for all  $g \in GL(d)$  and  $b \in B(d)$ . **TODO: Check to signs in this footnote to make sure they are correct.**

weight vector is simply that the  $B(d)$ -orbit of  $v_\lambda$ ,  $B(d) \cdot v_\lambda = \{\pi_\lambda(b)v_\lambda\}$ , is contained within the ray  $[v_\lambda]$ . In addition, as each  $g \in \mathrm{GL}(d)$  admits of a decomposition,  $g = ub$ , for  $(u, b) \in \mathrm{U}(d) \times B(d)$ , the  $\mathrm{U}(d)$ -orbit of  $[v_\lambda]$  is equivalent to the  $\mathrm{GL}(d)$ -orbit of  $[v_\lambda]$ . Moreover, there is a well-defined  $\mathrm{U}(d)$ -equivariant diffeomorphism<sup>2</sup> from  $\mathcal{O}_\lambda$  to  $\mathrm{U}(d) \cdot [v_\lambda]$  sending  $u \cdot D_\lambda \in \mathcal{O}_\lambda$  to  $u \cdot [v_\lambda] = [v_\lambda^u] \in \mathrm{U}(d) \cdot [v_\lambda] \subseteq \mathbb{P}(L_\lambda)$  where

$$v_\lambda^u := \pi_\lambda(u)v_\lambda. \quad (27)$$

Furthermore, if  $X \in \mathfrak{gl}(d)$  is a  $d \times d$  matrix, then it can be shown that  $\mathrm{Tr}(XD_\lambda) = \langle v_\lambda, \Pi_\lambda(X)v_\lambda \rangle$  where  $\Pi_\lambda$  is the induced representation of  $\mathfrak{gl}(d)$  on  $L_\lambda$ ,  $\langle \cdot, \cdot \rangle$  is a  $\mathrm{U}(d)$ -invariant inner product on  $L_\lambda$  and  $v_\lambda$  is a highest weight vector of unit norm in  $L_\lambda$ . Using the aforementioned diffeomorphism, one can also show that for all  $d \times d$  matrices  $X \in \mathfrak{gl}(d)$ ,

$$\mathrm{Tr}(X(u \cdot D_\lambda)) = \langle v_\lambda^u, \Pi_\lambda(X)v_\lambda^u \rangle. \quad (28)$$

Throughout this section, the following result serves as the basis for establishing a number of asymptotic results for large  $k$ .

**Lemma 6.** *Let  $\lambda \in \Lambda_{\geq d}^{\mathbb{Z}}$  be a dominant weight for  $\mathrm{GL}(d)$ , let  $L_\lambda$  be the irreducible representation of  $\mathrm{GL}(d)$  with highest weight  $\lambda$  and let  $k \in \mathbb{N}_{>0}$  be a positive integer. Then  $\dim(\mathrm{Hom}_{\mathrm{GL}(d)}(L_{k\lambda}, L_\lambda^{\otimes k})) = 1$ . Let*

$$\iota_\lambda^k : L_{k\lambda} \rightarrow L_\lambda^{\otimes k} \quad (29)$$

*be a  $\mathrm{GL}(d)$ -intertwining isometry in  $\mathrm{Hom}_{\mathrm{GL}(d)}(L_{k\lambda}, L_\lambda^{\otimes k})$ . If  $v_\lambda \in L_\lambda$  is a highest weight vector, then there exists a highest weight vector  $v_{k\lambda} \in L_{k\lambda}$  satisfying*

$$\iota_\lambda^k v_{k\lambda} = v_\lambda^{\otimes k}. \quad (30)$$

*Moreover, for all  $g \in \mathrm{GL}(d)$ ,*

$$\|\pi_{k\lambda}(g)v_{k\lambda}\| = \|\pi_\lambda(g)v_\lambda\|^k. \quad (31)$$

## 3.2 Reformulating the cycle condition

[sec:reform\\_cycle](#)

It will be shown how the cycle condition, Eq. (2), is equivalent to the existence of certain vectors which are minimal in norm with respect to a representation of  $\mathrm{GL}(d)$ . Now consider a polygonal graph consisting of a single connected cycle of length  $s$  such that  $\Gamma$  is a  $\{\pm 1\}$ -valued  $1 \times s$  matrix. Let  $\lambda : [s] \rightarrow \Lambda_{\geq}^{\mathbb{R}}(d)$  be an eigenvalue map for such a  $\Gamma$ , we consider the symplectic manifold

$$\mathcal{M}_\ell := \prod_{e \in \mathcal{E}} \mathcal{O}_{\lambda(e)\gamma_\ell(e)} \quad (32)$$

It can be shown that the moment map  $\Phi_\ell : \mathcal{M}_\ell \rightarrow \mathcal{H}(d)$  associated to this action satisfies

$$\Phi_\ell(\gamma_\ell(e)H(e) \mid e \in \ell) = \sum_{e \in \ell} \gamma_\ell(e)H(e) \quad (33)$$

and thus vanishes precisely when the cycle condition, Eq. (2), is satisfied.

**Theorem 7.**

---

<sup>2</sup>This map is well-defined because the  $\mathrm{U}(d)$ -stabilizers of  $D_\lambda$  and  $v_\lambda$  necessarily agree.

### 3.3 Trigonal spectra & Littlewood-Richardson coefficients

sec:trigonal\_lr\_coeff

**Definition 2.** Let  $\alpha, \beta, \gamma \in \Lambda_{\geq}^{\mathbb{Z}}(d)$  be a triple of dominant weights for  $\mathrm{GL}(d)$  and let  $L_{\alpha}, L_{\beta}, L_{\gamma}$  be the triple of irreducible representations of  $\mathrm{GL}(d)$  with corresponding highest weights. Let  $C_{\alpha, \beta}^{\gamma}$  denote the multiplicity space of  $\mathrm{GL}(d)$ -intertwiners from  $L_{\gamma}$  to  $L_{\alpha} \otimes L_{\beta}$ :

$$C_{\alpha, \beta}^{\gamma} := \mathrm{Hom}_{\mathrm{GL}(d)}(L_{\gamma}, L_{\alpha} \otimes L_{\beta}) = \mathrm{Inv}_{\mathrm{GL}(d)}(L_{\gamma}^* \otimes L_{\alpha} \otimes L_{\beta}). \quad (34)$$

The *Littlewood-Richardson coefficient* is defined as the dimension of this multiplicity space:

$$c_{\alpha, \beta}^{\gamma} := \dim(C_{\alpha, \beta}^{\gamma}). \quad (35)$$

Our first result can be viewed as a simple condition which is sufficient to conclude that a given triple of integer eigenvalues are trigonal.

**Lemma 8.** *If  $c_{\alpha, \beta}^{\gamma} > 0$ , then there exists Hermitian matrices  $A \in \mathcal{O}_{\alpha}$ ,  $B \in \mathcal{O}_{\beta}$ , and  $C \in \mathcal{O}_{\gamma}$ , satisfying  $A + B - C = 0$ .*

*Proof.* Our proof proceeds in the style of [BVW18, Proposition 2.3]. Given any dominant weight  $\lambda \in \Lambda_{\geq}^{\mathbb{Z}}(d)$ , let  $P_{\lambda}$  denote the Hermitian projection operator onto the invariant  $\square$

Using the theory of spectrum estimation developed in the context of quantum information theory [ARS88; KW01; HM02], the following pair of theorems were proven in [Chr08]. Recall that  $(\alpha, \beta, \gamma) \in \Lambda_{\geq}^{\mathbb{R}}(d)^{\times 3}$  are said to be *trigonal* if there exists Hermitian matrices  $A, B, C \in \mathcal{H}(d)$  satisfying (i)  $A + B + C = 0$  and (ii)

$$(\mathrm{eig}(A), \mathrm{eig}(B), \mathrm{eig}(C)) = (\alpha, \beta, \gamma). \quad (36)$$

**Theorem 9.** *If  $c_{\alpha, \beta, \gamma} > 0$  for some  $\alpha, \beta, \gamma \in \Lambda_{\geq}^{\mathbb{Z}}(d)$ , then  $(\alpha, \beta, \gamma)$  are trigonal.*

**Theorem 10.** *If  $(\alpha, \beta, \gamma) \in \Lambda_{\geq}^{\mathbb{R}}(d)^{\times 3}$  are triangular, then there exists a graded sequence  $(\tilde{\alpha}(n), \tilde{\beta}(n), \tilde{\gamma}(n)) \in \Lambda_{\geq}^{\mathbb{Z}}(d)^{\times 3}$  such that (i) for all  $n \in \mathbb{N}_{>0}$ ,*

$$c_{\tilde{\alpha}(n), \tilde{\beta}(n), \tilde{\gamma}(n)} > 0, \quad (37)$$

*and (ii)*

$$\lim_{n \rightarrow \infty} \frac{\tilde{\alpha}(n)}{n} = \alpha, \quad \lim_{n \rightarrow \infty} \frac{\tilde{\beta}(n)}{n} = \beta, \quad \lim_{n \rightarrow \infty} \frac{\tilde{\gamma}(n)}{n} = \gamma. \quad (38)$$

Consider the following well-known result, for instance see [BVW18, Proposition 2.3] or [Chr08].

prop:LR\_pos\_implies\_herm

**Proposition 11.** *Let  $\lambda(1), \dots, \lambda(\ell) \in \Lambda_{\geq}^{\mathbb{Z}}(d)$  be dominant weights for  $\mathrm{GL}(d)$  such that*

$$(L_{\lambda(1)} \otimes \dots \otimes L_{\lambda(\ell)})^{\mathrm{GL}(d)} \neq 0. \quad (39)$$

*Then there exists Hermitian matrices,  $H(e) \in \mathcal{O}_{\lambda(e)}$  for all  $e \in \{1, \dots, \ell\}$ , such that  $\sum_{e=1}^{\ell} H(e) = 0$ .*

*Proof.* Summarize the direct and pedagogical Kempf-Ness style proof in [BVW18].  $\square$

### 3.4 Tetrahedral spectra & recoupling operators

sec:recouple

TODO: Show how to generalize the construction due to Roberts [Rob99] by taking the Cartesian product of these manifolds over all cycles and then imposing additional group actions and moment-map conditions to stitch everything together.

**Remark 2.** As an immediate corollary of Proposition 11, we conclude that if

$$(L_\alpha \otimes L_\gamma^* \otimes L_\beta)^{\text{GL}(d)} \otimes (L_\beta^* \otimes L_\phi \otimes L_\delta^*)^{\text{GL}(d)} \otimes (L_\delta \otimes L_\gamma \otimes L_{\epsilon^*})^{\text{GL}(d)} \otimes (L_\epsilon \otimes L_\phi^* \otimes L_\alpha^*)^{\text{GL}(d)} \neq 0, \quad (40)$$

{eq:face\_inv}

then there exists Hermitian operators

$$A, A' \in \mathcal{O}_\alpha, \quad B, B' \in \mathcal{O}_\beta, \quad C, C' \in \mathcal{O}_\gamma, \quad D, D' \in \mathcal{O}_\delta, \quad E, E' \in \mathcal{O}_\epsilon, \quad F, F' \in \mathcal{O}_\phi, \quad (41)$$

{eq:prime\_unprime}

satisfying the following four conditions

$$A - C' + B = 0, \quad -B' + F - D' = 0, \quad D + C - E' = 0, \quad E - F' - A' = 0. \quad (42)$$

From this perspective, it becomes clear that Eq. (40) is a necessary condition for  $\alpha, \beta, \gamma, \delta, \epsilon$ , and  $\phi$  to be tetrahedral spectra. Unfortunately, it can be shown that this condition is indeed insufficient; the insufficiency can be attributed to the fact that Eq. (40) does not require that the primed and unprimed Hermitian matrices in Eq. (41) are the same, i.e., that  $A = A'$ ,  $B = B'$ , etc.

### 3.5 The $U(d)$ -recoupling operator as a $S_d$ -recoupling operator

sec:recoupling-correspondence

The purpose of this section is to describe how to view certain aspects of the recoupling theory for the unitary group  $U(d)$  as a special case of the recoupling theory for the symmetric group  $S_d$ , at least when  $d$  is taken to be sufficiently large. This connection between  $U(d)$  and  $S_d$  is derived from considering the symmetric group  $S_d$  as a *subgroup* of  $U(d)$  with respect to the faithful representation of  $S_d$  as  $d \times d$  permutation matrices (taken with respect to the canonical basis). The critical theorem, due to [Nis00], describes the restriction of the irrep of  $U(d)$  labelled by  $\lambda \vdash d$  to the symmetric subgroup  $S_d \subseteq U(d)$ , namely [Nis00, Theorem 2.3].

**Theorem 12.** *Let  $d \in \mathbb{N}$  and let  $\lambda \vdash d$ . Let  $L_\lambda$  denote the irrep of  $U(d)$  with highest weight  $\lambda$  and  $V_\lambda$  the irrep of  $S_d$ . Given a partition  $\mu$  with length  $\ell(\mu) \leq d$ , let  $\Upsilon_\mu$  denote the Young subgroup  $\Upsilon_\mu = S_{\mu_1} \times \cdots \times S_{\mu_{\ell(\mu)}} \subseteq S_d$  and let  $\tilde{\mu}$  denote the (not necessarily sorted) partition of  $\ell(\mu)$  corresponding to the multiplicities of  $\mu$ . Alternatively,  $\tilde{\mu}$  can be characterised in terms of the normaliser of  $\Upsilon_\mu$  inside  $S_d$  via the formula*

$$N_{S_d}(\Upsilon_\mu) = \Upsilon_{\tilde{\mu}} \ltimes \Upsilon_\mu \quad (43)$$

Then the restriction of the irrep  $L_\lambda$  of  $U(d)$  to the symmetric group  $S_d$  decomposes as follows:

$$\text{Res}_{S_d}^{U(d)} L_\lambda \simeq \bigoplus_{\mu \vdash d} \text{Ind}_{S_{d-\ell(\mu)} \times \Upsilon_{\tilde{\mu}}}^{S_d} (1 \otimes V_\lambda^{\Upsilon_\mu}) \quad (44)$$

{eq:nishiyama-decomp}

where  $1$  denotes the trivial representation of  $S_{d-\ell(\mu)}$  and  $V_\lambda^{\Upsilon_\mu}$  is the  $\Upsilon_\mu$ -invariant subspace of  $V_\lambda$  supporting a representation of  $\Upsilon_{\tilde{\mu}}$ .

While the summands appearing on the right-hand side of Eq. (44) are quite complicated in general, it was shown in [CSW18] that the irreducible representations of  $S_d$ , labelled by  $\nu \vdash d$ , which appear with non-zero multiplicity in the decomposition of  $\text{Res}_{S_d}^{\text{U}(d)} L_\lambda$  are constrained in a particular way for  $d$  sufficiently large.

**Definition 3.** Let  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_{\ell(\lambda)})$  be a Young diagram of size  $|\lambda|$ . Let  $d \geq \lambda_1 + |\lambda|$  and define the (*size  $d$* ) *augmentation* of  $\lambda$  as

$$\lambda^{+d} := (d - |\lambda|, \lambda_1, \lambda_2, \dots, \lambda_{\ell(\lambda)}). \quad (45)$$

Note that  $\lambda^{+d}$  also a Young diagram (because  $d - |\lambda| \geq \lambda_1$ ) of size  $d = |\lambda^{+d}|$  and length  $\ell(\lambda^{+d}) = \ell(\lambda) + 1$ .

The following is [CSW18, Lemma 13]:

*lem:largest\_first\_rows*

**Lemma 13.** *Let  $\lambda$  be a positive integer partition and let  $d \in \mathbb{N}$  be given. Let  $L_\lambda$  be the irrep of  $\text{U}(d)$  with highest weight  $\lambda$  and let  $\mu \vdash d$  be an arbitrary Young diagram of size  $d$ . Define  $M_\mu^\lambda$  as the multiplicity space*

$$M_\mu^\lambda := \text{Hom}_{S_d}(V_\mu, \text{Res}_{S_d}^{\text{U}(d)} L_\lambda). \quad (46)$$

*If  $M_\mu^\lambda$  is non-zero, then the first row of  $\mu$  cannot be smaller than  $d - \lambda$ :*

$$M_\mu^\lambda \neq 0 \implies \mu_1 \geq d - |\lambda|. \quad (47)$$

*If  $d < \lambda_1 + |\lambda|$ , then the above bound is strict*

$$M_\mu^\lambda \neq 0 \implies \mu_1 > d - |\lambda|. \quad (48)$$

*Otherwise  $d \geq \lambda_1 + |\lambda|$  and the above bound is saturated only when  $\mu = \lambda^{+d}$  and moreover  $\dim(M_{\lambda^{+d}}^\lambda) = 1$  in this case.*

At this stage, we note that the Littlewood-Richardson coefficients for  $\text{U}(d)$ ,

$$c_{\alpha,\beta}^\gamma = \dim \text{Hom}(L_\gamma, L_\alpha \otimes L_\beta), \quad (49)$$

are known to be independent of  $d$  (whenever  $d \geq \max\{\ell(\alpha), \ell(\beta), \ell(\gamma)\}$ ), while the Kronecker coefficients

$$g_{\mu,\nu}^\lambda = \dim \text{Hom}(V_\lambda, V_\mu \otimes V_\nu) \quad (50)$$

are implicitly dependent on  $d$  because  $d = |\mu| = |\nu| = |\lambda|$ .

As an immediate corollary of Lemma 13, one obtains a method (originally suggest by Murnaghan [Mur55] and later formalized by Littlewood [Lit58]) for viewing Littlewood-Richardson coefficients as a special types of Kronecker coefficients by appropriately augmenting the Young diagrams involved:

**Corollary 14.** *Let  $\alpha, \beta, \gamma$  be Young diagrams and let  $d$  be sufficiently large such that  $\alpha^{+d}, \beta^{+d}$  and  $\gamma^{+d}$  are all well-defined. Then the multiplicity space of  $S_d$ -intertwiners from  $V_{\gamma^{+d}}$  to  $V_{\alpha^{+d}} \otimes V_{\beta^{+d}}$  is obtained from the multiplicity space of  $\text{U}(d)$ -intertwiners from  $L_\gamma$  to  $L_\alpha \otimes L_\beta$  by restriction from  $\text{U}(d)$  to  $S_d$ :*

$$C_{\alpha,\beta}^\gamma = \text{Hom}_{\text{U}(d)}(L_\gamma, L_\alpha \otimes L_\beta) \simeq \text{Hom}_{S_d}(V_{\gamma^{+d}}, V_{\alpha^{+d}} \otimes V_{\beta^{+d}}) = G_{\alpha^{+d}, \beta^{+d}}^{\gamma^{+d}}. \quad (51)$$

*In particular, their dimensions are equal:  $c_{\alpha,\beta}^\gamma = g_{\alpha^{+d}, \beta^{+d}}^{\gamma^{+d}}$ .*

*Proof.* The following two representations of  $S_d$  are isomorphic (because restriction is a monoidal functor):

$$\text{Res}_{S_d}^{U(d)}(L_\alpha \otimes L_\beta) \simeq \text{Res}_{S_d}^{U(d)}(L_\alpha) \otimes \text{Res}_{S_d}^{U(d)}(L_\beta). \quad \text{\textcolor{blue}{\{eq:res_monoidal\}}} \quad (52)$$

Applying Lemma 13 (noting  $d$  is sufficiently large), yields

$$\begin{aligned} \text{Res}_{S_d}^{U(d)}(L_\alpha \otimes L_\beta) &= \bigoplus_{|\gamma|=|\alpha|+|\beta|} c_{\alpha,\beta}^\gamma \text{Res}_{S_d}^{U(d)}(L_\gamma) = \bigoplus_{\substack{|\gamma|=|\alpha|+|\beta| \\ d-|\gamma| \geq \gamma_1}} c_{\alpha,\beta}^\gamma V_{\gamma+d} \oplus \cdots, \quad (53) \\ \text{Res}_{S_d}^{U(d)}(L_\alpha) \otimes \text{Res}_{S_d}^{U(d)}(L_\beta) &= (V_{\alpha+d} \oplus \cdots) \otimes (V_{\beta+d} \oplus \cdots) = \bigoplus_{\substack{|\gamma|=|\alpha|+|\beta| \\ d-|\gamma| \geq \gamma_1}} g_{\alpha+d,\beta+d}^{\gamma+d} V_{\gamma+d} \oplus \cdots. \end{aligned} \quad (54)$$

In both formulas above, the “ $\cdots$ ” refers to a direct sum of irreps of  $S_d$  labelled by  $\lambda \vdash d$  whose first part,  $\lambda_1$ , is larger than the first part,  $d - |\gamma|$ , of  $\gamma^+d$ . By Eq. (52), one concludes there exists an isomorphism between all multiplicity spaces, and in particular,  $c_{\alpha,\beta}^\gamma = g_{\alpha+d,\beta+d}^{\gamma+d}$ .  $\square$

**Remark 3.** Note that this result indirectly implies a stability result for the Kronecker coefficients, namely that the value of  $g_{\lambda+(d)}^{\mu+(d),\nu+(d)}$  is eventually independent of  $d$  and moreover reaches the value  $c_{\lambda_{-1}}^{\mu_{-1},\nu_{-1}}$  where  $\lambda_{-1}$  is obtained from  $\lambda$  by removal of the first row of  $\lambda$ . \textcolor{blue}{cor:ud\_sd\_recouple\_correspondence}

**Corollary 15.** Let  $\alpha, \beta, \gamma, \mu, \nu, \lambda$  be Young diagrams. Furthermore, take  $d$  be sufficiently large such that  $\alpha^+d, \beta^+d, \gamma^+d, \mu^+d, \nu^+d, \lambda^+d$  are all well-defined. Then the following recoupling operator for  $S_d$  is obtained from the corresponding recoupling operator for  $U(d)$  by restriction from  $U(d)$  to  $S_d$ :

$$\left\{ \begin{matrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{matrix} \right\} \simeq \left[ \begin{matrix} \alpha^+d & \beta^+d & \mu^+d \\ \gamma^+d & \lambda^+d & \nu^+d \end{matrix} \right]. \quad (55)$$

### 3.6 An entropic inequality for tetrahedral spectra

In Section 3.4, it was shown that the realizability of a given sextuple of candidate eigenvalues,  $(\alpha, \beta, \gamma, \mu, \nu, \lambda)$ , for the tetrahedral Horn problem is encoded both the asymptotics of the a recoupling operator for  $U(d)$  as well as a recoupling operator for the Young subgroups of the symmetric group. The purpose of this section is to prove an upper bound for the norm of the  $U(d)$  recoupling operator whose asymptotic limit gives rise to an entropic inequality which is necessarily satisfied by realizable spectra.

**Lemma 16.** Let  $(\alpha, \beta, \gamma, \mu, \nu, \lambda) \in \Upsilon_k^{\times 6}$  be a sextuple of Young diagrams and let  $\left[ \begin{matrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{matrix} \right]$  be the corresponding  $S_k$ -recoupling operator. Then \textcolor{blue}{lem:sk\_recouple\_bound}

$$\left\| \left[ \begin{matrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{matrix} \right] \right\|_{HS}^2 \leq (g_{\alpha,\beta}^\mu g_{\beta,\gamma}^\nu g_{\mu,\gamma}^\lambda g_{\alpha,\nu}^\lambda)^{\frac{1}{2}} \frac{\dim(V_\nu) \dim(V_\mu)}{\dim(V_\lambda) \dim(V_\beta)} \quad (56)$$

*Proof.* See Appendix A. \(\square\)

**Corollary 17.** Let  $(\alpha, \beta, \gamma, \mu, \nu, \lambda) \in \Upsilon_a \times \Upsilon_b \times \Upsilon_c \times \Upsilon_{a+b} \times \Upsilon_{b+c} \times \Upsilon_{a+b+c}$  be Young diagrams, let  $d \in \mathbb{N}$  be sufficiently large (as in Corollary 17), and let  $\begin{Bmatrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{Bmatrix}$  be the corresponding  $U(d)$ -recoupling operator. Then

$$\left\| \begin{Bmatrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{Bmatrix} \right\|_{HS}^2 \leq (c_{\alpha,\beta}^\mu c_{\beta,\gamma}^\nu c_{\mu,\gamma}^\lambda c_{\alpha,\nu}^\lambda)^{\frac{1}{2}} \frac{\dim(V_{\nu+d}) \dim(V_{\mu+d})}{\dim(V_{\lambda+d}) \dim(V_{\beta+d})} \quad (57)$$

**Theorem 18.** If  $(\alpha, \beta, \gamma, \mu, \nu, \lambda) \in \Lambda_{\geq}^{\mathbb{R}}(d)$  are realizable eigenvalues for the tetrahedral Horn problem, then the following inequality holds:

$$S_p^{a,b,c}(\mu) + S_p^{a,b,c}(\nu) - S_p^{a,b,c}(\beta) - S_p^{a,b,c}(\lambda) \quad (58)$$

*TODO: Figure out how to take the limit of  $\dim(V_\lambda)$  when  $\lambda \vdash pn$  as  $n \rightarrow \infty$ ; the exact form of the entropic inequality is sensitive to how this is done.*

*Proof.* Let  $n \mapsto \lambda(n)$  be a sequence of Young diagrams satisfying  $\lim_{n \rightarrow \infty} \frac{\lambda(n)}{|\lambda(n)|} = r$  and  $\lim_{n \rightarrow \infty} \frac{|\lambda(n)|}{n} = p$  for some  $p \in [0, 1]$ . Then **TODO: prove this**

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \dim V_{\lambda(n)} = pS(r). \quad (59)$$

□

## 4 The spectral quantum marginal problem

The purpose of this section will be to re-articulate the spectral embedding defined in [CŞW18] and state the sense in which the tetrahedral spectra problem is an instance of the quantum marginal problem. From here we can discuss the extent to which known inequalities coming from quantum information theory yield constraints necessarily satisfied by tetrahedral spectra.

### 4.1 A spectral embedding

**Definition 4.** Let  $(H_1, \dots, H_k) \in \mathcal{H}(d)^{\times k}$  be a  $k$ -tuple of  $d \times d$  positive-semidefinite Hermitian matrices and let  $X$  be a subset of  $\{1, \dots, k\}$ . Let  $\{|1\rangle, \dots, |d\rangle\}$  be an orthonormal basis for  $\mathbb{C}^d$  and  $\{|0\rangle, |1\rangle, \dots, |d\rangle\}$  be an orthonormal basis for  $\mathbb{C}^{d+1}$ . Define  $H_X := \sum_{i \in X} H_i$  as the partial sum. Define a Hermitian operator,  $\Phi_X$ , acting on  $(\mathbb{C}^{d+1})^{\otimes k}$  with coefficients with respect to the computational bases  $|i_1 \dots i_k\rangle$  be given by the following formula:<sup>3</sup>

$$\langle i_1 \dots i_k | \Phi_X | j_1 \dots j_k \rangle = \begin{cases} \langle i_p | H_p^{\frac{1}{2}} H_q^{\frac{1}{2}} | j_q \rangle & \exists! i_p, j_q > 0, p, q \in X \\ 0 & \text{otherwise.} \end{cases} \quad (60)$$

<sup>3</sup>Specifically,  $\langle i_1 \dots i_k | \Phi_X | j_1 \dots j_k \rangle$  is non-zero only if there exists a unique  $p \in [k]$  such that  $i_p > 0$  and a unique  $q \in [k]$  such that  $j_q > 0$  (with all other indices  $i_{p'}$  and  $j_{q'}$  for  $p' \neq p$  and  $q' \neq q$  being zero). Furthermore, if  $p, q \in [k]$  are the unique such indices, then  $\langle i_1 \dots i_k | \Phi_X | j_1 \dots j_k \rangle$  is zero if either  $p \notin X$  or  $q \notin X$ ; otherwise  $p, q \in X$  and then  $\langle i_1 \dots i_k | \Phi_X | j_1 \dots j_k \rangle$  takes the value  $\langle i_p | H_p^{\frac{1}{2}} H_q^{\frac{1}{2}} | j_q \rangle$ , the inner product between the  $i_p$ th column of  $H_p^{\frac{1}{2}}$  and the  $j_q$ th column of  $H_q^{\frac{1}{2}}$ .

Note that, up to shuffling the  $k+1$  copies of  $\mathbb{C}^{d+1}$  in  $(\mathbb{C}^{d+1})^{\otimes k}$ ,  $\Phi_X$  is of the form  $\tilde{\Phi}_X \otimes Z_{[k] \setminus X}$  where

$$Z_X = |0 \dots 0\rangle \langle 0 \dots 0|. \quad (61)$$

and where  $\tilde{\Phi}_X$  acts on  $(\mathbb{C}^{d+1})^{\otimes |X|}$ . Also define the operator  $\Gamma_X$  which is  $\tilde{\Phi}_X$  shifted by  $Z_X$

$$\Gamma_X := \tilde{\Phi}_X + (1 - \text{Tr}(H_X))Z_X, \quad (62)$$

and in particular  $\Gamma_\emptyset := 1$ .

**Example 6.** Given two Hermitian operators  $(H_1, H_2) \in \mathcal{H}(d)^{\times 2}$ , the matrix  $\Gamma_{1,2}$  has the following coefficients for all  $i, j, k, l \in \{1, \dots, d\}$ .

$$\langle 00 | \Gamma_{1,2} | 00 \rangle = 1 - \text{Tr}(H_{1,2}), \quad (63)$$

$$\langle i0 | \Gamma_{1,2} | j0 \rangle = \langle i | H_1 | j \rangle, \quad (64)$$

$$\langle 0i | \Gamma_{1,2} | 0j \rangle = \langle i | H_2 | j \rangle, \quad (65)$$

$$\langle i0 | \Gamma_{1,2} | 0j \rangle = \langle i | H_1^{\frac{1}{2}} H_2^{\frac{1}{2}} | j \rangle, \quad (66)$$

$$\langle 0i | \Gamma_{1,2} | j0 \rangle = \langle i | H_2^{\frac{1}{2}} H_1^{\frac{1}{2}} | j \rangle, \quad (67)$$

$$\langle ik | \Gamma_{1,2} | jl \rangle = 0. \quad (68)$$

*lem:spec\_embed*

**Lemma 19.** *Let everything be defined as in Definition 4. The follow statements hold for all subsets  $X \subseteq [k]$ .*

(i) *Up to padding by zeros, the eigenvalues of  $\tilde{\Phi}_X$ ,  $\Phi_X$  and  $H_X$  are all identical.*

$$\lambda(\tilde{\Phi}_X) \simeq \lambda(\Phi_X) \simeq \lambda(H_X). \quad (69)$$

(ii) *The eigenvalues of  $\Gamma_X$  are equal to the eigenvalues of  $\tilde{\Phi}_X$  plus the additional value of  $1 - \text{Tr}(H_X)$ .*

$$\lambda(\Gamma_X) \simeq \lambda(\tilde{\Phi}_X) \oplus \{1 - \text{Tr}(H_X)\}. \quad (70)$$

(iii) *The partial trace of  $\Gamma_{[k]}$  over  $[k] \setminus X$  is equal to  $\Gamma_X$ :*

$$\text{Tr}_{[k] \setminus X}(\Gamma_{[k]}) = \Gamma_X. \quad (71)$$

*Proof.* (i) From the definition of  $\Phi_X$ , one observes that  $\langle i_1 \dots i_k | \Phi_X | i_1 \dots i_k \rangle$  is only non-zero if there exists a unique non-zero  $i_p \neq 0$  where  $p \in X$  and thus

$$\text{Tr}(\Phi_X) = \sum_{p \in X} \sum_{j=1}^d \langle j | H_p | j \rangle = \sum_{p \in X} \text{Tr}(H_p) = \text{Tr}(H_X). \quad (72)$$

More generally, for any positive integer  $\ell \in \mathbb{N}_{>0}$ , one observes that  $\text{Tr}(\Phi_X^\ell) = \text{Tr}(H_X^\ell)$ . **(TODO: Prove this directly.)** As the eigenvalues of any matrix  $X$  are determined (up to padding by zeros) by the values of  $\text{Tr}(X^\ell)$ , we conclude that  $\Phi_X$  and  $H_X$  have the same eigenvalues (up to padding by zeros) **(TODO: Is there a more direct way to prove this.)**

(ii) Note that  $\tilde{\Phi}_X$  and its eigenvectors are supported in the subspace of  $(\mathbb{C}^{d+1})^{\otimes |X|}$  spanned by  $|i_1 \cdots i_{|X|}\rangle$  for which exactly one  $i_p$ 's is non-zero. Therefore  $|0 \cdots 0\rangle$  is orthogonal to all eigenvectors of  $\tilde{\Phi}_X$  and thus  $|0 \cdots 0\rangle$  is an eigenvector of  $\Gamma_X$  with eigenvalue  $1 - \text{Tr}(H_X)$ .

(iii) The partial trace of  $\Phi_{[k]}$  over  $[k] \setminus X$ , when performed in the computational basis  $|i_1 \cdots i_k\rangle$ , can be divided into three portions depending on the number of non-zero entries in  $(i_p \mid p \in [k] \setminus X)$ ; first, is all indices  $i_p$  are zero, second, where exactly one  $i_p$  is non-zero, and third where more than one index  $i_p$  is non-zero. The respective contributions are as follows:

$$\text{Tr}_{[k] \setminus X}(\Phi_{[k]}) = \tilde{\Phi}_X + \text{Tr}(H_{[k] \setminus X})Z_X + 0 \quad (73)$$

Recalling the definition of  $\Gamma_{[k]}$ ,

$$\Gamma_{[k]} := \Phi_{[k]} + (1 - \text{Tr}(H_{[k]}))Z_{[k]} \quad (74)$$

One concludes

$$\begin{aligned} \text{Tr}_{[k] \setminus X}(\Gamma_{[k]}) &= \text{Tr}_{[k] \setminus X}(\Phi_{[k]} + (1 - \text{Tr}(H_{[k]}))Z_{[k]}), \\ &= \tilde{\Phi}_X + \text{Tr}(H_{[k] \setminus X})Z_X + (1 - \text{Tr}(H_{[k]}))Z_X, \\ &= \tilde{\Phi}_X + (1 - \text{Tr}(H_X))Z_X, \\ &= \Gamma_X. \end{aligned} \quad (75)$$

□

**Corollary 20.** *By Lemma 19, one has  $\text{Tr}(\Gamma_X) = 1$ . If the eigenvalues of  $H_X$  satisfy*

$$\text{Tr}(H_X) \leq 1, \quad (76)$$

*then  $\Gamma_X$  is positive semidefinite, and thus  $\Gamma_X$  is a density matrix. In such a case, assuming  $H_X \neq 0$  so that  $\text{Tr}(H_X) \in (0, 1]$ , the von Neumann entropy of  $\Gamma_X$  is related to the von Neumann entropy of  $H_X / \text{Tr}(H_X)$  by the equation <sup>4</sup>*

$$S(\Gamma_X) = h(\text{Tr}(H_X)) + \text{Tr}(H_X)S\left(\frac{H_X}{\text{Tr}(H_X)}\right). \quad (77)$$

*Moreover, if the eigenvalues of  $H_X$  satisfy the stronger condition*

$$\lambda_1(H_X) + \text{Tr}(H_X) \leq 1, \quad (78)$$

*then the maximum eigenvalue of  $\Gamma_X$  is  $1 - \text{Tr}(H_X)$ .*

*Proof.* The only non-trivial claim involves the relation between the von Neumann entropies of  $\Gamma_X$  and  $H_X / \text{Tr}(H_X)$ , respectively given by

$$S(\Gamma_X) = -(1 - \text{Tr}(H_X)) \log(1 - \text{Tr}(H_X)) - \sum_{i=1}^d \lambda_i(H_X) \log \lambda_i(H_X), \quad (79)$$

$$S\left(\frac{H_X}{\text{Tr}(H_X)}\right) = -\frac{1}{\text{Tr}(H_X)} \sum_{i=1}^d \lambda_i(H_X) \log \lambda_i(H_X) + \log(\text{Tr}(H_X)). \quad (80)$$

---

<sup>4</sup>Here  $h(p) = -p \log p - (1-p) \log(1-p)$  denotes the binary entropy of  $p \in [0, 1]$ .

Multiplying the von Neumann entropy of  $H_X/\text{Tr}(H_X)$  by  $\text{Tr}(H_X)$  yields

$$\text{Tr}(H_X)S\left(\frac{H_X}{\text{Tr}(H_X)}\right) = S(\Gamma_X) + \text{Tr}(H_X)\log(\text{Tr}(H_X)) + (1 - \text{Tr}(H_X))\log(1 - \text{Tr}(H_X)), \quad (81)$$

which is the claimed result. □

## 5 Discussion

## Reference notes

- A popular textbook on quantum groups with an emphasis on the connection to Poisson manifolds, Lie bialgebras and their quantizations by Chari and Pressley [CP95]
- The general subject of multivariate trace inequalities involving exponentials of sums of Hermitian and products of their exponentials is, of course, intimately related. See for instance the paper by Sutter, Berta, and Tomamichel [SBT17].
- Also the article [FT14] by Forrester and Thompson which describes the connection to hyperbolic geometry quite explicitly.
- Course notes on Poisson geometry with an eye toward quantization by Ciccoli and Witkowski [CW06]
- An excellent expository overview of the big picture by Knutson [Knu00]
- The masters essay/thesis of Backens (TODO: check if the pdf Matthias sent to me is the final version) [Bac10]
- An excellent introduction to these ideas is the recent masters thesis on various aspects of additive and multiplicative Horn problems in connection to Vinnikov curves and the Lax conjecture due to Hess and supervised by S. Kuhlmann and M. Infusino [Hes20]
- The PhD thesis of Esposito (from the maths department here in Copenhagen) serves as a introduction to the world of Poisson momentum maps and the notion of a “quantum” momentum map as a deformation of the ordinary “classical” momentum map (where deformation is in the sense of deformation quantization) [Esp12].
- Alekseev [Ale97]
- Papers contained results pertaining to the generalization of Duistermaat-Heckman measures for Hamiltonian group actions with group-valued moment maps by Alekseev, Meinrenken, and Woodward [AMW99; AMW00]
- The proof of Thompson’s conjecture in the style of Alekseev, Meinrenken, and Woodward is the “linearization paper” due to [AMW01].
- See also the related article due to Lamb which summarizes the ideas of [AMW01] while also providing more technical background [Lam05]
- Lu and Evens extend the proof technique of [AMW01] to a larger class of Lie groups [LE03]
- Alekseev, Podkopaeva, and Szenes [APS17a]
- Alekseev, Podkopaeva, and Szenes [APS17b]
- Alekseev and Meinrenken [AM07]

- Deforming the symplectic volume by the  $s$  parameter Alekseev et al. [Ale+18]
- Klyachko [Kly00]
- Christandl [Chr08]
- Christandl, Şahinoğlu, and Walter [CŞW18]
- Kirillov and Reshetikhin [KR90]
- Usage of Wigner’s  $6j$  symbol to compute asymptotic errors rates for the promise problem of distinguishing between two possible “change points” for a sequence of unknown pure states to transition from one unknown type to another [AH11]
- A great summary by Roberts [Rob02]; also the original result for the asymptotics of the  $6j$  symbols due to Roberts [Rob99]
- Coquereaux and Zuber explicitly outline the link between the orbital measures (i.e. the induced distributions in the sense of Klyachko’s random walk approach) and the magnitude of the stretched Littlewood-Richardson coefficients [CZ18]. See also the papers [CMZ20; CMZ19; Zub18].
- A recent article by McSwiggen is an excellent introduction to the Harish-Chandra integral [McS21]. The proof of the Harish-Chandra formula provide by McSwiggen using the heat equation methods from Fegan’s 1978 paper [Feg78] seems very promising for deriving a connection to the character formula for the norm of the recoupling operator!
- See also Coquereaux’s later paper which promotes the alternative, albeit equivalent, usage of the integer-valued gon and tet functions in place of the rational-valued theta (triangular) and irrational-valued  $6j$  symbols [Coq23]. Most relevant for this project is [Coq23, Section IV D] which discusses the apparent lack of substantial generalization from  $SU(2)$  to  $SU(N)$  (“*apart from very few special cases (low dimension of irreps and no multiplicity) worked out by physicists at the time of the eightfold way, we are not aware of any published general result*”).
- The recent paper by Alekseev, Morozov, and Sleptsov (different Alekseev!) covers the case of some multiplicity free formulas for the quantum  $6j$  symbols for  $U_q(\mathfrak{sl}_N)$  [AMS20]
- Boalch [Boa07]
- Taylor and Woodward [TW06]
- Henriques and Kamnitzer [HK05]
- Octahedron Recurrence/Rule Robbins and Rumsey Jr [RR86] applied to the “Hive” ring Knutson, Tao, and Woodward [KTW03], and also Henriques and Kamnitzer [HK06]

- The integral formula for the square of the  $SU(2)$   $6j$  symbol can be performed by an appropriate change of variables [FL03]. See also other papers by Laurent Freidel and collaborators that are relevant [FK00; FNR07; FKL10].
- Speyer [Spe05]
- Generalizations of Horn’s problem to the product of unitary matrices [AW97; Bel08; BK16]
- Bhatia [Bha01]
- History account offered by Daftuar and Hayden [DH05], and technical review article by Fulton [Ful00].
- Connecting group-valued moment maps to spherical, flat, and hyperbolic tetrahedron via Minkowski’s theorem from the reconstruction of flat tetrahedron Haggard, Han, and Riello. See also the comments in [HHR16, Section 11] on the utility of *quasi-poisson* (controlled failure of the Jacobi identity) manifolds and group-valued moment maps.
- Foundational results concerning the relationships between the edge lengths, face areas, dihedral and solid angles of a tetrahedron by Richardson [Ric02]
- Potential application to inverse kinematics of 3-link arms in three dimensions? (Thanks Harold!)
- TODO: track down pdfs for the original triple of papers by Thompson and Therianos titled “On the singular values of matrix products” I, II, and III from 1973
- TODO: track down pdfs for the original triple of papers by Thompson and Freede titled “On the eigenvalues of sums of Hermitian matrices” I, II, and III from 1973
- TODO: track down a pdf for Thompson’s 1988 “Matrix Spectral Inequalities” referenced by Klyachko
- Thompson and Therianos [TT73] present a number of interesting counter-examples to the question of whether the inequalities pertaining to the spectra of general matrices can be deduced from looking at the spectra of diagonal matrices (both in the additive and multiplicative cases).
- Elementary introduction to toy models of three-dimensional quantum space [Bar03]
- Horn’s original paper where they conjectured the form of a necessary and sufficient set of inequalities via an inductive scheme for increasing dimension [Hor62]
- Berline, Vergne, and Walter a pedagogical account of the triple role played by Horn’s inequalities in characterizing the existence of invariant vectors in tensor products, the eigenvalues of sums of Hermitian matrices, and intersections of Schubert varieties [BVW18]

- Regge and Williams [RW00] a relatively recent review of discrete approaches to gravity. The essential idea, in contrast to the differential geometry of general relativity, is to model curvature discretely instead of continuously. In particular, one can think of a triangulation of the surface of a sphere; the curvature of the sphere is encoded in the triangulation when a configuration of three (or more) triangles meet at a common vertex with sum of interior angles deviating from  $\pi$ . It is in this sense that Regge and Williams refers to these types of models as “piecewise-flat”. Contained is an overview of various attempts to generalize discrete theories of three-dimensional gravity to four dimensions [RW00, Section III].

## A Proof of symmetric group recoupling operator upper bound

sec:sk\_recouple\_bound\_proof

The purpose of this section is to prove Lemma 16.

TODO: Make the notation and diagrammatic conventions used here match the rest of the document, also resolve ambiguities introduced by the the “dot” notation.

**Definition 5** ( $S_k$  recoupling operators). Given six Young diagrams  $\alpha, \beta, \gamma, \mu, \nu$  and  $\lambda$  (each with fixed size  $k \in \mathbb{N}$ ), the *recoupling operator* is a linear map

$$\begin{bmatrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{bmatrix} : G_{\alpha,\beta}^{\mu} \otimes G_{\mu,\gamma}^{\lambda} \rightarrow G_{\alpha,\nu}^{\lambda} \otimes G_{\beta,\gamma}^{\nu}, \quad (82)$$

between the multiplicity spaces of irreps of the symmetric group defined as


$$\begin{array}{c} G_{\alpha,\nu}^{\lambda} \quad G_{\beta,\gamma}^{\nu} \\ \uparrow \quad \uparrow \\ \boxed{\begin{array}{ccc} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{array}} \\ \uparrow \quad \uparrow \\ G_{\alpha,\beta}^{\mu} \quad G_{\mu,\gamma}^{\lambda} \end{array} := \frac{1}{\textcircled{V_{\lambda}}} \begin{array}{c} G_{\alpha,\nu}^{\lambda} \quad G_{\beta,\gamma}^{\nu} \\ \nearrow V_{\nu} \quad \nwarrow V_{\gamma} \\ \begin{array}{ccc} & V_{\beta} & \\ \nearrow V_{\alpha} & & \nwarrow V_{\lambda} \\ \searrow V_{\mu} & & \nearrow V_{\gamma} \end{array} \\ \nwarrow G_{\alpha,\beta}^{\mu} \quad \nearrow G_{\mu,\gamma}^{\lambda} \end{array}. \quad (83)$$

The recoupling operator arises when applying Schur’s lemma (with multiplicities) to diagram on the left-hand side of the subsequent equation:

$$\begin{array}{c} \lambda \quad M_{\lambda}^{\alpha,\nu} \quad \nu \quad M_{\nu}^{\beta,\gamma} \\ \nearrow \quad \nwarrow \\ \begin{array}{ccc} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{array} \\ \nwarrow \quad \nearrow \\ \lambda \quad M_{\mu}^{\alpha,\beta} \quad \mu \quad M_{\lambda}^{\mu,\gamma} \end{array} = \begin{array}{c} M_{\lambda}^{\alpha,\nu} \quad M_{\nu}^{\beta,\gamma} \\ \uparrow \quad \uparrow \\ \boxed{\begin{array}{ccc} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{array}} \\ \uparrow \quad \uparrow \\ \lambda \quad M_{\mu}^{\alpha,\beta} \quad M_{\lambda}^{\mu,\gamma} \end{array}. \quad (84)$$


The recoupling operator derives its name from the recoupling theory of angular momenta and serves the purpose of a change of basis between two distinct (de-)coupling schemes.

Specifically, the recoupling operator satisfies the following equation:




$$\text{Diagram} = \sum_{\nu} \text{Diagram} \cdot \begin{bmatrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{bmatrix} \quad \text{\texttt{\{eq:recoupling-trees\}}} \quad (85)$$

As a shorthand for this equation, we simply write:




$$\text{Diagram} \left[ \begin{array}{ccc} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{array} \right] \text{Diagram} \quad (86)$$

The next result is a consequence of an application of Schur's lemma to the identity on  $V_{\alpha}$ :



$$\text{Diagram} = \frac{1}{\alpha} \text{Diagram} = \frac{\mu}{\alpha} \text{Diagram} \quad (87)$$

Combining two instances of the above result yields:



$$\text{Diagram} = \frac{\mu \nu}{\beta} \text{Diagram} \quad (88)$$

*Proof of Lemma 16.* Applying the Cauchy-Schwarz inequality along a particular edge-cut

indicated below, and applying Schur's lemma multiple times yields

$$\begin{aligned}
 & \left( \lambda \right)^2 \left[ \begin{array}{c} \boxed{\begin{matrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{matrix}} \\ G_{\alpha,\nu}^\lambda \\ \boxed{\begin{matrix} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{matrix}} \\ G_{\mu,\gamma}^\lambda \end{array} \right] = \left[ \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} \right]^{\frac{1}{2}} = \left[ \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array} \right]^{\frac{1}{2}} \\
 & \leq \left[ \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} \right]^{\frac{1}{2}} = \frac{1}{(\beta)(\lambda)} \left[ \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array} \right]^{\frac{1}{2}} \\
 & = \frac{(\lambda)^2 (\mu)(\nu)}{(\beta)(\lambda)} \left[ \begin{array}{c} \text{Diagram 9} \end{array} \right]^{\frac{1}{2}}
 \end{aligned} \tag{89}$$

□

## B Proof of recoupling operator norm equivalences

[sec:recouple\\_norms](#)

TODO: Figure out if a “Young recoupling operator” is even a useful object to consider now that my entropy inequality proof failed.

[defn:young\\_recouple](#)

**Definition 6.** Let  $a, b, c \in \mathbb{N}_{>0}$  be given and let

$$\alpha \in \Upsilon_a, \quad \beta \in \Upsilon_b, \quad \gamma \in \Upsilon_c, \quad \mu \in \Upsilon_{a+b}, \quad \nu \in \Upsilon_{b+c}, \quad \lambda \in \Upsilon_{a+b+c}. \tag{90}$$

The *Young recoupling operator* is the operator

$$\left[ \begin{array}{ccc} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{array} \right] : C_{\alpha,\beta}^\mu \otimes C_{\mu,\gamma}^\lambda \rightarrow C_{\alpha,\nu}^\lambda \otimes C_{\beta,\gamma}^\nu, \tag{91}$$

between the multiplicity spaces  $C_{\alpha,\beta}^\mu := \text{Hom}_{S_a \times S_b}(V_\alpha \otimes V_\beta, V_\mu)$  defined as

$$\left[ \begin{array}{ccc} \alpha & \beta & \mu \\ \gamma & \lambda & \nu \end{array} \right] := \frac{\text{Tr}_{V_\alpha \otimes V_\beta \otimes V_\gamma}[(\text{id}_{V_\alpha} \otimes Y_{\beta\gamma}^{\nu;\dagger}) Y_{\alpha\nu}^{\lambda;\dagger} Y_{\mu\gamma}^\lambda (Y_{\alpha\beta}^\mu \otimes \text{id}_{V_\gamma})]}{\dim(V_\alpha \otimes V_\beta \otimes V_\gamma)} \tag{92}$$



The remaining two isometries,  $\iota_{III}$  and  $\iota_{IV}$ , involve decompositions of tensor products of irreps of the unitary group:

$$\begin{aligned}
 & \text{Diagram 1} = \text{Diagram 2} = \text{Diagram 3} \quad (98)
 \end{aligned}$$

□

# References

- [AH11] Daiki Akimoto and Masahito Hayashi. “Discrimination of the change point in a quantum setting”. In: *Physical Review A* 83.5 (2011), p. 052328.
- [Ale+18] Anton Alekseev et al. “Concentration of symplectic volumes on Poisson homogeneous spaces”. In: *arXiv preprint arXiv:1808.06975* (2018).
- [Ale97] A Yu Alekseev. “On Poisson actions of compact Lie groups on symplectic manifolds”. In: *Journal of Differential Geometry* 45.2 (1997), pp. 241–256.
- [AM07] Anton Alekseev and Eckhard Meinrenken. “Ginzburg-Weinstein via Gelfand-Zeitlin”. In: *Journal of Differential Geometry* 76.1 (2007), pp. 1–34.
- [AMS20] Victor Alekseev, Andrey Morozov, and Alexey Sleptsov. “Multiplicity-free  $U_q(\mathfrak{sl}_N)$  6-j symbols: relations, asymptotics, symmetries”. In: *Nuclear Physics B* 960 (2020), p. 115164.
- [AMW00] Anton Alekseev, Eckhard Meinrenken, and Christopher Woodward. “Group-valued equivariant localization”. In: *Inventiones mathematicae* 140.2 (2000), pp. 327–350.
- [AMW01] Anton Alekseev, Eckhard Meinrenken, and Chris Woodward. “Linearization of Poisson actions and singular values of matrix products”. In: *Annales de l’institut Fourier*. Vol. 51. 6. 2001, pp. 1691–1717.
- [AMW99] Anton Alekseev, Eckhard Meinrenken, and Chris Woodward. “Duistermaat-Heckman distributions for group valued moment maps”. In: *arXiv preprint math/9903087* (1999).
- [APS17a] Anton Alekseev, Masha Podkopaeva, and Andras Szenes. “A symplectic proof of the Horn inequalities”. In: *Advances in Mathematics* 318 (2017), pp. 711–736.
- [APS17b] Anton Alekseev, Masha Podkopaeva, and Andras Szenes. “The Horn problem and planar networks”. In: *Advances in Mathematics* 318 (2017), pp. 678–710.
- [ARS88] Robert Alicki, Slawomir Rudnicki, and Slawomir Sadowski. “Symmetry properties of product states for the system of  $N$   $n$ -level atoms”. In: *Journal of mathematical physics* 29.5 (1988), pp. 1158–1162.
- [AW97] Sharad Agnihotri and Chris Woodward. “Eigenvalues of products of unitary matrices and quantum Schubert calculus”. In: *arXiv preprint alg-geom/9712013* (1997).
- [Bac10] Miriam Backens. “The 6j symbols and an extended version of Horn’s problem”. In: (2010).
- [Bar03] John W Barrett. “Geometrical measurements in three-dimensional quantum gravity”. In: *International Journal of Modern Physics A* 18.suppl2 (2003), pp. 97–113.
- [Bel08] Prakash Belkale. “Quantum generalization of the Horn conjecture”. In: *Journal of the American Mathematical Society* 21.2 (2008), pp. 365–408.

- [Bha01] Rajendra Bhatia. “[Linear algebra to quantum cohomology: the story of Alfred Horn’s inequalities](#)”. In: *The American Mathematical Monthly* 108.4 (2001), pp. 289–318.
- [BK16] Prakash Belkale and Shrawan Kumar. “The multiplicative eigenvalue problem and deformed quantum cohomology”. In: *Advances in Mathematics* 288 (2016), pp. 1309–1359.
- [Boa07] Philip P Boalch. “[Regge and Okamoto symmetries](#)”. In: *Communications in mathematical physics* 276.1 (2007), pp. 117–130.
- [BVW18] Nicole Berline, Michèle Vergne, and Michael Walter. “The Horn inequalities from a geometric point of view”. In: *L’Enseignement Mathématique* 63.3 (2018), pp. 403–470.
- [Chr08] Matthias Christandl. “A Quantum Information-Theoretic Proof of the Relation between Horn’s Problem and the Littlewood-Richardson Coefficients”. In: *Lecture Notes in Computer Science*. Springer Berlin Heidelberg, 2008, pp. 120–128.
- [CMZ19] Robert Coquereaux, Colin McSwiggen, and Jean-Bernard Zuber. “Revisiting Horn’s problem”. In: *Journal of Statistical Mechanics: Theory and Experiment* 2019.9 (2019), p. 094018.
- [CMZ20] Robert Coquereaux, Colin McSwiggen, and Jean-Bernard Zuber. “On Horn’s problem and its volume function”. In: *Communications in Mathematical Physics* 376.3 (2020), pp. 2409–2439.
- [Coq23] Robert Coquereaux. “About integer-valued variants of the theta and 6j symbols”. In: *Journal of Mathematical Physics* 64.3 (2023).
- [CP95] Vyjayanthi Chari and Andrew N Pressley. *A guide to quantum groups*. Cambridge university press, 1995.
- [CŞW18] Matthias Christandl, M Burak Şahinoğlu, and Michael Walter. “Recoupling coefficients and quantum entropies”. In: *Annales Henri Poincaré*. Vol. 19. 2. Springer. 2018, pp. 385–410.
- [CW06] Nicola Ciccoli and Pawel Witkowski. “From Poisson to quantum geometry”. In: *Lecture Notes, available at <http://toknotes.mimuw.edu.pl>* (2006).
- [CZ18] Robert Coquereaux and Jean-Bernard Zuber. “From orbital measures to Littlewood–Richardson coefficients and hive polytopes”. In: *Annales de l’Institut Henri Poincaré D* 5.3 (2018), pp. 339–386.
- [DH05] Sumit Daftuar and Patrick Hayden. “[Quantum state transformations and the Schubert calculus](#)”. In: *Annals of Physics* 315.1 (2005), pp. 80–122.
- [Esp12] Chiara Esposito. “On the classical and quantum momentum map”. In: (2012). 1203.4112: [arxiv](#).
- [Feg78] HD Fegan. “The heat equation on a compact Lie group”. In: *Transactions of the American Mathematical Society* 246 (1978), pp. 339–357.
- [FK00] Laurent Freidel and Kirill Krasnov. “Simple spin networks as Feynman graphs”. In: *Journal of Mathematical Physics* 41.4 (2000), pp. 1681–1690.

- [FKL10] Laurent Freidel, Kirill Krasnov, and Etera R Livine. “Holomorphic factorization for a quantum tetrahedron”. In: *Communications in Mathematical Physics* 297.1 (2010), pp. 45–93.
- [FL03] Laurent Freidel and David Louapre. “Asymptotics of 6j and 10j symbols”. In: *Classical and Quantum Gravity* 20.7 (2003), p. 1267.
- [FNR07] L Freidel, K Noui, and Ph Roche. “6J symbols duality relations”. In: *Journal of mathematical physics* 48.11 (2007).
- [FT14] Peter J Forrester and Colin J Thompson. “The Golden-Thompson inequality: Historical aspects and random matrix applications”. In: *Journal of Mathematical Physics* 55.2 (2014).
- [Ful00] William Fulton. “Eigenvalues, invariant factors, highest weights, and Schubert calculus”. In: *Bulletin of the American Mathematical Society* 37.3 (2000), pp. 209–249.
- [Hes20] Sarah Tanja Hess. “Examination of the connection between the Horn Problem and the Lax Conjecture”. In: (2020).
- [HHR16] Hal M Haggard, Muxin Han, and Aldo Riello. “Encoding curved tetrahedra in face holonomies: phase space of shapes from group-valued moment maps”. In: *Annales Henri Poincaré*. Vol. 17. Springer. 2016, pp. 2001–2048.
- [HK05] Andre Henriques and Joel Kamnitzer. *Crystals and coboundary categories*. 2005. arXiv: [math/0406478](https://arxiv.org/abs/math/0406478) [[math.QA](https://arxiv.org/archive/math)].
- [HK06] André Henriques and Joel Kamnitzer. “The octahedron recurrence and gl<sub>n</sub> crystals”. In: *Advances in Mathematics* 206.1 (2006), pp. 211–249.
- [HM02] Masahito Hayashi and Keiji Matsumoto. “Quantum universal variable-length source coding”. In: *Physical Review A* 66.2 (2002), p. 022311.
- [Hor62] Alfred Horn. “Eigenvalues of sums of Hermitian matrices.” In: (1962).
- [Kly00] Alexander A Klyachko. “Random walks on symmetric spaces and inequalities for matrix spectra”. In: *Linear Algebra and its Applications* 319.1-3 (2000), pp. 37–59.
- [Kly98] Alexander A Klyachko. “Stable bundles, representation theory and Hermitian operators”. In: *Selecta Mathematica* 4 (1998), pp. 419–445.
- [Knu00] Allen Knutson. “The symplectic and algebraic geometry of Horn’s problem”. In: *Linear Algebra and its Applications* 319.1-3 (2000), pp. 61–81.
- [KR90] AN Kirillov and N Yu Reshetikhin. “Representations of the Algebra  $U_q(\mathfrak{sl}(2))$ ,  $q$ -Orthogonal Polynomials and Invariants of Links”. In: *New developments in the theory of knots* 11 (1990), p. 202.
- [KT99] Allen Knutson and Terence Tao. “The honeycomb model of  $GL_n(\mathbb{C})$  tensor products I: Proof of the saturation conjecture”. In: *Journal of the American Mathematical Society* 12.4 (1999), pp. 1055–1090.

- [KTW03] Allen Knutson, Terence Tao, and Christopher T Woodward. “A positive proof of the Littlewood-Richardson rule using the octahedron recurrence”. In: *arXiv preprint math/0306274* (2003).
- [KW01] Michael Keyl and Reinhard F Werner. “[Estimating the spectrum of a density operator](#)”. In: *Physical Review A* 64.5 (2001), p. 052311.
- [Lam05] McKenzie Lamb. “A proof of the Thompson conjecture using Poisson geometry”. In: (2005).
- [LE03] Jiang-Hua Lu and Sam Evens. “Thompson’s conjecture for real semi-simple Lie groups”. In: (2003). arXiv: [math/0310098](#).
- [Lit58] Dudley E Littlewood. “Products and plethysms of characters with orthogonal, symplectic and symmetric groups”. In: *Canadian Journal of Mathematics* 10 (1958), pp. 17–32.
- [McS21] Colin McSwiggen. “The Harish-Chandra integral: an introduction with examples”. In: *L’Enseignement Mathématique* 67.3 (2021), pp. 229–299.
- [Mur55] Francis D Murnaghan. “On the analysis of the Kronecker product of irreducible representations of  $S_n$ ”. In: *Proceedings of the National Academy of Sciences* 41.7 (1955), pp. 515–518.
- [Nis00] Kyo Nishiyama. “Restriction of the irreducible representations of  $GL_n$  to the symmetric group  $S_n$ ”. In: *preprint* (2000).
- [Ric02] G Richardson. “The trigonometry of the tetrahedron”. In: *The Mathematical Gazette* 2.32 (1902), pp. 149–158.
- [Rob02] Justin Roberts. “Asymptotics and 6j-symbols”. In: *Geom. Topol. Monogr* 4 (2002), pp. 245–261.
- [Rob99] Justin Roberts. “[Classical 6j-symbols and the tetrahedron](#)”. In: *Geometry & Topology* 3.1 (1999), pp. 21–66.
- [RR86] David P Robbins and Howard Rumsey Jr. “Determinants and alternating sign matrices”. In: *Advances in Mathematics* 62.2 (1986), pp. 169–184.
- [RW00] Tullio Regge and Ruth M Williams. “Discrete structures in gravity”. In: *Journal of Mathematical Physics* 41.6 (2000), pp. 3964–3984.
- [SBT17] David Sutter, Mario Berta, and Marco Tomamichel. “[Multivariate trace inequalities](#)”. In: *Communications in Mathematical Physics* 352.1 (2017), pp. 37–58. arXiv: [1604.03023v3](#).
- [Spe05] David E Speyer. “Horn’s problem, Vinnikov curves, and the hive cone”. In: (2005).
- [TT73] RC Thompson and S Therianos. “The Eigenvalues and Singular Values of Matrix Sums and Products. VII (1)”. In: *Canadian Mathematical Bulletin* 16.4 (1973), pp. 561–569.
- [TW06] Yuka U Taylor and Christopher T Woodward. “6 j symbols for and non-Euclidean tetrahedra”. In: *Selecta Mathematica* 11.3-4 (2006), p. 539.

- [Zub18] Jean-Bernard Zuber. “Horn’s problem and Harish-Chandra’s integrals. Probability density functions”. In: *Annales de l’Institut Henri Poincaré D* 5.3 (2018), pp. 309–338.