

कृषि-मौसम संबंधी सलाहकार सेवा बुलेटिन

Agromet Advisory Service Bulletin for the State of Karnataka

(Advisories mentioned in the bulletin are prepared by different AMFUs and DAMUs in Karnataka)

बुलेटिन संख्या /Bulletin No. 69/2026

28.08.2026 को जारी किया गया

Issued on 28.08.2026

*FORECAST & WARNING FOR KARNATAKA FOR NEXT SEVEN DAYS: *

Very Heavy /Heavy Rainfall warning: Nil

***FORECAST & WARNING FOR KARNATAKA FOR NEXT SEVEN DAYS: ***

Day 1, 28-Aug-2026

COASTAL KARNATAKA

Light to moderate rain with sustained wind (30-40 Kmph) likely to occur at most places over Dakshina Kannada, Udupi & Uttara Kannada districts.

NORTH INTERIOR KARNATAKA

Light to moderate rain with sustained wind (40-50 Kmph) likely to occur at one or two places over Belagavi, Dharwad, Haveri districts.

SOUTH INTERIOR KARNATAKA

Light to moderate rain with sustained wind (30-40 Kmph) likely to occur at one or two places over Shivamogga, Chikkamagaluru, Kodagu, Hassan districts. Light rain with sustained wind (30-40 Kmph) likely to occur at a one or two places over Bengaluru (Rural), Bengaluru (Urban), Tumakuru, Kolar, Chikaballapura districts.

Day 2, 29-Aug-2026

COASTAL KARNATAKA

Light to moderate rain with sustained wind (30-40 Kmph) likely to occur at many places over Dakshina Kannada, Udupi and Uttara Kannada districts.

NORTH INTERIOR KARNATAKA

Light to moderate rain with sustained wind (40-50 Kmph) likely to occur at one or two places over Belagavi, Dharwad, Haveri districts.

SOUTH INTERIOR KARNATAKA

Light to moderate rain with sustained wind (30-40 Kmph) likely to occur at one or two places over Shivamogga, Chikkamagaluru, Kodagu, Hassan districts. Light rain with

sustained wind (30-40 Kmph) likely to occur at a one or two places over Bengaluru (Rural), Bengaluru (Urban), Tumakuru, Kolar, Chikkaballapura districts.

Day 3, 30-Aug-2026

COASTAL KARNATAKA

Light to moderate rain likely to occur at many places over Dakshina Kannada, Udupi and Uttara Kannada districts.

NORTH INTERIOR KARNATAKA

Light to moderate rain likely to occur at one or two places over Belagavi, Dharwad, Haveri districts.

SOUTH INTERIOR KARNATAKA

Light to moderate rain likely to occur at one or two places over Shivamogga, Chikkamagaluru, Kodagu, Hassan districts. Light rain likely to occur at a one or two places over Bengaluru (Rural), Bengaluru (Urban), Tumakuru, Kolar, Chikkaballapura districts.

Day 4, 31-Aug-2026

COASTAL KARNATAKA

Light to moderate rain likely to occur at many places over Dakshina Kannada, Udupi and Uttara Kannada districts.

NORTH INTERIOR KARNATAKA

Light to moderate rain likely to occur at one or two places over Belagavi, Dharwad, Haveri districts.

SOUTH INTERIOR KARNATAKA

Light to moderate rain likely to occur at one or two places over Shivamogga, Chikkamagaluru, Hassan and Kodagu districts.

Day 5, 01-Sept-2026

COASTAL KARNATAKA

Light to moderate rain likely to occur at many places over Dakshina Kannada, Udupi and Uttara Kannada districts.

NORTH INTERIOR KARNATAKA

Light to moderate rain likely to occur at one or two places over Belagavi, Dharwad, Haveri districts.

SOUTH INTERIOR KARNATAKA

Light to moderate rain likely to occur at one or two places over Shivamogga, Chikkamagaluru, Hassan and Kodagu districts.

Day 6, 02-Sept-2026

COASTAL KARNATAKA

Light to moderate rain likely to occur at most places over Dakshina Kannada, Udupi and Uttara Kannada districts.

NORTH INTERIOR KARNATAKA

Light to moderate rain likely to occur at a few places over Bagalkot, Belagavi, Bidar, Dharwad, Gadag, Haveri, Kalburgi, Koppal, Raichur, Vijayapura and Yadgir districts.

SOUTH INTERIOR KARNATAKA

Light to moderate rain likely to occur at a few places over Shivamogga, Chikkamagaluru, Hassan, Kodagu, Ballari, Bengaluru (Rural), Bengaluru (Urban), Chamarajanagara, Chikkaballapura, Chitradurga, Davanagere, Kolar, Mandya, Mysuru, Ramanagara, Tumakuru and Vijayanagara districts.

Day 7, 03-Sept-2026

COASTAL KARNATAKA

Light to moderate rain likely to occur at most places over Dakshina Kannada, Udupi and Uttara Kannada districts.

NORTH INTERIOR KARNATAKA

Light to moderate rain likely to occur at a few places over Bagalkot, Belagavi, Bidar, Dharwad, Gadag, Haveri, Kalburgi, Koppal, Raichur, Vijayapura and Yadgir districts.

SOUTH INTERIOR KARNATAKA

Light to moderate rain likely to occur at a few places over Shivamogga, Chikkamagaluru, Hassan, Kodagu, Ballari, Bengaluru (Rural), Bengaluru (Urban), Chamarajanagara, Chikkaballapura, Chitradurga, Davanagere, Kolar, Mandya, Mysuru, Ramanagara, Tumakuru and Vijayanagara districts.

Extended Range forecast at Met-Subdivision Level (04.09.2026 to 10.09.2026):

Meteorological Sub-Division	Rainfall	Max Temp	Min Temp
Coastal Karnataka	Below Normal..	Normal.	Normal.
North Interior Karnataka	Below Normal.	Normal.	Normal.
South Interior Karnataka	Below Normal.	Normal.	Normal

Likely impacts of Heavy rainfall/Gusty wind (General)

Temporary disruption of electricity in some areas. Minor traffic snarls. Possible uprooting of weak tree branches

Visibility may become poor due to intense spell of rainfall leading to traffic congestion. •

Temporary Disruption of traffic due to water logging in roads/ uprooting of trees/ breaking of tree branches leading to increased travel time.

- Uprooting of trees may cause damages to power sector
- Possibility of damages to vulnerable structures due to heavy to very heavy rainfall
- Partial Damages to Kutcha Houses and Huts due to uprooting of trees.
- Possibilities of Flash floods due to intense spell of rainfall.
- Water logging / flooding in many parts of low-lying areas.
- Landslides/mud slide/land slip very likely over Coast and ghat areas.

Impact based advisories (General)

Stay indoors, close windows & doors and avoid travel if possible. Take safe shelters; Keep away from all the objects that conduct electricity. Drive carefully if you are travelling.

- Avoid staying in vulnerable structure.
- Avoid going to the areas that face water logging problems/ river fronts.
- People in the vulnerable area are advised to move to safer places.
- Be Updated

Agromet advisories

Drain out excess water from standing crops.

Cover the harvested produce (if in the field) with polythene sheet.

Avoid contact with electrical equipment or cords.

Keep away contact with anything metal - tractors, farm equipment and bicycles. As metal or other surfaces that conduct electricity.

Waterlogging may occur in standing crops and fields. Pest and diseases may occur.

Monitor for occurrence of pest and diseases and take plant protection measures after heavy rainfall.

Raised nursery beds for nursery sowing of rice may be prepared. Provide staking to avoid lodging of Banana plant.

Summary of Seven days rainfall over the State (28th August –03rd September 2026):

Meteorological Subdivision	28/08/2026	29/08/2026	30/08/2026	31/08/2026	01/09/2026	02/09/2026	03/09/2026
Coastal Karnataka	WS	FWS	FWS	FWS	FWS	WS	WS
North Interior Karnataka	ISOL	ISOL	ISOL	ISOL	ISOL	SCT	SCT
South Interior Karnataka	ISOL	ISOL	ISOL	ISOL	ISOL	SCT	SCT
	ISOL:	Isolated <i>i.e.</i> Rain at 1 or 2 places			SCT:	Scattered <i>i.e.</i> Rain at a few places	
	FWS:	Fairly Widespread <i>i.e.</i> Rain at many places					
	WS:	Widespread <i>i.e.</i> Rain at most places			DRY:	No rain	

Crop stages

Districts	Paddy	Sugarcane	Green gram	Red gram	Sorghum
Coastal Karnataka					
Dakshina Kannada	Ti				
Udupi	Ti				
Uttara Kannada	LP/S	V			
North Interior Karnataka					
Bagalkote		Gg	H	V	
Belagavi		D	H		
Bidar			F/Pf	V	
Dharwad		D	H		
Gadag		V	V	V	
Haveri		D	H		
Kalaburgi	T		D		
Koppal	N				
Raichur	T		D		
Vijayapura		D			
Yadgir	T		D		
South Interior Karnataka					
Ballari	Pi		M		
Vijayanagara	T		D		
Bengaluru Rural				V	

Bengaluru Urban				V	
Chamarajanagar	Ti/Pi/DT		V to F	V to F	
Chikaballapura				V	
Chikkamagaluru	T/SD				
Chitradurga			F/EPF	B to FI	
Davangere			F/EPF	B to FI	
Hassan	T/SD				
Kodagu	Pi to F				
Kolar				S	
Mandya	Ti/Pi/DT		V to F	V to F	
Mysuru	Ti/Pi/DT		V to F	V to F	
Ramanagara				V	
Shivamogga	TSD				
Tumakuru	ETi		F/EPF	B to FI	

N: Nursery, V : Vegetative, D: Development; S-Sowing; M-Maturity; H-Harvesting; Gg: Grand growth; G: Germination; Ti-Tillering; SD- Seedling development; EV- Early vegetative; T-Transplanting; growth-g; E-Emergence; F: Flowering; Pf: Pod formation; GF: grain filling; LP: Land Preparation; Pd: Pod development ; Td: Top dressing; Pm: Pod maturity; H: Harvesting; FI: Flower Initiation; SE: Stem Elongation; E: Establishment; PF: Pod Filling, P: Pegging; PS: Pre Sowing; P: Pegging ; NP: New Plantation; CD: Cane Development; PaF:Panicle formation; B: Branching; PH: Post Harvest; St: Storage; B/F : Boot /Flag leaf; Pi: Panicle initiation; Fa: Fallow; R:Ratooning; GM in: Green manure incorporation stage; ATi: Active tillering stage; DT: Delayed Transplanting; PS: Pod Setting ETi: Early Tillering; PM: Pod Maturity

Agromet Advisories issued by different Agromet Field Units (AMFUs)

Hill Zone

Agromet Advisory for Uttara Kannada district

Crop	Stage/ Condition	Pest and Disease	Agro advisories
Paddy	Land preparation nursery raising And Transpalnting	-	- As a basal dose apply N:P:K fertilizer @ 37.5:75:43.8 kg per ha respectively to Paddy before transplanting. - In paddy fields for the control of weeds apply butachlor 1lit / ha by mixing with 20 to 25 kg sand maintaining 1 cm water. After application see that the water should not go from one field to other field at least 24 hrs.

Arecanut,	Flowering	Mites	• It is the right time to clean and repair of the all drainage channels, as it should be completed before the start of the rain. • It is the right time to clean up the inter drainage channels so that rain water should flow easily. • This is the right time for transplanting of arecanut. Use healthy seedlings. • Raising of Arecanut seedlings can be continued. • To prevent the nut drop spray Carbendazim @ 1 g / lit. of water and Planofix @ 0.25 m.l./lit. of water. • For control of mites and spider, spray Dicofal @ 2.5 ml /Lit. of water or wettable sulphur @ 4 g / Lit. of water.
<u>Maize</u>	Vegetative stage		• Farmers are advised provide the drainage to remove the excess water from the field. • If the early sown maize leaves turn yellow or white, advise to take up the spray of water soluble fertilizer 19-19-19 @ 5 g/ lit. of water. • Intercultivation or Hand weeding can be done. • After intercultivation or hand weeding 50 Kg Nitrogen (110 kg Urea) per hectare should be given as atop dressing.
Black Pepper			• For control of katte roga as a precautionary measure drench copper sulphate solution @ of 2 to 4 lit per vine. (@ 3g/lit. of water). After that spray 1% Bordeaux mixture solution. • For control of katte roga oraganically use Trichoderma @ 50 g/vine mix it with neem cake and apply to the base of vine.
Ginger	Tube rot		• Sufficient drainage should be given. • For the control of tube rot in ginger and turmeric, treat the tubers with 3 gm Copper Oxychloride per litre of water.
Sugarcane			• Attend Intercultivation at 50, 65, 80 and 95 Days after planting and earthing up can be done at 120 days after planting. • For control of woolly aphid in sugarcane spray 1.0 gm Acepate 75 WP or 2.0 ml Chlorpyriphos 20 EC or Dimethoate 30 EC @1.7 ml per litre of Water
Cotton	Vegetative stage		• Right time for Intercultivation or hand weeding. • 25 kg Nitrogen (50 kg Urea) and 25 kg potash (30 kg MOP) per hectare should be given as atop dressing.

			White fly, Jassids and Aphids is there then go for spraying of 0.5 ml Imidacloprid 178 SL or 1.5 ml Oxydemeton Methyl per litre of water.
Veterinary and animal husbandry			
Animals			- Animals to be screened for nasal discharge, wound and injuries regularly and the services of veterinary doctor be taken for treatment - Animals to be dewormed with suitable anti-helmentic drug and be checked and treated for ecto-parasites, if any

Agromet Advisory for Dakshina Kannada / Udupi districts

Crop/enterprise	Crop Stage	Weather Risk	Agro Advisory
Paddy	Tillering Stage	• Moisture stress for mid and up land crop	• Conserve available soil moisture by strengthening field bunds and minimizing water losses. Provide life-saving irrigation, wherever feasible, particularly at this critical crop growth stage and avoid moisture stress for yield reduction.
		• Likelihood of caseworm and leaf folder infestation	• If case worm infestation is observed in paddy, spray Chlorantraniliprole @ 0.5 ml/litre or Chlorpyrifos @ 2 ml/litre or Neem oil (3000 ppm) @ 3 ml/litre of water. • If leaf folder infestation is observed in paddy, spray Chlorantraniliprole @ 0.5 ml/litre or Chlorpyrifos @ 2 ml/litre or Neem oil (3000 ppm) @ 3 ml/litre of water
		General Advisory • 1st Top Dressing (25-30 Days after transplating : Apply Niotrogen 6 kg / ac (i.e Urea : 13kg) and Apply Potassium 7.5 kg /ac (i.e Muriate of Potash : 15 kg) • 2nd Top Dressing (55-60 Days after transplating : Apply Niotrogen 6 kg / ac (i.e Urea : 13kg) and Apply Potassium 7.5 kg /ac (i.e Muriate of Potash : 15 kg)	
Horticultural crops			
Arecanut	Nut Formation	Probability of Koleroga infestation	Due to continous rainfall there may be chances of Kole roga so, farmers can spray 1%

			Bordo mixture in clear weather condition to avoid yield losses
		Excess moisture	Farmers can provide drainage facility to drain excess rain water
		Likelihood of Spindle bug and Pentatomid Bug	If spindle or Pentatomid bug infestation is there spray Imidacloprid 1ml/ltr or Thiomethaxome 0.5gm/ltr
		Likelihood of Root Grub	Drenching of Chloropyriphos 3-5ml/ltr (Minimum 5 litre of this solution should be drenched to palm)
		General Advisory: In Nurseries farmers can take up hand weeding. In younger gardens the seedlings should be given anchorage with sticks and provide drainage facilities. In older gardens : Sanitation must be maintained	
Coconut	Physiological Maturity	Excess moisture	Farmers can provide drainage facility to drain excess rain water
		Likelihood of Rhinoceros Beetle	• Collect and destroy the various stages of the beetle from the manure pits (breeding ground of the pest) whenever manure is lifted from the pits. • Incorporate the Entomopathogen i.e, fungus (Metarrhizium anisopliae) in manure pits to check the perpetuation of the pest. • Examine the crowns of tree at every harvest and hook out and kill the adults. • Apply mixture of either phorate 10 G + sand (1:2) @150 g per palm or phorate 10 G + sand (1:2) @150 g per palm in the base of the 3 inner most leaves in the crown. • Place Phorate 10 G 5gm in perforated sachets in two inner most leaf axils for 2 times at 6 months intervals. • Set up one Rhinolure pheromone trap for half hectare area to trap and kill the beetles.
Black Pepper	Formation of Berries	Excess moisture	Farmers can provide drainage facility to drain excess rain water

		Chance of Quick wilt	As a precautionary measure go for pruning of severely affected leaves and destroy them followed by application of FYM enriched with neem cake or Soil drenching with 3gram Copper Oxychloride or 1% Bordeaux mixture.
Cashew	Grand growth	Likelihood of T-Mosquito Bug	To manage T-Mosquito bug farmers can spray Monocrotophos @ of 1.7ml/ltr or Lamdacyalothrin @ 0.5ml/ltr.
		Likelihood of Stem Borer	Clean the infected place. After cleaning fill the infected part with Chloropyriphos granules mixed with sand(1:1 ratio) or Prepare a bordaeux paste and mix 5ml of Choloropyriphos per kg of paste and this paste can be used to fill the infected part
		General advisory	• New growth on planted saplings start in 2 to 3 weeks time. Protect the new growth from the attack of leaf eating caterpillar or beetles etc., by spraying Quinolphos @ 2 ml./lt. of water at every 15 to 20 days interval. • Weeding should be done at least once in a month during monsoon season to control weeds. Simultaneously the basins around the plant should be made to retain water. • Remove the side shoots which appear on the root stock.
Jasmine	Flowering stage	Likelihood of Leaf Spot	Farmers can spray Hexaconazole @ of 1ml/ltr or Copper Oxy-Chloride @ 3gm/ltr
Animal Husbandry	----	Wet weather	Ensure that the cattle sheds remains dry to avoid respiratory related diseases
		General Advisory General Advisory	Animals should be vaccinated against FMD, Enterotoximea, Black quarter if not done so far., Farmers can add 30-50 g m of mineral mixture to animal feed to get good milk yield Ecto Parasite: Keep the shed/area where animals are kept, clean. Bouquets of Basil, Neem or Lemon grass may be hung up in the animal

			shed, the scent of which keeps the ecto-parasites away Alternatively a lemon oil or Neem oil or Eucalyptus oil based disinfectant may also used for this purpose.
Fisheries	----	General Advisory	Farmers can make preparation of fish pond for fingerling production The pH of fish fry ponds can be adjusted to 7.5 by adding lime. Then, by adding manure, the natural fish food production can be done and when the pond water turns from light green to brown, freshwater fish fry can be stocked. These chicks will be available for sale from authorized suppliers in June and July.
Rain harvesting structure	-----	Shortage of water in upcoming days	Make necessary arrangements for established rain water harvesting structures as Monsoon is progressing. Ex: Farm pond: By storing excess water during rainy season may help farmers to get rid of water scarcity in summer & farmers can earn extra income by rearing fish.

Agromet Advisory for Bidar district

Crops	Stages	Pest/ disease	Agro advisory
Red gram	Vegetative		1. Farmers are advised to take up clipping activity (removal of apical portion of the plant) which helps in profuse lateral branching and also helps in increasing red gram yields, when the crop is at around 40-50 days old.
Green gram	Flowering / Pod formation	Aphids/thrips	Aphids menace in green gram crop can be managed by spraying of systemic insecticides such as dimethoate (1.7ml) or imidacloprid (0.3ml) or acephate (1 gm) or acetamiprid (0.2gm) in one litre of water. Provide proper drainage every 10 mtrs distance to discharge the excess water from the field due to rains.

			To overcome the nutrient deficiency in green gram, spray the crop with 19:19:19 NPK (10 gram/lit of water) as foliar spray followed by second spray at an interval of 15 days at the time of flowering along with 0.5ml planofix per litre of water which results higher yields.
Black gram	Flowering/ Pod formation	Aphids/thrips	Aphids menace in black gram crop can be managed by spraying of systemic insecticides such as dimethoate (1.7ml) or imidacloprid (0.3ml) or acephate (1 gm) or acetamiprid (0.2gm) in one litre of water. Provide proper drainage every 10 mtrs distance to discharge the excess water from the field due to rains. To overcome the nutrient deficiency in green gram, spray the crop with 19:19:19 NPK (10 gram/lit of water) as foliar spray followed by second spray at an interval of 15 days at the time of flowering along with 0.5ml planofix per litre of water which results higher yields.
Soybean	Flowering/ Pod formation		At 50% flowering, spray soybean crop with 2% urea solution. Provide proper drainage every 10 mtrs distance to discharge the excess water from the field due to rains. In 25 days old crop, spray the crop with Imezatpyr 10SL that is 1000ml required for 1 ha, to overcome the weed menace.
Mango	Newly planted	-	1. Intercultivation: In a newly planted mango orchard, growing in between the rows with horse gram, cowpea & sun hemp improves fertility status & also increases the yield levels. 2 Pruning: (In 5-6 years old mango orchard) 5-6 years old high density mango plantation, due to profused shoot growth, pruning is done for easy access of sunrays, & the pruned part are applied with fumigants plus insecticide. Which improves yield as well as quality of the fruits
Animal Husbandry/ Poultry	1. Animals must be vaccinated against Foot & Mouth disease, HS, Black Quarter & enterotoxaemia etc. 2. To protect animals from ectoparasites, there sheds should be kept clean, bouquets of nirgundi, basil leaves, or lemon grass must be hung in the cattle shed. Neem oil based disinfectants can also be used. 3. Vaccinate Goat/Sheep against PPR, Goat pox/sheep pox & Enterotoxaemia. 4. Provision of fresh drinking water as for as possible. 5. Feed the animals with mineral supplements based on the deficiency		

Agromet Advisory for Kalburgi/ Raichur/ Ballari/Vijayanagara/ Yadgiri districts

Crops	Stage	Probable pest/diseases	Agricultural Advisories
General			- Split application of fertilizer may be done for crops like cotton, pigeon pea, maize & other long duration crops. - Thinning should be done to maintain optimum plant population and soil moisture conservation.
Green gram	Maturity	Powdery mildew	When green gram /black gram reaches physiological maturity, spray the crop with Paraquat (5ml/L of water). It helps to easy pod picking by shredding of leaves,
Groundnut	Developmental stage	leaf spot and rust	Spray Chlorothalonil (Kavach) (2g/l) twice with an interval of 10 days for the management of leaf spot and rust diseases in groundnut.
Cotton	vegetative stage	thrips and leaf hopper	- to manage leaf reddening in cotton Spraying 19:19:19 @ 1% + MgSO₄@ 1% - Spray 250 ml Fipronil 5 SC in 250 L of water for one acre controlling thrips and green leaf hopper during early stage of cotton crop.
Pigeon pea	Developmental stage		Farmers are advised to go for nipping of 5-6 cm apical bud and spray 10 gm 19:19:19 during 50-55 days old crop for enhancing branches and improving yield in pigeon pea.
Paddy	Panicle initiation		Apply 10 kg N as a top dressing at 3rd and 6th weeks after transplanting and during panicle initiation stage.
Chilli	Flowering stage		Spray 5 ml Nephthalene Acetic Acid (NAA) in 100 liter of water for controlling flower dropping in chilli.
sheep and goat			Do not allow sheep and goat to graze on young misty grass in early morning hours. This may lead to enterotoxaemia (ET) in unprotected animals.
	Livestock		Feed the drumstick leaves to animals because it is having high protein and mineral content so it will enhance the immunity as well as milk yield and quality in livestock.

Agromet Advisory for Koppal district

Major crops	Stage	Pest/disease	Agricultural Advisories
General			❖ Spraying of 1% KNO₃ or 1% 19:19:19 or Nano Urea @ 4 ml/litre to overcome moisture stress in non-rainfall areas. ❖ Carryout repeated intercultivation in already sown crop to overcome evaporative loss of soil moisture and to manage weeds effectively. ❖ Soaking seeds in a 2% calcium chloride solution or 10% cow urine before sowing enables the seedlings to withstand drought conditions; however, the soaking duration varies for each crop ❖ Repeated intercultivation and weed management and also Crop thinning in rainfed areas to reduce moisture stress ❖ To adopt short duration/drought tolerant varieties ❖ Wherever its possible go for life saving irrigation from farm ponds ❖ Looking to the weather and soil type irrigate the crop once in 12-15 days in black soil and 8-10 days in red soil ❖ Take up the repairs of bunds and check dam so as reduce the runoff to check the erosion ❖ Farmers are advised to take up plant protection measures during morning hours by mixing with adhesive gum, mainly enables uniform pesticide spread over crop foliage and also enhances the activity of the pesticide
Paddy	Transplanting		❖ Apply 5-7 tonnes of FYM or 2.0 tonnes of poultry manure per hectare should be applied three weeks before transplanting of paddy ❖ Transplanting of paddy seedlings at the optimum stage, 20-25 days old seedlings for short and

			medium duration varieties and 30-35 days old seedlings for long duration varieties. ❖ Before transplanting of paddy seedlings dipped in <i>Azospirillum</i> slurry for 15-20 minutes after that go for transplanting and which will helps to reduce 25 per cent of recommended nitrogen. ❖ Transplanting of paddy is carried out in different places, so that maintain optimum irrigation to avoid transplanting shock. ❖ Transplant two-three seedlings per hill at 20 cm x 10 cm distance under normal conditions. Increase the number of seedlings per hill if old seedlings are used. ❖ In drum seeder sown paddy fields, carry out gap filling and first weeding. Maintain a 2–3 cm water layer or adopt Alternate Wetting and Drying (AWD) under limited water. ❖ Under water scarcity conditions sowing of DSR paddy with short to medium duration varieties.
Maize	Vegetative /flowering	Fall army worm	❖ In 20–25 DAS crop, spray 1% MOP (25–30 DAS) or 2% KCl during dry spells, provide protective irrigation at critical stages, and undertake shallow intercultivation with mulching. ❖ Top dressing of remaining 50% Nitrogen @ 50 kg N/ha in two splits after receipt of rainfall. ❖ Provide one life-saving irrigation during tasseling and silking wherever irrigation facilities are available. ❖ Spraying of Nano Urea @ 4 ml/litre or 19:19:19 @ 1% or KNO₃ @ 1% to overcome moisture stress in non-rainfall areas. ❖ For Broad leaved weeds in Maize, Spray of 2,4-D Sodium salt (0.5 kg/acre) at 20-25 DAS. ❖ For management of Fall army worm in Maize, spraying of Emamectin Benzoate @0.2 g per liter of water. ❖ For control of fall army worm, Install 10 pheromone traps per hectare

Redgram	Flowering		❖ Nipping in Pigeon pea at 50 days after sowing to enhance branches, pods per plant and yield. ❖ Dropping of flower in pigeon pea due to continuous cloudy condition go for spraying of NAA @ 0.5 ml/lit of water ❖ Spraying of pulse magic (10 g/lit of water) to pigeon pea at flowering stage to enhance yield of crop ❖ Thinning of excess seedlings to maintain optimum plant population. ❖ Carry out intercultivation and weeding for soil moisture conservation. ❖ Foliar spray of 2% DAP during flowering under moisture stress.
Groundnut	Vegetative	Thrips and Aphids	❖ For control of Thrips and Aphids, farmers can take up spraying of Dimethoate 30 EC per liter of water ❖ If possible, farmers can take up one hand weeding or intercultivation at 15-20 DAS for control of weeds and better aeration to crops ❖ If iron and zinc deficiency observed during flowering stage, take up spraying of 0.5 % Iron Sulphate or 0.2 per cent borax or 0.1% Ammonium molybdate. Repeat the same spray at 10 days interval ❖ Apply gypsum at 200 kg per acre at 35 to 45 DAS on either side of the crop rows during last intercultural activity.
Cotton	Flowering	Thrips and Aphids leaf reddening	❖ Spray Nano Urea @ 4 ml/litre or 1% MgSO₄ + 1% 19:19:19 under moisture stress. ❖ Fertilizer application: Apply 25% of N, K and 100% P at the time sowing and remaining 75% of N and K apply @ 30, 60 & 90 DAS ❖ For control of Thrips and Aphids, farmers can take up spraying based on ETL. Fipronil 5 SC @ 1.0 ml or Thiamethoxam 25 WG @0.2 g in one liter of water. ❖ Management of leaf reddening in cotton, spraying of Magnesium sulphate @ 1.0% along with 2 % urea during 70 and 90 DAS.

Mango	Harvesting	Fruit fly	❖ For control of fruit fly: 10 Pheromone traps per hectare. Chemical used for Pheromone trap (1 ml methyl ugenol and 1 ml Melothion 50 EC per litre of water
Guava	Flowering /fruiting	Kujji Bug	❖ For management of Kujji Bug, spraying of Lamda cyhalothrin @ 0.5 ml per liter of water ❖ After 3-4 years of transplanting, intercrops like vegetables and pulses can be cultivated
Pomogranate	Harvesting	Sucking pests Bacteria l blight	❖ To manage sucking pests, take up spraying of 0.3 ml Imidacloprid or 0.25 ml Thiamethaxam dissolved in one litre of water ❖ To manage bacterial blight disease, take up spraying of Streptomycin Sulphate 0.5 g. + Copper oxychloride 2 g. dissolved in one liter water. ❖ In Pomegranate, intercrops like Cowpea, cucumber, Onion, Horsegram can be cultivated upto the year of 5-6 years.
Poultry	❖ Maintain the ambient room temperature of the poultry house. ❖ Poultry litter should be dried to avoid fly menace		
Animals (Cows & Buffalo)	❖ Proper ventilation should be maintained for free circulation of air in the sheds. ❖ For Milch animals fodder should maintain in the ratio of 1:3 (leguminous:non-legumious fodder) to get good quantity and quality of milk. ❖ Green fodder can be conserved and minimizing wastage (Silage making). It will enhance livestock productivity by ensuring fodder supply especially during the lean period. ❖ In case of irrigation facilities available go for growing of fodder crops like maize, jawar and green fodder especially during the lean period for higher milk production.		

Agromet Advisory for Gadag district

Crop	Stage	Management of pest /Disease
Sugarcane	Cane development stage	Top dressing of Nitrogenous and Potash fertilizers. To manage root grub menace application/spreading of 10 kg of Metarhizium anisopliae bio-insecticide mixing in dry organic manure or vermicompost. To manage early shoot borer : Spraying of Imidacloprid @ 0.5 ml /lit of water To manage scale insect spraying of Acephate 1 ml / lit of water
Pigeonpea	Vegetative stage	To manage Ash weevil : Spraying of 2 ml Quinalphos @ 2 ml/liter of water. To manage root grub menace application/spreading of 10 Kg of Metarhizium anisopliae bio-insecticide mixing in dry organic manure or vermicompost To manage leaf webber: Spraying of Propenophos @ 1 ml/liter of water. Nipping: 45-50 days after sowing nipping of terminal buds up to 5-6 cm will helps to get more branches and achieve good yield.
Groundnut	Vegetative stage to flowering stage	To manage leaf minor in groundnut, spraying Quinalphos 25 EC @ 2 ml or Dimethoate 30 EC @ 2 ml/lit.
Maize	Vegetative stage	Fall army worm (> 20% damaged plants): Spray Spinetoram 11.7 % SC @ 0.5 ml/l or Chlorantraniliprole 18.5 @ 0.3 ml/lit of water or Thiamethoxam 12.6 % + Lambda cyhalothrin 9.5% ZC @ 0.25 ml/l. Use high volume sprayer, the nozzle directed towards the whorls. To manage spotted stem borer : Take up spraying of Chlorantraniliprole @ 0.2ml/litre
Sunflower	Before sowing	To manage necrosis virus before sowing treat the seeds with 5 gm Imidacloprid 70 WS in 1 kg of seeds.
Onion	Vegetative stage	To manage onion bulb rot : Take up spraying of carbendazim @ 1 gm/lit of water.

		To manage thrips, whitefly (upside Leaf curling) take up spraying of 0.5 ml Imidacloprid 17.8 % SL dissolved in one liter of water at seedling stage & 19:19:19 @ 1 g/litre of water.
Chilli	Vegetative stage	To manage thrips, whitefly (upside leaf curling) : Take up spraying of 0.5 ml Imidacloprid 17.8 % SL dissolved in one litre of water at seedling stage.
Grapes	Under pruning/vegetative	Due to high cloudy condition and rainfall, there is chance of occurrence of Anthracnose disease . To manage Anthracnose disease, spraying of Thiophanate methyl @ 1 gram/liter of water. To manage mealy bug take up spraying of Imidacloprid 17.8 S.L. @ 0.4 ml/litre of water
Citrus	Fruit development to harvest	To control citrus canker disease cut the affected parts then spray 3gm copper oxychloride 50 W.P and 0.5 g streptomycin sulphate or 1% Boardo mixture in every liter of water.
Pomogranate	Fruit development to harvest	To manage Bacterial blight disease in pomogranate, spray the crop with Streptocyclin sulphate 0.5 g. + Copper Oxychloride @ 2 g / liter of water. To manage sucking pests spraying of Imidacloprid @ 0.3 ml or 0.25 g Thiamethoxam / liter of water.
Livestock management	Deworming should be done in all livestock regularly once in six months As a preventive measure of Lumpy skin disease (LSD), keep the livestock shed clean and take control measures for flies and ticks in order to control spreading of LSD. For every milch animals regularly follow schedule of 1 kg concentrated feed per 2 litres of milk yield Vaccinate sheep and goat against blue tongue and Haemorrhagic septicemia disease.	

Agromet Advisory for Vijayapura district

Crop	Stage	Management of pest /Disease
Sugarcane	Cane development stage	• Top dressing of Nitrogenous and Potash fertilizers. • To manage root grub menace application/spreading of 10 kg of Metarhizium anisopliae bio-insecticide mixing in dry organic manure or vermicompost. • To manage early shoot borer : Spraying of Imidacloprid @ 0.5 ml /lit of water • To manage scale insect spraying of Acephate 1 ml / lit of water
Pigeonpea	Vegetative stage	• To manage ash weevil : Spraying of 2 ml Quinalphos @ 2 ml/liter of water. • To manage root grub menace application/spreading of 10 Kg of Metarhizium anisopliae bio-insecticide

		mixing in dry organic manure or vermicompost - To manage leaf webber: Spraying of Propanophos @ 1 ml/liter of water. - Nipping: 45-50 days after sowing nipping of terminal buds up to 5-6 cm will helps to get more branches and achieve good yield.
Groundnut	Vegetative stage to flowering stage	To manage leaf minor in groundnut, spraying Quinalphos 25 EC @ 2 ml or Dimethoate 30 EC @ 2 ml/lit.
Maize	Vegetative stage	- Fall army worm (> 20% damaged plants): Spray Spinetoram 11.7 % SC @ 0.5 ml/l or Chlorantraniliprole 18.5 @ 0.3 ml/lit of water or Thiamethoxam 12.6 % + Lambda cyhalothrin 9.5% ZC @ 0.25 ml/l. Use high volume sprayer, the nozzle directed towards the whorls. - To manage spotted stem borer : Take up spraying of Chlorantraniliprole @ 0.2ml/litre
Sunflower	Before sowing	To manage necrosis virus before sowing treat the seeds with 5 gm Imidacloprid 70 WS in 1 kg of seeds.
Onion	Vegetative stage	- To manage onion bulb rot : Take up spraying of carbendazim @ 1 gm/lit of water. - To manage thrips, whitefly (upside Leaf curling) take up spraying of 0.5 ml Imidacloprid 17.8 % SL dissolved in one liter of water at seedling stage & 19:19:19 @ 1 g/litre of water.
Chilli	Vegetative stage	To manage thrips, whitefly (upside leaf curling) : Take up spraying of 0.5 ml Imidacloprid 17.8 % SL dissolved in one litre of water at seedling stage.
Grapes	Under pruning/vegetative	- Due to high cloudy condition and rainfall, there is chance of occurrence of Anthracnose disease . To manage Anthracnose disease, spraying of Thiophanate methyl @ 1 gram/liter of water. - To manage mealy bug take up spraying of Imidacloprid 17.8 S.L. @ 0.4 ml/litre of water
Citrus	Fruit development to harvest	To control citrus canker disease cut the affected parts then spray 3gm copper oxychloride 50 W.P and 0.5 g streptomycin sulphate or 1% Boardo mixture in every liter of water.
Pomogranate	Flowering to fruit development	During flowering to fruit setting stage Apply 25-30 kg Farm Yard Manure (FYM) per

		plant OR 15-20 kg FYM + 2 kg vermi-compost + 2 kg neem-cake per plant OR 7.5 kg well decomposed poultry manures + 2 kg neem-cake per plant. - To manage Bacterial blight disease in pomogranate, spray the crop with Streptocyclin sulphate 0.5 g. + Copper Oxychloride @ 2 g / liter of water. - To manage sucking pests spraying of Imidacloprid @ 0.3 ml or 0.25 g Thiamethoxam / liter of water.
Livestock management	- Deworming should be done in all livestock regularly once in six months - As a preventive measure of Lumpy skin disease (LSD), keep the livestock shed clean and take control measures for flies and ticks in order to control spreading of LSD. - For every milch animals regularly follow schedule of 1 kg concentrated feed per 2 litres of milk yield - Vaccinate sheep and goat against blue tongue and Haemorrhagic septicemia disease.	

Agromet Advisory for Bagalkote district

Crop	Stage of crop	Pest & Disease	Agro-Advisories
Sugarcane	Grand growth	Striga and Rootgrub	➤ To manage root grub infestation, mix 5-10 kg Metarhizium anisopliae in 500 kg of Compost fertilizer for one acre. It will help to control early stage of root grub infestation. ➤ Spread the trash on field every line after line for better moisture conservation. ➤ To manage leaf reddening, take up spraying of complex fertilizer 19:19:0 at the rate of 5 to 10 gram dissolved in one liter of water. ➤ Tie up the leaves of sugarcane (Propping) to withstand high wind speed effect in field. ➤ To manage striga weed, irrigate the crop, spread sugarcane trash in between the rows and take up spraying of 1 kg of 2,4 D dissolved in 500 liters on the weed.
Greengram	Harvest	-	➤ Harvest early sown mature crop during early morning hours.

Pigeon pea	Vegetative	-	➤ For efficient nutrient uptake and for better crop growth, 35-45 days after sowing apply 3 kg/ha Zinc Sulphate with 6 t FYM on the soil. ➤ In Pigeon pea 45 to 50 days after sowing carryout nipping operation (Pinching/Removal of terminal buds up to 5-6 cm) that will helps to get more side branches and helps to achieve higher pod set and good yield.
Soybean	flowering	Leaf eating Caterpillar	➤ For the control of leaf eating caterpillars in soybean spray Chlorpyriphos 20 EC @ 2 ml or Alphamethrin @ 1 ml or Diflubenzuron 25 WP @ 0.6 g or Flubendiamide 20% WG @ 0.5 ml per liter of water. ➤ To overcome micronutrient deficiency takes up spray of 19:19:19 to crop. ➤ In already sown crop take up one intercultivation operation to remove weeds and and to maintain optimum soil moisture condition.
Maize	flowering	Fall army worm	➤ In already sown crop, carry out one intercultivation operation to control weeds and maintain optimum soil moisture. ➤ In early sown Maize crop is affected with fall army worm to control eggs/larvae spray neem seed kernel extract 5 ml/ lit of water. ➤ Take up spray of Chlorantraniliprole 18.5 SC 0.4ml/lit or Emamectin benzoate 5 SG 0.4 g/lit.
Sunflower	Vegetative	Alternaria blight, Necrosis	➤ To manage Alternaria leaf blight spray 2g of zineb 80 W.P or 2 gm Mancozeb 75 W.P or 1 ml Hexaconazole in every liter of water. ➤ To manage necrosis virus disease, treat the seeds with imidacloprid 70 W.S/ KG of seeds. ➤ Destroy the disease affected plants. ➤ After 30 days after sowing spray Imidacloprid 17.8 S. L in every liter of water. ➤ Destroy weeds like Parthenium and Xanthium.

HORTICULTURAL CROPS :

Grape	Vegetative to Fruit bud differentiation stage	Anthracnose and Mealy bug	➤ Possibility of leaf curling, check the leaf margins, if slight to more yellow, possibility of potassium deficiency. Foliar spray of SOP @ 3-4g/L followed by fertigation of 20-25 kg SOP/acre in 2 to 3 splits. ➤ During humid conditions take up spray of Bio-agents, it will help to control in the incidence of disease to vines. ➤ Spraying of Imidacloprid 17.8 S.L @ 0.4ml/lit of water will help in controlling mealy bug infestation on new shoot growth. ➤ Due to humid and cloudy condition, one preventive spray of Thiophenate methyl/carbendazim@1g/L should be given for anthracnose control.
Chilli	Vegetative	Sucking pest	➤ To control Sucking pests like thrips after three weeks of sowing spray 1.7 ml. Dimethoate 30 E.C or 0.5 ml or 1 ml. Methyl Parathion 50 E.C in every liter of water.
Onion	Vegetative	Spodoptera, Thrips	➤ To control spodoptera litura apply chlorantraniliprole 18.5 SC @ 0.15 ml or Spinosad 45 SC @ 0.12 ml or chlorpyrifos 20 EC @ 2 ml per liter. ➤ To control thrips spray 0.3 g thiamethoxam 25 WP or 0.3 ml imidacloprid 17.8 SL or 1.7 ml Dimethoate 30 EC or fipronil 5 SL @ 1 ml per liter of water.
Citrus	Fruiting	Citrus canker	➤ To manage Bacterial blight disease, spray the crop with 3 gram Copper Oxychloride or 1 per cent Bordo mixture dissolved in one liter of water or 300 ppm Streptomycin 3 gram in 10 liters of water. ➤ To control citrus canker disease cut the affected parts then spray 3gm copper oxychloride 50 W.P and 0.5 gm streptomycin sulphate or 1% Boardo mixture in every liter of water.

Pomegranate	Harvest	weeds	➤ Harvest Matured Fruits. ➤ Plant not under proper stress due to intermittent rains or other reasons, take 2 to 3 sprays of Ethephon 39% SL, first spray @ 0.5 ml/lit followed by second spray after 5-8 days with @ 0.5 ml/lit depending on yellowing of leaves third spray can be taken after 7 days of 2nd spray with Ethephon @ 1 to 1.5 ml/L. Mix 18:46:00 or 12:61:00 or 00:52:34 @ 5 g/L with each Ethephon spray for more efficiency. ➤ Plants under proper stress with yellow leaves, take spray of Ethephon 39 % SL @ 1 ml /lit + 18:46:00 or 12:61:00 @ 5 g/L. ➤ Remove all weeds surrounding the plant and mulch the soil with crop debris to conserve the soil moisture.
--------------------	----------------	--------------	--

Livestock management	➤ Livestock animals should be vaccinated for Hemorrhage Septecemia (HS), Black quarter (BQ) and foot and mouth disease. ➤ For milch animals regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 litre of milk yield. ➤ Farmers are advised to take spray on animals frequently for removal of ectoparasites. And cut all bushes near their sheds. ➤ Vaccinate sheep/goat for blue tongue, enterotoxaemia and Haemorrhagic Septicemia diseases. ➤ Vaccinate chickens for every six months to avoid incidence of Stork disease.
-----------------------------	---

North Transition Zone

Agromet Advisory for Dharwad / Haveri/ Belagavi districts

Crop	Stage	Pest & disease	Agro advisories
Green gram	Harvesting	Yellow mosaic:	• Yellow mosaic: Uproot and destroy the affected plant, to reduce further spread spray the crop with 5.0 g 13:0:45 potassium nitrate & 0.4 g Thiamethoxam per litre of water. • Crop is at harvesting stage harvesting may be attended with rainfall precautions.

Black gram	Maturity to harvesting	Yellow mosaic:	- Yellow mosaic: Uproot and destroy the affected plant, to reduce further spread spray the crop with 5.0 g 13:0:45 potassium nitrate & 0.4 g Thiamethoxam per litre of water. - Early sown crop is at harvesting stage harvesting may attended with rainfall precautions.
Soybean	Grain filling	Pod borer	- To manage stem fly in soybean spray Thiamethoxam 25 WG @ 0.5 g/lit OR (Thiamethoxam 12.6% + Lambda-cyhalothrin 9.5% ZC) @ 0.5 ml per litre of water. - For control of defoliators in Soybean Spray 0.6 g Diflubenzuron 25% WP or 0.5 ml Flubendiamide 20% WG per litre of water. - For control of Pod borer in Soybean Spray: 0.2 ml Chlorantraniliprole 18.5% SC OR Chlorantraniliprole + Lambda-cyhalothrin 0.5 ml OR Flubendiamide 0.5 ml per litre of water.
Ground nut	Flowering to pod initiation		- advised to attend inter cultivation in 25-30 days crop to control the weeds and conserve the soil moisture in rain fed situation. - In Groundnut apply 200 kg of Gypsum per acre when crop is 35-40 days old crop. - Spry 2% urea (20 g urea in one litre of water) at 30 and 60 days after sowing.
Maize	vegetative (4-5 leaf stage)	fall armyworm	- advised to attend inter cultivation in 25-30 days crop to control the weeds and conserve the soil moisture in rain fed situation. - Top dressing of nutrients and inter cultivation may be attended at 30-35 days crop (4-6 week stage), looking to the soil moisture condition. - For the control of fall armyworm spray 0.3 g Imamectine benzoate 5% SG (Proclame)/ Chlorantraniliprole 18.5 SC per litre of water.
Cotton	Development		- Sowing May be taken up where moisture is favorable and irrigation for sowing desi cotton. - Treat the Seeds with Seed polymer @ 8 g/kg and micronutrients like Copper sulphate, Magnesium Sulphate, Iron Sulphate @ 4 gm per kg of Seed. - pre emergent herbicide spry: on the same day or next day after sowing Spray 500 g Diuron 80 WP or 2.0 litre Pendimethlin 30 EC mix in 300 litre of water spray over one acre area.

			• Top dressing of fertilizer may be attended in 30-35 days crop looking to the soil moisture condition.
Sugarcane	Development	-	• Sugarcane development stages (101-270 days) irrigate the crop at 7 days interval. • Attend Intercultivation at 50, 65, 80 and 95 Days after planting and urthing up can be done at 120 days after planting. • Management of rust in Sugarcane: spray Hexaconazole/Propiconazole @1.0 ml per litre of water.
Horticulture/Vegetable Crops			•
Chilli/ Tomato Brinjol	Flowering	Sucking pest	<i>Pest and Diseases management</i> • Looking to the soil moisture condition provide light irrigation to vegetable crops. • For control of flower dropping in brinjol spray planofix 1ml/4liter of water + boran 1g/litre of water. • looking to the weather and soil type irrigate the crop once in 15 days for black soil and 8-10days for red soils. • To manage leaf curl disease in tomato take up spraying of 1 ml Acetamaprid dissolved in 4 liter of water. • Under irrigated situation chilli crop can be sown during January - February.
Veterinary			
Animals			• For milch animals regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 litres of milk yield. • Feed animals with a mixture of green grass + hay + minerals + dry feed like a khichri. • Livestock owners are advised to vaccinate the animals against the foot & mouth (FMD) and Lumpy Skin Disease (LSD). • It is advised to provide de worming medicine by consulting nearest veterinary hospital. • Provide pure drinking water during day time and Keep the animal shed clean and dry.

Agromet Advisory for Chitradurga/Davangere districts

Crops	Stage / Condition	Agro advisories
Maize	Grand Growth to Tassel Initiation / Pre-Tasseling	➤ Utilize expected light rainfall and avoid irrigation immediately before or after showers. ➤ Maintain adequate soil moisture because tassel initiation is a critical water-demand period ➤ Complete intercultivation and earthing-up where required without damaging roots. ➤ Apply remaining nitrogen top dressing only when soil has adequate moisture; avoid excess nitrogen. ➤ Maintain drainage in low-lying fields and monitor for lodging after wind-associated rain.
Maize Pest and Disease:		➤ Fall Armyworm: Monitor whorls and tassels; destroy egg masses and early larvae, and use neem oil/azadirachtin or biological control where appropriate. ➤ Turcicum leaf blight: Inspect lower leaves under humid weather; where disease is established, mancozeb or propiconazole may be used as per approved label recommendation. ➤ Apply chlorantraniliprole or another recommended insecticide for FAW only when infestation crosses the economic threshold, and only during a rain-free, low-wind period. Conserve natural enemies and use neem-based/IPM measures first. Apply insecticide/fungicide only during a clear rain-free, low-wind period when required.
Ragi	Active Tillering to Panicle Initiation	➤ Use light rainfall efficiently and avoid unnecessary irrigation. ➤ Complete intercultivation and weed management before canopy closure. ➤ Maintain adequate moisture during active tillering and panicle initiation. ➤ Apply recommended nitrogen top dressing when soil moisture is adequate. ➤ Avoid water stagnation in heavier soils.
Ragi Pest and Disease:		➤ Blast: Regularly inspect leaves and collar region under cloudy, humid weather; maintain balanced nitrogen management. ➤ For confirmed blast, use tricyclazole or mancozeb as per approved recommendation and only during dry weather.

		➤ Monitor leaf-feeding caterpillars and sucking pests; conserve natural enemies and use neem oil/azadirachtin first, with chlorantraniliprole used only when necessary and during a rain-free window.
Redgram	Branching to Flowering Initiation	➤ Utilize available rainfall for moisture conservation and avoid irrigation immediately following showers. ➤ Complete intercultivation and weed control without damaging feeder roots; maintain a clean field. ➤ Maintain good drainage and avoid prolonged water stagnation around the root zone. ➤ Apply recommended micronutrients only where deficiency symptoms are observed; avoid excess nitrogen near flowering.
Redgram Pest and Disease:		➤ Pod borer: Start close monitoring of flowers, eggs and early larvae; install pheromone traps and conserve parasitoids. ➤ For significant pod-borer infestation, emamectin benzoate or chlorantraniliprole may be used as per label recommendation during clear, calm weather. ➤ Monitor leaf webber and defoliators; use neem oil/azadirachtin or suitable biological control first and avoid unnecessary broad-spectrum insecticides.
Green gram	Flowering to Early Pod Formation	➤ Maintain uniform soil moisture during flowering and pod setting; avoid irrigation immediately after light rainfall. ➤ Keep fields weed-free and avoid heavy intercultivation that may damage the root zone at this stage. ➤ Remove badly diseased or virus-affected plants and maintain good field sanitation. ➤ Apply foliar nutrient correction only where deficiency is evident and weather remains dry and calm.
Green gram Pest and Disease:		➤ Whitefly/Yellow Mosaic Virus: Monitor the crop frequently; remove severely affected plants and control vector populations using neem oil/azadirachtin where appropriate. ➤ Pod borer: Inspect flowers and young pods; install pheromone traps and use emamectin benzoate when economic threshold is reached. ➤ Under humid weather, monitor Cercospora leaf spot/powdery mildew; use mancozeb or sulfur only when disease is confirmed and during a dry-weather interval.

Sunflower	Bud Initiation to Flowering	➤ Use forecast rainfall efficiently and avoid excess irrigation; maintain adequate soil moisture during bud initiation and flowering. ➤ Complete essential weed management without disturbing the root zone. ➤ Avoid unnecessary late nitrogen application; maintain balanced nutrition for proper head development. ➤ Ensure proper drainage after showers to prevent prolonged root-zone saturation.
Sunflower Pest and Disease:		➤ Monitor developing heads and foliage for caterpillars/head feeders; remove early infestations and conserve natural enemies. ➤ Where caterpillar infestation exceeds threshold, emamectin benzoate or chlorantraniliprole may be used as per approved recommendation in clear weather. ➤ Monitor Alternaria leaf spot under humid conditions; where necessary use mancozeb or propiconazole during a rain-free period.
Arecanut	Inflorescence & Nut Formation Stage	➤ Maintain effective drainage around palm basins after intermittent rainfall and avoid prolonged water stagnation. ➤ Retain organic mulch to conserve moisture during dry intervals and remove weeds regularly. ➤ Avoid fertilizer application immediately before rainfall; apply nutrients only when soil is moist but not waterlogged. ➤ Remove fallen and diseased nuts and maintain clean orchard conditions.
Arecanut Pest and Disease:		➤ Mahali/Fruit rot: Remove infected nuts promptly and maintain sanitation; under confirmed disease, copper oxychloride can be used during a suitable dry interval as per recommendation. ➤ Spindle bug: Inspect emerging spindle leaves and use neem-based management where infestation is low; use a recommended insecticide only when economically justified. ➤ Bud rot: Inspect crown region after humid periods; where required, mancozeb or copper oxychloride may be used during dry weather as per approved recommendation.
Coconut	Post-Nut Setting to Pre-Harvest	➤ Maintain good drainage around basins while conserving sufficient soil moisture during dry intervals. ➤ Remove dried fronds, dead material and weeds to improve orchard sanitation and airflow. ➤ Avoid water stagnation around the root zone and irrigate only according to actual soil moisture requirement.

		➤ Apply organic manure/compost when soil is moist but not waterlogged and avoid application immediately before rainfall.
Coconut Pest and Disease:		➤ Rhinoceros beetle: Maintain crown sanitation and pheromone-trap monitoring; use neem cake/biological measures where appropriate. ➤ Black-headed caterpillar: Monitor leaflets and conserve parasitoids; where economically required, use azadirachtin or a recommended insecticide during dry weather. ➤ Bud rot: Inspect spindle/crown region under prolonged humidity; where disease is confirmed, copper oxychloride may be used according to approved recommendation and only in a dry period.
Pomegranate	Fruit Development	➤ Maintain uniform soil moisture and avoid excess irrigation after light rainfall to reduce fruit cracking and root-zone problems. ➤ Improve canopy ventilation through removal of excessively dense or diseased growth without heavy pruning. ➤ Remove cracked, diseased and damaged fruits immediately and maintain orchard sanitation. ➤ Maintain good drainage and use organic mulch to reduce rapid soil-moisture fluctuations.
Pomegranate Pest and Disease:		➤ Bacterial blight: Remove infected plant parts and affected fruits promptly; where recommended, use copper oxychloride during dry weather with proper coverage. ➤ Anthracnose/fruit spot: Remove infected tissues and monitor closely after humid weather; mancozeb may be used when required during a rain-free window. ➤ Fruit borer: Install pheromone traps and collect damaged fruits; where ETL is reached, use emamectin benzoate during clear, calm weather.
Livestock Management		
➤ Keep animal sheds, calf pens and poultry houses clean, dry and well ventilated; remove dung and stagnant water regularly after rainfall. ➤ Ensure continuous availability of clean drinking water, green fodder, dry fodder, mineral mixture and common salt; avoid feeding wet or mouldy fodder. ➤ Avoid grazing animals in waterlogged or very muddy areas immediately after rainfall and provide a dry resting place to prevent hoof infections and parasitic problems. ➤ Monitor livestock for FMD, Haemorrhagic Septicaemia (HS), Black Quarter (BQ), mastitis and parasitic infections; isolate suspected animals and consult a veterinarian promptly. ➤ Protect calves, pregnant animals and poultry from prolonged wet and humid conditions and provide adequate ventilation.		

- Follow the recommended **vaccination and deworming schedule** and avoid indiscriminate use of antibiotics or other veterinary medicines.

Agromet Advisory for Tumakuru

Crops	Stage / Condition	Agro advisories
Maize	Grand Growth to Tassel Initiation / Pre-Tasseling	➤ Utilize expected light rainfall and avoid irrigation immediately before or after showers. ➤ Maintain adequate soil moisture because tassel initiation is a critical water-demand period ➤ Complete intercultivation and earthing-up where required without damaging roots. ➤ Apply remaining nitrogen top dressing only when soil has adequate moisture; avoid excess nitrogen. ➤ Maintain drainage in low-lying fields and monitor for lodging after wind-associated rain.
Maize Pest and Disease:		➤ Fall Armyworm: Monitor whorls and tassels; destroy egg masses and early larvae, and use neem oil/azadirachtin or biological control where appropriate. ➤ Turcicum leaf blight: Inspect lower leaves under humid weather; where disease is established, mancozeb or propiconazole may be used as per approved label recommendation. ➤ Apply chlorantraniliprole or another recommended insecticide for FAW only when infestation crosses the economic threshold, and only during a rain-free, low-wind period. Conserve natural enemies and use neem-based/IPM measures first. Apply insecticide/fungicide only during a clear rain-free, low-wind period when required.
Ragi	Active Tillering to Panicle Initiation	➤ Use light rainfall efficiently and avoid unnecessary irrigation. ➤ Complete intercultivation and weed management before canopy closure. ➤ Maintain adequate moisture during active tillering and panicle initiation. ➤ Apply recommended nitrogen top dressing when soil moisture is adequate. ➤ Avoid water stagnation in heavier soils.
Ragi Pest and Disease:		➤ Blast: Regularly inspect leaves and collar region under cloudy, humid weather; maintain balanced nitrogen management.

		➤ For confirmed blast, use tricyclazole or mancozeb as per approved recommendation and only during dry weather. ➤ Monitor leaf-feeding caterpillars and sucking pests; conserve natural enemies and use neem oil/azadirachtin first, with chlorantraniliprole used only when necessary and during a rain-free window.
Redgram	Branching to Flowering Initiation	➤ Utilize available rainfall for moisture conservation and avoid irrigation immediately following showers. ➤ Complete intercultivation and weed control without damaging feeder roots; maintain a clean field. ➤ Maintain good drainage and avoid prolonged water stagnation around the root zone. ➤ Apply recommended micronutrients only where deficiency symptoms are observed; avoid excess nitrogen near flowering.
Redgram Pest and Disease:		➤ Pod borer: Start close monitoring of flowers, eggs and early larvae; install pheromone traps and conserve parasitoids. ➤ For significant pod-borer infestation, emamectin benzoate or chlorantraniliprole may be used as per label recommendation during clear, calm weather. ➤ Monitor leaf webber and defoliators; use neem oil/azadirachtin or suitable biological control first and avoid unnecessary broad-spectrum insecticides.
Green gram	Flowering to Early Pod Formation	➤ Maintain uniform soil moisture during flowering and pod setting; avoid irrigation immediately after light rainfall. ➤ Keep fields weed-free and avoid heavy intercultivation that may damage the root zone at this stage. ➤ Remove badly diseased or virus-affected plants and maintain good field sanitation. ➤ Apply foliar nutrient correction only where deficiency is evident and weather remains dry and calm.
Green gram Pest and Disease:		➤ Whitefly/Yellow Mosaic Virus: Monitor the crop frequently; remove severely affected plants and control vector populations using neem oil/azadirachtin where appropriate. ➤ Pod borer: Inspect flowers and young pods; install pheromone traps and use emamectin benzoate when economic threshold is reached. ➤ Under humid weather, monitor Cercospora leaf spot/powdery mildew; use mancozeb or sulfur only

		when disease is confirmed and during a dry-weather interval.
Sunflower	Bud Initiation to Flowering	➤ Use forecast rainfall efficiently and avoid excess irrigation; maintain adequate soil moisture during bud initiation and flowering. ➤ Complete essential weed management without disturbing the root zone. ➤ Avoid unnecessary late nitrogen application; maintain balanced nutrition for proper head development. ➤ Ensure proper drainage after showers to prevent prolonged root-zone saturation.
Sunflower Pest and Disease:		➤ Monitor developing heads and foliage for caterpillars/head feeders; remove early infestations and conserve natural enemies. ➤ Where caterpillar infestation exceeds threshold, emamectin benzoate or chlorantraniliprole may be used as per approved recommendation in clear weather. ➤ Monitor Alternaria leaf spot under humid conditions; where necessary use mancozeb or propiconazole during a rain-free period.
Paddy	Early Tillering	➤ Complete transplanting where adequate soil moisture is available and avoid unnecessary delay under suitable field conditions. ➤ Maintain shallow standing water after establishment rather than excessive flooding; provide drainage following rainfall. ➤ Complete gap filling and early weed management; avoid prolonged deep water stagnation. ➤ Apply the first nitrogen top dressing after establishment when soil moisture is suitable and avoid excess nitrogen under humid weather.
Paddy Pest and Disease:		➤ Leaf folder: Monitor folded leaves and early infestation; conserve natural enemies and avoid unnecessary spraying. ➤ Blast: Inspect leaves and collar region during humid/cloudy weather; maintain balanced nitrogen and proper water management. ➤ Where economic threshold is reached, use chlorantraniliprole for leaf folder or tricyclazole for blast according to approved label recommendations, only during rain-free weather.
Arecanut	Inflorescence & Nut	➤ Maintain effective drainage around palm basins after intermittent rainfall and avoid prolonged water stagnation.

	Formation Stage	➤ Retain organic mulch to conserve moisture during dry intervals and remove weeds regularly. ➤ Avoid fertilizer application immediately before rainfall; apply nutrients only when soil is moist but not waterlogged. ➤ Remove fallen and diseased nuts and maintain clean orchard conditions.
Arecanut Pest and Disease:		➤ Mahali/Fruit rot: Remove infected nuts promptly and maintain sanitation; under confirmed disease, copper oxychloride can be used during a suitable dry interval as per recommendation. ➤ Spindle bug: Inspect emerging spindle leaves and use neem-based management where infestation is low; use a recommended insecticide only when economically justified. ➤ Bud rot: Inspect crown region after humid periods; where required, mancozeb or copper oxychloride may be used during dry weather as per approved recommendation.
Coconut	Post-Nut Setting to Pre-Harvest	➤ Maintain good drainage around basins while conserving sufficient soil moisture during dry intervals. ➤ Remove dried fronds, dead material and weeds to improve orchard sanitation and airflow. ➤ Avoid water stagnation around the root zone and irrigate only according to actual soil moisture requirement. ➤ Apply organic manure/compost when soil is moist but not waterlogged and avoid application immediately before rainfall.
Coconut Pest and Disease:		➤ Rhinoceros beetle: Maintain crown sanitation and pheromone-trap monitoring; use neem cake/biological measures where appropriate. ➤ Black-headed caterpillar: Monitor leaflets and conserve parasitoids; where economically required, use azadirachtin or a recommended insecticide during dry weather. ➤ Bud rot: Inspect spindle/crown region under prolonged humidity; where disease is confirmed, copper oxychloride may be used according to approved recommendation and only in a dry period.
Pomegranate	Fruit Development	➤ Maintain uniform soil moisture and avoid excess irrigation after light rainfall to reduce fruit cracking and root-zone problems. ➤ Improve canopy ventilation through removal of excessively dense or diseased growth without heavy pruning. ➤ Remove cracked, diseased and damaged fruits immediately and maintain orchard sanitation.

		➤ Maintain good drainage and use organic mulch to reduce rapid soil-moisture fluctuations.
Pomegranate Pest and Disease:		➤ Bacterial blight: Remove infected plant parts and affected fruits promptly; where recommended, use copper oxychloride during dry weather with proper coverage. ➤ Anthracnose/fruit spot: Remove infected tissues and monitor closely after humid weather; mancozeb may be used when required during a rain-free window. ➤ Fruit borer: Install pheromone traps and collect damaged fruits; where ETL is reached, use emamectin benzoate during clear, calm weather.
Livestock Management		
➤ Keep animal sheds, calf pens and poultry houses clean, dry and well ventilated; remove dung and stagnant water regularly after rainfall. ➤ Ensure continuous availability of clean drinking water, green fodder, dry fodder, mineral mixture and common salt; avoid feeding wet or mouldy fodder. ➤ Avoid grazing animals in waterlogged or very muddy areas immediately after rainfall and provide a dry resting place to prevent hoof infections and parasitic problems. ➤ Monitor livestock for FMD, Haemorrhagic Septicaemia (HS), Black Quarter (BQ), mastitis and parasitic infections; isolate suspected animals and consult a veterinarian promptly. ➤ Protect calves, pregnant animals and poultry from prolonged wet and humid conditions and provide adequate ventilation. ➤ Follow the recommended vaccination and deworming schedule and avoid indiscriminate use of antibiotics or other veterinary medicines.		

Eastern Dry Zone

Agromet Advisory for Bengaluru Urban district

Crop	Stage	Pest & Diseases	Weather-Based Agromet Advisory
Late Kharif			Procure high-quality seeds and fertilizers for the Kharif sowing season. Ploughing across the slope will make insitu soil moisture conservation. Land preparation for sowing of kharif crops is advised to farmers
Finger millet	Sowing, Early vegetative		The following varieties are recommended for sowing: GPU-28, 45, 48 and 66, ML-365 Postpone sowing operation until sufficient soil moisture available in crop field.

Cowpea	Sowing, Early vegetative		e following varieties are recommended for sowing: C-152, TVX-944, KBC-1, Green gram: PS-16, PDM-84,178 Before sowing seeds treat with Rhizobium biofertilizer @ 200g/acre recommendation.
Black gram	Sowing, Early vegetative		The following varieties are recommended for sowing: Karagao-3, T-9, Rashmi (LBG-625)
Redgram	Vegetative		Intercultural operation is recommended. Thinning plant population Mulching is recommended for conserve the soil moisture. One or two protective irrigations is advisable.
Maize			
Groundnut			
Sunflower			
Baby corn	Maturity stage	Fall Armyworm	Spray Emamectin Benzoate @ 0.4 g/ l of water. Maintain regular irrigation at 4–5 day intervals. Remove weeds at regular interval. Top dressing of nitrogen fertilizer is recommended.
Bhendi	Fruit formation stage	Whitefly	Install Yellow Sticky Traps (15 per acre). Spray Neem Oil (3000 ppm)
Papaya	Vegetative	Mealybug	Maintain regular irrigation at 3–4-day intervals. Neem Oil + Liquid Soap: High-concentration Neem oil (1500-3000 ppm) mixed with a surfactant (shampoo) Apply recommended nitrogen fertilizers in split doses for vigorous vegetative growth.
Chilli	Maturity stage	Thrips	For thrips, use Fipronil 1 g/l of water or Neem oil 3 ml/l of water Apply recommended dose of potash to improve fruit quality and pungency. Foliar spray of micronutrients (Zn + Boron) if deficiency symptoms appear.
Rose	Flowering stage	Aphids	Maintain regular irrigation at 4–6 day intervals. Avoid moisture stress to ensure good bud size and quality. To control of aphids in rose to spray Dimethoate 30 EC @ 1.7 ml/litre of water.
Guava	Fruit development and maturity stage	Fruit fly	Maintain regular irrigation. Adopt mulching (dry leaves/straw) to conserve soil moisture. Apply recommended dose of potash to improve fruit size and quality. Foliar spray of micronutrients (Zn + Boron) if deficiency symptoms appear. Fruit fly (<i>Bactrocera spp.</i>) spray entomopathogenic fungus (<i>Beauveria bassiana</i>) @ 10 g/litre water on infested fruits. Use Methyl eugenol traps (10/acre) for fruit fly.

Agromet Advisory for Bengaluru Rural district

Crop	Stage	Pest and Diseases	Weather-Based Agromet Advisory
Late Kharif land preparation			Procure high-quality seeds and fertilizers for the late kharif sowing season. Ploughing across the slope will make insitu soil moisture conservation. Land preparation for sowing of kharif crops is advised to farmers
Finger millet	Sowing, Early vegetative		The following varieties are recommended for sowing: GPU-28, 45,48 and 66, ML-365 Postpone sowing operation until sufficient soil moisture available in crop field.
Cowpea	Sowing, Early vegetative		The following varieties are recommended for sowing: C-152, TVX-944, KBC-1, Green gram: PS-16, PDM-84,178 Before sowing seeds treat with Rhizobium biofertilizer @ 200g/acre recommendation.
Black gram	Sowing, Early vegetative		The following varieties are recommended for sowing: Karagao-3, T-9, Rashmi (LBG-625)
Redgram	Vegetative		Intercultural operation is recommended. Thinning plant population Mulching is recommended for conserve the soil moisture. One or two protective irrigations is advisable.
Maize			
Groundnut			
Sunflower			
Chrysanthemum	Flowering	Aphids	Avoid overhead sprinkler irrigation to protect the open flower blooms from rotting. Instead, apply light drip irrigation at the base, and lay down organic mulch (crop residues/dry straw) to prevent soil moisture evaporation. Spraying a 1% Potassium Nitrate solution (10 g per liter of water) increases the flower size and enhances its post-harvest shelf life
Grapes	Vegetative stage	Thrips	Install Blue Sticky Traps (15/acre). Spray Spinosad @ 0.3ml/ l of water or Neem Oil @ 3ml/ l of water Apply recommended nitrogen and magnesium sulphate fertilizers in split doses for vigorous vegetative growth.

Tomato	Fruit development and maturity stage	Mites and fruit borer	Continue irrigation at regular intervals. Watch for mites and fruit borer incidence under dry conditions. Spray calcium nitrate (0.5%) to reduce blossom end rot incidence. Avoid excessive nitrogen during fruiting stage.
Rose	Flowering stage	Aphids	Maintain regular irrigation at 4-6 days intervals. Avoid moisture stress to ensure good bud size and quality. To control, of aphids in rose to spray Dimethoate 30 EC@ 1.7 ml/ litre of water.
Guava	Fruit development and maturity stage	Fruit fly	Apply recommended dose of potash to improve fruit size and quality. Foliar spray of micronutrients (Zn + Boron) if deficiency symptoms appear. Maintain uniform soil moisture by providing regular drip irrigation. Avoid sudden heavy watering after a dry spell, as fluctuating moisture levels cause fruit cracking. Use Methyl eugenol traps (10/acre) for fruit fly.

Agromet Advisory for Kolar district

Crop	Stage	Pest & Diseases	Weather-Based Agromet Advisory
Late Kharif land preparation			Procure high-quality seeds and fertilizers for the Kharif sowing season. Ploughing across the slope will make insitu soil moisture conservation. Land preparation for sowing of kharif crops is advised to farmers
Finger millet	Sowing, Early vegetative		The following varieties are recommended for sowing: GPU-28, 45, 48 and 66, ML-365. Postpone sowing operation until sufficient soil moisture available in crop field.
Cowpea	Sowing, Early vegetative		The following varieties are recommended for sowing: C-152, TVX-944, KBC-1, Green gram: PS-16, PDM-84, 178. Before sowing seeds treat with Rhizobium biofertilizer @ 200g/acre recommendation.
Black gram	Sowing, Early vegetative		The following varieties are recommended for sowing: Karagao-3, T-9, Rashmi (LBG-625)
Redgram	Vegetative		Intercultural operation is recommended. Thinning plant population
Maize			
Groundnut			

Sunflower			Mulching is recommended for conserve the soil moisture. One or two protective irrigations is advisable.
Brinjal	Flowering and fruit formation stage	Shoot Borer	Spray Flubendiamide @ 0.5ml/l of water and Chlorantraniliprole 0.4ml/ l of water. Avoid excess nitrogen application during fruiting stage.
Beetroot	Marble stage	Aphids	Spray Borax @ 1g/liter or Solubor when the roots are about the size of a marble. Use Yellow Sticky Traps. Spray Thiamethoxam @ 0.5g/ l of water
Chilli	Maturity stage	Thrips	For thrips, use Fipronil 1 g/l of water or Neem oil 3 ml/l of water Apply recommended dose of potash to improve fruit quality and pungency. Foliar spray of micronutrients (Zn + Boron) if deficiency symptoms appear.
Rose	Flowering stage	Aphids	Maintain regular irrigation at 4–6-day intervals. Avoid moisture stress to ensure good bud size and quality. To control of aphids in rose to spray Dimethoate 30 EC @ 1.7 ml/litre of water.
Guava	Fruit development	Fruit fly	Maintain regular irrigation. Adopt mulching (dry leaves/straw) to conserve soil moisture. Apply recommended dose of potash to improve fruit size and quality. Foliar spray of micronutrients (Zn + Boron) if deficiency symptoms appear. Fruit fly (<i>Bactrocera spp.</i>) spray entomopathogenic fungus (<i>Beauveria bassiana</i>) @ 10 g/litre water on infested fruits. Use Methyl eugenol traps (10/acre) for fruit fly.
Mango			Recommended for pruning of dried twigs and branches done in June-July month to facilitate the sunlight to bear the more fruits in next seasons.

Agromet Advisory for Chikkaballapura district

Crop	Stage	Pest & Diseases	Weather-Based Agromet Advisory
Late Kharif land preparation			Procure high-quality seeds and fertilizers for the Kharif sowing season. Ploughing across the slope will make insitu soil moisture conservation. Land preparation for sowing of kharif crops is advised to farmers

Finger millet	Sowing, Early vegetative		The following varieties are recommended for sowing: GPU-28, 45,48 and 66, ML-365 Postpone sowing operation until sufficient soil moisture available in crop field.
Cowpea	Sowing, Early vegetative		The following varieties are recommended for sowing: C-152, TVX-944, KBC-1, Green gram: PS-16, PDM-84,178 Before sowing seeds treat with Rhizobium biofertilizer @ 200g/acre recommendation.
Black gram	Sowing, Early vegetative		The following varieties are recommended for sowing: Karagao-3, T-9, Rashmi (LBG-625)
Redgram	Vegetative		Intercultural operation is recommended. Thinning plant population Mulching is recommended for conserve the soil moisture. One or two protective irrigations is advisable.
Maize			
Groundnut			
Sunflower			
Capsicum	Flowering and Fruiting stage	Thrips & Mites	High humidity management; use overhead misters for 5-10 mins at noon to cool the canopy. Spray Spinosad @ 0.5 ml/ litre of water. or Fipronil @ 2 ml/ litre of water.
Chilli	Maturity stage	Thrips	For thrips, use Fipronil 1 g/l of water or Neem oil 3 ml/l of water Apply recommended dose of potash to improve fruit quality and pungency. Foliar spray of micronutrients (Zn + Boron) if deficiency symptoms appear.
Rose	Flowering stage	Aphids	Maintain regular irrigation at 4–6-day intervals. Avoid moisture stress to ensure good bud size and quality. To control of aphids in rose to spray Dimethoate 30 EC @ 1.7 ml/litre of water.
Guava	Fruit development and maturity stage	Fruit fly	Maintain regular irrigation. Adopt mulching (dry leaves/straw) to conserve soil moisture. Apply recommended dose of potash to improve fruit size and quality. Foliar spray of micronutrients (Zn + Boron) if deficiency symptoms appear. Fruit fly (<i>Bactrocera spp.</i>) spray entomopathogenic fungus (<i>Beauveria bassiana</i>) @ 10 g/litre water on infested fruits. Use Methyl eugenol traps (10/acre) for fruit fly.
Mango			Recommended for pruning of dried twigs and branches done in June-July month to facilitate the sunlight to bear the more fruits in next seasons.

Agromet Advisory for Ramanagara district

Crop	Stage	Pest and Diseases	Weather-Based Agromet Advisory
Late Kharif land preparation			Procure high-quality seeds and fertilizers for the Kharif sowing season. Ploughing across the slope will make insitu soil moisture conservation. Land preparation for sowing of kharif crops is advised to farmers
Finger millet	Sowing, Early vegetative		The following varieties are recommended for sowing: GPU-28, 45,48 and 66, ML-365 Postpone sowing operation until sufficient soil moisture available in crop field.
Cowpea	Sowing, Early vegetative		The following varieties are recommended for sowing: C-152, TVX-944, KBC-1, Green gram: PS-16, PDM-84,178 Before sowing seeds treat with Rhizobium biofertilizer @ 200g/acre recommendation.
Black gram	Sowing, Early vegetative		The following varieties are recommended for sowing: Karagao-3, T-9, Rashmi (LBG-625)
Redgram	Vegetative		Intercultural operation is recommended. Thinning plant population Mulching is recommended for conserve the soil moisture. One or two protective irrigations is advisable.
Maize			
Groundnut			
Sunflower			
French beans	Flowering and fruit formation stage	Mites	Avoid water stress during flowering and frequent irrigation is required for better growth.
Chilli	Maturity stage	Thrips	For thrips, use Fipronil 1 g/l of water or Neem oil 3 ml/l of water Apply recommended dose of potash to improve fruit quality and pungency. Foliar spray of micronutrients (Zn + Boron) if deficiency symptoms appear.
Ridge guard	Flowering and fruit formation stage	Fruit Fly	Install Cue-lure pheromone traps (10/acre). Wrap young fruits in paper/cloth bags. Irrigate at 3–4 day intervals in summer.
Tomato	Fruit formation stage	Whitefly / Aphids	Install Yellow Sticky Traps (10/acre). Spray Imidacloprid (0.3 ml/L) or Neem Oil.
Mango			Recommended for pruning of dried twigs and branches done in June-July month to facilitate the sunlight to bear the more fruits in next seasons.
Arecanut		Inflorescence Die-back	Apply dry leaves, coconut husk or organic mulch in basins to conserve moisture.

			Spray Hexaconazole (1 ml/liter) or 1 % Bordeaux mixture.
--	--	--	---

Southern Dry Zone

Agromet Advisory for Kodagu district

Crop / Enterprise	Crop Stage	Weather-based Advisory	Severe Pests / Diseases Likely	Recommended Management
Coffee	Berry maturation to harvesting	Low rainfall and dry spells may affect berry filling and maturation. Conserve soil moisture through mulching and provide protective irrigation wherever available.	Coffee Berry Borer, Leaf Rust, Black Rot	Maintain shade and soil moisture. Collect and destroy fallen berries. Monitor berry borer and fungal diseases and undertake need-based management during suitable weather conditions.
Black Pepper	Berry development to maturity	Moisture stress may affect berry development and cause premature berry drop. Maintain organic mulch around vines and provide protective irrigation during prolonged dry periods.	Pollu Beetle, Quick Wilt, Foot Rot	Maintain mulch and proper drainage. Monitor spikes and vines regularly. Remove severely affected plant parts and undertake need-based pest and disease management.
Arecanut	Nut development to maturity	Low rainfall may cause moisture stress and affect nut development. Conserve soil moisture through mulching and provide protective irrigation wherever available.	Koleroga (Fruit Rot), Spindle Bug, Mites	Maintain basin mulch and clean basins. Monitor bunches and spindle region regularly and undertake need-based plant protection.
Ginger	Rhizome development to maturity	Rainfall deficit may restrict rhizome development. Maintain mulch and provide protective irrigation during prolonged dry periods while avoiding excessive irrigation.	Rhizome Rot, Leaf Spot	Maintain adequate soil moisture and drainage. Remove infected clumps and undertake need-based disease management during suitable weather.
Paddy	Panicle initiation to flowering	Very low rainfall may cause moisture stress during critical reproductive stages and affect grain formation.	Blast, Stem Borer, Leaf Folder	Give priority to protective irrigation wherever water is available. Maintain adequate soil moisture during critical stages.

				Monitor for blast, stem borer and leaf folder and undertake need-based management.
Maize	Pre-tasseling to tasseling	Moisture stress during tasseling may seriously affect pollination and grain formation.	Fall Armyworm, Stem Borer, Downy Mildew	Provide protective irrigation at critical stages and conserve moisture through intercultivation and earthing-up. Monitor Fall Armyworm. If severe moisture stress persists and crop recovery is poor, utilize the crop as green fodder or silage .
Butterfruit (Avocado)	Fruit development to maturity	Dry spell may reduce fruit enlargement and increase fruit drop. Conserve soil moisture through organic mulching and provide protective irrigation wherever available.	Anthraxnose, Fruit Rot, Fruit Fly	Maintain mulch and adequate soil moisture. Harvest physiologically mature fruits. Collect fallen fruits and monitor for fruit fly and fungal diseases.
Cattle & Buffalo	Maintenance / Lactation	Dry conditions may reduce green fodder availability and increase water requirement.	Heat stress, Dehydration, Reduced feed intake	Ensure continuous access to clean drinking water and adequate green and dry fodder. Provide shade and good ventilation. Avoid strenuous activities during hotter periods.
Sheep & Goat	Maintenance / Growing	Reduced availability of green fodder and water may affect feed intake and body condition.	Heat stress, Dehydration, Reduced feed intake	Provide adequate drinking water, green/dry fodder and shade. Avoid overcrowding and maintain clean, well-ventilated shelters.
Poultry	Growing / Laying	Warm and dry conditions may increase heat stress and reduce feed intake and production.	Heat stress, Reduced feed intake, Reduced egg production	Ensure continuous clean drinking water and good ventilation. Avoid overcrowding, maintain dry litter and follow proper shed hygiene.
Sericulture – Mulberry	Leaf production	Rainfall deficit may reduce mulberry leaf yield and quality.	Moisture stress, Leaf spot	Provide protective irrigation wherever available. Maintain mulch and weed-free conditions. Apply nutrients only when adequate soil moisture is available.

Sericulture – Silkworm	Late larval stage to cocooning	Poor-quality mulberry leaves due to moisture stress may affect larval growth and cocoon quality.	Grasserie, Muscardine, Flacherie	Provide fresh, tender and good-quality mulberry leaves. Maintain proper ventilation, hygiene and suitable rearing conditions. Avoid overcrowding.
Sericulture – Cocoon	Cocooning to harvesting	Unfavourable rearing conditions may affect cocoon formation and quality.	Poor cocoon formation, Cocoon loss	Maintain clean, hygienic and well-ventilated conditions. Avoid disturbance during cocoon formation and harvest mature cocoons promptly.

Agromet Advisory for Chamarajanagara district

Crop / Enterprise	Crop Stage	Likely Impact of Rainfall Deficit	Recommended Management
Field Bean	Vegetative to Flowering	Moisture stress may restrict vegetative growth, flowering and pod development.	Provide protective irrigation wherever available. Conserve soil moisture through intercultivation and mulching. Apply fertilizer only when adequate soil moisture is available.
Cowpea	Vegetative to Flowering	Moisture stress may reduce growth, flowering and pod formation.	Provide protective irrigation during prolonged dry spells. Maintain mulch and undertake intercultivation to conserve moisture.
Black Gram	Vegetative to Flowering	Moisture stress may reduce branching, flowering and pod formation.	Provide protective irrigation at critical stages. Maintain weed-free conditions to reduce competition for moisture. Monitor for Yellow Mosaic Virus and whitefly.
Green Gram	Vegetative to Flowering	Inadequate rainfall may affect vegetative growth and reproductive development.	Provide protective irrigation wherever possible. Conserve soil moisture through intercultivation and mulching. Monitor for whitefly and leaf spot.
Red Gram	Vegetative to Flowering	Prolonged dry spell may reduce growth and flowering.	Provide protective irrigation at critical stages. Undertake intercultivation and maintain soil cover to conserve moisture.
Maize	Vegetative to Pre-tasseling	Moisture stress before tasseling can severely affect crop growth and subsequent grain formation.	Give priority to protective irrigation . Undertake earthing-up and moisture conservation. If severe moisture stress persists and recovery is poor, utilize the crop as green fodder or silage .
Chilli	Flowering to Fruiting	Moisture stress may cause flower drop, poor fruit set and reduced fruit size.	Provide life-saving irrigation , preferably through drip. Maintain mulch and avoid moisture fluctuations. Avoid excessive nitrogen under moisture stress.

Tomato	Fruit Development	Moisture stress may affect fruit growth and quality.	Maintain drip irrigation and mulch. Provide protective irrigation based on soil-moisture status. Harvest mature fruits regularly and remove diseased fruits.
Tobacco	Vegetative to Grand Growth	Moisture stress may restrict leaf expansion and affect leaf yield and quality.	Provide protective irrigation wherever available. Conserve moisture through intercultivation. Avoid fertilizer application when the soil is dry.
Ginger	Rhizome Development	Moisture deficit may restrict rhizome development and reduce yield.	Maintain mulch and provide protective irrigation during prolonged dry periods. Avoid excessive irrigation and maintain suitable soil moisture around the rhizome zone.
Paddy	Tillering to Panicle Initiation / Delayed Transplanting	Very low rainfall may cause inadequate water availability and moisture stress.	Provide protective irrigation wherever irrigation is assured. For delayed transplanting, avoid long-duration varieties without assured water and prefer short-duration varieties where suitable.
Livestock	Maintenance / Lactation / Growing	Dry conditions may reduce green fodder availability and increase water requirement.	Ensure continuous access to clean drinking water , adequate green and dry fodder, shade and well-ventilated housing.
Poultry	Growing / Laying	Warm conditions may increase heat stress, reduce feed intake and affect production.	Ensure continuous clean drinking water, adequate ventilation and proper housing hygiene. Avoid overcrowding and maintain dry litter.
Sericulture – Mulberry	Vegetative / Leaf Production	Moisture stress may reduce mulberry leaf yield and quality.	Provide protective irrigation wherever available. Maintain mulch and weed-free conditions. Apply nutrients only when adequate soil moisture is available.
Sericulture – Silkworm	Larval to Cocooning	Poor-quality mulberry leaves may affect larval growth and cocoon quality.	Provide fresh, tender and good-quality mulberry leaves. Maintain proper ventilation, hygiene and suitable rearing conditions. Avoid overcrowding.

Agromet Advisory for Mysore/ Mandya districts

Crop / Enterprise	Crop Stage	Likely Impact of Rainfall Deficit	Recommended Management
Field Bean	Vegetative to Flowering	Moisture stress may restrict vegetative growth, flowering and pod development.	Provide protective irrigation wherever available. Conserve soil moisture through intercultivation and mulching. Apply fertilizer only when adequate soil moisture is available.

Cowpea	Vegetative to Flowering	Moisture stress may reduce growth, flowering and pod formation.	Provide protective irrigation during prolonged dry spells. Maintain mulch and undertake intercultivation to conserve moisture.
Black Gram	Vegetative to Flowering	Moisture stress may reduce branching, flowering and pod formation.	Provide protective irrigation at critical stages. Maintain weed-free conditions to reduce competition for moisture. Monitor for Yellow Mosaic Virus and whitefly.
Green Gram	Vegetative to Flowering	Inadequate rainfall may affect vegetative growth and reproductive development.	Provide protective irrigation wherever possible. Conserve soil moisture through intercultivation and mulching. Monitor for whitefly and leaf spot.
Red Gram	Vegetative to Flowering	Prolonged dry spell may reduce growth and flowering.	Provide protective irrigation at critical stages. Undertake intercultivation and maintain soil cover to conserve moisture.
Maize	Vegetative to Pre-tasseling	Moisture stress before tasseling can severely affect crop growth and subsequent grain formation.	Give priority to protective irrigation . Undertake earthing-up and moisture conservation. If severe moisture stress persists and recovery is poor, utilize the crop as green fodder or silage .
Chilli	Flowering to Fruiting	Moisture stress may cause flower drop, poor fruit set and reduced fruit size.	Provide life-saving irrigation , preferably through drip. Maintain mulch and avoid moisture fluctuations. Avoid excessive nitrogen under moisture stress.
Tomato	Fruit Development	Moisture stress may affect fruit growth and quality.	Maintain drip irrigation and mulch. Provide protective irrigation based on soil-moisture status. Harvest mature fruits regularly and remove diseased fruits.
Tobacco	Vegetative to Grand Growth	Moisture stress may restrict leaf expansion and affect leaf yield and quality.	Provide protective irrigation wherever available. Conserve moisture through intercultivation. Avoid fertilizer application when the soil is dry.
Ginger	Rhizome Development	Moisture deficit may restrict rhizome development and reduce yield.	Maintain mulch and provide protective irrigation during prolonged dry periods. Avoid excessive irrigation and maintain suitable soil moisture around the rhizome zone.
Paddy	Tillering to Panicle Initiation / Delayed Transplanting	Very low rainfall may cause inadequate water availability and moisture stress.	Provide protective irrigation wherever irrigation is assured. For delayed transplanting, avoid long-duration varieties without assured water and prefer short-duration varieties where suitable.
Livestock	Maintenance / Lactation / Growing	Dry conditions may reduce green fodder	Ensure continuous access to clean drinking water , adequate green and

		availability and increase water requirement.	dry fodder, shade and well-ventilated housing.
Poultry	Growing / Laying	Warm conditions may increase heat stress, reduce feed intake and affect production.	Ensure continuous clean drinking water, adequate ventilation and proper housing hygiene. Avoid overcrowding and maintain dry litter.
Sericulture – Mulberry	Vegetative / Leaf Production	Moisture stress may reduce mulberry leaf yield and quality.	Provide protective irrigation wherever available. Maintain mulch and weed-free conditions. Apply nutrients only when adequate soil moisture is available.
Sericulture – Silkworm	Larval to Cocooning	Poor-quality mulberry leaves may affect larval growth and cocoon quality.	Provide fresh, tender and good-quality mulberry leaves. Maintain proper ventilation, hygiene and suitable rearing conditions. Avoid overcrowding.

Southern Transition Zone

Agromet Advisory Service Bulletin for Shivamogga district

Crop	Stage	Advisory
Paddy	Transplanting stage/Seedling stage	- Go for transplanting of paddy seedlings. - Apply recommended dose of fertilizer@ 40:20:20:8 N. P. K, Zn Per acre. Recommended P, K, Zn and 50% N apply at the time of transplanting and remaining 50% of N should apply in two splits. - Dip seedlings in Bavistin powder @ 0.1% for 10 – 20 minutes. - Apply pre emergent herbicide like Butachlor 50 EC @ 2 litre a.i/ha, within 3-5 days after transplanting.
Arecanut	Nut development	Weather favours Mahali (fruit rot). Remove fallen infected nuts and spray 1% Bordeaux mixture or Copper oxychloride @ 3 g/L during a rain-free period.
Maize	Flowering stage	Provide adequate irrigation and fertilizers when the crop is at the flowering stage; otherwise, the grains will shrivel.
Ginger	Vegetative stage	Continuous rainfall and high humidity favour rhizome rot and leaf spot. Ensure proper drainage and drench with Metalaxyl + Mancozeb @ 2 g/L or Copper oxychloride @ 3 g/L if symptoms appear

		For managing root rot provide adequate drainage to remove excess of water from the field.
Black pepper	Fruiting stage	Conditions favour quick wilt (Phytophthora foot rot). Ensure drainage and spray 1% Bordeaux mixture or Copper oxychloride @ 3 g/L around vines and the collar region during a rain-free period.
Cattle	-	Early diagnosis of mastitis in dairy animals by farmers during milking by observing the swelling of udder and change in the smell of the milk and immediately drain out the milk from affected teat infuse intra mammary Pendistrin Infusion.

Agromet Advisory Service Bulletin for Hassan district






Crop	Stage	Advisory
Arecanut	Nut development	Weather favours Mahali (fruit rot) . Remove fallen infected nuts and spray 1% Bordeaux mixture or Copper oxychloride @ 3 g/L during a rain-free period.
Maize	Flowering stage	Provide adequate irrigation and fertilizers when the crop is at the flowering stage; otherwise, the grains will shrivel.
Paddy	Transplanting stage/Seedling stage	- Go for transplanting of paddy seedlings. - Apply recommended dose of fertilizer@ 40:20:20:8 N. P. K, Zn Per acre. Recommended P, K, Zn and 50% N apply at the time of transplanting and remaining 50% of N should apply in two splits. - Dip seedlings in Bavistin powder @ 0.1% for 10 – 20 minutes. - Apply pre emergent herbicide like Butachlor 50 EC @ 2 litre a.i/ha, within 3-5 days after transplanting.
Ginger	Vegetative stage	- Continuous rainfall and high humidity favour rhizome rot and leaf spot. Ensure proper drainage and drench with Metalaxyl + Mancozeb @ 2 g/L or Copper oxychloride @ 3 g/L if symptoms appear - For managing root rot provide adequate drainage to remove excess of water from the field.
Coconut	Maturity /harvesting stage	For managing Rhinoceros beetle remove the beetle from infected part with the long iron rod by piercing in whorl and fill 2 % Quinolphos or 5 % Melathion in sand @ 1:1 ratio.

Black pepper	Fruiting stage	Conditions favour quick wilt (Phytophthora foot rot). Ensure drainage and spray 1% Bordeaux mixture or Copper oxychloride @ 3 g/L around vines and the collar region during a rain-free period.
Cattle	-	Early diagnosis of mastitis in dairy animals by farmers during milking by observing the swelling of udder and change in the smell of the milk and immediately drain out the milk from affected teat infuse intra mammary Pendistrin Infusion.

Agromet Advisory Service Bulletin for Chikkamagaluru district

Crop	Stage	Advisory
Maize	Flowering stage	Provide adequate irrigation and fertilizers when the crop is at the flowering stage; otherwise, the grains will shrivel.
Arecanut	Nut development	Weather favours Mahali (fruit rot). Remove fallen infected nuts and spray 1% Bordeaux mixture or Copper oxychloride @ 3 g/L during a rain-free period.
Paddy	Transplanting stage/Seedling stage	- Go for transplanting of paddy seedlings. - Apply recommended dose of fertilizer@ 40:20:20:8 N. P. K, Zn Per acre. Recommended P, K, Zn and 50% N apply at the time of transplanting and remaining 50% of N should apply in two splits. - Dip seedlings in Bavistin powder @ 0.1% for 10 – 20 minutes. - Apply pre emergent herbicide like Butachlor 50 EC @ 2 litre a.i/ha, within 3-5 days after transplanting.
Ginger	Vegetative stage	- Continuous rainfall and high humidity favour rhizome rot and leaf spot. Ensure proper drainage and drench with Metalaxyl + Mancozeb @ 2 g/L or Copper oxychloride @ 3 g/L if symptoms appear - For managing root rot provide adequate drainage to remove excess of water from the field.
Black pepper	Fruiting stage	Conditions favour quick wilt (Phytophthora foot rot). Ensure drainage and spray 1% Bordeaux mixture or Copper oxychloride @ 3 g/L around vines and the collar region during a rain-free period.
Cattle	-	Early diagnosis of mastitis in dairy animals by farmers during milking by observing the swelling of udder and change in the smell of the milk and

		immediately drain out the milk from affected teat infuse intra mammary Pendistrin Infusion.
--	--	--

5-day Weather Warning maps :

[illegible]

KARNATAKA STATE FORECAST ISSUED AT 1300 HOURS IST OF 28/Aug/26															
Met Sub.Division/ DISTRICTS	Distribution	Rainfall forecast for next 7 days													
		Day 1		Day 2		Day 3		Day 4		Day 5		Day 6		Day 7	
		valid from 1300	valid from 0830	valid from 0830	valid from 0830	valid from 0830	valid from 0830	valid from 0830	valid from 0830	valid from 0830	valid from 0830	valid from 0830	valid from 0830	valid from 0830	
		Hrs IST of 28/Aug/2026 to 0830 hrs IST of 29/Aug/2026	Hrs IST of 29/Aug/2026 to 0830 hrs IST of 30/Aug/2026	Hrs IST of 30/Aug/2026 to 0830 hrs IST of 31/Aug/2026	Hrs IST of 31/Aug/2026 to 0830 hrs IST of 01/Sep/2026	Hrs IST of 01/Sep/2026 to 0830 hrs IST of 02/Sep/2026	Hrs IST of 02/Sep/2026 to 0830 hrs IST of 03/Sep/2026	Hrs IST of 03/Sep/2026 to 0830 hrs IST of 04/Sep/2026	Hrs IST of 04/Sep/2026 to 0830 hrs IST of 05/Sep/2026	Hrs IST of 05/Sep/2026 to 0830 hrs IST of 06/Sep/2026	Hrs IST of 06/Sep/2026 to 0830 hrs IST of 07/Sep/2026	Hrs IST of 07/Sep/2026 to 0830 hrs IST of 08/Sep/2026	Hrs IST of 08/Sep/2026 to 0830 hrs IST of 09/Sep/2026		
		FORECAST	WARNING	FORECAST	WARNING	FORECAST	WARNING	FORECAST	WARNING	FORECAST	WARNING	FORECAST	WARNING	FORECAST	WARNING
CHIKKAMAGALURU	Intensity	L/M	GW	L/M	GW	L/M	NIL	L/M	NIL	L/M	NIL	L/M	NIL	L/M	NIL
	Probability	VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY	
CHITRADURGA	Intensity	DRY	G'W	DRY	G'W	DRY	NIL	DRY	NIL	DRY	NIL	L/M	NIL	L/M	NIL
	Probability	VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY	
DAVANGERE	Intensity	DRY	G'W	DRY	G'W	DRY	NIL	DRY	NIL	DRY	NIL	L/M	NIL	L/M	NIL
	Probability	VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY	
HASSAN	Intensity	L/M	GW	L/M	GW	L/M	NIL	L/M	NIL	L/M	NIL	L/M	NIL	L/M	NIL
	Probability	VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY	
KODAGU	Intensity	L/M	GW	L/M	GW	L/M	NIL	L/M	NIL	L/M	NIL	L/M	NIL	L/M	NIL
	Probability	VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY	
KOLAR	Intensity	L	GW	L	GW	L	NIL	DRY	NIL	DRY	NIL	L/M	NIL	L/M	NIL
	Probability	VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY	
MANDYA	Intensity	DRY	G'W	DRY	G'W	DRY	NIL	DRY	NIL	DRY	NIL	L/M	NIL	L/M	NIL
	Probability	VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY	
MYSURU	Intensity	DRY	GW	DRY	GW	DRY	NIL	DRY	NIL	DRY	NIL	L/M	NIL	L/M	NIL
	Probability	VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY	
RAMANAGARA	Intensity	DRY	GW	DRY	GW	DRY	NIL	DRY	NIL	DRY	NIL	L/M	NIL	L/M	NIL
	Probability	VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY	
SHIVAMOGGA	Intensity	L/M	GW	L/M	GW	L/M	NIL	L/M	NIL	L/M	NIL	L/M	NIL	L/M	NIL
	Probability	VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY	
TUMAKURU	Intensity	L	GW	L	GW	L	NIL	DRY	NIL	DRY	NIL	L/M	NIL	L/M	NIL
	Probability	VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY	
VIJAYANAGARA	Intensity	DRY	G'W	DRY	GW	DRY	NIL	DRY	NIL	DRY	NIL	L/M	NIL	L/M	NIL
	Probability	VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY		VERY LIKELY	

KARNATAKA STATE FORECAST ISSUED AT 1300 HOURS IST OF 28/Aug/26																
Met Sub.Division/ DISTRICTS	Distribution	Rainfall forecast for next 7 days														
		Day 1		Day 2		Day 3		Day 4		Day 5		Day 6		Day 7		
		valid from 1300	valid from 0830	valid from 0830	valid from 0830	valid from 0830	valid from 0830	valid from 0830	valid from 0830	valid from 0830	valid from 0830	valid from 0830	valid from 0830			
		Hrs IST of 28/Aug/2026	Hrs IST of 29/Aug/2026	Hrs IST of 30/Aug/2026	Hrs IST of 31/Aug/2026	Hrs IST of 1/Sep/2026	Hrs IST of 2/Sep/2026	Hrs IST of 3/Sep/2026	Hrs IST of 4/Sep/2026	Hrs IST of 5/Sep/2026	Hrs IST of 6/Sep/2026	Hrs IST of 7/Sep/2026				
		to 0830 hrs IST of 29/Aug/2026	to 0830 hrs IST of 30/Aug/2026	to 0830 hrs IST of 31/Aug/2026	to 0830 hrs IST of 01/Sep/2026	to 0830 hrs IST of 02/Sep/2026	to 0830 hrs IST of 03/Sep/2026	to 0830 hrs IST of 04/Sep/2026								
		FORECAST	WARNING	FORECAST	WARNING	FORECAST	WARNING	FORECAST	WARNING	FORECAST	WARNING	FORECAST	WARNING			
Sustained wind with speed 30-40 kmph likely over South Interior Karnataka from 28th to 29th August 2026.																
VL – Very Light Rainfall (Trace to 2.4 mm)		L/M – Light to Moderate Rain (2.5 mm to 64.4 mm)					ISL – Isolated places (<=25%)									
L – Light Rainfall (2.5 mm to 15.5 mm)		M/M – Moderate to Heavy Rain (15.6 mm to 115.5 mm)					SCT – At a few places (>25% & <=50%)									
M – Moderate Rainfall (15.6 mm to 64.4 mm)		GR – Hail Storm					GW – Gusty wind									
H (R+) – Heavy Rainfall (64.5 to 115.5)		SQ – Squall					SW – Sustained Wind									
VH (R++) – Very Heavy Rainfall (115.6 mm to 204.4 mm)		TSGW – Thunderstorm with Gusty wind					WS – Wide spread – At most places (>75%)									
Ext(R+++)-Extremely Heavy Rainfall (>= 204.5mm)		TSH – Thunderstorm with Lightning					HH – Hot & Humid HW –Heat Wave CW: Cold Wave									
SEVERE COLD WAVE		TSGR- Thundershower with hail					S HW – Severe Heat Wave upto 2days; S CW – Severe Cold Wave upto 2days									
							S HW* – Severe Heat Wave > 2days; S CW* – Severe Cold Wave > 2days									
							COLD WAVE									
Probability		Likely (26%-50%);					Very Likely (51%-75%)					Most Likely (>75%)				