

# Pure Mathematics for Engineers and Scientists

Anthony Daniell

June 3, 2025



# Preface

The basic aim is to provide exposure to a solid foundation of mathematics based on proofs. The exit goals would be:

- Readers will be able to write valid proofs using the standard techniques of modern mathematics.

- Readers will be able to read and understand proofs at some reasonable level.

- Readers will have working knowledge of broad foundations of mathematics, and be able to confidently take dedicated courses in each area, or pursue independent studies.

From my experience, I certainly never got this stuff explicitly, but on retrospect, it certainly seems like it would help.

My original concept was that the course would be targeted from someone maybe at the junior or senior level in physics/engineering. So, the typical prerequisite would be calculus, maybe differential equations, and experience applying this math to physics, engineering problems like in mechanics, electricity and magnetism, etc. That's essentially my background, so that's why I thought of this cohort.

The main reason is that this person, in my concept, would have some context and exposure to different math, and may be better motivated to wanting to understand the underpinnings. Also, they may be more interested to branch out into the other topics on the list as these will likely be important as they take more advanced classes in their careers.

However, that being said, I think the main prerequisite is being interested in pure mathematics, perhaps with some future application in mind, but not necessarily.

Topics to be covered:

Mathematical Foundations - Mathematical logic, basic notation, axiomatic set theory (Zermelo-Fraenkel), Peano Axioms of Arithmetic, Gödel theorems.

Methods of Proof: Induction, Contradiction, Direct Proofs. Examples of classic proofs - infinite primes, irrationality of square root of two, etc. Plenty of practice examples.

Modern Algebra: Groups, Rings, Fields, Transformations, Representation Theory, Modules, Galois Theory

Real Analysis - Study of real numbers, including construction by Dedekind Cuts etc, point set topology, continuity, metric spaces (Basically a compressed version of Rudin or equivalent).

Complex Analysis - Study of functions of a complex variable with connections to other areas of mathematics including number theory. Riemman Zeta Function and Riemann Hypothesis.

Non-euclidean geometry and topology: Various spaces and properties. Invariants, knot theory, homotopy, homology.

Number theory:

Category theory:

Computational complexity theory: Finite Automata, lambda calculus, NP and other spaces.

# Contents

|          |                                                 |           |
|----------|-------------------------------------------------|-----------|
| <b>1</b> | <b>Mathematical Foundations</b>                 | <b>7</b>  |
| 1.1      | Basic Notation . . . . .                        | 7         |
| 1.2      | Mathematical Logic . . . . .                    | 8         |
| 1.3      | Zermelo-Fraenkel Axiomatic Set Theory . . . . . | 12        |
| 1.3.1    | The Zermelo-Fraenkel Axioms . . . . .           | 12        |
| 1.3.2    | Set Operations . . . . .                        | 14        |
| 1.4      | Peano Axioms of Arithmetic . . . . .            | 23        |
| 1.5      | Gödel Theorems . . . . .                        | 25        |
| <b>2</b> | <b>Methods of Proof</b>                         | <b>39</b> |
| 2.1      | Induction . . . . .                             | 39        |
| 2.1.1    | Weak Induction . . . . .                        | 39        |
| 2.1.2    | Strong Induction . . . . .                      | 41        |
| 2.2      | Proof by Contradiction . . . . .                | 41        |
| 2.3      | Direct Proof . . . . .                          | 42        |
| <b>3</b> | <b>Modern Algebra</b>                           | <b>43</b> |
| <b>4</b> | <b>Real Analysis</b>                            | <b>45</b> |
| <b>5</b> | <b>Complex Analysis</b>                         | <b>47</b> |
| <b>6</b> | <b>Non-Euclidean Geometry and Topology</b>      | <b>49</b> |
| <b>7</b> | <b>Number Theory</b>                            | <b>51</b> |
| <b>8</b> | <b>Category Theory</b>                          | <b>53</b> |
| <b>9</b> | <b>Computational Complexity Theory</b>          | <b>55</b> |



# Chapter 1

## Mathematical Foundations

The objective of this chapter is to present some basic concepts that underlie much of the remaining material in this book.

### 1.1 Basic Notation

The reader may be familiar with some if not all of the following notation, but it is presented here for concreteness and review.

- $\forall$  is read as “for all”
- $\exists$  is read as “exists”
- $\subset$  is read as “subset of”
- $\in$  is read as “in”
- $\mathbb{Z}$  is the set of integers:  $\dots, -2, -1, 0, 1, 2, 3, \dots$
- $\mathbb{N}$  is the set of natural numbers:  $0, 1, 2, 3, \dots$  (note: Sometimes these start with 0, and sometimes with 1. Usually doesn’t matter to the argument in question, but the reader should be aware.)
- $\mathbb{Q}$  is the set of rational numbers. These are numbers of the form  $\frac{p}{q}$  where  $p, q$  are integers, and  $q \neq 0$
- $\mathbb{R}$  is the set of real numbers. These will later be defined precisely in terms of *Dedekind Cuts*, but for now we’ll just use the informal notion of all the points on a coordinate line. However, hopefully the reader can see this is not good enough for precise work.
- We use curly braces to indicate sets in general. For example,  $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, 3, \dots\}$  defines the set of integers using the curly braces and examples. It is hoped the reader is able to abstract that the dots to the left imply the list has

no smallest negative value and the dots to the right imply that there is no largest positive value.

- **Set builder notation:** There are other ways to specify the set within the braces. For example, we can indicate the property that each member of the set has:  $S = \{p | p = 2m + 1, m \in \mathbb{Z}\}$ . This is read as the set  $S$  consists of those values  $p$  of the form  $2m + 1$  where  $m$  is an integer. In other words,  $S$  the set of odd integers. The vertical bar in the braces is read as “such that” and the comma is read as “and.”

#### *Exercise*

Write the definition of the set of rational numbers using set builder notation.

#### *Solution*

$$\mathbb{Q} = \left\{ r \mid r = \frac{p}{q}, p, q \in \mathbb{Z}, q \neq 0 \right\}$$

## 1.2 Mathematical Logic

It should be remarked that a good portion of this section (as well as some others) was strongly inspired by the YouTube video series “Math Major Basics” by MathDoctorBob (<https://www.youtube.com/playlist?list=PLF2DF6C3C8015DF5F>). We cover some key aspects here. It is very helpful to understand *Truth Tables* in this context. For those who have some experience with digital logic or computer programming, much of this will be quite familiar.

We have the following logical relations, with symbols, defined by the truth tables shown below: Negation ( $\neg$ ), Conjunction/And ( $\wedge$  - note that this symbol looks like the outline of the letter “A”, to help you remember “And”), Disjunction/or ( $\vee$ ), Implication ( $\implies$ ), If and Only If ( $\iff$ ):

#### **Negation:**

| $A$ | $\neg A$ |
|-----|----------|
| $T$ | $F$      |
| $F$ | $T$      |

#### **Conjunction/And:**

| $A$ | $\wedge$ | $B$ |
|-----|----------|-----|
| $T$ | $T$      | $T$ |
| $T$ | $F$      | $F$ |
| $F$ | $F$      | $T$ |
| $F$ | $F$      | $F$ |

#### **Disjunction/Or:**

| $A$ | $\vee$ | $B$ |
|-----|--------|-----|
| $T$ | $T$    | $T$ |
| $T$ | $T$    | $F$ |
| $F$ | $T$    | $T$ |
| $F$ | $F$    | $F$ |



**Implication:**

| $A$ | $\implies$ | $B$ |
|-----|------------|-----|
| $T$ | $T$        | $T$ |
| $T$ | $F$        | $F$ |
| $F$ | $T$        | $T$ |
| $F$ | $T$        | $F$ |

**If and only if (Iff):**

| $A$ | $\iff$ | $B$ |
|-----|--------|-----|
| $T$ | $T$    | $T$ |
| $T$ | $F$    | $F$ |
| $F$ | $F$    | $T$ |
| $F$ | $T$    | $F$ |

In the above,  $A$  and  $B$  are statements which can be either true (T) or false (F). The result of the logical operation is shown in the column under the particular symbol. For example, if  $A$  is true, and  $B$  is true, then  $A \wedge B$  is true. However, either either  $A$  or  $B$  is false (or both) then  $A \wedge B$  is false. It is important to refer back to the truth table definitions of these logical operations to avoid confusion.

Sometimes, implication is called “if-then” as in “If  $A$  then  $B$ .” However, note that it is possible for  $A$  to be false and  $B$  to be true, and the implication  $A \implies B$  to be true. This is somewhat at odds with how this if-then might be expressed in common usage: How could something false imply something true? This seems confusing. Therefore, always refer to the truth table for clarity on this point.

Here is an illustration regarding implication ( $\implies$ ).

Let  $A = (a > 5)$ ,  $B = (a > 3)$  where  $a$  is some integer. Now consider the following cases:

If  $a = 6$ ,  $(6 > 5) = T$ ,  $(6 > 3) = T$

If  $a = 4$ ,  $(4 > 5) = F$ ,  $(4 > 3) = T$

If  $a = 2$ ,  $(2 > 5) = F$ ,  $(2 > 3) = F$

If we compare this to the truth table for implication, we see that implication is always true. Note that with this particular definition of the statements  $A$  and  $B$  we cannot have a case where  $A$  is true and  $B$  is false.

*Exercise* Show  $A \iff B$  is equivalent to  $(A \implies B) \wedge (B \implies A)$ .

*Solution*

| $(A \implies B)$ | $\wedge$ | $(B \implies A)$ |
|------------------|----------|------------------|
| $T$              | $T$      | $T$              |
| $T$              | $F$      | $T$              |
| $F$              | $T$      | $F$              |
| $F$              | $F$      | $F$              |

We now consider some important terminology:

- *Tautology*: Any statement that is true for all values of its components is a tautology.
- *Contradiction*: Any statement that is false for all values of its components is a contradiction. The negation of a contradiction is a tautology and the negation of a tautology is a contradiction.
- *Logically Equivalent*: When  $A \iff B$  is a tautology. This means  $A$  and  $B$  have the same truth table. The notation for this is  $A \equiv B$ .

We illustrate logical equivalence by considering “De Morgan’s Laws” for disjunction (or) and conjunction (and). De Morgan’s Law’s are useful for manipulating logical expressions:

$$\neg(A \vee B) \equiv (\neg A) \wedge (\neg B)$$

$$\neg(A \wedge B) \equiv (\neg A) \vee (\neg B)$$

In words, the first law says that the negation ( $\neg$ ) of a disjunction ( $A \vee B$ ) is logically equivalent to the conjunction of the negations of the individual statements  $A$  and  $B$ . The second law is similar, except we exchange conjunction for disjunction and vice-versa.

*Exercise* Verify the De Morgan law for disjunction by showing the logical equivalence of the left and right hand sides using a truth table:

*Solution*

| $\neg$ | $(A \vee B)$ | $\iff$ | $(\neg A) \wedge (\neg B)$ |
|--------|--------------|--------|----------------------------|
| $F$    | $T$          | $T$    | $F$                        |
| $F$    | $T$          | $T$    | $F$                        |
| $F$    | $F$          | $T$    | $T$                        |
| $T$    | $F$          | $T$    | $T$                        |

We note that for any value of statements  $A$  and  $B$ , the values under the column for  $\iff$  are all true, indicating that both expressions  $\neg(A \vee B)$ ,  $(\neg A) \wedge (\neg B)$  have the same truth tables, as required.

Another useful logical equivalence is the following “contrapositive” form of implication, which we explore in the next exercise:

*Exercise* Show that  $(A \implies B) \iff (\neg B \implies \neg A)$  is a tautology, i.e. this means that  $A \implies B$  is logically equivalent (LE) to  $\neg B \implies \neg A$ .

*Solution*

| $(A \implies B)$ | $\iff$ | $(\neg B \implies \neg A)$ |
|------------------|--------|----------------------------|
| $T$              | $T$    | $T$                        |
| $T$              | $F$    | $F$                        |
| $F$              | $T$    | $T$                        |
| $F$              | $F$    | $F$                        |

To show a situation where expressions are not logically equivalent, the following exercise may be considered:

*Exercise* Show  $A \implies B$  is not logically equivalent to  $B \implies A$

*Solution*

| $(A \implies B)$ | $\iff$ | $(B \implies A)$ |
|------------------|--------|------------------|
| $T$              | $T$    | $T$              |
| $T$              | $F$    | $F$              |
| $F$              | $T$    | $T$              |
| $F$              | $F$    | $F$              |

Notice the middle column  $\iff$  has false entries, so we don't have logical equivalency.

Another item of terminology is *Logical Implication*. Logical implication is when the expression  $A \implies B$  is a tautology. If the statements  $A$  and  $B$  are such that we only get true statements from  $A \implies B$ , then  $A$  logically implies (**LI**)  $B$ . This was the situation with the earlier illustration where we had defined statements  $A$  and  $B$  such that  $A = (a > 5), B = (a > 3)$  where  $a$  is some integer.

If  $A$  is logically equivalent (LE) to  $B$ , then  $A$  LI  $B$  and  $B$  LI  $A$ .

*Exercise:* Show  $A \wedge (A \implies B)$  logically implies  $B$ . This is called "Modus Ponens."

*Solution*

| $(A \wedge (A \implies B))$ | $\implies$ | $B$ |
|-----------------------------|------------|-----|
| $T$                         | $T$        | $T$ |
| $T$                         | $F$        | $F$ |
| $F$                         | $T$        | $T$ |
| $F$                         | $F$        | $F$ |

### 1.3 Zermelo-Fraenkel Axiomatic Set Theory

Set theory is at the foundation of mathematics. The reader should have some awareness of the axioms of set theory, although in practice it seems that one rarely sees a mathematical proof that explicitly cites these axioms. These axioms are referred to as the Zermelo-Fraenkel Axioms.

#### 1.3.1 The Zermelo-Fraenkel Axioms

The following so-called Zermelo-Fraenkel Axioms of set theory are presented to help provide the reader a more comprehensive picture of mathematics foundations. This section is inspired by the set of YouTube lectures by Richard Borchers (ZF Axioms)

([https://www.youtube.com/playlist?list=PL8yHsr3EFj52EKVgPi-p50fRP2\\_SbG2oi](https://www.youtube.com/playlist?list=PL8yHsr3EFj52EKVgPi-p50fRP2_SbG2oi)) and also the book *Axiomatic Set Theory* by Paul Bernays. The definitions are adapted from Bernays.

We note that in the axioms below, the notion of *set* is undefined, as is the inclusion relation,  $\in$ . Furthermore, all elements or members of sets are sets themselves.

##### *Axiom of Extensionality*

If  $s \subseteq t$  and  $t \subseteq s$ , then  $s = t$  where  $s, t$  are sets and  $\subseteq$  means *subset*.

We define subset by the following: If  $s$  and  $t$  are sets such that, for all  $x$ ,  $x \in s$  implies  $x \in t$ ,  $s$  is called a subset of  $t$ , where  $x$  is a set element or member. Occasionally, we might use the notation  $s \subset t$  to indicate a *proper subset* whereby there is an element in  $t$  that is not in  $s$ .

To recap: The axiom states what it means for two sets to be equal - namely they contain the same elements.

##### *Axiom of Foundation*

Every non-empty set  $s$  contains an element  $t$  such that  $s$  and  $t$  have no common element.

This axiom prevents an element of a set having an element which appears in the original set. For example, the set  $s = \{s\}$  is not allowed.

##### *Axiom of Pairing*

For any two different sets  $a$  and  $b$ , the pair  $\{a, b\}$ , or  $\{b, a\}$ , exists.

##### *Axiom of Union*

For any set  $s$  which contains at least two elements, there exists the set whose elements are the elements of the elements of  $s$ .

This set is called the union of the elements of  $s$  and is denoted by  $\cup s$ .

*Axiom of Infinity*

There exists at least one set  $W$  with the properties:

- a)  $\emptyset \in W$ , where  $\emptyset$  refers to the “empty set.” - the set with no elements.
- b) if  $x \in W$ , then  $\{x\} \in W$ .

The interpretation of the above is that a set exists with a non-ending sequence of elements. For example, if we think of the infinite sequence of non-negative integers being encoded as follows:  $0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$ , then we can have the corresponding infinite set  $S = \{\emptyset, \{\emptyset\}, \{\{\emptyset\}\}, \dots\}$  which is justified by the axiom.

*Axiom of Power-set*

For any set  $s$ , there exists the set whose elements are all subsets of  $s$ .

To take an example, consider the set  $s = \{1, 2, 3\}$ . Then the power-set is  $P = \{\{\}, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}$ . The number of elements, also called the *cardinality*, in this case is given by  $2^N$ , where  $N$  is the number of elements in the original set  $s$ . Thus, we have  $2^{N=3} = 8$ . We note that the empty set,  $\emptyset$ , is considered a subset of every set.

*Axiom of Subsets* (Note: Borchers refers to this as “separation”)

For any set  $s$  and any predicate  $P$  which is meaningful for all elements of  $s$ , there exists the set  $y$  that contains just those elements  $x$  of  $s$  which satisfy the predicate  $P$ .

For example, let  $s = \{1, 2, 3, 4, 5\}$  and the predicate  $P(x)$  be true whenever  $x$  is even. Then, we have the set  $y = \{x \in s \mid P(x) \text{ true}\} = \{2, 4\}$ .

*Axiom of Replacement*

For every set  $s$  and every single-valued function  $f$  of one argument which is defined for all the elements of  $s$ , there exists the set that contains all  $f(x)$  with  $x \in s$ .

*Axiom of Choice*

For every disjointed set  $t$  for which  $\emptyset \notin t$ , the Cartesian product  $\mathfrak{P}t$  differs from the  $\emptyset$ .

We define a *disjointed set* by the following: If a set  $s$  contains at least two elements, and any two elements of  $s$  are mutually exclusive (i.e. these elements have no common elements themselves).

**Theorem** The *Cartesian product* of the elements of a disjointed set  $t$ , denoted  $\mathfrak{P}t$  is the set whose elements are the sets which contain a single element

from each element of  $t$ . If  $a, b, \dots$  are the elements of  $t$ , the Cartesian product is also denoted by  $a \times b \times \dots$ . *Note:* For those interested, this theorem is proved in Bernays.

To illustrate the applications of the above axioms to a proof, consider the following exercise.

*Exercise* Complete the proof of the following theorem (adapted from Bernays):

**Theorem** For every non-empty set  $t$ , there exists the set whose elements are common to all elements of  $t$ .

This set is called the *intersection* of the elements of  $t$  and is denoted by  $\cap t$ .

**Proof:** By the Axiom of Union,  $s = \cup t$  exists and contains the elements of the elements of  $t$ . The predicate “ $x$  is contained in each element of  $t$ ” defines, by the Axiom of fill in the blank, a subset of  $s$  which is the intersection  $\cap t$ .

*Solution*

Axiom of **Subsets**

The following exercise explores the Cartesian product.

*Exercise* According to Bernays, for a disjointed set  $t$ , one has the Cartesian product,  $\mathfrak{P}t = \emptyset$  if  $\emptyset \in t$ . Prove this using the Cartesian Product Theorem stated above in Axiom of Choice section.

*Solution* According to the theorem, the Cartesian product is the set whose elements are the sets which contain a single element from each element of  $t$ . Since  $\emptyset \in t$  and contains no elements, the Cartesian product must be empty since there is no set possible that contains the required one element from  $\emptyset$ .

### 1.3.2 Set Operations

In this section, we present some set operations that are useful in much of the remainder of this book. The reader may be familiar with some of the material, although perhaps not necessarily in the formalism presented. Concepts and proofs in mathematics are very frequently expressed in terms of set language to be precise, so utility with set operations is important. For example, in later sections, we will discuss algebraic structures such as “groups” which will be defined in terms of sets with certain properties.

The emphasis here will be to continue the development following the axiomatic approach, largely adapted from *Axiomatic Set Theory* by Patrick Suppes. Hopefully, this will illustrate some of the notation and concepts we have been developing in the previous sections.

### Binary Relations

Definition:  $R$  is a binary relation  $\iff (\forall x)(x \in R \implies (\exists y)(\exists z)(x = \langle y, z \rangle))$ .

We note that in the above,  $\langle y, z \rangle$  is called an *ordered pair* and is formally defined in set notation as  $\langle y, z \rangle = \{\{y\}, \{y, z\}\}$ . Note how the ordering is encoded. So we have that a relation is a set of ordered pairs.

The definition is read as “ $R$  is a binary relation if and only if for all  $x$  if  $x$  is in  $R$ , then there exists a  $y$  and there exists a  $z$  such that  $x$  is equal to the ordered pair  $\langle y, z \rangle$ .”

It should be remembered that the phrases “if and only if” and the “if ... then” are used to refer to the truth tables for the corresponding symbols and not the casual english meaning. If desired, we could express the definition by using a truth table as we did previously to be completely clear.

Sometimes the following notation is used discussing relations:  $xRy$ . This is formally defined as  $xRy \iff \langle x, y \rangle \in R$ .

Consider the following example set which is a relation:

$$R = \{\langle 1, 2 \rangle, \langle 2, 1 \rangle, \langle 1, 3 \rangle, \langle 3, 1 \rangle, \langle 2, 3 \rangle, \langle 3, 2 \rangle\}.$$

We have utilized the angle bracket notation for ordered pairs for simplicity. Observe that this relation,  $R$ , has some special properties. To begin with, this relation is called *symmetric* because if  $xRy$  is true then we have  $yRx$  true. For example,  $\langle 1, 2 \rangle \in R$  and  $\langle 2, 1 \rangle \in R$ .

*Exercise* For the relation  $R$  defined above, verify that it is symmetric by explicitly listing the ordered pairs in question and their corresponding symmetric partners.

*Solution*

$\langle 1, 2 \rangle, \langle 2, 1 \rangle$   
 $\langle 2, 1 \rangle, \langle 1, 2 \rangle$   
 $\langle 1, 3 \rangle, \langle 3, 1 \rangle$   
 $\langle 3, 1 \rangle, \langle 1, 3 \rangle$   
 $\langle 2, 3 \rangle, \langle 3, 2 \rangle$   
 $\langle 3, 2 \rangle, \langle 2, 3 \rangle$

We consider as an exercise, the proof that  $\emptyset$  is a relation.

*Exercise* Prove that  $\emptyset$  (the empty set) is a relation by appealing to the definition of relation given earlier.

*Solution*

Recalling the definition:  $R$  is a binary relation  $\iff (\forall x)(x \in R \implies (\exists y)(\exists z)(x =$

$\langle y, z \rangle \rangle$ ), we substitute  $\emptyset$  for  $R$  and obtain:

$$\emptyset \text{ is a binary relation} \iff (\forall x)(x \in \emptyset \implies (\exists y)(\exists z)(x = \langle y, z \rangle)).$$

Now, since  $x \in \emptyset$  is false (the empty set contains no members), the statement  $(x \in \emptyset \implies (\exists y)(\exists z)(x = \langle y, z \rangle))$  is true. Recall the truth table for  $\implies$  whereby if the first term is false, the entire statement is true no matter what the second term is. Therefore, we have shown the  $\emptyset$  is a relation because it satisfies the definition.

### Ordering Relations

In the previous section, we introduced the idea of a *symmetric* relation. We now present a more complete list of such definitions:

- $R$  is reflexive in  $A \iff (\forall x)(x \in A \implies xRx)$ .
- $R$  is irreflexive in  $A \iff (\forall x)(x \in A \implies \neg(xRx))$ .
- $R$  is symmetric in  $A \iff (\forall x)(\forall y)(x, y \in A \wedge xRy \implies yRx)$ .
- $R$  is asymmetric in  $A \iff (\forall x)(\forall y)(x, y \in A \wedge xRy \implies \neg(yRx))$ .
- $R$  is antisymmetric in  $A \iff (\forall x)(\forall y)(x, y \in A \wedge xRy \wedge yRx \implies x = y)$ .
- $R$  is transitive in  $A \iff (\forall x)(\forall y)(\forall z)(x, y, z \in A \wedge xRy \wedge yRz \implies xRz)$ .
- $R$  is connected in  $A \iff (\forall x)(\forall y)(x, y \in A \wedge x \neq y \implies xRy \vee yRx)$ .
- $R$  is strongly connected in  $A \iff (\forall x)(\forall y)(x, y \in A \implies xRy \vee yRx)$ .

A couple of items regarding notation. The notation  $x, y \in A$  means  $x \in A \wedge y \in A$ . Also, by convention, the  $\wedge$  and  $\vee$  operations are performed before  $\implies$  which saves writing extra parentheses in the definitions. For example, if we explicitly included the parentheses in the definition of symmetric to indicate order of operations, it would appear as  $A(\forall x)(\forall y)((x, y \in A \wedge xRy) \implies yRx)$ .

If we consider the definition of *symmetric* above, we see how this corresponds to the example symmetric relation in the previous section. The reader should be cautioned to carefully observe the difference between *asymmetric* and *antisymmetric* relations.

*Exercise* Prove that the relation from the previous section is irreflexive by appealing to the definition. Define the set  $A = \{1, 2, 3\}$  for purposes of this exercise.

*Solution*

Recalling  $R$ :



$$R = \{\langle 1, 2 \rangle, \langle 2, 1 \rangle, \langle 1, 3 \rangle, \langle 3, 1 \rangle, \langle 2, 3 \rangle, \langle 3, 2 \rangle\}.$$

we see that  $\forall x \in A$ , namely 1, 2, 3, we are missing the corresponding ordered pairs  $\langle 1, 1 \rangle, \langle 2, 2 \rangle, \langle 3, 3 \rangle$  in the relation  $R$ . Now if we consider the definition of *irreflexive* we have:  $R$  is *irreflexive* in  $A \iff (\forall x)(x \in A \implies \neg(xRx))$ . Clearly,  $xRx$  is false for any choice of  $x \in A$ , thus we have  $\neg(xRx)$  true for all choices of  $x \in A$  and we see that  $R$  is therefore indeed *irreflexive*.

*Exercise* Show by counterexample, that the relation from the previous section is not *transitive*. Once again, consider the set  $A = \{1, 2, 3\}$  for purposes of this exercise.

*Solution*

Recalling the definition:  $R$  is *transitive* in  $A \iff (\forall x)(\forall y)(\forall z)(x, y, z \in A \wedge xRy \wedge yRz \implies xRz)$ , we see that if we can find an instance where  $x, y, z \in A \wedge xRy \wedge yRz \implies xRz$  is false, we are done. Consider  $1R2$  and  $2R1$  which are both true. However,  $1R1$  is false because  $\langle 1, 1 \rangle \notin R$ . Therefore, we conclude that  $R$  is not transitive.

The following two definitions are important:

- $R$  is a *partial ordering* of  $A \iff R$  is *reflexive*, *anti-symmetric* and *transitive* in  $A$ .
- $R$  *well-orders*  $A \iff R$  is *connected* in  $A \wedge (\forall B)(B \subseteq A \wedge B \neq \emptyset \implies B$  has an  $R$ -*minimal* element).

The definition of *R-minimal element* mentioned above is as follows:  $x$  is an *R-minimal element* of  $A \iff x \in A \wedge (\forall y)(y \in A \implies \neg(yRx))$ .

Let us examine the above two ordering definitions with some examples. Firstly, let us consider *partial ordering*. We will take  $A = \{1, 2, 3\}$  and construct an appropriate relation. We can begin with taking all possible ordered pairs which gives us the following provisional relation:

$$R = \{\langle 1, 1 \rangle, \langle 1, 2 \rangle, \langle 1, 3 \rangle, \langle 2, 1 \rangle, \langle 2, 2 \rangle, \langle 2, 3 \rangle, \langle 3, 1 \rangle, \langle 3, 2 \rangle, \langle 3, 3 \rangle\}$$

The above relation is symmetric, because it contains  $\langle 1, 1 \rangle, \langle 2, 2 \rangle, \langle 3, 3 \rangle$ . The anti-symmetric requirement restricts the ordered pair partners. For example, if we have  $\langle 1, 2 \rangle$  we cannot have  $\langle 2, 1 \rangle$ . So, let us remove those ordered pairs where the second element is smaller than the first:

$$R = \{\langle 1, 1 \rangle, \langle 1, 2 \rangle, \langle 1, 3 \rangle, \langle 2, 2 \rangle, \langle 2, 3 \rangle, \langle 3, 3 \rangle\}$$

Lastly, we need to verify transitivity:

$$1R1 \wedge 1R1 \implies 1R1$$

$$1R1 \wedge 1R2 \implies 1R2$$

$$\begin{aligned}
1R1 \wedge 1R3 &\implies 1R3 \\
1R2 \wedge 2R2 &\implies 1R2 \\
1R2 \wedge 2R3 &\implies 1R3 \\
1R3 \wedge 3R3 &\implies 1R3 \\
2R2 \wedge 2R2 &\implies 2R2 \\
2R2 \wedge 2R3 &\implies 2R3 \\
2R3 \wedge 3R3 &\implies 2R3 \\
3R3 \wedge 3R3 &\implies 3R3
\end{aligned}$$

So, our revised  $R$  is a partial ordering of the set  $A$ . When dealing with real numbers, the symbol  $\leq$  (less than or equal to) represents a partial ordering relation. The relation  $R$  we constructed above parallels this.

Now, let us do a similar construction to obtain a well-ordering. Again, consider the set  $A = \{1, 2, 3\}$  and let us construct a relation  $R$  that satisfies the definition. To begin with, we can take as provisional all possible ordered pairs:

$$R = \{\langle 1, 1 \rangle, \langle 1, 2 \rangle, \langle 1, 3 \rangle, \langle 2, 1 \rangle, \langle 2, 2 \rangle, \langle 2, 3 \rangle, \langle 3, 1 \rangle, \langle 3, 2 \rangle, \langle 3, 3 \rangle\}$$

Now, we first need to make sure that  $R$  is *connected* in  $A$  by appealing to the definition of connected:  $R$  is connected in  $A \iff (\forall x)(\forall y)(x, y \in A \wedge x \neq y \implies xRy \vee yRx)$ . We have as members of  $R$ ,  $\langle 1, 2 \rangle, \langle 1, 3 \rangle, \langle 2, 3 \rangle$ , so  $R$  is connected.

Next, we need to verify the second part of the definition:  $(\forall B)(B \subseteq A \wedge B \neq \emptyset \implies B \text{ has an } R\text{-minimal element})$ . There are seven subsets of  $A$  (we don't need to consider  $\emptyset$ ), which are listed here:

$$\begin{aligned}
&\{1\} \\
&\{2\} \\
&\{3\} \\
&\{1, 2\} \\
&\{1, 3\} \\
&\{2, 3\} \\
&\{1, 2, 3\}
\end{aligned}$$

For 1 to be an  $R$ -minimal in  $B$ , we must remove  $\langle 1, 1 \rangle$  from  $R$ . Similar, arguments apply to  $\langle 2, 2 \rangle$  and  $\langle 3, 3 \rangle$  corresponding to  $\{2\}$  and  $\{3\}$ . Now, for  $\{1, 2\}$  we must remove  $\langle 2, 1 \rangle$  from  $R$  so that 1 is  $R$ -minimal in  $\{1, 2\}$ , and likewise for  $\{1, 3\}$  we remove  $\langle 3, 1 \rangle$ . Similarly, we remove  $\langle 3, 2 \rangle$  corresponding to  $\{2, 3\}$ . Now,  $\{1, 2, 3\}$  has an  $R$ -minimal element already, so we are done. Our revised relation  $R$  is now:

$$R = \{\langle 1, 2 \rangle, \langle 1, 3 \rangle, \langle 2, 3 \rangle\}$$

This revised relation  $R$  well-orders  $A$ . This parallels the  $<$  (less than) relation in real numbers.

*Exercise* There is a theorem in Suppes (Theorem 62) as follows:  $R$  well-orders  $A \implies R$  is asymmetric and transitive in  $A$ . Verify the relation  $R$  above satisfies this theorem.

*Solution.* We first verify that  $R$  is asymmetric. We have  $\langle 1, 2 \rangle$  but not  $\langle 2, 1 \rangle$ ,  $\langle 1, 3 \rangle$  but not  $\langle 3, 1 \rangle$ , and lastly  $\langle 2, 3 \rangle$  but not  $\langle 3, 2 \rangle$ . Now, we verify transitivity: The only statement that makes sense is  $1R2 \wedge 2R3 \implies 1R3$  which is true. Therefore, we have verified that  $R$  satisfies the theorem.

### Equivalence Relations

We now discuss some important concepts that will appear in some form later, notably in abstract algebra. The reader will benefit by getting comfortable with these topics as early as possible, so we present them here.

- Definition:  $R$  is an equivalence relation  $\iff R$  is a relation  $\wedge R$  is reflexive, symmetric, and transitive.
- Definition: the  $R$ -coset of  $x$  is defined by  $R[x] = \{y | xRy\}$ .
- Definition:  $\Pi$  is a partition of  $A \iff \cup \Pi = A \wedge (\forall B)(\forall C)(B \in \Pi \wedge C \in \Pi \wedge B \neq C \implies B \cap C = \emptyset) \wedge (\forall x)(x \in \Pi \implies (\exists y)(y \in x))$ .

A *partition* of a set is an exhaustive, mutually exclusive decomposition of a set into subsets, whose union is the set itself. For example, if our set is  $A = \{1, 2, 3\}$ , then a partition might be  $\Pi = \{\{1\}, \{2\}, \{3\}\}$  or perhaps  $\Pi = \{\{1\}, \{2, 3\}\}$ .

We can see the partitioning through an example. Let us consider a set  $A = \{1, 2, 3, 4, 5\}$  and construct an equivalence relation,  $R$ . Firstly, we know that  $R$  must be reflexive, so we provisionally begin with:

$$R = \{\langle 1, 1 \rangle, \langle 2, 2 \rangle, \langle 3, 3 \rangle, \langle 4, 4 \rangle, \langle 5, 5 \rangle\}$$

We could actually stop here, because  $R$  happens to also be symmetric and transitive. However, to make the example slightly more complex, let us add some more members that are symmetric:

$$R = \{\langle 1, 1 \rangle, \langle 1, 2 \rangle, \langle 2, 1 \rangle, \langle 2, 2 \rangle, \langle 3, 3 \rangle, \langle 4, 4 \rangle, \langle 5, 5 \rangle\}$$

where it can be seen that  $\langle 1, 2 \rangle$  and  $\langle 2, 1 \rangle$  have been added.

Now we need to verify that transitivity is satisfied:

$$1R1 \wedge 1R1 \implies 1R1$$

$$1R1 \wedge 1R2 \implies 1R2$$

$$1R2 \wedge 2R1 \implies 1R1$$

$$1R2 \wedge 2R2 \implies 1R2$$

$$2R1 \wedge 1R1 \implies 2R1$$

$$2R1 \wedge 1R2 \implies 2R2$$

$$2R2 \wedge 2R1 \implies 2R1$$

$$\begin{aligned}
2R2 \wedge 2R2 &\implies 2R2 \\
3R3 \wedge 3R3 &\implies 3R3 \\
4R4 \wedge 4R4 &\implies 4R4 \\
5R5 \wedge 5R5 &\implies 5R5
\end{aligned}$$

Now, let us determine the  $R$ -coset of  $x$  for each  $x \in A$ :

$$\begin{aligned}
R[1] &= \{1, 2\} \\
R[2] &= \{1, 2\} \\
R[3] &= \{3\} \\
R[4] &= \{4\} \\
R[5] &= \{5\}
\end{aligned}$$

Now, we can form a set of all the cosets, which will be a partition of  $A$ :  
 $\Pi = \{\{1, 2\}, \{3\}, \{4\}, \{5\}\}.$

### Functions

To help motivate the discussion of *functions*, which presumably many readers will have encountered previously and have a fair degree of experience with, within the set theoretical framework, the following directly quoted passage from Suppes (p.86) may be helpful:

“Since the eighteenth century, clarification and generalization of the concept of a function have attracted much attention. Fourier’s representation of ‘arbitrary’ functions (actually piecewise continuous ones) by trigonometric series encountered much opposition; and later when Weierstrass and Riemann gave examples of continuous functions without derivatives, mathematicians refused to consider them seriously. Even today many textbooks of the differential and integral calculus do not give a mathematically satisfactory definition of functions. An exact and completely general definition is immediate within our set-theoretical framework. A function is simply a many-one relation, that is, a relation which to any element in its domain relates exactly one element in its range.”

We now present the formal definition:

Definition:  $f$  is a function  $\iff f$  is a relation  $\wedge (\forall x)(\forall y)(\forall z)(x f y \wedge x f z \implies y = z).$

Here is an example from Suppes which we use as an exercise.

*Exercise* Argue that the following relation is not a function:  $f = \{\langle 1, 1 \rangle, \langle 1, 2 \rangle, \langle 3, 4 \rangle\}.$

*Solution:*

Appealing to the definition of function, we see that the statement  $1 f 1 \wedge 1 f 2 \implies 1 = 2$  is false, therefore we conclude  $f$  is not a function. In other words, the

ordered pairs  $\langle 1, 1 \rangle$  and  $\langle 1, 2 \rangle$  which appear in  $f$  have the same first element, 1, but different second elements, 1 and 2, which is not allowed for a function.

*Composition* of functions is an important concept that appears quite frequently in some of the later topics. The notation for this is  $(f \circ g)(x) = f(g(x))$ , where  $f$  and  $g$  are functions,  $x$  is an element of a set, and the circle indicates composition. In words, we can think of this as a chain, whereby, firstly the mapping is applied using the  $g$  function, then the result of this is mapped via the  $f$  function. Suppes states a formal definition of composition using something called the *relative product*, so for completeness we provide the definitions correspondingly:

Definition:  $f \circ g = g/f$ .

The relative product is given by the notation  $g/f$  and is defined by the following:

Definition:  $g/f = \{\langle x, y \rangle \mid (\exists z)(xgz \wedge zfy)\}$ .

*Exercise* Given the two functions  $f = \{\langle 3, 2 \rangle, \langle 4, 5 \rangle, \langle 6, 7 \rangle\}$  and  $g = \{\langle 1, 3 \rangle, \langle 2, 6 \rangle\}$ , determine the composition  $f \circ g$  using the definition  $g/f$ .

*Solution:*

Let us pair up the terms  $xgz$  and  $zfy$  (in that order), where  $z$  is common to both:

$\langle 1, 3 \rangle, \langle 3, 2 \rangle$   
 $\langle 2, 6 \rangle, \langle 6, 7 \rangle$

Then we have  $g/f = \{\langle 1, 2 \rangle, \langle 2, 7 \rangle\}$ .

*Exercise* Prove the following theorem (which is actually part of “Theorem 82” in Suppes):  $f$  and  $g$  are functions  $\implies f \circ g$  is a function .

*Solution:*

From the definition of a function provided earlier, we need to establish that  $h = f \circ g$  is a relation, and also prove that  $(\forall x)(\forall y)(\forall z)(xhy \wedge xhz \implies y = z)$ . For the first part, we appeal to the definition  $f \circ g = g/f$  and subsequently note that the definition of  $g/f$  is a set of ordered pairs, which is a (binary) relation, also by definition.

For the second part, we need to show:  $(\forall x)(\forall y)(\forall z)(xhy \wedge xhz \implies y = z)$ . Now consider  $xhy$ . From the definition of relative product, this means that  $(\exists w)(xgw \wedge wfy)$ , and likewise for  $xhz$  this means that  $(\exists w')(xgw' \wedge w'fz)$ . Rewriting the hypothesis, we obtain:  $(\forall x)(\forall y)(\forall z)((\exists w)(xgw \wedge wfy) \wedge (\exists w')(xgw' \wedge w'fz))$ . Now, since  $g$  is a function, this implies that  $w = w'$ . Then, since  $f$  is a function and  $w = w'$ , this implies  $y = z$  thereby completing the proof.

We now define the concept of a converse relation, denoted  $\widetilde{R}$ , which will be

immediately useful in what comes next.

Definition:  $\widetilde{R} = \{\langle x, y \rangle | yRx\}$ , where  $R$  is a relation.

The following definitions are key:

- Definition:  $f$  is 1-1  $\iff f$  and  $\widetilde{f}$  are functions .
- Definition:  $f$  is 1-1  $\implies f^{-1} = \widetilde{f}$ , where  $f^{-1}$  is the *inverse* of  $f$ .

We note that “1-1” is read “one-to-one.” We now illustrate a counter-example to a 1-1 function.

*Exercise* Show that the following function,  $f = \{\langle 1, 2 \rangle, \langle 2, 2 \rangle, \langle 3, 4 \rangle\}$  is not 1-1.

*Solution:* Let us first determine the converse relation to  $f$ :  $\widetilde{f} = \{\langle 2, 1 \rangle, \langle 2, 2 \rangle, \langle 4, 3 \rangle\}$ . We see that this relation is not a function because the elements  $\langle 2, 1 \rangle, \langle 2, 2 \rangle$  have the same first entry, but different second entries. Therefore, we conclude  $f$  is not 1-1.

We now introduce some additional terminology:

- Definition: Domain of a relation  $R$ :  $\mathcal{D}R = \{x | (\exists y)(xRy)\}$
- Definition: Range of a relation  $R$ :  $\mathcal{R}R = \{y | (\exists x)(xRy)\}$ . This is also referred to as the *counterdomain* or *converse domain*.
- Definition:  $f$  is a function from set  $A$  into set  $B \iff f$  is a function  $\wedge \mathcal{D}f = A \wedge \mathcal{R}f \subseteq B$ .
- Definition:  $f$  is a function from set  $A$  onto set  $B \iff f$  is a function  $\wedge \mathcal{D}f = A \wedge \mathcal{R}f = B$ .
- Definition: *Restriction* of the domain of a relation  $R$  to a given set  $A$ :  $R|A = R \cap (A \times \mathcal{R}R)$ .
- Definition: the *image* of the set  $A$  under  $R$ :  $R^{\circ}A = \mathcal{R}(R|A)$ .

*Exercise* What is the image of  $A = \{1, 2, 4, 6\}$  under  $R = \{\langle 1, 10 \rangle, \langle 2, 11 \rangle, \langle 5, 15 \rangle\}$ ?

*Solution:* First, let us determine the restriction  $R|A$ . We need to obtain the Cartesian product  $A \times \mathcal{R}R$  which is:

$$\{\langle 1, 10 \rangle, \langle 1, 11 \rangle, \langle 1, 15 \rangle, \langle 2, 10 \rangle, \langle 2, 11 \rangle, \langle 2, 15 \rangle, \langle 4, 10 \rangle, \langle 4, 11 \rangle, \langle 4, 15 \rangle, \langle 6, 10 \rangle, \langle 6, 11 \rangle, \langle 6, 15 \rangle\}$$

Next, we determine the intersection with  $R$ :

$$R|A = R \cap (A \times \mathcal{R}R) = \{\langle 1, 10 \rangle, \langle 2, 11 \rangle\}$$

Lastly, we obtain the image of  $A$  under  $R$ :

$$R^{\ast}A = \mathcal{R}(R|A) = \{10, 11\}$$

It should be kept in mind that we are always dealing with sets when we refer to functions now and later in the material, so if there is any confusion, the reader should always consult the pertinent definitions to see exactly what sets or subsets are being referenced in any manipulations or results. For example, does a particular operation involve the entire set or only some portion of a set. Typically, engineers and scientists may normally think of functions in terms of graphs on a say two-dimensional plot, but in the pure mathematical context, keeping in mind the underlying set framework as has been outlined in the above material is important.

## 1.4 Peano Axioms of Arithmetic

In the previous section, we built up a basic framework for set theory based on the Zermelo-Frankel axioms. We now establish the foundations of the natural numbers, founded on top of that set theory, by stating the Peano Axioms. Much of this material is based directly on *Schaum's Outline of Theory and Problems of Abstract Algebra* by Frank Ayres and Lloyd R. Jaisingh.

Let there exist a non-empty set  $\mathbb{N}$  such that:

- Axiom I:  $1 \in \mathbb{N}$ .
- Axiom II: For each  $n \in \mathbb{N}$  there exists a unique  $n^* \in \mathbb{N}$ , called the *successor* of  $n$ .
- Axiom III: For each  $n \in \mathbb{N}$  we have  $n^* \neq 1$ .
- Axiom IV: If  $m, n \in \mathbb{N}$  and  $m^* = n^*$ , then  $m = n$ .
- Axiom V: Any subset  $K$  of  $\mathbb{N}$  having the properties (a)  $1 \in K$ , (b)  $k^* \in K$  whenever  $k \in K$  is equal to  $\mathbb{N}$ .

We now define addition and multiplication on  $\mathbb{N}$ .

Addition:

- (i)  $n + 1 = n^*$ , for every  $n \in \mathbb{N}$
- (ii)  $n + m^* = (n + m)^*$ , whenever  $n + m$  is defined.

We then have the following laws for addition. For all  $m, n, p \in \mathbb{N}$ :

- Closure:  $n + m \in \mathbb{N}$
- Commutative:  $n + m = m + n$

- Associative:  $m + (n + p) = (m + n) + p$
- Cancellation: If  $m + p = n + p$ , then  $m = n$

*Exercise* Suppose we have two natural numbers,  $n, m$ , whereby  $n = 1, m = 1^*$ . For this restricted case, prove the commutative law for addition:  $n + m = m + n$ .

*Solution:*

We have that  $n + m = 1 + 1^*$ . Now, from the definition of addition (ii), we obtain  $1 + 1^* = (1 + 1)^*$ . From the definition of addition (i), we obtain  $(1 + 1)^* = (1^*)^*$ . Finally,  $(1^*)^* = 1^* + 1$  from the definition of addition (i) which is equal to  $m + n$ , thus completing the proof.

Multiplication:

- (i)  $n \cdot 1 = n$
- (ii)  $n \cdot m^* = n \cdot m + n$ , whenever  $n \cdot m$  is defined.

Similar to addition, we have laws for multiplication. For all  $m, n, p \in \mathbb{N}$ :

- Closure:  $n \cdot m \in \mathbb{N}$
- Commutative:  $m \cdot n = n \cdot m$
- Associative:  $m \cdot (n \cdot p) = (m \cdot n) \cdot p$
- Cancellation: If  $m \cdot p = n \cdot p$ , then  $m = n$ .

If we consider both addition and multiplication, there are so-called distributive laws. For all  $m, n, p \in \mathbb{N}$ :

- $D_1$ :  $m \cdot (n + p) = m \cdot n + m \cdot p$
- $D_2$ :  $(n + p) \cdot m = n \cdot m + p \cdot m$ .

The reader should be cautioned or reminded at this point that the order of the variables and operations in laws such as in the above are important to observe in general. With certain mathematical objects that we will consider, such as *groups*, swapping the positions of variables may not necessarily yield the same results. Therefore, one must carefully consider the definitions relevant to each mathematical object in question.

*Exercise* How are we justified in asserting that  $m \cdot n = n \cdot m$ , when  $m, n \in \mathbb{N}$ ?

*Solution:* The commutative law for multiplication under  $\mathbb{N}$  provides the justification.

*Exercise* Suppose we have  $m, n \in \mathbb{N}$  and form the product  $p = n \cdot m$ . Is the result  $p \in \mathbb{N}$ ?



*Solution:* Yes. The closure law for multiplication under  $\mathbb{N}$  provides the justification.

Note in the above exercises, even though we may have some intuitive sense for the reasons, we must appeal to an exact statement which justifies the assertion. This may be an axiom, definition, theorem, law etc. which has been established for the objects in question. This habit will be useful to form as we consider other contexts in pure mathematics.

## 1.5 Gödel Theorems

To provide a more complete picture to the reader regarding the underpinnings of mathematics and logic, it is useful to consider the important results by Gödel which are cited in this context. The section is largely adapted from entries in *Stanford Encyclopedia of Philosophy* (<https://plato.stanford.edu/entries/goedel/>) and *Logic for Mathematicians* by A.G. Hamilton.

There are three theorems that we will consider. The first is called “the Completeness Theorem”, the second is “the First Incompleteness Theorem”, and the third is “The Second Incompleteness Theorem.”

We begin with some definitions adapted from (<https://plato.stanford.edu/entries/goedel-incompleteness/>):

- *Formal system:* (Roughly) A system of axioms equipped with rules of inference, which allow one to generate new theorems.
- *Complete:* A formal system for which every statement of the system, either the statement or its negation can be derived (“proved”) is complete.
- *Consistent:* A formal system for which there is no statement such that the statement and its negation are both derivable is consistent.

### The Completeness Theorem

*Every valid logical expression is provable. Equivalently, every logical expression is either satisfiable or refutable.*

An illustration of how this works is provided in the Stanford reference mentioned earlier, which is reproduced here. Suppose we have a logical expression  $\phi = \forall x_0 \exists x_1 \psi(x_0, x_1)$ . We can transform that expression with quantifiers into a list of expressions without quantifiers:

$$\begin{aligned}
\phi_0 &= \psi(x_0, x_1) \\
\phi_1 &= \psi(x_0, x_1) \wedge \psi(x_1, x_2) \\
&\dots \\
\phi_n &= \psi(x_0, x_1) \wedge \dots \wedge \psi(x_n, x_{n+1})
\end{aligned}$$

Then, there are two cases:

Case 1: For some  $n$ ,  $\phi_n$  is not satisfiable. This implies the original  $\phi$  is refutable (its negation is provable).

Case 2: Each  $\phi_n$  is satisfiable. This implies the original  $\phi$  is satisfiable.

The Stanford reference states that “logical expression” in the terminology of the theorem is “a well-formed first order formula without identity.” Gödel’s 1929 dissertation *On the Completeness of the Calculus of Logic* which is the source of the theorem, states the following:

“The main object of the following investigations is the proof of the completeness of the axiom system for what is called the restricted functional calculus, namely the system given in *Whitehead and Russell 1910*, Part I, \*1 and \*10, and, in a similar way, in *Hilbert and Ackermann 1928* (hereafter cited as H.A.) , III, §5.” (Source: Kurt Gödel *Collected Works* Vol.1 by Feferman, et al. (1986) - Page 61.)

The notation and axioms for this system are then specified in the same reference. Briefly, the logic symbols are the same we have seen previously, namely:  $\vee, \wedge, \implies, \iff, \neg$  and the axioms are as follows:

- (1)  $X \vee X \implies X$
- (2)  $X \implies X \vee Y$
- (3)  $X \vee Y \implies Y \vee X$
- (4)  $(X \implies Y) \implies (Z \vee X \implies Z \vee Y)$
- (5)  $(\forall x)F(x) \implies F(y)$
- (6)  $(\forall x)[X \vee F(x)] \implies X \vee (\forall x)F(x)$

The notation  $F(x)$  above refers to a “predicate” that can be true or false depending on the value of  $x$ . For example,  $F(x)$  can be true if  $x \in A$ , where  $A$  is some set.

There are also rules of inference, one of which is the “modus ponens” introduced previously. Namely,  $(p \wedge (p \implies q)) \implies q$  where  $p$  and  $q$  are statements that can be true or false.

*Exercise* In the text of Gödel’s dissertation, it is stated that  $\implies$  is used as an abbreviation. Also, it is implied that only  $\vee$  and  $\neg$  are necessary in order to represent the needed logic. Substantiate this by showing that it is possible to obtain the truth table for  $\implies$  only using  $\vee$  and  $\neg$ .

*Solution:*

Recalling the truth table for  $\implies$ :

| $A$ | $\implies$ | $B$ |
|-----|------------|-----|
| $T$ | $T$        | $T$ |
| $T$ | $F$        | $F$ |
| $F$ | $T$        | $T$ |
| $F$ | $T$        | $F$ |

we can see by inspection that if we negate the  $A$  column, and use  $\vee$  instead of  $\implies$  we will reproduce the same truth table. To confirm, let us do this explicitly:

| $(\neg A)$ | $\vee$ | $B$ |
|------------|--------|-----|
| $F$        | $T$    | $T$ |
| $F$        | $F$    | $F$ |
| $T$        | $T$    | $T$ |
| $T$        | $F$    | $F$ |

*Exercise* Is the following logical expression provable (within the above axiom system):  $(X \implies Y) \vee (X \implies Y) \implies (X \implies Y)$ ?

*Solution:* Yes, according to Gödel’s Completeness Theorem, since the formula is valid (taking “valid” to mean true for any combination of true or false for the variables  $X$  and  $Y$ ). In fact, in this case it was obtained from Axiom (1) by substituting  $X \implies Y$  for  $X$ , immediately verifying the completeness theorem.

It is worth emphasizing the following point, raised by Gödel in his dissertation. Namely, since the logical axioms are valid and the rules of inference preserve truth, this means that every provable formula (i.e. derived from the axioms using the rules of inference) is valid. Gödel’s Completeness Theorem actually states the converse: *Every valid logical expression is provable*. Again, this is within the specific system described above.

### The First Incompleteness Theorem

To introduce this section and the next, the following quote from the *Stanford Encyclopedia of Philosophy* entry on Gödel (<https://plato.stanford.edu/entries/goedel/>) may be illuminating:

“The First Incompleteness Theorem provides a counterexample to completeness by exhibiting an arithmetic statement which is neither provable nor refutable in Peano arithmetic, though true in the standard model. The Second Incompleteness Theorem shows that the consistency of arithmetic cannot be proved

in arithmetic itself.”

It is hoped that the previous sections on Peano Axioms and the Completeness Theorem will provide a good starting point for considering these incompleteness theorems.

We now present the First Incompleteness Theorem, based on the entry in the *Stanford Encyclopedia of Philosophy* for Gödel Incompleteness (<https://plato.stanford.edu/entries/goedel-incompleteness/>):

*Any consistent formal system  $F$  within which a certain amount of elementary arithmetic can be carried out is incomplete; i.e., there are statements of the language of  $F$  which can neither be proved nor disproved in  $F$ .*

At this point, it is useful to introduce the concept of *Gödel numbering* (ref: <https://plato.stanford.edu/entries/goedel-incompleteness/sup1.html>), which provides a procedure for encoding components of a formal system as natural numbers. The idea is that certain statements can be represented by a number and reasoned about as a number. For example, the statement (Axiom (1) from previous section)  $X \vee X \implies X$  might be represented as a natural number like 1329487 and so forth.

To see how Gödel numbers are determined in an unambiguous fashion, we proceed in two steps. Firstly, each symbol of the “language” of the formal system is given a natural number code. For example: the symbol ‘0’ might be assigned code number 1, the plus sign ‘+’ might be 3, the equals sign, ‘=’ might be 4, a variable  $x_1$  might be 13, and  $>$  is 14.

The next step is to encode a particular formula, which is a sequence of code numbers, into a final encoded natural number. This is done (apparently following Gödel’s original method), by leveraging the prime numbers and the fundamental theorem of arithmetic, which states that every natural number greater than one has a unique prime factorization (up to order of the factors). This means that a given sequence of symbols will have a unique representation in this prime encoding scheme.

Let us represent the primes as  $p_1, p_2, p_3, \dots, p_n, \dots$ . The first few primes in order are: 2, 3, 5, 7, 11. The next step is to raise each prime to the power of the corresponding symbol code at that position in the symbol sequence. For example, let us take the symbol sequence:  $0 + 0 = 0$ . The code sequence for this, using our above mapping, is: 1, 3, 1, 4, 1. To do the final encoding with the primes, we compute  $p_1^1 \times p_2^3 \times p_3^1 \times p_4^4 \times p_5^1 = 2^1 \times 3^3 \times 5^1 \times 7^4 \times 11^1 = 2 \times 27 \times 5 \times 2401 \times 11 = 7130970$ . So this final result, 7130970, would be the Gödel number for the formula  $0 + 0 = 0$ .

*Exercise* Using the above symbol encoding, give an argument for why just representing a given formula by the symbol code sequence is not enough to guarantee a unique Gödel number.

*Solution:* Consider the formulas  $0 + 0 = 0$  vs.  $x_1 > 0$ . Supposing that both of these are meaningful formulas, we can see that the code sequence for both would be the same, namely: 13141. Therefore, if we left the encoding this way, given the number 13141 we would not be able to determine which of those two formulas was being represented. However, if we do the second step of the process of encoding involving the primes, we would obtain for  $x_1 > 0$  the following Gödel number:  $2^{13} \times 3^{14} \times 5^1 = 8192 \times 4782969 \times 5 = 195910410240$ , which is obviously distinct from 7130970.

Now we have established the concept of Gödel Numbering, we can continue the discussion. In particular, in addition to assigning Gödel numbers to individual formulas, we can imagine assigning a Gödel number to the proof of a formula. If we think of a proof as a sequence of formulas, starting from axioms, applying the inference rules and other logical operations defined in the formal system, we can convert this proof to a Gödel number. For example, taking the *modus ponens* inference rule, suppose we might have something like this  $X, X \implies Y, Y, (\neg Y) \implies Z$ , where we are trying to prove  $(\neg Y) \implies Z$ . Suppose that  $X$  and  $X \implies Y$  are our axioms, then the above sequence could be considered a proof of  $(\neg Y) \implies Z$ .

As we did before, we can convert each symbol of the eleven symbol sequence to its code number, say: 123, 123, 11, 125, 125, 15, 13, 125, 17, 11, 127 and then encode to a Gödel number using the primes:  $2^{123} \times 3^{123} \times \dots p_{11}^{127}$ , where  $p_{11}$  is the eleventh prime, 31.

With this in mind, we can then construct the formula which Gödel showed could not be proved or disproved in the particular formal system at hand. Following the Hamilton logic book reference, we define a formula  $\mathcal{U}$  in words as follows: for every natural number  $n$ ,  $n$  is not the Gödel number of a proof of the formula  $\mathcal{U}$ . To clarify, Gödel's theorem states that  $\mathcal{U}$  is not a theorem in the formal system, and neither is  $\neg(\mathcal{U})$ .

Another concept that is useful in this context is *recursive functions*. For those with experience in computer science and software development, the idea of recursive functions may be familiar. According to the Hamilton reference, recursive functions can be constructed from three basic function and three rules which we provide below. Note that the convention used in Hamilton's reference that the set of natural numbers starts with 0 as opposed to 1, and the symbol  $D_N$  is used to represent this set of natural numbers starting from 0. The symbol  $D_N^k$  then represents a  $k$ -dimensional cartesian product set of natural numbers. For example,  $D_N^2 = \{\langle 0, 0 \rangle, \langle 1, 0 \rangle, \langle 2, 0 \rangle, \dots\}$ , and then  $\langle 2, 3 \rangle$  and  $\langle 5, 11 \rangle$  are two particular elements of that set.

The basic functions are:

- 1. The zero function  $z : D_N \rightarrow D_N$ , given by  $z(n) = 0$  for every  $n \in D_N$ . The notation  $z : D_N \rightarrow D_N$  is read as “ $z$  is the function which maps

elements in  $D_N$  to  $D_N$ ". Note that the arrow is not the same and the logical implication  $\implies$  previously discussed. So in this case, the function  $z$  maps (i.e. connects) every natural number in  $D_N$  to the natural number 0.

- 2. The successor function  $s : D_N \rightarrow D_N$ , given by  $s(n) = n + 1$  for every  $n \in D_N$ .
- 3. The projection functions  $p_i^k : D_N^k \rightarrow D_N$ , given by  $p_i^k(n_1, \dots, n_k) = n_i$ , for every  $n_1, \dots, n_k \in D_N$ .

The rules are as follows:

- R1. Composition. If  $g : D_N^j \rightarrow D_N$  and  $h_i : D_N^k \rightarrow D_N$  for  $1 \leq i \leq j$ , then  $f : D_N^k \rightarrow D_N$ , defined by  $f(n_1, \dots, n_k) = g(h_1(n_1, \dots, n_k), \dots, h_j(n_1, \dots, n_k))$  is obtained by composition from  $g$  and  $h_1, \dots, h_j$ .
- R2. Recursion. If  $g : D_N^k \rightarrow D_N$  and  $h : D_N^{k+2} \rightarrow D_N$ , then the function  $f : D_N^{k+1} \rightarrow D_N$ , defined by  $f(n_1, \dots, n_k, 0) = g(n_1, \dots, n_k)$ , and  $f(n_1, \dots, n_k, n+1) = h(n_1, \dots, n_k, n, f(n_1, \dots, n_k, n))$  is said to be obtained by recursion from  $g$  and  $h$ . We can also have recursion with all the  $n_i$  absent, as in  $f(0) = a$  and  $f(n+1) = h(n, f(n))$ , where  $a$  is a fixed member of  $D_N$ .
- R3. Least Number Operator. Let  $g : D_N^{k+1} \rightarrow D_N$  be any function with the property that for each  $n_1, \dots, n_k \in D_N$  there is at least one  $n \in D_N$  such that  $g(n_1, \dots, n_k, n) = 0$ . Then the function  $f : D_N^k \rightarrow D_N$ , defined by  $f(n_1, \dots, n_k) = \text{least number } n \in D_N \text{ such that } g(n_1, \dots, n_k, n) = 0$ , is said to be obtained from  $g$  by the use of the least number operator.

Let us consider some examples to illustrate the above functions and rules.

Taking the projection function  $p_2^3$ , we can apply it to  $\langle 4, 6, 9 \rangle$  and obtain  $p_2^3(4, 6, 9) = 6$ . This is the second value in the triplet  $\langle 4, 6, 9 \rangle$  extracted (or "projected").

Now, moving to composition, take  $g(n_1, n_2, n_3) = n_1 n_2 + n_3$  and  $h_1(m_1, m_2) = m_1 + m_2$ ,  $h_2(m_1, m_2) = m_1 m_2$ ,  $h_3(m_1, m_2) = m_1 + 2m_2$ . Then, the composition is  $f(m_1, m_2) = g(h_1(m_1, m_2), h_2(m_1, m_2), h_3(m_1, m_2)) = (m_1 + m_2)(m_1 m_2) + (m_1 + 2m_2)$ .

Lastly, let us consider recursion. Let  $h(m_1, m_2) = m_1 + m_2$  and  $f(0) = 1$ . Then,  $f(1) = h(0, f(0)) = 0 + 1 = 1$ ,  $f(2) = h(1, f(1)) = 1 + 1 = 2$ ,  $f(3) = h(2, f(2)) = 2 + 2 = 4$ ,  $f(4) = h(3, f(3)) = 3 + 4 = 7$ , and so forth.

We now show that the "addition function", which is one of the basic operators in our formal system for arithmetic, is recursive. In other words, it can be constructed from the above basic functions and finite application of the rules.

We show how this example (taken from Hamilton) is done.

We want to show that the addition function  $f_1(p, q) = p + q$  can be constructed as recursive. If we take the projection functions  $p_1^1, p_3^3$ , and the successor function  $s$  we can form the following:

$$\begin{aligned} f_1(m, 0) &= p_1^1(m) \\ f_1(m, n+1) &= s(p_3^3(m, n, f_1(m, n))) \end{aligned}$$

Let's check this for  $f_1(2, 3)$  which we anticipate should be  $2 + 3 = 5$ . We start by working backwards from the final function:

$$\begin{aligned} f_1(2, 3) &= s(p_3^3(2, 2, f_1(2, 2))) \\ f_1(2, 2) &= s(p_3^3(2, 1, f_1(2, 1))) \\ f_1(2, 1) &= s(p_3^3(2, 0, f_1(2, 0))) \\ f_1(2, 0) &= p_1^1(2) = 2 \end{aligned}$$

We have reached the “base case” and now we can evaluate the function working in the forward direction:

$$\begin{aligned} f_1(2, 0) &= p_1^1(2) = 2 \\ f_1(2, 1) &= s(p_3^3(2, 0, f_1(2, 0))) = s(p_3^3(2, 0, 2)) = 3 \\ f_1(2, 2) &= s(p_3^3(2, 1, f_1(2, 1))) = s(p_3^3(2, 1, 3)) = 4 \\ f_1(2, 3) &= s(p_3^3(2, 2, f_1(2, 2))) = s(p_3^3(2, 2, 4)) = 5 \end{aligned}$$

*Exercise* Show how to construct the multiplication function,  $f_2(m, n) = mn$  as a recursive, using the addition function,  $f_1$  defined above recursively, as well as the appropriate projection functions.

*Solution* If we consider what  $mn$  means as an addition problem, this can give us a clue how to proceed. Namely, we know that  $mn = \overbrace{m + m + \dots + m}^n$ . So, if we start with a base case of 0, and then add  $m$  to that base value  $n$  times, we should get the desired result. Putting this into the recursive format, and leveraging the addition function,  $f_1$  defined earlier, we obtain:

$$\begin{aligned} f_2(m, 0) &= z(m) = 0, \text{ where } z \text{ is the zero function defined earlier} \\ f_2(m, n+1) &= f_1(p_3^3(m, n, f_2(m, n)), p_1^1(m, n, f_2(m, n))) \end{aligned}$$

Let's check this for a simple case to see if it works. Take  $f_2(3, 5)$  which we expect should be 15. As before let us start by working backwards from the final function:

$$\begin{aligned} f_2(3, 5) &= f_1(f_2(3, 4), 3) \\ f_2(3, 4) &= f_1(f_2(3, 3), 3) \\ f_2(3, 3) &= f_1(f_2(3, 2), 3) \\ f_2(3, 2) &= f_1(f_2(3, 1), 3) \\ f_2(3, 1) &= f_1(f_2(3, 0), 3) \\ f_2(3, 0) &= 0 \end{aligned}$$

We have reached the “base case”, so now we can evaluate the function working in the forward direction:

$$\begin{aligned} f_2(3, 0) &= 0 \\ f_2(3, 1) &= f_1(f_2(3, 0), 3) = 0 + 3 = 3 \\ f_2(3, 2) &= f_1(f_2(3, 1), 3) = 3 + 3 = 6 \\ f_2(3, 3) &= f_1(f_2(3, 2), 3) = 6 + 3 = 9 \\ f_2(3, 4) &= f_1(f_2(3, 3), 3) = 9 + 3 = 12 \\ f_2(3, 5) &= f_1(f_2(3, 4), 3) = 12 + 3 = 15 \end{aligned}$$

We now expand the idea of recursion to apply to relations more generally. First we define a characteristic function (adapted from Hamilton Def 6.17):

*Definition* Let  $R$  be a  $k$ -place relation on  $D_N$ . The *characteristic function* of  $R$ , denoted  $C_R$  is defined by

$$C_R(n_1, \dots, n_k) = \begin{cases} 0 & \text{if } R(n_1, \dots, n_k) \\ 1 & \text{if } \neg R(n_1, \dots, n_k) \end{cases}$$

Note that previously we only considered binary relations in detail (i.e. two place), but we can extend that to three place, four place, etc. For example, we can think of an entry in a three place relations using the ordered pair notation as  $\langle\langle x, y \rangle, z\rangle$  and describe it generally as the set  $R(x, y, z)$ . If a particular triplet, in this case, is present in the set  $R(x, y, z)$ , then the relation is true otherwise it is false.

Given the definition of a characteristic function of a relation, we make the following definition (Hamilton Def. 6.18):

*Definition* A relation  $R$  on  $D_N$  is *recursive* if its characteristic function is a recursive function.



We adapt an example from Hamilton to illustrate this. Define a binary relation  $R(m, n)$  on  $D_N$  to be true whenever  $m + n$  is odd. If we can show that the associated characteristic function is recursive, then by definition, so is the relation  $R$ . The characteristic function is given as:

$$C_R(m, n) = \begin{cases} 0 & \text{if } R(m, n) \text{ is true, which means } m + n \text{ is odd} \\ 1 & \text{if } \neg R(m, n) \text{ is true, which means } m + n \text{ is even} \end{cases}$$

In order to proceed, we assume a recursive function exists which computes the remainder after division by 2. Call this function  $rm(n)$ . For example,  $rm(5) = 1$  and  $rm(8) = 0$ . In other words, this function returns 1 for odd arguments and 0 for even. This function  $rm$  is actually shown to be recursive in Hamilton, so we leverage that result. We know that the addition function  $f_2(m, n)$  is recursive, since we showed that earlier. We almost have all the necessary building blocks. We will need a recursive function to represent a constant. It is also stated in Hamilton that such a function  $f(n) = k$ , where  $k \in D_N$  is some constant, is indeed recursive.

However, we still need a recursive function which returns 0 when the argument is non-zero and returns 1 when the argument is 0, so that we can construct our characteristic function  $C_R(m, n)$  to have the correct result for when  $R(m, n)$  is true (i.e. when  $m + n$  is odd). We now construct such a recursive function, and call it  $\overline{sg}$ , as is done in Hamilton. We now define it as follows:

$$\begin{aligned} \overline{sg}(0) &= f(0), \text{ where } f(n) = 1, \text{ the constant function } 1 \\ \overline{sg}(n + 1) &= z(p_1^2(n, \overline{sg}(n))) = 0 \end{aligned}$$

Given all these functions, we can now construct the characteristic function by using “composition” (i.e. Rule R1.):

$$C_R(m, n) = \overline{sg}(rm(f_2(m, n)))$$

Since each individual function is recursive (i.e. constructed using the base functions and rules R1-R3), and we constructed  $C_R$  using the composition rule, the resulting function  $C_R$  is also recursive. This then implies that the relation  $R(m, n)$  is also recursive (by the earlier definition) and we are done.

We note that the definition of recursiveness applies to sets of numbers. In particular, if  $A \subseteq D_N$ , the set  $A$  is recursive if the characteristic function of the relation  $\in A$  is recursive. An example is given to show that the set of even numbers is recursive. The remainder function  $rm$  discussed above, which is recursive, can act as the characteristic function in this case. In particular,  $n \in A$ , where  $A$  is the set of even natural numbers, then  $rm(n) = 0$  otherwise  $rm(n) = 1$ .

An important connection exist between recursive relations and the formal system we are considering, namely the system of first order arithmetic. In particular, a recursive relation is “expressible” in this system, which is referred to as  $\mathcal{N}$ .

To continue this discussion towards a proof of Gödel’s First Incompleteness Theorem, let us consider the following recursive relation  $W(m, n)$  defined on  $D_N$ , (the natural numbers), which is expressible in  $\mathcal{N}$  (adapted from Proposition 6.29 in Hamilton)

$W(m, n)$  holds if and only if  $m$  is the Gödel number of a wf  $\mathcal{A}(x_1)$  in which  $x_1$  occurs free, and  $n$  is the Gödel number of a proof in  $\mathcal{N}$  of  $\mathcal{A}(0^{(m)})$ .

Then, since  $W$  is expressible in  $\mathcal{N}$ , this means there is a well-formed formula (wf)  $\mathcal{W}(x_1, x_2)$  such that:

If  $W(m, n)$  holds then  $\vdash_{\mathcal{N}} \mathcal{W}(0^{(m)}, 0^{(n)})$ , where  $\vdash_{\mathcal{N}}$  means that the formula is derivable or is a theorem of the system,  $\mathcal{N}$ . The symbol  $0^{(m)}$  corresponds to the natural number  $m$ . Conversely, if  $W(m, n)$  does not hold then  $\vdash_{\mathcal{N}} \neg \mathcal{W}(0^{(m)}, 0^{(n)})$ .

Let us examine expressibility with an example adapted from Hamilton (Example 6.8). The function  $f : D_N \rightarrow D_N$  given by  $f(m) = 2m$  is expressible in  $\mathcal{N}$ . Let  $\mathcal{A}(x_1, x_2)$  be the wf  $x_2 = x_1 \times 0^{(2)}$ . By using the axioms of  $\mathcal{N}$  (which is done in Hamilton) the two following required statements can be established.

$$\begin{aligned} \text{if } n = 2m \text{ then } \vdash_{\mathcal{N}} 0^{(n)} &= 0^{(m)} \times 0^{(2)} \\ \text{if } n \neq 2m \text{ then } \vdash_{\mathcal{N}} \neg(0^{(n)} &= 0^{(m)} \times 0^{(2)}) \end{aligned}$$

To illustrate the first couple of steps in the proof, we have:

$$\begin{aligned} 0^{(m)} \times 0^{(2)} &= 0^{(m)} \times 0'' && \text{(Reason: notation)} \\ &= (0^{(m)} \times 0') + 0^{(m)} && \text{(Reason: Axiom N6*)} \\ &\dots \\ &= 0^{(n)} \end{aligned}$$

Axiom N6\* is given in Hamilton as:  $(\forall x_1)(\forall x_2)(x_1 \times x'_2 = (x_1 \times x_2) + x_1)$ . We note that the prime on  $x_2$  means the “successor” of  $x_2$ . For example, 3 is the successor of 2 in the natural numbers.

We now state Gödel’s First Incompleteness Theorem in precise language, followed by the proof, based on Hamilton (Proposition 6.32):

*Under the assumption that  $\mathcal{N}$  is  $\omega$ -consistent, the wf  $\mathcal{U}$  is not a theorem of  $\mathcal{N}$ , nor is its negation. Hence, if  $\mathcal{N}$  is  $\omega$ -consistent, then  $\mathcal{N}$  is not complete.*

The well-formed formula  $\mathcal{U}$  is defined as  $(\forall x_2)\neg\mathcal{W}(0^{(p)}, x_2)$ , where  $p$  is the Gödel number of the wf  $(\forall x_2)\neg\mathcal{W}(x_1, x_2)$ .

The term  $\omega$ -consistent asserts that if each well-formed formula  $\mathcal{A}(0^{(n)})$  is a theorem of  $\mathcal{N}$  (where  $n \in D_N$ ), then  $\neg(\forall x_1)\mathcal{A}(x_1)$  is not a theorem. We note that in Hamilton, Remark 6.33 states that “We have explicitly stated the assumption of  $\omega$ -consistency, although there is an obvious demonstration that  $\mathcal{N}$  is  $\omega$ -consistent, using the model  $N$  ( $N$  refers to the an interpretation of arithmetic in natural numbers, including the distinguished element 0, addition and multiplication, the successor function, and the equality relation  $=$ ).

The proof is below:

Part I: We show that  $\mathcal{U}$  is not a theorem of  $\mathcal{N}$ .

Suppose that  $\mathcal{U}$  is a theorem of  $\mathcal{N}$ . Let  $q$  be the Gödel number of a proof of  $\mathcal{U}$  in  $\mathcal{N}$ . Let  $p$  be the Gödel number of  $(\forall x_2)\neg\mathcal{W}(x_1, x_2)$ . Thus  $W(p, q)$  holds.  $W$  is expressible in  $\mathcal{N}$  so we have  $\vdash_{\mathcal{N}} \mathcal{W}(0^{(p)}, 0^{(q)})$ . In other words,  $\mathcal{W}$  is derivable from the axioms of  $\mathcal{N}$ .

But we assumed  $\mathcal{U}$  is theorem, which means  $\vdash_{\mathcal{N}} (\forall x_2)\neg\mathcal{W}(0^{(p)}, x_2)$ , and so  $\vdash_{\mathcal{N}} \neg\mathcal{W}(0^{(p)}, 0^{(q)})$ , using Axiom K5 and modus ponens. This contradicts the consistency of  $\mathcal{N}$  (note that  $\omega$ -consistency implies consistency), so  $\mathcal{U}$  cannot be a theorem of  $\mathcal{N}$ .

Axiom K5 of the system is:  $((\forall x_i)\mathcal{A}(x_i) \implies \mathcal{A}(t))$ .

Part II: We show that  $\neg\mathcal{U}$  is not a theorem of  $\mathcal{N}$ .

From Part I, we know that  $\mathcal{U}$  is not a theorem of  $\mathcal{N}$ , i.e. there is no proof in  $\mathcal{N}$  of  $\mathcal{U}$ , so for no number  $q$  is  $q$  the Gödel number of a proof in  $\mathcal{N}$  of  $\mathcal{U}$ , i.e. of  $(\forall x_2)\neg\mathcal{W}(0^{(p)}, x_2)$ , thus  $W(p, q)$  does not hold for any number  $q$ . Therefore

$$\vdash_N \neg\mathcal{W}(0^{(p)}, 0^{(q)}) \text{ for every } q.$$

By  $\omega$ -consistency, then,

$$\neg(\forall x_2)\neg\mathcal{W}(0^{(p)}, x_2)$$

is not a theorem of  $\mathcal{N}$  i.e.  $(\neg\mathcal{U})$  is not a theorem of  $\mathcal{N}$ .

End of proof.

Since we have shown that a formula exists in  $\mathcal{N}$ , called  $\mathcal{U}$  which is not a theorem, and the negation  $(\neg\mathcal{U})$  is not a theorem, this means that  $\mathcal{N}$  is not complete.

We now consider some exercises to better understand the proof.

*Exercise* In Part I of the proof, it was stated that there was a contradiction to the consistency of  $\mathcal{N}$ . Please explain why this statement could be made.

*Solution* First recall the definition of consistency: *A formal system for which there is no statement such that the statement and its negation are both derivable is consistent.* In Part I of the proof, under the assumption that  $\mathcal{U}$  was a theorem of  $\mathcal{N}$ , we obtained  $\vdash_{\mathcal{N}} \mathcal{W}(0^{(p)}, 0^{(q)})$  and  $\vdash_{\mathcal{N}} \neg \mathcal{W}(0^{(p)}, 0^{(q)})$ . Since we know that  $\mathcal{N}$  is consistent, obtaining both these results, as one is the negation of the other, violates consistency. From this contradiction, we conclude that our original assumption that  $\mathcal{U}$  is a theorem of  $\mathcal{N}$  must be false.

*Exercise* In Part II of the proof, please explain how  $\omega$ -consistency implies that if we have  $\vdash_{\mathcal{N}} \neg \mathcal{W}(0^{(p)}, 0^{(q)})$  for every  $q$ , we get that  $\neg(\forall x_2) \neg \mathcal{W}(0^{(p)}, x_2)$  is not a theorem of  $\mathcal{N}$ .

*Solution* Recalling that the term  $\omega$ -consistent asserts that if each well-formed formula  $\mathcal{A}(0^{(q)})$  is a theorem of  $\mathcal{N}$  (where  $q \in D_N$ ), then  $\neg(\forall x_1) \mathcal{A}(x_1)$  is not a theorem, we first let  $\mathcal{A}(0^{(q)}) = \neg \mathcal{W}(0^{(p)}, 0^{(q)})$ . (Note that we substituted  $q$  for  $n$  in the original statement of  $\omega$ -consistency). Then, we have that  $\neg(\forall x_2) \mathcal{A}(x_2)$  is not a theorem. (Note that we have substituted  $x_2$  for  $x_1$ ). Substituting in for  $\mathcal{A}$ , we obtain finally that  $\neg(\forall x_2) \neg \mathcal{W}(0^{(p)}, x_2)$  is not a theorem of  $\mathcal{N}$ .

### The Second Incompleteness Theorem

We now present Gödel Second Incompleteness Theorem, largely following the Stanford Encyclopedia of Philosophy entry on Kurt Gödel (<https://plato.stanford.edu/entries/goedel/>). We note that in this reference, the formal system is called  $P$  and “contains” Peano arithmetic. (Note: Please be aware that in this particular case of Peano arithmetic, the minimal value is 0 as opposed to 1).

In summary, the Second Incompleteness Theorem establishes that the consistency of  $P$  cannot be *proved* in  $P$ . Here, “consistency” is meant in the technical term we defined earlier, namely in a consistent formal system, you cannot derive both a statement and its negation. We state the theorem below in precise terms:

*If  $P$  is consistent, then  $\neg \text{Prov}(\ulcorner 0 = 1 \urcorner)$  is not provable from  $P$ .*

In the above statement of the theorem, the relation  $\text{Prov}(y)$  is true whenever the “sentence” (formula) with Gödel number  $y$  is provable in the system  $P$ . The sequence  $\ulcorner x \urcorner$  means the Gödel number of the sentence  $x$ . The underline signifies the formal term corresponding to the natural number code of the argument. For example, assume the Gödel number of the statement  $0 = 1$  is 57, i.e.  $\ulcorner 0 = 1 \urcorner = 57$ . Then,  $\underline{57}$  would be how that number is represented in the formal system  $P$ . In particular, it would be whatever formula represents 57 applications of the successor function on 0, say  $0^{(57)}$ . It should be emphasized that the sentence  $\neg \text{Prov}(\ulcorner 0 = 1 \urcorner)$  is a statement which asserts the consistency

of  $P$ .

In order to give the proof of the Second Incompleteness Theorem, we define a “sentence”  $\phi$  in the following manner:

$$P \vdash (\phi \iff \neg \text{Prov}(\ulcorner \phi \urcorner))$$

In words, the  $\phi$  in the above definition is not provable in  $P$ . It is shown in the Stanford reference that  $P \not\vdash \phi$ , meaning  $\phi$  cannot be derived in  $P$ , (i.e. is not a theorem in  $P$ .)

We now continue with the proof of the Second Incompleteness Theorem (again following the Stanford reference):

Utilize the definition of  $\phi$  stated above. It is first assumed that the reasoning to obtain the statement “if  $P \vdash \phi$ , then  $P \vdash 0 \neq 1$ ” can be stated in  $P$ . This yields:  $P \vdash (\text{Prov}(\ulcorner \phi \urcorner) \implies \text{Prov}(\ulcorner 0 = 1 \urcorner))$ . Then we also have, by the definition of  $\phi$ ,  $P \vdash (\neg \text{Prov}(\ulcorner 0 = 1 \urcorner) \implies \phi)$ . Since we know that actually  $P \not\vdash \phi$ , this means that  $P \not\vdash \neg \text{Prov}(\ulcorner 0 = 1 \urcorner)$ . This means that a statement which asserts the consistency of  $P$  cannot be proved in  $P$ .

*End of proof.*

*Exercise* It was stated in the proof from the definition of  $\phi$  we can state  $P \vdash (\neg \text{Prov}(\ulcorner 0 = 1 \urcorner) \implies \phi)$ . Please explain how this is possible using logical manipulations.

*Solution* We can rewrite  $P \vdash (\phi \iff \neg \text{Prov}(\ulcorner \phi \urcorner))$  as  $P \vdash ((\phi \implies \neg \text{Prov}(\ulcorner \phi \urcorner)) \wedge (\neg \text{Prov}(\ulcorner \phi \urcorner) \implies \phi))$  and as both terms must be true for the  $\wedge$  to be true, the second term corresponds to part of what we seek. In the proof, we also have  $P \vdash (\text{Prov}(\ulcorner \phi \urcorner) \implies \text{Prov}(\ulcorner 0 = 1 \urcorner))$ . Writing this expression in the contrapositive, we obtain  $P \vdash (\neg \text{Prov}(\ulcorner 0 = 1 \urcorner) \implies \neg \text{Prov}(\ulcorner \phi \urcorner))$ . So, in summary, we have

$$\begin{aligned} P \vdash (\neg \text{Prov}(\ulcorner 0 = 1 \urcorner) \implies \neg \text{Prov}(\ulcorner \phi \urcorner)) \\ P \vdash (\neg \text{Prov}(\ulcorner \phi \urcorner) \implies \phi) \end{aligned}$$

and thus we can finally write

$$P \vdash ((\neg \text{Prov}(\ulcorner 0 = 1 \urcorner) \implies \phi).$$

*Exercise* It was emphasized that the sentence  $\neg \text{Prov}(\ulcorner 0 = 1 \urcorner)$  is a statement which asserts the consistency of  $P$ . Does this make sense, since it is only a single formula and the definition of consistency is a statement about a formula and

the negation of that formula?

*Solution* This does makes sense, if we take consider the negation of  $0 = 1$ , which would be  $0 \neq 1$ . In the system  $P$ , which contains Peano arithmetic, there is an axiom (Axiom III in our earlier section on Peano Arithmetic) which states: “Axiom III: For each  $n \in \mathbb{N}$  we have  $n^* \neq 1$ .” If we replace 1 with 0 (since the system  $P$  we are using for the Second Incompleteness Theorem uses 0 as the minimal element), we see that  $n^*$ , the successor of any  $n$ , does not equal the minimal element. Thus,  $0 \neq 1$  is true by Axiom III and indeed derivable in one step, if we let  $n = 0$  and  $n^* = 1$ . Therefore, if  $P$  is consistent, then we must have  $\neg \text{Prov}(\ulcorner 0 = 1 \urcorner)$  since  $\text{Prov}(\ulcorner 0 \neq 1 \urcorner)$  as we just showed. However, note carefully that we cannot actual *prove*  $\neg \text{Prov}(\ulcorner 0 = 1 \urcorner)$  in  $P$ , as we learned by the Second Incompleteness Theorem.

## Chapter 2

# Methods of Proof

In this chapter, we will discuss various methods of proof and provide illustrations of each. In the first chapter, we saw some examples of proofs without dwelling on the process of doing the proof itself. The emphasis in this book is to gain experience with reading and writing proofs within various topics in pure mathematics. It is important to both read *and* write proofs in order to become comfortable with them. The exit goal of this chapter is that the reader will have a general knowledge of the various types of proofs and be able to read and write such proofs.

One may see proofs that are written as lists, say with the statement on the left and the justification on the right, or written more like prose, where more usual written language is combined with the mathematics and logic. In the previous chapter, there were examples of each type.

Much of the material in this chapter is based on the YouTube video series “Math Major Basics” by MathDoctorBob (<https://www.youtube.com/playlist?list=PLF2DF6C3C8015DF5F>).

We first remark that a *theorem* is a set of valid statements chained together by logical “ands” (i.e.  $\wedge$ ). If all the statements are true and the conclusion is true, then the theorem is *proved*.

## 2.1 Induction

In the *induction* type of proof, the statements in question will be associated with an index which is used to construct the proof.

### 2.1.1 Weak Induction

Let us begin by stating the “principle of mathematic induction”, based on the presentation in Schaum’s Outline of Theory and Problems of Set Theory and

Related Topics (2nd Ed), by Seymour Lipschutz.

Let  $A(n)$  be an assertion defined on the set of positive integers,  $\mathbb{P}$ .  $A(n)$  is true or false for each integer  $n \geq 1$ . Suppose  $A(n)$  has the following two properties:

- $A(1)$  is true (sometimes called the “base case”)
- $A(n+1)$  is true whenever  $A(n)$  is true (sometimes called the “inductive hypothesis”)

Then  $A(n)$  is true for every  $n \geq 1$ .

Let us see an example of this in practice.

We wish to prove that  $\sum_{i=1}^n i = \frac{n(n+1)}{2}$  using (weak) induction. That is, the sum of the first  $n$  positive integers is equal to  $\frac{n(n+1)}{2}$ . For example:  $1 + 2 + 3 + 4 + 5 = 15$ ,  $\frac{5(6)}{2} = \frac{30}{2} = 15\checkmark$ .

We can first check the base case:  $\sum_{i=1}^1 i = 1$ ,  $\frac{1(1+1)}{2} = 1\checkmark$ . Thus  $A(1)$  is true, where  $A(n)$  is the assertion that  $\sum_{i=1}^n i = \frac{n(n+1)}{2}$ .

Now, suppose that  $A(n)$  is true. We then check  $A(n+1)$ . This yields:  $\sum_{i=1}^{n+1} i = \frac{(n+1)(n+2)}{2}$ . If  $A(n+1)$  is true, then  $A(n+1) = A(n) + (n+1)$ . So we check:

$$\begin{aligned} \frac{(n+1)(n+2)}{2} &= \frac{n(n+1)}{2} + (n+1)? \\ \frac{n(n+1)}{2} + (n+1) &= \frac{n(n+1) + 2(n+1)}{2} \\ \frac{n(n+1) + 2(n+1)}{2} &= \frac{(n+1)(n+1)}{2} \checkmark \end{aligned}$$

Thus, we conclude that  $A(n)$  is true for  $n \geq 1$  which completes our proof.

We consider a second example, taken from Schaum's Outline of Theory and Problems of Abstract Algebra (2nd Ed.) by Frank Ayres, Jr. and Lloyd R. Jaisingh. This is associated with the Peano Postulates of Arithmetic that we considered in Chapter 1. We wish to prove the Closure Law for Addition, namely  $n + m \in \mathbb{N}$ , for all  $n, m \in \mathbb{N}$ , using induction. Recall that  $\mathbb{N}$  is the set of natural numbers (which starts at 1 in this case). We proceed as follows:



We use the two definitions associated with addition on  $\mathbb{N}$ , namely (i)  $n + 1 = n^*$ , for every  $n \in \mathbb{N}$ , where  $n^*$  is the successor of  $n$ , and (ii)  $n + m^* = (n + m)^*$ , whenever  $n + m$  is defined (i.e. whenever  $n + m$  is a natural number), as well as Peano Postulate II: For each  $n \in \mathbb{N}$ , there exists a unique  $n^* \in \mathbb{N}$ , called the successor of  $n$ .

We first fix  $n$  as some natural number, and define the proposition  $P(m)$  :  $n + m \in \mathbb{N}$  for every  $m \in \mathbb{N}$ . That is,  $P(m)$  is true, whenever  $n + m$  is a natural number. First, let us determine if  $P(1)$  is true (i.e. the base case). Since  $n + 1 = n^*$ , from Postulate II we know that  $n^* \in \mathbb{N}$ , so  $P(1)$  is true. Now, for the inductive hypothesis portion. Suppose that  $P(m)$  is true, then we want to see if  $P(m + 1)$  is true, which is the same as  $P(m^*)$  by Definition (i). Now,  $P(m^*)$  is true if  $n + m^* \in \mathbb{N}$ . From Definition (ii), we have that  $n + m^* = (n + m)^*$  whenever,  $n + m$  is defined (i.e. is a natural number). Since we assumed  $P(m)$  was true, this means that  $n + m$  is a natural number, and by Postulate II,  $(n + m)^*$  is also a natural number, so we are done.

### 2.1.2 Strong Induction

## 2.2 Proof by Contradiction

Proof by contradiction is a very useful method of proof which is used quite frequently. The usual pattern is one assumes the converse of what is being asserted, and then following a sequence of logical deductions arriving at a contradiction. Once this contradiction has been uncovered, then it implies the original assumption must have been false thus completing the proof. Here is an example to illustrate the points.

*Prove that there is no rational number whose square is 12.* (Note: This was originally an exercise in Rudin PMA 3rd Edition).

Proof:

**Suppose that there was a rational number whose square was 12.** This means that

$$\left(\frac{p}{q}\right)^2 = 12$$

for integers  $p, q$ . We assume that  $p$  and  $q$  are not both even. In other words, all common factors of 2 have been cancelled. This can always been done for a rational number.

Upon expansion we obtain:

$$p^2 = 12q^2 = 2(6q^2)$$

This implies that  $p^2$  and  $p$  are both even. Thus  $q$  is odd. So we write  $p = 2m$  for integer  $m$  and obtain:

$$(2m)^2 = 4m^2 = 12q^2$$

This implies that

$$m^2 = 3q^2$$

Since  $q$  is assumed to be odd,  $q^2$  is odd, and  $3q^2$  is also odd. Now,  $m^2$  must be therefore be odd, and likewise for  $m$ . Therefore, for integers  $r, n$  we can rewrite the above as:

$$(2r + 1)^2 = 3(2n + 1)^2$$

Now, expanding both sides yields:

$$4r^2 + 4r + 1 = 3(4n^2 + 4n + 1)$$

$$4r^2 + 4r + 1 = 12n^2 + 12n + 3$$

$$4r^2 + 4r = 12n^2 + 12n + 2$$

$$4r^2 + 4r = 12n^2 + 12n + 2$$

The left side is clearly divisible by 4.

$$r^2 + r = 3n^2 + 3n + \frac{2}{4}$$

The left side is an integer, but the right side is not, which is a contradiction. Therefore our assumption about both  $p, q$  not even is false. However, since this can always be done for the rational number  $p/q$  it implies that there is no rational  $p/q$  such that  $(p/q)^2 = 12$ .

### 2.3 Direct Proof

## Chapter 3

# Modern Algebra



## Chapter 4

# Real Analysis



## Chapter 5

# Complex Analysis





## Chapter 6

# Non-Euclidean Geometry and Topology



## Chapter 7

# Number Theory



## Chapter 8

# Category Theory



## Chapter 9

# Computational Complexity Theory

# Index

- antisymmetric, 16
- asymmetric, 16
- Cardinality, 13
- characteristic function, 32
- Complete, 25
- composition, 21
- connected, 16, 18
- Consistent, 25
- converse domain, 22
- Coset, 19
- counterdomain, 22
- De Morgan's Law's, 10
- Dedekind Cuts, 7
- domain, 22
- expressible, 34
- Formal system, 25
- function, 20
- Gödel numbering, 28
- image, 22
- induction, 39
- inverse, 22
- irreflexive, 16
- partial ordering, 17
- Partition, 19
- R-minimal element, 17
- range, 22
- recursive functions, 29
- recursive relation, 32
- reflexive, 16
- relative product, 21
- restriction, 22
- Set builder notation, 8
- strongly connected, 16
- symmetric, 16
- transitive, 16
- well-order, 17