

# TOPICS IN ANALYTIC NUMBER THEORY

## EXERCISE SHEET 1

- (1) Prove that

$$N_{\mathbb{P}^n, H}(B) \sim \frac{2^n}{\zeta(n+1)} B^{n+1},$$

as  $B \rightarrow \infty$ .

- (2) Let  $\nu(B)$  denote the number of  $(a, b, c, d) \in \mathbb{N}^4$  such that  $a^3 + b^3 = c^3 + d^3$  and  $a, b, c, d \leq B$ . Show that  $\nu(B) = O_\varepsilon(B^{2+\varepsilon})$ , for any  $\varepsilon > 0$ . You may assume the standard estimate for the divisor function.
- (3) Let  $e_q(\cdot) = \exp(\frac{2\pi i}{q} \cdot)$  and let

$$S_q = \sum_{a \in (\mathbb{Z}/q\mathbb{Z})^*} \sum_{\mathbf{x} \in (\mathbb{Z}/q\mathbb{Z})^n} e_q(aF(\mathbf{x})),$$

where  $F \in \mathbb{Z}[x_1, \dots, x_n]$  is a homogeneous polynomial. Prove that  $S_q$  is a multiplicative function of  $q$ .

- (4) Let  $S$  be the set of unramified rational primes  $p$  that split in the cubic number field  $\mathbb{Q}(2^{1/3})$ . Use the Chebotarev density theorem to check the convergence of the Euler product

$$\prod_{\substack{p \equiv 1 \pmod{3} \\ p \in S}} \left(1 - \frac{1}{p}\right) \left(1 + \frac{6}{p} - \frac{6}{p^2}\right).$$

Similarly, check the convergence of

$$\prod_{\substack{p \equiv 1 \pmod{3} \\ p \notin S}} \left(1 - \frac{1}{p}\right) \left(1 - \frac{3}{p} + \frac{3}{p^2}\right).$$

- (5) Let  $p$  be a prime and let  $\mathbf{a}, \mathbf{b} \in \mathbb{Z}^n$  be vectors that are both coprime to  $p$ . Let  $\mathbf{M}$  denote the set of vectors  $\mathbf{x} \in \mathbb{Z}^n$  such that  $p^2 \mid \mathbf{a} \cdot \mathbf{x}$  and  $\mathbf{x} \equiv \lambda \mathbf{b} \pmod{p}$ , for some  $\lambda \in \mathbb{Z}$ .
- (a) Prove that  $\mathbf{M}$  is a lattice of rank  $n$ .
- (b) Prove that  $\det \mathbf{M} = p^n$ .