

Direct fold test: all 26 candidate intervals

10/26 pass all paired checks. All 16 finite-return curve families retained.

b = 0.2, c = 7.212 | Unresolved does not mean absent | No symbolic letters assigned

a = 0.21575

Region / depth / direction / candidate	DOP	Radau	Both	Fold input x
Left / 4 / 0 / 1	N	N	U	--
Left / 4 / 0 / 2	N	N	U	--
Left / 4 / 1 / 1	N	N	U	--
Left / 4 / 1 / 2	N	N	U	--
Left / 8 / 0 / 1	N	N	U	--
Left / 8 / 0 / 2	N	N	U	--
Left / 8 / 1 / 1	N	N	U	--
Left / 8 / 1 / 2	N	N	U	--
Right / 4 / 0 / 1	Q	Q	Q	-6.880758
Right / 4 / 1 / 1	Q	Q	Q	-6.880758
Right / 8 / 0 / 1	Q	Q	Q	-6.880758
Right / 8 / 1 / 1	Q	Q	Q	-6.880758

4/12 candidates qualify in both solvers

a = 0.21577

Region / depth / direction / candidate	DOP	Radau	Both	Fold input x
Left / 4 / 0 / 1	N	N	U	--
Left / 4 / 1 / 1	N	N	U	--
Left / 8 / 0 / 1	N	N	U	--
Left / 8 / 1 / 1	N	N	U	--
Right / 4 / 0 / 1	Q	Q	Q	-6.880532
Right / 4 / 1 / 1	Q	Q	Q	-6.880532
Right / 8 / 0 / 1	N	N	U	--
Right / 8 / 0 / 2	Q	Q	Q	-6.880532
Right / 8 / 0 / 3	Q	Q	Q	-6.880532
Right / 8 / 0 / 4	E	E	U	--
Right / 8 / 1 / 1	N	N	U	--
Right / 8 / 1 / 2	Q	Q	Q	-6.880532
Right / 8 / 1 / 3	Q	Q	Q	-6.880532
Right / 8 / 1 / 4	E	E	U	--

6/14 candidates qualify in both solvers

Q Qualified N No qualified shooting/census E Event or input projection invalid

F A fold check failed U Paired result unresolved * Outside the prior observed x region

Input x is the DOP853 value only when both solvers qualify; both unrounded values are retained in the receipt.

Depth is the number of prior returns; directions 0 and 1 are the two frozen initial tangents. Candidate numbers are local to each family.