

### **Activity Overview**


- Using SPENVIS to generate some environment results for a few reference orbits
- Guided example for Mid-Latitude LEO (ex. ISS)
  - Looking at trapped fluxes, dose-depth curves
- On your own: generate environments for couple other orbits
  - Walking around to help with issues and answer questions

1

# **Guided Activity**

Mid Latitude LEO (ISS, Starlink)	
<b>Mission Duration</b>	10 years
Apogee	420 km
Perigee	420 km
Inclination	52.0 deg
RAAN <sup>1</sup>	0 deg
Argument of Perigee	0 deg
True Anomaly	0 deg

<sup>&</sup>lt;sup>1</sup>Right Ascension of the Ascending Node



### On your own



#### By yourself or in a small group, generate results for these other reference orbits

Polar LEO (POES, IRIDIUM)		
<b>Mission Duration</b>	7 years	
Apogee	825 km	
Perigee	825 km	
Inclination	98.8 deg	
RAAN <sup>1</sup>	0 deg	
Argument of Perigee	0 deg	
True Anomaly	0 deg	

HEO (Van Allen Probes, MMS)		
<b>Mission Duration</b>	2 years	
Apogee	70000 km	
Perigee	2500 km	
Inclination	28 deg	
RAAN	0 deg	
Argument of Perigee	0 deg	
True Anomaly	0 deg	


How do the environments at these orbits differ?


<sup>&</sup>lt;sup>1</sup>Right Ascension of the Ascending Node




### **Mid Latitude LEO**












### **Polar LEO**









## **HEO**



