

# Liftings of Elliptic and Hyperelliptic Curves

Luís Finotti

University of Tennessee

AMS Joint Meeting  
Jan 10, 2013

# Witt Vectors

Let  $\mathbb{k}$  be a *perfect* field of characteristic  $p > 0$  and  $W = W(\mathbb{k})$  be the ring of Witt vectors over  $\mathbb{k}$ .

# Witt Vectors

Let  $\mathbb{k}$  be a *perfect* field of characteristic  $p > 0$  and  $W = W(\mathbb{k})$  be the ring of Witt vectors over  $\mathbb{k}$ .

Remember also that we have a Frobenius map, which we denote by  $\sigma$  on  $W$ , defined by

$$\sigma(a_0, a_1, \dots) = (a_0^p, a_1^p, \dots).$$

# Witt Vectors

Let  $\mathbb{k}$  be a *perfect* field of characteristic  $p > 0$  and  $W = \mathbf{W}(\mathbb{k})$  be the ring of Witt vectors over  $\mathbb{k}$ .

Remember also that we have a Frobenius map, which we denote by  $\sigma$  on  $W$ , defined by

$$\sigma(a_0, a_1, \dots) = (a_0^p, a_1^p, \dots).$$

So, if  $\sigma$  denotes the Frobenius in characteristic  $p$  (i.e.,  $\sigma(a) = a^p$ ), we have a *lifting of the Frobenius*.

# Witt Vectors

Let  $\mathbb{k}$  be a *perfect* field of characteristic  $p > 0$  and  $W = \mathbf{W}(\mathbb{k})$  be the ring of Witt vectors over  $\mathbb{k}$ .

Remember also that we have a Frobenius map, which we denote by  $\sigma$  on  $W$ , defined by

$$\sigma(a_0, a_1, \dots) = (a_0^p, a_1^p, \dots).$$

So, if  $\sigma$  denotes the Frobenius in characteristic  $p$  (i.e.,  $\sigma(a) = a^p$ ), we have a *lifting of the Frobenius*. More precisely, if  $\pi : W \rightarrow \mathbb{k}$  is the reduction modulo  $p$ , we have the following diagram on multiplicative groups:

$$\begin{array}{ccc} W^\times & \xrightarrow{\sigma} & W^\times \\ \pi \downarrow & & \downarrow \pi \\ \mathbb{k}^\times & \xrightarrow{\sigma} & \mathbb{k}^\times \end{array}$$

# Witt Vectors (cont.)

Moreover, the *Teichmüller lift*  $\tau : a \mapsto (a, 0, 0, \dots)$  (a *group homomorphism*) yields the following diagram:

$$\begin{array}{ccc}
 W^\times & \xrightarrow{\sigma} & W^\times \\
 \pi \updownarrow \tau & & \tau \updownarrow \pi \\
 \mathbb{k}^\times & \xrightarrow{\sigma} & \mathbb{k}^\times
 \end{array}$$

# Witt Vectors (cont.)

Moreover, the *Teichmüller lift*  $\tau : a \mapsto (a, 0, 0, \dots)$  (a *group homomorphism*) yields the following diagram:

$$\begin{array}{ccc}
 W^\times & \xrightarrow{\sigma} & W^\times \\
 \pi \updownarrow \tau & & \tau \updownarrow \pi \\
 \mathbb{k}^\times & \xrightarrow{\sigma} & \mathbb{k}^\times
 \end{array}$$

## Question

Can we also lift the Frobenius for curves over  $\mathbb{k}$ ?

# Curves

More precisely, given a curve  $C/\mathbb{k}$  and if  $\phi : C \rightarrow C^\sigma$  is the Frobenius map, is there a lifting  $C/W$  for which we can lift the Frobenius:

$$\begin{array}{ccc}
 C(W(\bar{\mathbb{k}})) & \xrightarrow{\phi} & C^\sigma(W(\bar{\mathbb{k}})) \\
 \pi \downarrow & & \downarrow \pi \\
 C(\bar{\mathbb{k}}) & \xrightarrow{\phi} & C^\sigma(\bar{\mathbb{k}})
 \end{array}$$



# Curves

More precisely, given a curve  $C/\mathbb{k}$  and if  $\phi : C \rightarrow C^\sigma$  is the Frobenius map, is there a lifting  $C/W$  for which we can lift the Frobenius:

$$\begin{array}{ccc}
 C(W(\bar{\mathbb{k}})) & \xrightarrow{\phi} & C^\sigma(W(\bar{\mathbb{k}})) \\
 \pi \downarrow & & \downarrow \pi \\
 C(\bar{\mathbb{k}}) & \xrightarrow{\phi} & C^\sigma(\bar{\mathbb{k}})
 \end{array}$$

**Answer:** Yes, for *ordinary* elliptic curves and Abelian varieties (Deuring and Serre-Tate),

# Curves

More precisely, given a curve  $C/\mathbb{k}$  and if  $\phi : C \rightarrow C^\sigma$  is the Frobenius map, is there a lifting  $C/W$  for which we can lift the Frobenius:

$$\begin{array}{ccc}
 C(W(\bar{\mathbb{k}})) & \xrightarrow{\phi} & C^\sigma(W(\bar{\mathbb{k}})) \\
 \pi \downarrow & & \downarrow \pi \\
 C(\bar{\mathbb{k}}) & \xrightarrow{\phi} & C^\sigma(\bar{\mathbb{k}})
 \end{array}$$

**Answer:** Yes, for *ordinary* elliptic curves and Abelian varieties (Deuring and Serre-Tate), but no for higher genus curves (Raynaud).

# Curves

More precisely, given a curve  $C/\mathbb{k}$  and if  $\phi : C \rightarrow C^\sigma$  is the Frobenius map, is there a lifting  $C/W$  for which we can lift the Frobenius:

$$\begin{array}{ccc}
 C(W(\bar{\mathbb{k}})) & \xrightarrow{\phi} & C^\sigma(W(\bar{\mathbb{k}})) \\
 \pi \downarrow & & \downarrow \pi \\
 C(\bar{\mathbb{k}}) & \xrightarrow{\phi} & C^\sigma(\bar{\mathbb{k}})
 \end{array}$$

**Answer:** Yes, for *ordinary* elliptic curves and Abelian varieties (Deuring and Serre-Tate), but no for higher genus curves (Raynaud). In the case of elliptic curves we also have a *Teichmüller lift*.

# Curves

More precisely, given a curve  $C/\mathbb{k}$  and if  $\phi : C \rightarrow C^\sigma$  is the Frobenius map, is there a lifting  $C/W$  for which we can lift the Frobenius:

$$\begin{array}{ccc}
 C(W(\bar{\mathbb{k}})) & \xrightarrow{\phi} & C^\sigma(W(\bar{\mathbb{k}})) \\
 \pi \downarrow & & \downarrow \pi \\
 C(\bar{\mathbb{k}}) & \xrightarrow{\phi} & C^\sigma(\bar{\mathbb{k}})
 \end{array}$$

**Answer:** Yes, for *ordinary* elliptic curves and Abelian varieties (Deuring and Serre-Tate), but no for higher genus curves (Raynaud). In the case of elliptic curves we also have a *Teichmüller lift*.

Also, **Mochizuki** showed that one can lift the Frobenius for some curves of genus  $g \geq 2$  if we allow **singularities** (at  $(p-1)(g-1)$  points).

# Ordinary Elliptic Curve

An elliptic curve (given by  $y^2 = x^3 + ax + b$ ) over a field  $\mathbb{k}$  of characteristic  $p > 3$  is *ordinary* if  $E[p] \cong \mathbb{Z}/p$ .

# Ordinary Elliptic Curve

An elliptic curve (given by  $y^2 = x^3 + ax + b$ ) over a field  $\mathbb{k}$  of characteristic  $p > 3$  is *ordinary* if  $E[p] \cong \mathbb{Z}/p$ . (Or, equivalently, if the coefficient of  $x^{p-1}$  in  $(x^3 + ax + b)^{(p-1)/2}$  is non-zero.)

# Ordinary Elliptic Curve

An elliptic curve (given by  $y^2 = x^3 + ax + b$ ) over a field  $\mathbb{k}$  of characteristic  $p > 3$  is *ordinary* if  $E[p] \cong \mathbb{Z}/p$ . (Or, equivalently, if the coefficient of  $x^{p-1}$  in  $(x^3 + ax + b)^{(p-1)/2}$  is non-zero.) Otherwise, the elliptic curve is said to be *supersingular*.

# Ordinary Elliptic Curve

An elliptic curve (given by  $y^2 = x^3 + ax + b$ ) over a field  $\mathbb{k}$  of characteristic  $p > 3$  is *ordinary* if  $E[p] \cong \mathbb{Z}/p$ . (Or, equivalently, if the coefficient of  $x^{p-1}$  in  $(x^3 + ax + b)^{(p-1)/2}$  is non-zero.) Otherwise, the elliptic curve is said to be *supersingular*.

**Note:** Only finitely many elliptic curves (up to isomorphism) are supersingular.



# Ordinary Elliptic Curve

An elliptic curve (given by  $y^2 = x^3 + ax + b$ ) over a field  $\mathbb{k}$  of characteristic  $p > 3$  is **ordinary** if  $E[p] \cong \mathbb{Z}/p$ . (Or, equivalently, if the coefficient of  $x^{p-1}$  in  $(x^3 + ax + b)^{(p-1)/2}$  is non-zero.) Otherwise, the elliptic curve is said to be **supersingular**.

**Note:** Only finitely many elliptic curves (up to isomorphism) are supersingular.

We can lift the Frobenius for ordinary elliptic curves, i.e., if  $\mathbb{k}$  is a perfect field with  $\text{char}(\mathbb{k}) = p$  and  $E/\mathbb{k} : y_0^2 = x_0^3 + a_0x_0 + b_0$ , then there exists  $\mathbf{a} = (a_0, a_1, \dots), \mathbf{b} = (b_0, b_1, \dots) \in W$  such that  $E/W : y^2 = x^3 + ax + b$  has a lifting of the Frobenius:

$$\begin{array}{ccc}
 E(W(\bar{\mathbb{k}})) & \xrightarrow{\phi} & E^\sigma(W(\bar{\mathbb{k}})) \\
 \pi \downarrow & & \downarrow \pi \\
 E(\bar{\mathbb{k}}) & \xrightarrow{\phi} & E^\sigma(\bar{\mathbb{k}})
 \end{array}$$

# Elliptic Teichmüller Lift

Moreover, the curve  $E$  above is unique up to isomorphism and it is called the *canonical lifting of  $E$* .

# Elliptic Teichmüller Lift

Moreover, the curve  $E$  above is unique up to isomorphism and it is called the *canonical lifting of  $E$* .

As with Witt vectors, we also have a section of the reduction modulo  $p$ , the so called *elliptic Teichmüller lift  $\tau$* :

$$\begin{array}{ccc}
 E(W(\bar{\mathbb{k}})) & \xrightarrow{\phi} & E^\sigma(W(\bar{\mathbb{k}})) \\
 \pi \downarrow \uparrow \tau & & \pi \downarrow \uparrow \tau^\sigma \\
 E(\bar{\mathbb{k}}) & \xrightarrow{\phi} & E^\sigma(\bar{\mathbb{k}})
 \end{array}$$

# Elliptic Teichmüller Lift

Moreover, the curve  $E$  above is unique up to isomorphism and it is called the *canonical lifting of  $E$* .

As with Witt vectors, we also have a section of the reduction modulo  $p$ , the so called *elliptic Teichmüller lift  $\tau$* :

$$\begin{array}{ccc}
 E(W(\bar{\mathbb{k}})) & \xrightarrow{\phi} & E^\sigma(W(\bar{\mathbb{k}})) \\
 \pi \downarrow \uparrow \tau & & \pi \downarrow \uparrow \tau^\sigma \\
 E(\bar{\mathbb{k}}) & \xrightarrow{\phi} & E^\sigma(\bar{\mathbb{k}})
 \end{array}$$

Also,  $\tau$  is a *group homomorphism*

# Elliptic Teichmüller Lift

Moreover, the curve  $E$  above is unique up to isomorphism and it is called the *canonical lifting of  $E$* .

As with Witt vectors, we also have a section of the reduction modulo  $p$ , the so called *elliptic Teichmüller lift  $\tau$* :

$$\begin{array}{ccc}
 E(W(\bar{\mathbb{k}})) & \xrightarrow{\phi} & E^\sigma(W(\bar{\mathbb{k}})) \\
 \pi \downarrow \uparrow \tau & & \pi \downarrow \uparrow \tau^\sigma \\
 E(\bar{\mathbb{k}}) & \xrightarrow{\phi} & E^\sigma(\bar{\mathbb{k}})
 \end{array}$$

Also,  $\tau$  is a *group homomorphism*, and one can show that:

$$\tau(x_0, y_0) = ((F_0, F_1, F_2, \dots), (y_0, y_0 G_1, y_0 G_2, \dots)),$$

where  $F_i, G_i \in \mathbb{k}[x_0]$ .

# Error Correcting Codes

Voloch and Walker used canonical liftings and the elliptic Teichmüller lift to create error-correcting codes.

# Error Correcting Codes

Voloch and Walker used canonical liftings and the elliptic Teichmüller lift to create error-correcting codes. The bounds for the parameters (which measure “how good” the resulting codes are likely to be) depend on the degrees of  $F_i$ 's and  $G_i$ 's,

# Error Correcting Codes

Voloch and Walker used canonical liftings and the elliptic Teichmüller lift to create error-correcting codes. The bounds for the parameters (which measure “how good” the resulting codes are likely to be) depend on the degrees of  $F_i$ 's and  $G_i$ 's, with lower degrees giving better bounds.



# Error Correcting Codes

Voloch and Walker used canonical liftings and the elliptic Teichmüller lift to create error-correcting codes. The bounds for the parameters (which measure “how good” the resulting codes are likely to be) depend on the degrees of  $F_i$ ’s and  $G_i$ ’s, with lower degrees giving better bounds. They showed that  $F_1$  and  $G_1$  had minimal degrees, making the canonical lifting the natural choice.

# Error Correcting Codes

Voloch and Walker used canonical liftings and the elliptic Teichmüller lift to create error-correcting codes. The bounds for the parameters (which measure “how good” the resulting codes are likely to be) depend on the degrees of  $F_i$ ’s and  $G_i$ ’s, with lower degrees giving better bounds. They showed that  $F_1$  and  $G_1$  had minimal degrees, making the canonical lifting the natural choice.

On the other hand,  $F_i$  and  $G_i$  for  $i \geq 2$  are *not* minimal.

# Error Correcting Codes

Voloch and Walker used canonical liftings and the elliptic Teichmüller lift to create error-correcting codes. The bounds for the parameters (which measure “how good” the resulting codes are likely to be) depend on the degrees of  $F_i$ ’s and  $G_i$ ’s, with lower degrees giving better bounds. They showed that  $F_1$  and  $G_1$  had minimal degrees, making the canonical lifting the natural choice.

On the other hand,  $F_i$  and  $G_i$  for  $i \geq 2$  are *not* minimal.

One should note that, one can construct codes with more general liftings of curves in a very similar way.

# Error Correcting Codes (cont.)

With elliptic curves, we have:

## Theorem

Let  $E/\mathbb{k}$  as above and  $\tilde{E}/W_3(\mathbb{k})$  be a lifting for which we have a lifting of points  $\nu : E(\bar{\mathbb{k}}) \rightarrow \tilde{E}/W_3(\bar{\mathbb{k}})$  having “minimal degrees”. Then  $\tilde{E}$  is the canonical lifting of  $E$  (modulo  $p^3$ )

# Error Correcting Codes (cont.)

With elliptic curves, we have:

## Theorem

Let  $E/\mathbb{k}$  as above and  $\tilde{E}/W_3(\mathbb{k})$  be a lifting for which we have a lifting of points  $\nu : E(\bar{\mathbb{k}}) \rightarrow \tilde{E}/W_3(\bar{\mathbb{k}})$  having “minimal degrees”. Then  $\tilde{E}$  is the canonical lifting of  $E$  (modulo  $p^3$ ) and we have a lifting of the Frobenius on the affine part of  $E$  so that the following diagram commutes:

$$\begin{array}{ccc}
 \tilde{E}(W_3(\bar{\mathbb{k}})) & \xrightarrow{\tilde{\phi}} & \tilde{E}^\sigma(W_3(\bar{\mathbb{k}})) \\
 \pi \left( \begin{array}{c} \uparrow \\ \downarrow \end{array} \right) \nu & & \pi \left( \begin{array}{c} \uparrow \\ \downarrow \end{array} \right) \nu^\sigma \\
 E(\bar{\mathbb{k}}) & \xrightarrow{\phi} & E^\sigma(\bar{\mathbb{k}})
 \end{array}$$

# Error Correcting Codes (cont.)

With elliptic curves, we have:

## Theorem

Let  $E/\mathbb{k}$  as above and  $\tilde{E}/W_3(\mathbb{k})$  be a lifting for which we have a lifting of points  $\nu : E(\bar{\mathbb{k}}) \rightarrow \tilde{E}/W_3(\bar{\mathbb{k}})$  having “minimal degrees”. Then  $\tilde{E}$  is the canonical lifting of  $E$  (modulo  $p^3$ ) and we have a lifting of the Frobenius on the affine part of  $E$  so that the following diagram commutes:

$$\begin{array}{ccc}
 \tilde{E}(W_3(\bar{\mathbb{k}})) & \xrightarrow{\tilde{\phi}} & \tilde{E}^\sigma(W_3(\bar{\mathbb{k}})) \\
 \pi \left( \begin{array}{c} \uparrow \\ \downarrow \end{array} \right) \nu & & \pi \left( \begin{array}{c} \uparrow \\ \downarrow \end{array} \right) \nu^\sigma \\
 E(\bar{\mathbb{k}}) & \xrightarrow{\phi} & E^\sigma(\bar{\mathbb{k}})
 \end{array}$$

Moreover, any supersingular elliptic curve will yield larger degrees.

# Minimal Degree Liftings

Therefore, the notions of *ordinary elliptic curve* and its *canonical lifting* (at least modulo  $p^3$ ) can be defined strictly from the point of view of minimal degree liftings:

# Minimal Degree Liftings

Therefore, the notions of *ordinary elliptic curve* and its *canonical lifting* (at least modulo  $p^3$ ) can be defined strictly from the point of view of minimal degree liftings:

- $E$  is ordinary if there is a lifting satisfying the lower bound on the degrees of the lifting map;



# Minimal Degree Liftings

Therefore, the notions of *ordinary elliptic curve* and its *canonical lifting* (at least modulo  $p^3$ ) can be defined strictly from the point of view of minimal degree liftings:

- $E$  is ordinary if there is a lifting satisfying the lower bound on the degrees of the lifting map;
- $E$  is the canonical lifting of  $E$  if there is a lifting map satisfying the lower bound.

# Minimal Degree Liftings

Therefore, the notions of *ordinary elliptic curve* and its *canonical lifting* (at least modulo  $p^3$ ) can be defined strictly from the point of view of minimal degree liftings:

- $E$  is ordinary if there is a lifting satisfying the lower bound on the degrees of the lifting map;
- $E$  is the canonical lifting of  $E$  if there is a lifting map satisfying the lower bound.

On the other hand, in this way, these notions can be generalized to higher genus curves, and in a very similar way, one can obtain very similar results for *hyperelliptic* curves!

# Mochizuki Liftings

For genus 2 curves (and so hyperelliptic) in characteristic 3, one can have a Mochizuki lifting of the Frobenius if one removes (some) 2 points from the curve.

# Mochizuki Liftings

For genus 2 curves (and so hyperelliptic) in characteristic 3, one can have a Mochizuki lifting of the Frobenius if one removes (some) 2 points from the curve. These two points are invariant by the hyperelliptic involution and thus can be put at “infinity”.

# Mochizuki Liftings

For genus 2 curves (and so hyperelliptic) in characteristic 3, one can have a Mochizuki lifting of the Frobenius if one removes (some) 2 points from the curve. These two points are invariant by the hyperelliptic involution and thus can be put at “infinity”.

We then have:

## Theorem (F.-Mochizuki)

*The notions of “ordinary” and “canonical lifting” (modulo  $p^2$ ) from minimal degree lifting coincide with the ones coming from Mochizuki’s theory.*

# Mochizuki Liftings

For genus 2 curves (and so hyperelliptic) in characteristic 3, one can have a Mochizuki lifting of the Frobenius if one removes (some) 2 points from the curve. These two points are invariant by the hyperelliptic involution and thus can be put at “infinity”.

We then have:

## Theorem (F.-Mochizuki)

*The notions of “ordinary” and “canonical lifting” (modulo  $p^2$ ) from minimal degree lifting coincide with the ones coming from Mochizuki’s theory.*

Thus, we were able to give a concrete example of a family of Mochizuki liftings.

# The $J_n$ Functions

We now return to ordinary elliptic curves and their canonical liftings.

# The $J_n$ Functions

We now return to ordinary elliptic curves and their canonical liftings.

If  $\mathbb{k}^{ord}$  denotes the set of ordinary  $j$ -invariants in  $\mathbb{k}$ , we have functions  $J_i : \mathbb{k}^{ord} \rightarrow \mathbb{k}$  such that  $(j_0, J_1(j_0), J_2(j_0), \dots)$  is the  $j$ -invariant of the canonical lifting of the curve with  $j$ -invariant  $j_0 \in \mathbb{k}^{ord}$ .



# The $J_n$ Functions

We now return to ordinary elliptic curves and their canonical liftings.

If  $\mathbb{k}^{ord}$  denotes the set of ordinary  $j$ -invariants in  $\mathbb{k}$ , we have functions  $J_i : \mathbb{k}^{ord} \rightarrow \mathbb{k}$  such that  $(j_0, J_1(j_0), J_2(j_0), \dots)$  is the  $j$ -invariant of the canonical lifting of the curve with  $j$ -invariant  $j_0 \in \mathbb{k}^{ord}$ .

## Mazur's Question (to John Tate)

What kind of functions are these  $J_n$ ? Can one say anything about them?

# First Computations

## Examples:

$$p = 5 \quad \blacksquare \quad J_1 = 3j_0^3 + j_0^4;$$

# First Computations

## Examples:

$$p = 5 \quad \blacksquare \quad J_1 = 3j_0^3 + j_0^4;$$

$$p = 7 \quad \blacksquare \quad J_1 = 3j_0^5 + 5j_0^6;$$

# First Computations

## Examples:

$$p = 5 \quad \blacksquare \quad J_1 = 3j_0^3 + j_0^4;$$

**Question:** Can these functions all be polynomials?

$$p = 7 \quad \blacksquare \quad J_1 = 3j_0^5 + 5j_0^6;$$

# First Computations

## Examples:

$$p = 5 \quad \begin{array}{l} \blacksquare J_1 = 3j_0^3 + j_0^4; \\ \blacksquare J_2 = \\ \quad 3j_0^5 + 2j_0^{10} + 2j_0^{13} + 4j_0^{14} + 4j_0^{15} + 4j_0^{16} + j_0^{17} + 4j_0^{18} + j_0^{19} + j_0^{20} + 3j_0^{23} + j_0^{24}. \end{array}$$

**Question:** Can these functions all be polynomials?

$$p = 7 \quad \blacksquare J_1 = 3j_0^5 + 5j_0^6;$$

# First Computations

## Examples:

$$p = 5 \quad \begin{array}{l} \blacksquare J_1 = 3j_0^3 + j_0^4; \\ \blacksquare J_2 = \\ \quad 3j_0^5 + 2j_0^{10} + 2j_0^{13} + 4j_0^{14} + 4j_0^{15} + 4j_0^{16} + j_0^{17} + 4j_0^{18} + j_0^{19} + j_0^{20} + 3j_0^{23} + j_0^{24}. \end{array}$$

**Question:** Can these functions all be polynomials?

$$p = 7 \quad \begin{array}{l} \blacksquare J_1 = 3j_0^5 + 5j_0^6; \\ \blacksquare J_2 = \\ \quad (3j_0^{21} + 6j_0^{28} + 3j_0^{33} + 5j_0^{34} + 4j_0^{35} + 2j_0^{36} + 3j_0^{37} + 6j_0^{38} + 3j_0^{39} + 5j_0^{40} + 5j_0^{41} + \\ \quad 5j_0^{42} + 2j_0^{43} + 3j_0^{44} + 6j_0^{45} + 3j_0^{46} + 5j_0^{47} + 5j_0^{48} + 3j_0^{49} + 3j_0^{54} + 5j_0^{55}) / (1 + j_0^7). \end{array}$$

# First Computations

## Examples:

$$p = 5 \quad \begin{array}{l} \blacksquare J_1 = 3j_0^3 + j_0^4; \\ \blacksquare J_2 = \\ 3j_0^5 + 2j_0^{10} + 2j_0^{13} + 4j_0^{14} + 4j_0^{15} + 4j_0^{16} + j_0^{17} + 4j_0^{18} + j_0^{19} + j_0^{20} + 3j_0^{23} + j_0^{24}. \end{array}$$

**Question:** Can these functions all be polynomials?

$$p = 7 \quad \begin{array}{l} \blacksquare J_1 = 3j_0^5 + 5j_0^6; \\ \blacksquare J_2 = \\ (3j_0^{21} + 6j_0^{28} + 3j_0^{33} + 5j_0^{34} + 4j_0^{35} + 2j_0^{36} + 3j_0^{37} + 6j_0^{38} + 3j_0^{39} + 5j_0^{40} + 5j_0^{41} + \\ 5j_0^{42} + 2j_0^{43} + 3j_0^{44} + 6j_0^{45} + 3j_0^{46} + 5j_0^{47} + 5j_0^{48} + 3j_0^{49} + 3j_0^{54} + 5j_0^{55}) / (1 + j_0^7). \end{array}$$

**Note:** If  $j_0 = -1$ , then  $E$  is **supersingular**, i.e., no canonical lifting.

# Pseudo-Canonical Liftings

## (Superficial) Answer to Mazur's Question

For any  $p$ , we have that  $J_n \in \mathbb{F}_p(X)$ .



# Pseudo-Canonical Liftings

## (Superficial) Answer to Mazur's Question

For any  $p$ , we have that  $J_n \in \mathbb{F}_p(X)$ .

## Tate's Question

Is there a **supersingular** value of  $j_0$  (for some fixed characteristic  $p$ ) for which all functions  $J_n$  are regular at  $j_0$ . (E.g.,  $j_0 = 0$  for  $p = 5$  for  $J_1$  and  $J_2$ ?)

# Pseudo-Canonical Liftings

## (Superficial) Answer to Mazur's Question

For any  $p$ , we have that  $J_n \in \mathbb{F}_p(X)$ .

## Tate's Question

Is there a **supersingular** value of  $j_0$  (for some fixed characteristic  $p$ ) for which all functions  $J_n$  are regular at  $j_0$ . (E.g.,  $j_0 = 0$  for  $p = 5$  for  $J_1$  and  $J_2$ ?)

This lead us to define:

## Definition

The elliptic curve over  $\mathbf{W}(\mathbb{k})$  given by  $j \stackrel{\text{def}}{=} (j_0, J_1(j_0), J_2(j_0), \dots)$  for such a supersingular  $j_0$  is a **pseudo-canonical lifting** of the elliptic curve given by  $j_0$ .

# Pseudo-Canonical Liftings

## (Superficial) Answer to Mazur's Question

For any  $p$ , we have that  $J_n \in \mathbb{F}_p(X)$ .

## Tate's Question

Is there a **supersingular** value of  $j_0$  (for some fixed characteristic  $p$ ) for which all functions  $J_n$  are regular at  $j_0$ . (E.g.,  $j_0 = 0$  for  $p = 5$  for  $J_1$  and  $J_2$ ?)

This lead us to define:

## Definition

The elliptic curve over  $W(\mathbb{k})$  given by  $j \stackrel{\text{def}}{=} (j_0, J_1(j_0), J_2(j_0), \dots)$  for such a supersingular  $j_0$  is a **pseudo-canonical lifting** of the elliptic curve given by  $j_0$ . **Tate's question:** do they exist at all?

# Answer to Tate's Question

## Theorem

Let  $j_0 \notin \mathbb{k}^{ord}$  and  $p \geq 5$ . Then:

# Answer to Tate's Question

## Theorem

Let  $j_0 \notin \mathbb{k}^{ord}$  and  $p \geq 5$ . Then:

- 1  $J_1$  is regular at  $j_0$  if, and only if,  $j_0$  is either 0 or 1728.

# Answer to Tate's Question

## Theorem

Let  $j_0 \notin \mathbb{k}^{ord}$  and  $p \geq 5$ . Then:

- 1  $J_1$  is regular at  $j_0$  if, and only if,  $j_0$  is either 0 or 1728.
- 2  $J_2$  is regular at  $j_0$  if, and only if,  $j_0$  is 0.

# Answer to Tate's Question

## Theorem

Let  $j_0 \notin \mathbb{k}^{ord}$  and  $p \geq 5$ . Then:

- 1  $J_1$  is regular at  $j_0$  if, and only if,  $j_0$  is either 0 or 1728.
- 2  $J_2$  is regular at  $j_0$  if, and only if,  $j_0$  is 0.
- 3 For  $n \geq 3$ , we have that  $J_n$  is never regular at  $j_0$ .

# Answer to Tate's Question

## Theorem

Let  $j_0 \notin \mathbb{k}^{\text{ord}}$  and  $p \geq 5$ . Then:

- 1  $J_1$  is regular at  $j_0$  if, and only if,  $j_0$  is either 0 or 1728.
- 2  $J_2$  is regular at  $j_0$  if, and only if,  $j_0$  is 0.
- 3 For  $n \geq 3$ , we have that  $J_n$  is never regular at  $j_0$ .

For  $p = 2, 3$  (in which case only  $j_0 = 0$  is supersingular), we have that  $J_i$  is regular at 0 if, and only if,  $i \leq 11$  for  $p = 2$



# Answer to Tate's Question

## Theorem

Let  $j_0 \notin \mathbb{k}^{\text{ord}}$  and  $p \geq 5$ . Then:

- 1  $J_1$  is regular at  $j_0$  if, and only if,  $j_0$  is either 0 or 1728.
- 2  $J_2$  is regular at  $j_0$  if, and only if,  $j_0$  is 0.
- 3 For  $n \geq 3$ , we have that  $J_n$  is never regular at  $j_0$ .

For  $p = 2, 3$  (in which case only  $j_0 = 0$  is supersingular), we have that  $J_i$  is regular at 0 if, and only if,  $i \leq 11$  for  $p = 2$  or  $i \leq 5$  for  $p = 3$ .

# Answer to Tate's Question

## Theorem

Let  $j_0 \notin \mathbb{k}^{\text{ord}}$  and  $p \geq 5$ . Then:

- 1  $J_1$  is regular at  $j_0$  if, and only if,  $j_0$  is either 0 or 1728.
- 2  $J_2$  is regular at  $j_0$  if, and only if,  $j_0$  is 0.
- 3 For  $n \geq 3$ , we have that  $J_n$  is never regular at  $j_0$ .

For  $p = 2, 3$  (in which case only  $j_0 = 0$  is supersingular), we have that  $J_i$  is regular at 0 if, and only if,  $i \leq 11$  for  $p = 2$  or  $i \leq 5$  for  $p = 3$ .

So, (unrestricted) pseudo-canonical liftings don't exist.

# Answer to Mazur's Question

We need some notation: let

$$\mathrm{ss}_p(X) \stackrel{\mathrm{def}}{=} \prod_{j \text{ supers.}} (X - j)$$

(the *supersingular polynomial*)

# Answer to Mazur's Question

We need some notation: let

$$\mathrm{ss}_p(X) \stackrel{\mathrm{def}}{=} \prod_{j \text{ supers.}} (X - j)$$

(the *supersingular polynomial*) and

$$S_p(X) \stackrel{\mathrm{def}}{=} \prod_{\substack{j \text{ supers.} \\ j \neq 0, 1728}} (X - j).$$

# Answer to Mazur's Question

We need some notation: let

$$\mathrm{ss}_p(X) \stackrel{\mathrm{def}}{=} \prod_{j \text{ supers.}} (X - j)$$

(the *supersingular polynomial*) and

$$S_p(X) \stackrel{\mathrm{def}}{=} \prod_{\substack{j \text{ supers.} \\ j \neq 0, 1728}} (X - j).$$

One then has that  $\mathrm{ss}_p(X), S_p(X) \in \mathbb{F}_p[X]$ , and  $S_p(0), S_p(1728) \neq 0$ .

# Answer to Mazur's Question

We need some notation: let

$$\mathrm{ss}_p(X) \stackrel{\mathrm{def}}{=} \prod_{j \text{ supers.}} (X - j)$$

(the *supersingular polynomial*) and

$$S_p(X) \stackrel{\mathrm{def}}{=} \prod_{\substack{j \text{ supers.} \\ j \neq 0, 1728}} (X - j).$$

One then has that  $\mathrm{ss}_p(X), S_p(X) \in \mathbb{F}_p[X]$ , and  $S_p(0), S_p(1728) \neq 0$ . Also, let

$$\iota = \begin{cases} 8, & \text{if } p = 2; \\ 3, & \text{if } p = 3; \\ 2, & \text{if } p = 31; \\ 1, & \text{otherwise.} \end{cases}$$

# Answer to Mazur's Question

Then, we have:

## Theorem

Let  $J_i = F_i/G_i$ , with  $F_i, G_i \in \mathbb{F}_p[X]$ ,  $(F_i, G_i) = 1$ , and  $G_i$  monic.

# Answer to Mazur's Question

Then, we have:

## Theorem

Let  $J_i = F_i/G_i$ , with  $F_i, G_i \in \mathbb{F}_p[X]$ ,  $(F_i, G_i) = 1$ , and  $G_i$  monic. Also, let  $r_i = (i-1)p^{i-1}$ ,  $s_i = ((i-3)p^i + ip^{i-1})/3$  and  $s'_i = \max\{0, s_i\}$ . Then, for all  $i \in \mathbb{Z}_{>0}$  we have:



# Answer to Mazur's Question

Then, we have:

## Theorem

Let  $J_i = F_i/G_i$ , with  $F_i, G_i \in \mathbb{F}_p[X]$ ,  $(F_i, G_i) = 1$ , and  $G_i$  monic. Also, let  $r_i = (i-1)p^{i-1}$ ,  $s_i = ((i-3)p^i + ip^{i-1})/3$  and  $s'_i = \max\{0, s_i\}$ .

Then, for all  $i \in \mathbb{Z}_{>0}$  we have:

$$1 \quad \deg F_i - \deg G_i = p^i - \iota;$$

# Answer to Mazur's Question

Then, we have:

## Theorem

Let  $J_i = F_i/G_i$ , with  $F_i, G_i \in \mathbb{F}_p[X]$ ,  $(F_i, G_i) = 1$ , and  $G_i$  monic. Also, let  $r_i = (i-1)p^{i-1}$ ,  $s_i = ((i-3)p^i + ip^{i-1})/3$  and  $s'_i = \max\{0, s_i\}$ .

Then, for all  $i \in \mathbb{Z}_{>0}$  we have:

- 1  $\deg F_i - \deg G_i = p^i - \iota$ ;
- 2 if  $p \geq 5$ , then  $G_i = S_p(X)^{ip^{i-1} + (i-1)p^{i-2}} \cdot H_i$ , where  $H_i \mid X^{s'_i} \cdot (X - 1728)^{r_i}$ ;

# Answer to Mazur's Question

Then, we have:

## Theorem

Let  $J_i = F_i/G_i$ , with  $F_i, G_i \in \mathbb{F}_p[X]$ ,  $(F_i, G_i) = 1$ , and  $G_i$  monic. Also, let  $r_i = (i-1)p^{i-1}$ ,  $s_i = ((i-3)p^i + ip^{i-1})/3$  and  $s'_i = \max\{0, s_i\}$ .

Then, for all  $i \in \mathbb{Z}_{>0}$  we have:

- 1  $\deg F_i - \deg G_i = p^i - \iota$ ;
- 2 if  $p \geq 5$ , then  $G_i = S_p(X)^{ip^{i-1} + (i-1)p^{i-2}} \cdot H_i$ , where  $H_i \mid X^{s'_i} \cdot (X - 1728)^{r_i}$ ;
- 3 if  $p = 2, 3$ , then  $G_i = X^{t_i}$ , where  $t_i \leq p^i$ .

# Answer to Mazur's Question

Then, we have:

## Theorem

Let  $J_i = F_i/G_i$ , with  $F_i, G_i \in \mathbb{F}_p[X]$ ,  $(F_i, G_i) = 1$ , and  $G_i$  monic. Also, let  $r_i = (i-1)p^{i-1}$ ,  $s_i = ((i-3)p^i + ip^{i-1})/3$  and  $s'_i = \max\{0, s_i\}$ .

Then, for all  $i \in \mathbb{Z}_{>0}$  we have:

- 1  $\deg F_i - \deg G_i = p^i - \iota$ ;
- 2 if  $p \geq 5$ , then  $G_i = S_p(X)^{ip^{i-1} + (i-1)p^{i-2}} \cdot H_i$ , where  $H_i \mid X^{s'_i} \cdot (X - 1728)^{r_i}$ ;
- 3 if  $p = 2, 3$ , then  $G_i = X^{t_i}$ , where  $t_i \leq p^i$ .

Also, there is a formula for  $J_i(X)$  (which can be simplified if  $p \geq 3$ ) obtained from the *classical modular polynomial*.

# Modular Functions

Assume from now  $p \geq 5$ .

# Modular Functions

Assume from now  $p \geq 5$ . Another perspective: if  $E/\mathbb{k}$ , ordinary, is given by  $y_0^2 = x_0^3 + a_0x_0 + b_0$ , and  $\mathbf{E}/W$  is its canonical lifting and (after some “choice”) is given by  $\mathbf{y}^2 = \mathbf{x}^3 + \mathbf{a}\mathbf{x} + \mathbf{b}$ , then

# Modular Functions

Assume from now  $p \geq 5$ . Another perspective: if  $E/\mathbb{k}$ , ordinary, is given by  $y_0^2 = x_0^3 + a_0x_0 + b_0$ , and  $E/W$  is its canonical lifting and (after some “choice”) is given by  $y^2 = x^3 + ax + b$ , then

$$a = (A_0, A_1, A_2, \dots),$$

$$b = (B_0, B_1, B_2, \dots),$$

where  $A_i, B_i \in \mathbb{k}(a_0, b_0)$ .

# Modular Functions

Assume from now  $p \geq 5$ . Another perspective: if  $E/\mathbb{k}$ , ordinary, is given by  $y_0^2 = x_0^3 + a_0x_0 + b_0$ , and  $\mathbf{E}/W$  is its canonical lifting and (after some “choice”) is given by  $y^2 = x^3 + \mathbf{a}x + \mathbf{b}$ , then

$$\mathbf{a} = (A_0, A_1, A_2, \dots),$$

$$\mathbf{b} = (B_0, B_1, B_2, \dots),$$

where  $A_i, B_i \in \mathbb{k}(a_0, b_0)$ . In fact, if  $\mathcal{H}$  is the *Hasse invariant* of  $E$  (i.e., the coefficient of  $x_0^{p-1}$  is  $(x_0^3 + a_0x_0 + b_0)^{(p-1)/2}$ ), then  $A_i, B_i$  possibly have poles only at the zeros of  $\mathcal{H}$



# Modular Functions

Assume from now  $p \geq 5$ . Another perspective: if  $E/\mathbb{k}$ , ordinary, is given by  $y_0^2 = x_0^3 + a_0x_0 + b_0$ , and  $\mathbf{E}/W$  is its canonical lifting and (after some “choice”) is given by  $y^2 = x^3 + \mathbf{a}x + \mathbf{b}$ , then

$$\mathbf{a} = (A_0, A_1, A_2, \dots),$$

$$\mathbf{b} = (B_0, B_1, B_2, \dots),$$

where  $A_i, B_i \in \mathbb{k}(a_0, b_0)$ . In fact, if  $\mathcal{H}$  is the *Hasse invariant* of  $E$  (i.e., the coefficient of  $x_0^{p-1}$  is  $(x_0^3 + a_0x_0 + b_0)^{(p-1)/2}$ ), then  $A_i, B_i$  possibly have poles only at the zeros of  $\mathcal{H}$  (or  $\Delta = 4a_0^3 + 27b_0^2$ ).

# Modular Functions

Assume from now  $p \geq 5$ . Another perspective: if  $E/\mathbb{k}$ , ordinary, is given by  $y_0^2 = x_0^3 + a_0x_0 + b_0$ , and  $\mathbf{E}/W$  is its canonical lifting and (after some “choice”) is given by  $y^2 = x^3 + \mathbf{a}x + \mathbf{b}$ , then

$$\mathbf{a} = (A_0, A_1, A_2, \dots),$$

$$\mathbf{b} = (B_0, B_1, B_2, \dots),$$

where  $A_i, B_i \in \mathbb{k}(a_0, b_0)$ . In fact, if  $\mathcal{H}$  is the *Hasse invariant* of  $E$  (i.e., the coefficient of  $x_0^{p-1}$  is  $(x_0^3 + a_0x_0 + b_0)^{(p-1)/2}$ ), then  $A_i, B_i$  possibly have poles only at the zeros of  $\mathcal{H}$  (or  $\Delta = 4a_0^3 + 27b_0^2$ ).

## Question

What are the weights of the  $A_i$ 's and  $B_i$ 's?

# Modular Functions

Assume from now  $p \geq 5$ . Another perspective: if  $E/\mathbb{k}$ , ordinary, is given by  $y_0^2 = x_0^3 + a_0x_0 + b_0$ , and  $\mathbf{E}/W$  is its canonical lifting and (after some “choice”) is given by  $y^2 = x^3 + \mathbf{a}x + \mathbf{b}$ , then

$$\mathbf{a} = (A_0, A_1, A_2, \dots),$$

$$\mathbf{b} = (B_0, B_1, B_2, \dots),$$

where  $A_i, B_i \in \mathbb{k}(a_0, b_0)$ . In fact, if  $\mathcal{H}$  is the *Hasse invariant* of  $E$  (i.e., the coefficient of  $x_0^{p-1}$  is  $(x_0^3 + a_0x_0 + b_0)^{(p-1)/2}$ ), then  $A_i, B_i$  possibly have poles only at the zeros of  $\mathcal{H}$  (or  $\Delta = 4a_0^3 + 27b_0^2$ ).

## Question

What are the weights of the  $A_i$ 's and  $B_i$ 's? What are the order of the poles?



# Modular Functions (cont.)

## Conjecture

1  $A_i$  has weight  $4p^i$ .

# Modular Functions (cont.)

## Conjecture

- 1  $A_i$  has weight  $4p^i$ .
- 2  $B_i$  has weight  $6p^i$ .

# Modular Functions (cont.)

## Conjecture

- 1  $A_i$  has weight  $4p^i$ .
- 2  $B_i$  has weight  $6p^i$ .
- 3  $A_i$  and  $B_i$  have poles of order at most  $(i-1)p+1$  at the zeros of  $\mathcal{H}$ .

# Modular Functions (cont.)

## Conjecture

- 1  $A_i$  has weight  $4p^i$ .
- 2  $B_i$  has weight  $6p^i$ .
- 3  $A_i$  and  $B_i$  have poles of order at most  $(i-1)p+1$  at the zeros of  $\mathcal{H}$ .  
(At least for  $i \leq 2$ . Not enough data yet.)

# Modular Functions (cont.)

## Conjecture

- 1  $A_i$  has weight  $4p^i$ .
- 2  $B_i$  has weight  $6p^i$ .
- 3  $A_i$  and  $B_i$  have poles of order at most  $(i-1)p+1$  at the zeros of  $\mathcal{H}$ .  
(At least for  $i \leq 2$ . Not enough data yet.)
- 4  $A_i$  and  $B_i$  have no zeros at zeros of  $\Delta$ .



# Modular Functions (cont.)

## Conjecture

- 1  $A_i$  has weight  $4p^i$ .
- 2  $B_i$  has weight  $6p^i$ .
- 3  $A_i$  and  $B_i$  have poles of order at most  $(i-1)p+1$  at the zeros of  $\mathcal{H}$ .  
(At least for  $i \leq 2$ . Not enough data yet.)
- 4  $A_i$  and  $B_i$  have no zeros at zeros of  $\Delta$ .

So, if true, the isomorphism  $(a_0, b_0) \leftrightarrow (\lambda_0^4 a_0, \lambda_0^6 b_0)$  corresponds to the isomorphism  $(\mathbf{a}, \mathbf{b}) \leftrightarrow (\boldsymbol{\lambda}^4 \mathbf{a}, \boldsymbol{\lambda}^6 \mathbf{b})$ , where  $\boldsymbol{\lambda} = \tau(\lambda_0) = (\lambda_0, 0, 0, \dots)$ .

Thank you!

# Thank you!

And sorry for going over...