



Learning  
Variation  
via Optimal  
Control

Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

LVOC

New Insight  
LVOC  
 $\epsilon$ -Constrained  
Method

Other  
Variation or  
Functional  
Method

Active Contour  
...

Experimental  
Results

Pre-analysis  
Results

# Learning Variation via Optimal Control

CVBIOUC



# Contents

Learning  
Variation  
via Optimal  
Control

## 1 Introduction

- Numerical Methods and Applications in Total Variation Image Restoration
- L-PDE

## 2 LVOC

- New Insight
- LVOC
- $\epsilon$ -Constrained Method

## 3 Other Variation or Functional Method

- Active Contour
- ...

## 4 Experimental Results

- Pre-analysis
- Results

Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

LVOC

New Insight  
LVOC  
 $\epsilon$ -Constrained  
Method

Other  
Variation or  
Functional  
Method

Active Contour  
...

Experimental  
Results

Pre-analysis  
Results



# Learning Variation via Optimal Control

## Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

## LVOC

New Insight  
LVOC  
 $\epsilon$ -Constrained  
Method

Other  
Variation or  
Functional  
Method

Active Contour  
...


## Experimental Results

Pre-analysis  
Results



The total variation of an image  $u : \Omega \rightarrow \mathbb{R}$  is defined as:

$$\int_{\Omega} |\nabla u| dx \quad (1)$$



## Deblurring

Rudin L, Osher S, Fatemi E (1992) Nonlinear total variation based noise removal algorithms. Physica D 60:259–268



# Learning Variation via Optimal Control

## Introduction

Numerical Methods and Applications in Total Variation Image Restoration  
L-PDE

## LVOC

New Insight  
LVOC  
 $\epsilon$ -Constrained Method

Other Variation or Functional Method

Active Contour  
...

## Experimental Results

Pre-analysis Results



The total variation of an image  $u : \Omega \rightarrow \mathbb{R}$  is defined as:

$$\int_{\Omega} |\nabla u| dx$$



## Inpainting

Rudin L, Osher S, Fatemi E (1992) Nonlinear total variation based noise removal algorithms. Physica D 60:259–268



# Learning Variation via Optimal Control

## Introduction

Numerical Methods and Applications in Total Variation Image Restoration  
L-PDE

## LVOC


New Insight  
LVOC  
 $\epsilon$ -Constrained Method

Other Variation or Functional Method

Active Contour  
...

## Experimental Results

Pre-analysis Results



The total variation of an image  $u : \Omega \rightarrow \mathbb{R}$  is defined as:

$$\int_{\Omega} |\nabla u| dx$$



## Decomposition

Rudin L, Osher S, Fatemi E (1992) Nonlinear total variation based noise removal algorithms. Physica D 60:259–268



# Learning Variation via Optimal Control

## Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration

L-PDE

## LVOC

New Insight

LVOC

$\epsilon$ -Constrained  
Method

Other  
Variation or  
Functional  
Method

Active Contour

...

## Experimental Results

Pre-analysis

Results



The unconstrained total variation based image restoration model reads:

$$\min_u \int_{\Omega} \frac{1}{2} (u - f)^2 d\mathbf{x} + \beta \int_{\Omega} |\nabla u| d\mathbf{x} \quad (2)$$



# Learning Variation via Optimal Control

## Introduction

Numerical Methods and Applications in Total Variation Image Restoration  
L-PDE

## LVOC

New Insight  
LVOC  
 $\epsilon$ -Constrained Method


Other Variation or Functional Method

Active Contour

...

## Experimental Results

Pre-analysis Results



The corresponding Euler-Lagrange equation of (2):

$$K^T Ku - K^T f - \beta \operatorname{div} \left( \frac{\nabla u}{|\nabla u|} \right) = 0 \quad (3)$$

With the help of an auxiliary variable  $\mathbf{p} = \frac{\nabla u}{\sqrt{|\nabla u|^2 + \epsilon}}$ , the singularity will vanish.

$$\begin{aligned} \nabla u - \mathbf{p} \sqrt{|\nabla u|^2 + \epsilon} &= 0 \\ K^T Ku - K^T f - \beta \operatorname{div} \mathbf{p} &= 0 \end{aligned} \quad (4)$$

---

Chan T, Golub G, Mulet P (1999) A nonlinear primal-dual method for total variation-based image restoration. SIAM J Sci Comp 20:1964-1977



# Learning Variation via Optimal Control

## Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

## LVOC

New Insight

LVOC

$\epsilon$ -Constrained  
Method

Other  
Variation or  
Functional  
Method


Active Contour

...

Experimental  
Results

Pre-analysis

Results



A pur dual method which is equivalent to the ROF model avoids the singularity of  $|\nabla u| = 0$ .

$$\sup_{|p| \leq 1} \left\{ -\frac{\lambda^2}{2} \int_{\Omega} |K^{-T} \operatorname{div} p - \frac{f}{\lambda}|^2 dx \right\} \quad (5)$$

Semi-implicit scheme:

$$p^{n+1} = \frac{p^n + \tau H(p^n)}{p^n + \tau |H(p^n)|} \quad (6)$$

where  $H(p) := \nabla[(K^T K)^{-1} \operatorname{div} p - \frac{1}{\lambda} K^{-1} f]$

Chambolle A (2004) An algorithm for total variation minimization and applications. J Math Imaging Vis 20:89–97





# Learning Variation via Optimal Control

- Introduction
- Numerical Methods and Applications in Total Variation Image Restoration
- L-PDE
- LVOC
- New Insight
- LVOC
- $\epsilon$ -Constrained Method
- Other
- Variation or Functional Method
- Active Contour
- ...
- Experimental Results
- Pre-analysis
- Results



Consider the general quadratic problem:

$$\min_{y, y \leq \phi} \quad \langle y, Ay \rangle - \langle f, y \rangle \tag{7}$$

KKT conditions:

$$\begin{aligned} Ay + \lambda &= f \\ \lambda \odot (\phi - y) &= 0 \\ \lambda &\geq 0 \\ \phi - y &\geq 0 \end{aligned} \tag{8}$$

---

Hintermüller M, Stadler G (2006) A primal-dual algorithm for TV-based inf-convolution-type image restoration. SIAM J Sci Comput 28:1–23



# Learning Variation via Optimal Control

## Introduction

Numerical Methods and Applications in Total Variation Image Restoration  
L-PDE

## LVOC

New Insight  
LVOC  
 $\epsilon$ -Constrained Method


Other  
Variation or Functional Method

Active Contour

...

## Experimental Results

Pre-analysis  
Results



The non-negative constrained TV model:

$$\min_{u, u \geq 0} \int_{\Omega} \frac{1}{2} (Ku - f)^2 d\mathbf{x} + \beta \int_{\Omega} |\nabla u| d\mathbf{x} \quad (9)$$

The equivalent system:

$$\begin{aligned} \nabla u - \mathbf{p} \sqrt{|\nabla u|^2 + \epsilon} &= 0 \\ (K^T K + \alpha I)u - K^T f - \beta \operatorname{div} \mathbf{p} - \lambda &= 0 \\ \lambda - \max\{0, \lambda - cu\} &= 0 \end{aligned} \quad (10)$$

---

Krishnan D, Lin P, Yip A (2007) A primal-dual active-set method for non-negativity constrained total variation deblurring problems. IEEE Trans Image Process 16(11):2766–2777



# Learning Variation via Optimal Control

## Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration

L-PDE

LVOC

New Insight

LVOC

$\epsilon$ -Constrained  
Method

Other  
Variation or  
Functional  
Method

Active Contour

...

Experimental  
Results

Pre-analysis

Results



Associated with the Fisher-Burmeister function  $\phi$ , the KKT system is shown as follows.

$$\begin{aligned}\mu \mathbf{p} - H(\mathbf{p}) &= 0 \\ \phi(\mu, 1 - |\mathbf{p}|^2) &= 0\end{aligned}\tag{11}$$



# Learning Variation via Optimal Control

## Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

## LVOC

New Insight  
LVOC  
 $\epsilon$ -Constrained  
Method

Other  
Variation or  
Functional  
Method

Active Contour  
...

## Experimental Results

Pre-analysis  
Results



The primal-dual formulation:

$$F(u, \mathbf{p}) := \frac{1}{2} \int_{\Omega} (Ku - f)^2 d\mathbf{x} + \beta \int_{\Omega} u \operatorname{div} \mathbf{p} d\mathbf{x} \quad (12)$$

Two subproblems:

$$\sup_{|\mathbf{p}| \leq 1} F(u, \mathbf{p}) \quad \inf_u F(u, \mathbf{p}) \quad (13)$$



# Learning Variation via Optimal Control

## Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

## LVOC

New Insight  
LVOC  
 $\epsilon$ -Constrained  
Method

## Other Variation or Functional Method

Active Contour  
...

## Experimental Results

Pre-analysis

Results

## Numerical Methods:

- Dual and Primal-Dual Methods
- Bregman Iteration
- Graph Cut Methods
- Splitting Methods
- Quadratic Programming
- Second-Order Cone Programming
- Majorization-Minimization



# Learning Variation via Optimal Control

## Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration

### L-PDE

## LVOC

New Insight

LVOC

$\epsilon$ -Constrained  
Method

Other  
Variation or  
Functional  
Method

Active Contour

...

## Experimental Results

Pre-analysis

Results

## Optimal Control Governed by Evolutionary PDEs:

minimize  $J(f, u)$ , where  $u \in \mathcal{U}$  controls  $f$  via the following PDE:

$$\begin{cases} f_t = L(u, \beta), & (x, t) \in \Omega \\ f = 0, & (x, t) \in \Gamma \\ f|_{t=0} = f_0 & x \in \Omega \end{cases} \quad (14)$$


where  $L(u, \beta) = \kappa(u) + F(u, \beta)$ .



## Adaptive partial differential equation learning for visual saliency detection

Saliency diffusion is formulated as an evolutionary PDE:

$$\frac{\partial f(\mathbf{p}, t)}{\partial t} = \text{div}(K_{\mathbf{p}} \nabla f(\mathbf{p})) + \lambda(f(\mathbf{p}) - g(\mathbf{p})), \mathbf{p} \in \mathcal{S} \quad (15)$$



Liu R, Cao J, Lin Z, Shan S (2014) Adaptive partial differential equation learning for visual saliency detection. In Proceedings of IEEE conference on computer vision and pattern recognition



Learning  
Variation  
via Optimal  
Control

Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

LVOC

New Insight

LVOC

$\epsilon$ -Constrained  
Method

Other

Variation or  
Functional  
Method

Active Contour

...

Experimental  
Results

Pre-analysis

Results

The non-negative constrained TV model:

$$\min_{u, u \geq 0} \int_{\Omega} \frac{1}{2} (Ku - f)^2 d\mathbf{x} + \beta \int_{\Omega} |\nabla u| d\mathbf{x}$$

The equivalent system:

$$\begin{aligned} \nabla u - p \sqrt{|\nabla u|^2 + \epsilon} &= 0 \\ K^T Ku - K^T f - \beta \operatorname{div} p - \lambda &= 0 \\ \lambda - \max\{0, \lambda - cu\} &= 0 \end{aligned}$$





# Learning Variation via Optimal Control

## Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

## LVOC

New Insight

LVOC

$\epsilon$ -Constrained  
Method

Other  
Variation or  
Functional  
Method

Active Contour

...

Experimental  
Results

Pre-analysis

Results

What if beta Is adaptive?

The non-negative constrained TV model:

$$\min_{u, u \geq 0} \int_{\Omega} \frac{1}{2} (Ku - f)^2 d\mathbf{x} + \beta \int_{\Omega} |\nabla u| d\mathbf{x}$$

The equivalent system:

$$\begin{aligned} \nabla u - p \sqrt{|\nabla u|^2 + \epsilon} &= 0 \\ K^T Ku - K^T f - \beta \operatorname{div} p - \lambda &= 0 \\ \lambda - \max\{0, \lambda - cu\} &= 0 \end{aligned}$$



How?

The non-negative constrained TV model:

$$\min_{u, u \geq 0} \int_{\Omega} \frac{1}{2} (Ku - f)^2 d\mathbf{x} + \beta \int_{\Omega} |\nabla u| d\mathbf{x}$$

The semi-smooth Newton's update:

$$\begin{bmatrix} |\nabla u^k|_{\epsilon} & -(I - \frac{p^k (\nabla u^k)^T}{|\nabla u^k|_{\epsilon}}) \nabla & 0 \\ -\beta^k \operatorname{div} & K^T K & -I \\ 0 & \frac{\partial F_3}{\partial u} & \frac{\partial F_3}{\partial \lambda} \end{bmatrix} \begin{bmatrix} \delta p \\ \delta u \\ \delta \lambda \end{bmatrix} = - \begin{bmatrix} F_1^k \\ F_2^k \\ F_3^k \end{bmatrix}$$

$\beta^k$  reflects the difference between  $u^k$  and the groundtruth.



Train It!

The non-negative constrained TV model:

$$\min_{u, u \geq 0} \int_{\Omega} \frac{1}{2} (Ku - f)^2 d\mathbf{x} + \beta \int_{\Omega} |\nabla u| d\mathbf{x}$$

The semi-smooth Newton's update:

$$\begin{bmatrix} |\nabla u^k|_{\epsilon} & -(I - \frac{p^k (\nabla u^k)^T}{|\nabla u^k|_{\epsilon}}) \nabla & 0 \\ -\beta^k \operatorname{div} & K^T K & -I \\ 0 & \frac{\partial F_3}{\partial u} & \frac{\partial F_3}{\partial \lambda} \end{bmatrix} \begin{bmatrix} \delta p \\ \delta u \\ \delta \lambda \end{bmatrix} = - \begin{bmatrix} F_1^k \\ F_2^k \\ F_3^k \end{bmatrix}$$

$\beta^k$  reflects the difference between  $u^k$  and the groundtruth.



# Learning Variation via Optimal Control

## Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

## LVOC

New Insight

### LVOC

$\epsilon$ -Constrained  
Method

## Other

Variation or  
Functional  
Method

Active Contour

...

## Experimental Results

Pre-analysis

Results

The non-negative constrained variation:

$$\min_{u, u \geq 0} \int_{\Omega} \left[ \frac{1}{2} (Ku - f)^2 + V(\beta, u) \right] d\mathbf{x} \quad (16)$$

where  $V(\beta, u) = \sum_{i=1}^3 \beta_i v_i(u)$ .

- $v_2(u) = |u|_{TV}$
- $v_1(u) = |u|^2$
- $v_3(u) = |\nabla u|^2$



# Learning Variation via Optimal Control

## Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

## LVOC

New Insight  
LVOC  
 $\epsilon$ -Constrained  
Method

## Other Variation or Functional Method

Active Contour  
...

## Experimental Results

Pre-analysis  
Results

Optimal control:

$$\min \left\{ \frac{1}{2} \int_{\Omega} (u - \hat{f})^2 d\mathbf{x} + \frac{1}{2} \sum_{i=1}^3 a_i \beta_i^2 \right\} \quad (17)$$

Learning variation via optimal control:

$$\begin{aligned} \min \quad & \left\{ \frac{1}{2} \int_{\Omega} (u - \hat{f})^2 d\mathbf{x} + \frac{1}{2} \sum_{i=1}^3 a_i \beta_i^2 \right\} \\ \min_{u, u \geq 0} \quad & \left\{ \int_{\Omega} \left[ \frac{1}{2} (Ku - f)^2 + V(\beta, u) \right] d\mathbf{x} \right\} \end{aligned} \quad (18)$$



*Problem A:*

$$\begin{aligned} \min_{u, \alpha} \quad & [f(x, u, \alpha), G(\alpha)] \\ \text{s.t.} \quad & g(x, u, \alpha) \leq 0 \end{aligned} \tag{19}$$

*Problem B:*

$$\begin{aligned} \min_{u, \alpha} \quad & f(x, u, \alpha) \\ \text{s.t.} \quad & G(\alpha) \leq \epsilon \\ & g(x, u, \alpha) \leq 0 \end{aligned} \tag{20}$$

## Equivalence Theorem

*Let  $\epsilon \geq \min G(\alpha)$ , let  $V^*$  solve Problem B( $\epsilon$ ), and assume that, if  $V^*$  is not unique, then  $V^*$  is an optimal solution of Problem B( $\epsilon$ ) with minimal  $G(\alpha)$  value. Then  $V^*$  solves Problem A.*

Haimes Y, Lasdon L, Wismer D (1971) On a Bicriterion Formulation of the Problems of Integrated System Identification and System Optimization. IEEE Transactions on Systems, Man, and Cybernetics 1(3) 296–297

Learning  
Variation  
via Optimal  
Control

Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

LVOC

New Insight  
LVOC  
 $\epsilon$ -Constrained  
Method

Other  
Variation or  
Functional  
Method

Active Contour  
...

Experimental  
Results

Pre-analysis  
Results



# Learning Variation via Optimal Control

Learning variation via optimal control:

$$\begin{aligned} \min \quad & \left\{ \frac{1}{2} \int_{\Omega} (u - \hat{f})^2 d\mathbf{x} + \frac{1}{2} \sum_{i=1}^3 a_i \beta_i^2 \right\} \\ \min \quad & \left\{ \int_{\Omega} \left[ \frac{1}{2} (Ku - f)^2 + V(\beta, u) \right] d\mathbf{x} \right\} \quad s.t. \quad u \geq 0 \end{aligned}$$

Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

LVOC

New Insight

LVOC

$\epsilon$ -Constrained  
Method

Other  
Variation or  
Functional  
Method

Active Contour

...

Experimental  
Results

Pre-analysis

Results



## Learning variation via optimal control:

### Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

### LVOC

New Insight  
LVOC  
 $\epsilon$ -Constrained  
Method

Other  
Variation or  
Functional  
Method

Active Contour  
...

Experimental  
Results

Pre-analysis  
Results

$$\begin{aligned} \min \quad & \left\{ \frac{1}{2} \int_{\Omega} (u - \hat{f})^2 d\mathbf{x} + \frac{1}{2} \sum_{i=1}^3 a_i \beta_i^2 \right\} \\ \min \quad & \left\{ \int_{\Omega} \left[ \frac{1}{2} (Ku - f)^2 + V(\beta, u) \right] d\mathbf{x} \right\} \quad s.t. \quad u \geq 0 \end{aligned}$$



$$\begin{aligned} \min \quad & \left\{ \frac{1}{2} \int_{\Omega} (u - \hat{f})^2 d\mathbf{x} + \frac{1}{2} \sum_{i=1}^3 a_i \beta_i^2 \right\} \\ s.t. \quad & \int_{\Omega} \left[ \frac{1}{2} (Ku - f)^2 + V(\beta, u) \right] d\mathbf{x} \leq \epsilon, u \geq 0 \end{aligned}$$





## Learning variation via optimal control:

$$\begin{aligned} \min \quad & \left\{ \frac{1}{2} \int_{\Omega} (u - \hat{f})^2 d\mathbf{x} + \frac{1}{2} \sum_{i=1}^3 a_i \beta_i^2 \right\} \\ \min \quad & \left\{ \int_{\Omega} \left[ \frac{1}{2} (Ku - f)^2 + V(\beta, u) \right] d\mathbf{x} \right\} \quad s.t. \quad u \geq 0 \end{aligned}$$



$$\begin{aligned} \min \quad & \left\{ \frac{1}{2} \int_{\Omega} (u - \hat{f})^2 d\mathbf{x} + \frac{1}{2} \sum_{i=1}^3 a_i \beta_i^2 \right\} \\ s.t. \quad & \int_{\Omega} \left[ \frac{1}{2} (Ku - f)^2 + V(\beta, u) \right] d\mathbf{x} \leq \epsilon, u \geq 0 \end{aligned}$$



$$\begin{aligned} \min \quad & \left\{ \frac{1}{2} \int_{\Omega} (u - \hat{f})^2 d\mathbf{x} + \frac{1}{2} \sum_{i=1}^3 a_i \beta_i^2 \right\} \\ s.t. \quad & K^T Ku + \text{Lag}(V(\beta, u, p)) = 0 \\ & |\nabla u|_{\epsilon} p - |\nabla u| = 0 \\ & u, |p|^2 - 1 \geq 0 \end{aligned} \tag{21}$$

Learning  
Variation  
via Optimal  
Control

Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

LVOC

New Insight

LVOC

$\epsilon$ -Constrained  
Method

Other  
Variation or  
Functional  
Method

Active Contour

...

Experimental  
Results

Pre-analysis

Results



The corresponding minimization:

$$\begin{aligned} \min \quad & \left\{ \frac{1}{2} \int_{\Omega} (u - \hat{f})^2 d\mathbf{x} + \frac{1}{2} \sum_{i=1}^3 a_i \beta_i^2 \right. \\ & + \int_{\Omega} \phi_1 (K^T K u + \text{Lag}(V(\boldsymbol{\beta}, u, p))) d\mathbf{x} \\ & + \int_{\Omega} \phi_2 (|\nabla u|_{\epsilon} p - \nabla u) d\mathbf{x} \\ & \left. + \langle u, \lambda_1 \rangle + \langle |p|^2 - 1, \lambda \rangle \right\} \end{aligned} \quad (22)$$

The KKT conditions for the problem(21):

$$\begin{aligned} (u - \hat{f}) + \text{Lag}(\text{Lag}(V)) + \nabla(\phi_2 p^2) - \nabla \phi_2 - \lambda + \lambda_2 \odot p &= 0 \\ K^T K u + \text{Lag}(V(\boldsymbol{\beta}, u, p)) &= 0 \\ |\nabla u|_{\epsilon} p - \nabla u &= 0 \\ (|p|^2 - 1) \odot \lambda_2 &= 0 \\ u \odot \lambda_1 &= 0 \\ \lambda_1, \lambda_2 &\geq 0 \end{aligned} \quad (23)$$



The equivalent minimization:

$$\min \quad \left\{ \frac{1}{2} \int_{\Omega} (u - \hat{f})^2 d\mathbf{x} + \frac{1}{2} \sum_{i=1}^3 a_i \beta_i^2 \right. \\ \left. + \int_{\Omega} \phi_1(K^T Ku + \text{Lag}(\tilde{V}(\boldsymbol{\beta}, u, p))) d\mathbf{x} \right\} \quad (24)$$

The corresponding system:

$$(u - \hat{f}) + \text{Lag}(\text{Lag}(\tilde{V})) = 0 \quad (25)$$

$$K^T Ku + \text{Lag}(\tilde{V}(\boldsymbol{\beta}, u, p)) = 0 \quad (26)$$

$$|\nabla u|_{\epsilon} p - \nabla u = 0 \quad (27)$$

$$\lambda - \max\{0, \lambda - cu\} = 0 \quad (28)$$

where  $\tilde{V} = \beta_1 u - \beta_3 \Delta u - \beta_2 \text{div} p - \lambda - K^T f$ . The left hand sides are denoted by  $F_1, F_2, F_3, F_4$  respectively.



# Learning Variation via Optimal Control

## Introduction

Numerical Methods and Applications in Total Variation Image Restoration  
L-PDE

## LVOC


New Insight  
LVOC  
 $\epsilon$ -Constrained Method

## Other Variation or Functional Method

Active Contour  
...

## Experimental Results

Pre-analysis  
Results





# Active contours model

## Introduction

- Numerical Methods and Applications in Total Variation
- Image Restoration
- L-PDE

## LVOC

- New Insight
- LVOC
- $\epsilon$ -Constrained Method

## Other Variation or Functional Method

- Active Contour


...

## Experimental Results

- Pre-analysis Results

Snake model:

$$E_{snake} = \int_1^2 \frac{1}{2} [\alpha(s) |\frac{\partial v}{\partial s}|^2 + \beta(s) |\frac{\partial^2 v}{\partial s^2}|^2] ds + \int_1^2 E_{ext}(v(s)) ds \tag{29}$$




---

Kass M, Witkin A, Terzopoulos D (1988) Snake: Active contours models. Int. J. comput. Vis., vol. 1, pp. 321-331



# Active contours model

## Introduction

- Numerical Methods and Applications in Total Variation
- Image Restoration
- L-PDE

## LVOC

- New Insight
- LVOC
- $\epsilon$ -Constrained Method

## Other Variation or Functional Method

## Active Contour


...

## Experimental Results

- Pre-analysis Results

Snake model:

$$E_{snake} = \int_1^2 \frac{1}{2} [\alpha(s) |\frac{\partial v}{\partial s}|^2 + \beta(s) |\frac{\partial^2 v}{\partial s^2}|^2] ds + \int_1^2 E_{ext}(v(s)) ds$$






## Active contours model

**Why not associate  
optimal control  
with Snake model?**

Snake model:

$$E_{snake} = \int_1^2 \frac{1}{2} [\alpha(s) |\frac{\partial v}{\partial s}|^2 + \beta(s) |\frac{\partial^2 v}{\partial s^2}|^2] ds + \int_1^2 E_{ext}(v(s)) ds$$






# Active contours model

Future Work!

Snake model:

$$E_{snake} = \int_1^2 \frac{1}{2} [\alpha(s) |\frac{\partial v}{\partial s}|^2 + \beta(s) |\frac{\partial^2 v}{\partial s^2}|^2] ds + \int_1^2 E_{ext}(v(s)) ds$$







# Learning Variation via Optimal Control

## Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

## LVOC

New Insight  
LVOC  
 $\epsilon$ -Constrained  
Method

## Other Variation or Functional Method

Active Contour

...

## Experimental Results

Pre-analysis  
Results

Thanks!



# Learning Variation via Optimal Control

## Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

## LVOC

New Insight  
LVOC  
 $\epsilon$ -Constrained  
Method

## Other Variation or Functional Method

Active Contour  
...

## Experimental Results

Pre-analysis  
Results

此处将列举其他能量泛函方法, 敬请期待!



# Learning Variation via Optimal Control

## Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

## LVOC

New Insight  
LVOC  
 $\epsilon$ -Constrained  
Method

## Other Variation or Functional Method

Active Contour  
...

## Experimental Results

Pre-analysis  
Results

此处将列举  $V(u, \beta)$  的处理效果, 敬请期待!



# Learning Variation via Optimal Control

## Introduction

Numerical  
Methods and  
Applications in  
Total Variation  
Image  
Restoration  
L-PDE

## LVOC

New Insight  
LVOC  
 $\epsilon$ -Constrained  
Method

## Other Variation or Functional Method

Active Contour  
...

## Experimental Results

Pre-analysis  
Results

此处将列举 LVOC 的处理效果 (加大训练集), 并进一步分析, 包括线性拟合最优参数和评价指标的内容, 敬请期待!