

Reconsidered error analysis in the finite element methods

Hiroki ISHIZAKA

April 1, 2025

Abstract

We introduce new proof methods for interpolation error estimates.

Contents

1	Preliminaries	2
1.1	General Convention	2
1.2	Basic Notation	2
1.3	Vectors and Matrices	2
1.4	Function Spaces	3
1.5	Finite-Element-Methods-Related Symbols	3
1.5.1	Symbols	3
1.5.2	Meshes	3
1.5.3	Broken Sobolev Spaces, Mesh faces, Averages and Jumps	4
1.6	Useful Tools for Analysis	5
1.6.1	Jensen-type Inequality	5
1.6.2	Embedding Theorems	5
1.6.3	Trace Theorem	6
1.6.4	Bramble–Hilbert–type Lemma	6
1.6.5	Poincaré inequality	7
2	Geometric Conditions	7

1 Preliminaries

1.1 General Convention

Throughout this article, we denote by c a constant independent of h (defined later) and the angles and aspect ratios of simplices, unless specified otherwise all constants c are bounded if the maximum angle is bounded. These values vary across different contexts.

1.2 Basic Notation

d	The space dimension, $d \in \{2, 3\}$
\mathbb{R}^d	d -dimensional real Euclidean space
\mathbb{N}_0	$\mathbb{N}_0 := \mathbb{N} \cup \{0\}$
\mathbb{R}_+	The set of positive real numbers
$ \cdot _d$	d -dimensional Hausdorff measure
$v _D$	Restriction of the function v to the set D
$\dim(V)$	Dimension of the vector space V
δ_{ij}	Kronecker delta: $\delta_{ij} = 1$ if $i = j$ and 0 otherwise
$(x_1, \dots, x_d)^T$	Cartesian coordinates in \mathbb{R}^d

1.3 Vectors and Matrices

$(v_1, \dots, v_d)^T$	Cartesian components of the vector v in \mathbb{R}^d
$x \cdot y$	Euclidean scalar product in \mathbb{R}^d : $x \cdot y := \sum_{i=1}^d x_i y_i$
$ x _E$	Euclidean norm in \mathbb{R}^d : $ x _E := (x \cdot x)^{1/2}$
$\mathbb{R}^{m \times n}$	Vector space $m \times n$ matrices with real-valued entries
A, B	Matrices
A_{ij} or $[A]_{ij}$	Entry of A in the i th and the j th column
A^\top	Transpose of the matrix A
$\text{Tr}(A)$	Trace of A : For $A \in \mathbb{R}^{m \times n}$, $\text{Tr}(A) := \sum_{i=1}^d A_{ii}$
$\det(A)$	Determinant of A
$\text{diag}(A)$	Diagonal of A : For $A \in \mathbb{R}^{m \times n}$, $\text{diag}(A)_{ij} := \delta_{ij} A_{ij}$, $1 \leq i, j \leq d$
Ax	Matrix-vector product: For $A \in \mathbb{R}^{m \times n}$ and $x \in \mathbb{R}^n$, $(Ax)_i := \sum_{j=1}^d A_{ij} x_j$ for $1 \leq i \leq d$
$A : B$	Double contraction: For $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{m \times n}$, $A : B := \sum_{i=1}^m \sum_{j=1}^n A_{ij} B_{ij}$
$\ A\ _2$	Operator norm of A : For $A \in \mathbb{R}^{d \times d}$, $\ A\ _2 := \sup_{0 \neq x \in \mathbb{R}^d} \frac{ Ax _E}{ x _E}$
$\ A\ _{\max}$	Max norm of A : For $A \in \mathbb{R}^{d \times d}$, $\ A\ _{\max} := \max_{1 \leq i, j \leq d} A_{ij} $
$O(d)$	$O(d)$ consists of all orthogonal matrices of determinant ± 1

In this article, we use the following facts.

For $A \in \mathbb{R}^{m \times n}$, it holds that

$$\|A\|_{\max} \leq \|A\|_2 \leq \sqrt{mn} \|A\|_{\max}, \quad (1.1)$$

e.g., see [10, p. 56]. For $A, B \in \mathbb{R}^{m \times m}$, it holds that

$$\|AB\|_2 \leq \|A\|_2 \|B\|_2. \quad (1.2)$$

If $A^\top A$ is a positive definite matrix in $\mathbb{R}^{d \times d}$, the spectral norm of the matrix $A^\top A$ is the largest eigenvalue of $A^\top A$; i.e.,

$$\|A\|_2 = (\lambda_{\max}(A^\top A))^{1/2} = \sigma_{\max}(A), \quad (1.3)$$

where $\lambda_{\max}(A)$ and $\sigma_{\max}(A)$ are respectively the largest eigenvalues and singular values of A .

If $A \in O(d)$, because $A^\top = A^{-1}$ and

$$|Ax|_E^2 = (Ax)^\top (Ax) = x^\top A^\top A x = x^\top A^{-1} A x = |x|_E^2,$$

it holds that

$$\|A\|_2 = \sup_{0 \neq x \in \mathbb{R}^d} \frac{|Ax|_E}{|x|_E} = \sup_{0 \neq x \in \mathbb{R}^d} \frac{|x|_E}{|x|_E} = 1.$$

1.4 Function Spaces

In this article, we use standard Sobolev spaces with associated norms (e.g., see [2, 6, 7]).

1.5 Finite-Element-Methods-Related Symbols

1.5.1 Symbols

\mathbb{P}^k	Vector space of polynomials in the variables x_1, \dots, x_d of global degree at most $k \in \mathbb{N}_0$
$N^{(d,k)}$	$N^{(d,k)} := \dim(\mathbb{P}^k) = \binom{d+k}{k}$
\mathbb{RT}^k	The Raviart–Thomas polynomial space of order $k \in \mathbb{N}_0$ as $\mathbb{RT}^k := (\mathbb{P}^k)^d + x\mathbb{P}^k$ for any $x \in \mathbb{R}^d$
$N^{(RT)}$	$N^{(RT)} := \dim \mathbb{RT}^k$
T, \tilde{T}, \hat{T}, K	Closed simplices in \mathbb{R}^d
$\mathbb{P}^k(T), \mathbb{RT}^k(T)$	$\mathbb{P}^k(T)$ (or $\mathbb{RT}^k(T)$) is spanned by the restriction to T of polynomials in \mathbb{P}^k (or \mathbb{RT}^k)

1.5.2 Meshes

Let $\Omega \subset \mathbb{R}^d$, $d \in \{2, 3\}$, be a bounded polyhedral domain. Furthermore, we assume that Ω is convex if necessary. Let $\mathbb{T}_h = \{T\}$ be a simplicial mesh of $\bar{\Omega}$ made up of closed d -simplices, such as

$$\bar{\Omega} = \bigcup_{T \in \mathbb{T}_h} T,$$

with $h := \max_{T \in \mathbb{T}_h} h_T$, where $h_T := \text{diam}(T)$. We also use a symbol ρ_T which means the radius of the largest ball inscribed in T . We assume that each face of any d -simplex T_1 in \mathbb{T}_h is either

a subset of the boundary $\partial\Omega$ or a face of another d -simplex T_2 in \mathbb{T}_h . That is, \mathbb{T}_h is a simplicial mesh of $\bar{\Omega}$ without hanging nodes. Such mesh \mathbb{T}_h is said to be conformal. Let $\{\mathbb{T}_h\}$ be a family of conformal meshes.

Let T be a simplex of \mathbb{T}_h which is a convex hull of $d+1$ vertices, p_1, \dots, p_{d+1} , that do not belong to the same hyperplane. Let S_i be the face of a simplex T opposite to the vertex p_i . For $d=3$, angles between faces of a tetrahedron are called *dihedral*, whereas angles between its edges are called *solid*.

1.5.3 Broken Sobolev Spaces, Mesh faces, Averages and Jumps

Let \mathcal{F}_h^i be the set of interior faces, and \mathcal{F}_h^∂ be the set of faces on boundary $\partial\Omega$. We set $\mathcal{F}_h := \mathcal{F}_h^i \cup \mathcal{F}_h^\partial$. For any $F \in \mathcal{F}_h$, we define the unit normal n_F to F as follows: (i) If $F \in \mathcal{F}_h^i$ with $F = T_{\natural} \cap T_{\sharp}$, $T_{\natural}, T_{\sharp} \in \mathbb{T}_h$, $\natural > \sharp$, let n_F be the unit normal vector from T_{\natural} to T_{\sharp} . (ii) If $F \in \mathcal{F}_h^\partial$, n_F is the unit outward normal n to $\partial\Omega$. We also use the following set. For any $F \in \mathcal{F}_h$,

$$\mathbb{T}_F := \{T \in \mathbb{T}_h : F \subset T\}.$$

We consider \mathbb{R}^q -valued functions for some $q \in \mathbb{N}$. Let $p \in [1, \infty]$ and $s > 0$ be a positive real number. We define a broken (piecewise) Sobolev space as

$$W^{s,p}(\mathbb{T}_h; \mathbb{R}^q) := \{v \in L^p(\Omega; \mathbb{R}^q) : v|_T \in W^{s,p}(T; \mathbb{R}^q) \ \forall T \in \mathbb{T}_h\}$$

with the norms

$$\begin{aligned} \|v\|_{W^{s,p}(\mathbb{T}_h; \mathbb{R}^q)} &:= \left(\sum_{T \in \mathbb{T}_h} \|v\|_{W^{s,p}(T; \mathbb{R}^q)}^p \right)^{1/p} \quad \text{if } p \in [1, \infty), \\ \|v\|_{W^{s,\infty}(\mathbb{T}_h; \mathbb{R}^q)} &:= \max_{T \in \mathbb{T}_h} \|v\|_{W^{s,\infty}(T; \mathbb{R}^q)}. \end{aligned}$$

When $q=1$, we denote $W^{s,p}(\mathbb{T}_h) := W^{s,p}(\mathbb{T}_h; \mathbb{R})$. When $p=2$, we write $H^s(\mathbb{T}_h)^q := H^s(\mathbb{T}_h; \mathbb{R}^q) := W^{s,2}(\mathbb{T}_h; \mathbb{R}^q)$ and $H^s(\mathbb{T}_h) := W^{s,2}(\mathbb{T}_h; \mathbb{R})$. We use the norm

$$|\varphi|_{H^1(\mathbb{T}_h)} := \left(\sum_{T \in \mathbb{T}_h} \|\nabla \varphi\|_{L^2(T)^d}^2 \right)^{1/2} \quad \varphi \in H^1(\mathbb{T}_h).$$

Let $\varphi \in H^1(\mathbb{T}_h)$. Suppose that $F \in \mathcal{F}_h^i$ with $F = T_{\natural} \cap T_{\sharp}$, $T_{\natural}, T_{\sharp} \in \mathbb{T}_h$, $\natural > \sharp$. We set $\varphi_{\natural} := \varphi|_{T_{\natural}}$ and $\varphi_{\sharp} := \varphi|_{T_{\sharp}}$. The jump in φ across F is defined as

$$[\![\varphi]\!] := [\![\varphi]\!]_F := \varphi_{\natural} - \varphi_{\sharp}, \quad \natural > \sharp.$$

For a boundary face $F \in \mathcal{F}_h^\partial$ with $F = \partial T \cap \partial\Omega$, $[\![\varphi]\!]_F := \varphi|_T$. For any $v \in H^1(\mathbb{T}_h)^d$, the notations

$$\begin{aligned} [v \cdot n] &:= [v \cdot n]_F := v_{\natural} \cdot n_F - v_{\sharp} \cdot n_F, \quad \natural > \sharp, \\ [v] &:= [v]_F := v_{\natural} - v_{\sharp}, \quad \natural > \sharp, \end{aligned}$$

denote the jump in the normal component of v and the jump of v . Set two nonnegative real numbers $\omega_{T_{\natural}, F}$ and $\omega_{T_{\sharp}, F}$ such that

$$\omega_{T_{\natural}, F} + \omega_{T_{\sharp}, F} = 1.$$

The skew-weighted average of φ across F is then defined as

$$\{\{\varphi\}\}_{\bar{\omega}} := \{\{\varphi\}\}_{\bar{\omega},F} := \omega_{T_{\sharp},F}\varphi_{\natural} + \omega_{T_{\natural},F}\varphi_{\sharp}.$$

For a boundary face $F \in \mathcal{F}_h^{\partial}$ with $F = \partial T \cap \partial\Omega$, $\{\{\varphi\}\}_{\bar{\omega}} := \varphi|_T$. Furthermore,

$$\{\{v\}\}_{\omega} := \{\{v\}\}_{\omega,F} := \omega_{T_{\natural},F}v_{\natural} + \omega_{T_{\sharp},F}v_{\sharp},$$

for the weighted average of v . For any $v \in H^1(\mathbb{T}_h)^d$ and $\varphi \in H^1(\mathbb{T}_h)$,

$$\llbracket (v\varphi) \cdot n \rrbracket_F = \{\{v\}\}_{\omega,F} \cdot n_F \llbracket \varphi \rrbracket_F + \llbracket v \cdot n \rrbracket_F \{\{\varphi\}\}_{\bar{\omega},F}.$$

We define a broken gradient operator as follows. Let $p \in [1, \infty]$. For $\varphi \in W^{1,p}(\mathbb{T}_h)$, the broken gradient $\nabla_h : W^{1,p}(\mathbb{T}_h) \rightarrow L^p(\Omega)^d$ is defined by

$$(\nabla_h \varphi)|_T := \nabla(\varphi|_T) \quad \forall T \in \mathbb{T}_h,$$

and we define the broken $H(\operatorname{div}; T)$ space by

$$H(\operatorname{div}; \mathbb{T}_h) := \{v \in L^2(\Omega)^d; v|_T \in H(\operatorname{div}; T) \quad \forall T \in \mathbb{T}_h\},$$

and the broken divergence operator $\operatorname{div}_h : H(\operatorname{div}; \mathbb{T}_h) \rightarrow L^2(\Omega)$ such that, for all $v \in H(\operatorname{div}; \mathbb{T}_h)$,

$$(\operatorname{div}_h v)|_T := \operatorname{div}(v|_T) \quad \forall T \in \mathbb{T}_h.$$

1.6 Useful Tools for Analysis

1.6.1 Jensen-type Inequality

Let r, s be two nonnegative real numbers and $\{x_i\}_{i \in I}$ be a finite sequence of nonnegative numbers. It then holds that

$$\begin{cases} (\sum_{i \in I} x_i^s)^{\frac{1}{s}} \leq (\sum_{i \in I} x_i^r)^{\frac{1}{r}} & \text{if } r \leq s, \\ (\sum_{i \in I} x_i^s)^{\frac{1}{s}} \leq \operatorname{card}(I)^{\frac{r-s}{rs}} (\sum_{i \in I} x_i^r)^{\frac{1}{r}} & \text{if } r > s, \end{cases} \quad (1.4)$$

see [7, Exercise 12.1].

1.6.2 Embedding Theorems

The following is well known as the Sobolev embedding theorem.

Theorem 1.1. Let $d \geq 2$, $s > 0$, and $p \in [1, \infty]$. Let $D \subset \mathbb{R}^d$ be a bounded open subset of \mathbb{R}^d . If D is a Lipschitz set, we then have

$$W^{s,p}(D) \hookrightarrow \begin{cases} L^q(D) & \forall q \in [p, \frac{pd}{d-sp}] \text{ if } sp < d, \\ L^q(D) & \forall q \in [p, \infty), \text{ if } sp = d, \\ L^\infty(D) \cap \mathcal{C}^{0,\xi}(\bar{D}) & \xi = 1 - \frac{d}{sp} \text{ if } sp > d. \end{cases} \quad (1.5)$$

Furthermore,

$$W^{s,p}(D) \hookrightarrow L^\infty(D) \cap \mathcal{C}^0(\bar{D}) \quad (\text{case } s = d \text{ and } p = 1). \quad (1.6)$$

Proof. See, for example, [6, Corollary B.43, Theorem B.40] and [7, Theorem 2.31] and the references therein. \square

The following is the embedding theorem related to operator from $W^{s,p}(D)$ into $L^q(S_r)$, where S_r is some plane r -dimensional piece belonging to D with dimensions $r < d$.

Theorem 1.2. Let $p, q \in [1, +\infty]$ and $s \geq 1$ be an integer. Let $D \subset \mathbb{R}^d$ be a bounded open set having piecewise smooth boundaries. The following embeddings are then continuous:

$$W^{s,p}(D) \hookrightarrow \begin{cases} L^q(S_r) & \text{if } 1 \leq p < \frac{d}{s}, r > d - sp \text{ and } q \leq \frac{pr}{d-sp}, \\ L^q(S_r) & \text{if } p = \frac{d}{s} \text{ for } q < +\infty. \end{cases} \quad (1.7)$$

Proof. See, for example, [22, Theorem 2.1 (p. 61)] and the references therein. \square

1.6.3 Trace Theorem

Theorem 1.3 (Trace on low-dimensional manifolds). Let $p \in [1, \infty)$ and let D be a Lipschitz domain in \mathbb{R}^d . Let M be a smooth, or polyhedral, manifold of dimension r in \overline{D} , $r \in \{0, \dots, d\}$. Then, there exists a bounded trace operator from $W^{s,p}(D)$ to $L^p(M)$, provided $sp > d - r$, or $s \geq d - r$ if $p = 1$.

Proof. See [7, Theorem 3.15]. \square

1.6.4 Bramble–Hilbert–type Lemma

The Bramble–Hilbert–type lemma (e.g., see [5, 3]) plays a major role in interpolation error analysis. We use the following estimates on anisotropic meshes proposed in [1, Lemma 2.1].

Lemma 1.4. Let $D \subset \mathbb{R}^d$ be a connected open set that is star-shaped with respect to balls B . Let γ be a multi-index with $m := |\gamma|$ and $\varphi \in L^1(D)$ be a function with $\partial^\gamma \varphi \in W^{\ell-m,p}(D)$, where $\ell \in \mathbb{N}$, $m \in \mathbb{N}_0$, $0 \leq m \leq \ell$, $p \in [1, \infty]$. It then holds that

$$\|\partial^\gamma(\varphi - Q^{(\ell)}\varphi)\|_{W^{\ell-m,p}(D)} \leq C^{BH} |\partial^\gamma \varphi|_{W^{\ell-m,p}(D)}, \quad (1.8)$$

where C^{BH} depends only on $d, \ell, \text{diam } D$, and $\text{diam } B$, and $Q^{(\ell)}\varphi$ is defined as

$$(Q^{(\ell)}\varphi)(x) := \sum_{|\delta| \leq \ell-1} \int_B \eta(y) (\partial^\delta \varphi)(y) \frac{(x-y)^\delta}{\delta!} dy \in \mathbb{P}^{\ell-1}, \quad (1.9)$$

where $\eta \in \mathcal{C}_0^\infty(B)$ is a given function with $\int_B \eta dx = 1$.

To give local interpolation error estimates on isotropic meshes, we use the inequalities given in [4, Theorem 1.1] and [5, 3, 25] which are variants of the Bramble–Hilbert lemma.

Lemma 1.5. Let $D \subset \mathbb{R}^d$ be a bounded convex domain. Let $\varphi \in W^{m,p}(D)$ with $m \in \mathbb{N}$ and $1 \leq p \leq \infty$. There exists a polynomial $\eta \in \mathbb{P}^{m-1}$ such that

$$|\varphi - \eta|_{W^{k,p}(D)} \leq C^{BH}(d, m) \text{diam}(D)^{m-k} |\varphi|_{W^{m,p}(D)}, \quad k = 0, 1, \dots, m. \quad (1.10)$$

Proof. The proof is found in [4, Theorem 1.1]. \square

Remark 1.6. In [3, Lemma 4.3.8], the Bramble–Hilbert lemma is given as follows. Let B be a ball in $D \subset \mathbb{R}^d$ such that D is star-shaped with respect to B and its radius $r > \frac{1}{2}r_{\max}$, where $r_{\max} := \sup\{r : D \text{ is star-shaped with respect to a ball of radius } r\}$. Let $\varphi \in W^{m,p}(D)$ with $m \in \mathbb{N}$ and $1 \leq p \leq \infty$. There exists a polynomial $\eta \in \mathbb{P}^{m-1}$ such that

$$|\varphi - \eta|_{W^{k,p}(D)} \leq C^{BH}(d, m, \gamma) \text{diam}(D)^{m-k} |\varphi|_{W^{m,p}(D)}, \quad k = 0, 1, \dots, m. \quad (1.11)$$

Here, γ is called the chunkiness parameter of D , which is defined by

$$\gamma := \frac{\text{diam}(D)}{r_{\max}}.$$

The main drawback is that the constant $C^{BH}(d, m, \gamma)$ depends on the chunkiness parameter. Meanwhile, the constant $C^{BH}(d, m)$ of the estimate (1.10) does not depend on the geometric parameter γ .

Remark 1.7. For general Sobolev spaces $W^{m,p}(\Omega)$, the upper bounds on the constant $C^{BH}(d, m)$ are not given, as far as we know. However, when $p = 2$, the following result has been obtained by Verfürth [25].

Let $D \subset \mathbb{R}^d$ be a bounded convex domain. Let $\varphi \in H^m(D)$ with $m \in \mathbb{N}$. There exists a polynomial $\eta \in \mathbb{P}^{m-1}$ such that

$$|\varphi - \eta|_{H^k(D)} \leq C^{BH}(d, k, m) \text{diam}(D)^{m-k} |\varphi|_{H^m(D)}, \quad k = 0, 1, \dots, m-1. \quad (1.12)$$

Verfürth has given upper bounds on the constants in the estimates such that

$$C^{BH}(d, k, m) \leq \pi^{k-m} \binom{d+k-1}{k}^{1/2} \frac{\{(m-k)!\}^{1/2}}{\{[\frac{m-k}{d}]\}^{d/2}},$$

where $[x]$ denotes the largest integer less than or equal to x .

As an example, let us consider the case $d = 3$, $k = 1$, and $m = 2$. We then have

$$C^{BH}(3, 1, 2) \leq \frac{\sqrt{3}}{\pi},$$

thus on the standard reference element \hat{T} , we obtain

$$|\hat{\varphi} - \hat{\eta}|_{H^1(\hat{T})} \leq \frac{\sqrt{6}}{\pi} |\hat{\varphi}|_{H^2(\hat{T})} \quad \forall \hat{\varphi} \in H^2(\hat{T}),$$

because $\text{diam}(\hat{T}) = \sqrt{2}$.

1.6.5 Poincaré inequality

Theorem 1.8 (Poincaré inequality). Let $D \subset \mathbb{R}^d$ be a convex domain with diameter $\text{diam}(D)$. It then holds that, for $\varphi \in H^1(D)$ with $\int_D \varphi dx = 0$,

$$\|\varphi\|_{L^2(D)} \leq \frac{\text{diam}(D)}{\pi} |\varphi|_{H^1(D)}. \quad (1.13)$$

Proof. The proof is found in [23, Theorem 3.2], also see [24]. □

Remark 1.9. The coefficient $\frac{1}{\pi}$ of (1.13) may be improved.

2 Geometric Conditions

References

- [1] Apel, Th.: Anisotropic finite elements: Local estimates and applications. Advances in Numerical Mathematics. Teubner, Stuttgart, (1999)
- [2] Bernardi, C., Girault, V., Hecht, F., Raviart, P.-A., Rivière, B.: Mathematics and Finite Element Discretizations of Incompressible Navier–Stokes Flows. SIAM, (2024)
- [3] Brenner, S.C., Scott, L.R.: The Mathematical Theory of Finite Element Methods, Third Edition. Springer Verlag, New York (2008)
- [4] Dekel, S., Leviatan, D.: The Bramble–Hilbert Lemma for Convex Domains, SIAM Journal on Mathematical Analysis **35** No. 5, 1203–1212 (2004)
- [5] Dupont, T., Scott, R.: Polynomial approximation of functions in Sobolev spaces. Math. Comp. **34**, 441–463 (1980)
- [6] Ern, A., Guermond, J.L.: Theory and Practice of Finite Elements. Springer Verlag, New York (2004)
- [7] Ern, A., Guermond, J. L.: Finite Elements I: Approximation and Interpolation. Springer-Verlag, New York (2021)
- [8] Ern, A., Guermond, J. L.: Finite elements II: Galerkin Approximation, Elliptic and Mixed PDEs. Springer-Verlag, New York (2021)
- [9] Ern, A. and Guermond, J. L.: Finite elements III: First-Order and Time-Dependent PDEs. Springer-Verlag, New York (2021).
- [10] Golub, G. H., Loan, C. F. V.: Matrix Computations 3rd edition. The Johns Hopkins University Press, (1996)
- [11] Ishizaka H. Anisotropic interpolation error analysis using a new geometric parameter and its applications. Ehime University, Ph. D. thesis (2022)
- [12] Ishizaka, H.: Anisotropic Raviart–Thomas interpolation error estimates using a new geometric parameter. Calcolo **59** (4), (2022)
- [13] Ishizaka, H.: Anisotropic weakly over-penalised symmetric interior penalty method for the Stokes equation. Journal of Scientific Computing **100**, (2024)
- [14] Ishizaka, H.: Morley finite element analysis for fourth-order elliptic equations under a semi-regular mesh condition. Applications of Mathematics **69** (6), 769–805 (2024)
- [15] Ishizaka, H.: Hybrid weakly over-penalised symmetric interior penalty method on anisotropic meshes. Calcolo **61**, (2024)
- [16] Ishizaka, H.: Anisotropic modified Crouzeix-Raviart finite element method for the stationary Navier-Stokes equation. Numerische Mathematik, (2025)
- [17] Ishizaka, H.: Nitsche method under a semi-regular mesh condition. preprint (2025) <https://arxiv.org/abs/2501.06824>
- [18] Ishizaka, H., Kobayashi, K., Suzuki, R., Tsuchiya, T.: A new geometric condition equivalent to the maximum angle condition for tetrahedrons. Comput. Math. Appl. **99**, 323–328 (2021)

- [19] Ishizaka, H., Kobayashi, K., Tsuchiya, T.: General theory of interpolation error estimates on anisotropic meshes. *Jpn. J. Ind. Appl. Math.* 38 (1), 163–191 (2021)
- [20] Ishizaka, H., Kobayashi, K., Tsuchiya, T.: Crouzeix–Raviart and Raviart–Thomas finite element error analysis on anisotropic meshes violating the maximum-angle condition. *Jpn. J. Ind. Appl. Math.* 38 (2), 645–675 (2021)
- [21] Ishizaka, H., Kobayashi, K., Tsuchiya, T.: Anisotropic interpolation error estimates using a new geometric parameter. *Jpn. J. Ind. Appl. Math.* 40 (1), 475–512 (2023)
- [22] Ladyženskaja, O. A., Solonnikov, V. A., Ural’ceva, N. N.: *Linear and Quasi-linear Equations of Parabolic Type*. Translations of Mathematical Monographs **23**, AMS, Providence, (1968)
- [23] Mario, B.: A note on the Poincaré inequality for convex domains. *Z. Anal. ihre. Anwend.*, **22**, 751–756 (2003)
- [24] Payne, L.E., Weinberger, H.F.: An optimal Poincaré-inequality for convex domains, *Arch. Rational Mech. Anal.* **5**, 286–292 (1960)
- [25] Verfürth, R.: A note on polynomial approximation in Sobolev spaces, *Math. Modelling and Numer. Anal.* **33**, 715–719 (1999)