

NUEN 647
Uncertainty Quantification for Nuclear Engineering
Assignment 1

Due on Tuesday, October 4, 2016

Dr. McClarren

Paul Mendoza

Contents

Problem 1	3
Problem 2	5
Problem 3	6
Problem 4	7

Complete the exercises in the Chapter 2 notes. Be sure to include discussion of results where appropriate. You may use any tools that are appropriate to solving the problem.

Problem 1

Show that the transformation in equation 1 results in a standard normal random variable by computing the mean and variance of z .

$$z = \frac{x - \mu}{\sigma} \quad (1)$$

An important special case of the expectation value is the mean which is the expected value of x . It is often denoted as μ ,

$$\mu = E[x] = \int_{-\infty}^{\infty} x f(x) dx$$

where x is a realization of a random sample and $f(x)$ is the probability density function (PDF) for the random variable. For a normal distribution,

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

For the sake of the transformation, the value of z substitutes for x , the realization of a random sample (not the PDF because we are transforming that distribution). Therefore, the mean for z is:

$$\mu_z = \int_{-\infty}^{\infty} \frac{x - \mu}{\sigma} \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx$$

If $u = (x - \mu)^2$ and $\frac{du}{2} = (x - \mu)dx$ (note that the limits change from $(-\infty, \infty)$ to (∞, ∞) - but that seems fishy to me so I will change it back after integration).

$$\begin{aligned} \mu_z &= \int_{-\infty}^{\infty} \frac{1}{2\sigma^2\sqrt{2\pi}} e^{-\frac{u}{2\sigma^2}} du = \left| \frac{-1}{\sqrt{2\pi}} e^{-\frac{u}{2\sigma^2}} \right|_{-\infty}^{\infty} \\ \mu_z &= \left| \frac{-1}{\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} \right|_{-\infty}^{\infty} = \frac{-1}{\sqrt{2\pi}} (e^{-\infty} - e^{-\infty}) = \boxed{0} \end{aligned}$$

The variance is defined as:

$$\text{Var}(X) = E[(X - \mu)^2]$$

Substituting Eq. 1 for X , (but not for the pdf - I could be wrong about that)

$$\text{Var}(X) = E\left[\left(\frac{x - \mu}{\sigma} - \mu\right)^2\right] = E\left[\left(\frac{x - \mu - \mu\sigma}{\sigma}\right)^2\right] = \frac{1}{\sigma^2} (E[x^2] - 2\mu E[x] - 2\mu\sigma E[x] + \mu^2 E[1] + \mu^2\sigma^2 + 2\mu^2\sigma)$$

Noting that above it was proven that $E[x] = \mu$ and given that the definition of $E[1] = 1$ and assuming that $E[x^2] = \sigma^2 + \mu^2$ (will solve on next page)

$$\frac{1}{\sigma^2} (\sigma^2 + \mu^2 - 2\mu^2 - 2\mu^2\sigma + \mu^2 + \mu^2\sigma^2 + 2\mu^2\sigma) = \frac{1}{\sigma^2} (\sigma^2 + \mu^2\sigma^2) = \boxed{1 + \mu^2 = 1}$$

This is assuming that $\mu = 0$. Which was shown above.

$$E[x^2] = \int_{-\infty}^{\infty} \frac{x^2}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx$$

If $t = \frac{(x-\mu)}{\sqrt{2}\sigma}$ and $\sqrt{2}\sigma dt = dx$ and $x = t\sqrt{2}\sigma + \mu$ then (limits of integration don't change)

$$E[x^2] = \int_{-\infty}^{\infty} \frac{(t\sqrt{2}\sigma + \mu)^2}{\sqrt{\pi}} e^{-t^2} dt = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \left(2\sigma^2 (t^2 e^{-t^2}) + 2\sqrt{2}\sigma\mu (te^{-t^2}) + \mu^2 (e^{-t^2}) \right) dt$$

According to wolfram alpha

$$\int_{-\infty}^{\infty} t^2 e^{-t^2} dt = \frac{\sqrt{\pi}}{2}$$

$$\int_{-\infty}^{\infty} te^{-t^2} dt = 0$$

$$\int_{-\infty}^{\infty} e^{-t^2} dt = \sqrt{\pi}$$

Which simplifies the above to $\sigma^2 + \mu^2$.

Problem 2

Consider the random variables $X \sim U(-1, 1)$ and $Y \sim X^2$. Are these independent random variables? What is their covariance?

If two random variables, X and Y, are independent, they satisfy the following condition: [*link*](#)

- $P(X|Y) = P(X)$, for all values of X and Y.

The PDF for X is:

$$f_X(x) = \frac{1}{(1 - (-1))} = 0.5 \quad x \in [-1, 1]$$

The PDF for Y is: [*link*](#)

$$f_Y(y) = \frac{1}{2\sqrt{y}} \quad y \in [0, 1]$$

If the covariance is non zero, then these two variables are independent. The covariance of two random variables can be given by: [*link*](#)

$$\sigma_{XY} = E(XY) - \mu_X \mu_Y = \int_{-1}^1 dx \int_0^1 dy \quad xyf(x, y) - \mu_X \mu_Y$$

Because $\mu_X = 0$ this reduces to

$$\sigma_{XY} = E(XY) = \int_{-1}^1 dx \int_0^1 dy \quad xyf(x, y)$$

Where $f(x, y)$ is:

$$f(x, y) = f(y|x)f_X(x)$$

From the definition of Y, $f(y|x)$ is 0 except when $y = x^2$. I think this would be.

$$f(y|x) = \delta(y - x^2)$$

Which means,

$$f(x, y) = 0.5\delta(y - x^2)$$

I am not 100% sure to do here, but after using wolfram, I think its 0.25.

Problem 3

Show that a general covariance matrix must be positive definite, i.e. $\vec{x}^T \Sigma \vec{x} > 0$ for any vector \vec{x} that is not all zeros.

Given that \vec{Y} is a vector of random variables and $\vec{\mu}_Y$ is a vector of the mean values for the random variables found in \vec{Y} .

$$\begin{aligned}\vec{x}^T \Sigma \vec{x} &= \vec{x}^T E[(\vec{Y} - \vec{\mu}_Y)(\vec{Y} - \vec{\mu}_Y)^T] \vec{x} \\ &= E[\vec{x}^T (\vec{Y} - \vec{\mu}_Y)(\vec{Y} - \vec{\mu}_Y)^T \vec{x}]\end{aligned}$$

The last step above puts a constant inside the expectation value integral. Notice

$$\vec{x}^T (\vec{Y} - \vec{\mu}_Y) = (\vec{Y} - \vec{\mu}_Y)^T \vec{x}$$

and that both are scalar functions of the random variables. Therefore,

$$\begin{aligned}\vec{x}^T \Sigma \vec{x} &= E[(\vec{x}^T (\vec{Y} - \vec{\mu}_Y))^2] \\ &= E[g(Y)^2] = \sigma_f^2\end{aligned}$$

The expectation value for a multivariate distribution is defined as

$$E[g(Y)] = \int_{-\infty}^{\infty} dy_1 \int_{-\infty}^{\infty} dy_2 \dots \int_{-\infty}^{\infty} dy_p g(y) f(y)$$

Where $f(y)$ is the multivariate PDF for the random variables of \vec{Y} . To prove that the covariance matrix is positive definite the above integral must be proved to be positive with $g(x) = (\vec{x}^T (\vec{Y} - \vec{\mu}_Y))^2$. Explicitly,

$$\begin{aligned}E[g(Y)] &= \int_{-\infty}^{\infty} dy_1 \int_{-\infty}^{\infty} dy_2 \dots \int_{-\infty}^{\infty} dy_p (\vec{x}^T (\vec{Y} - \vec{\mu}_Y))^2 f(y) \\ &= \int_{-\infty}^{\infty} dy_1 \int_{-\infty}^{\infty} dy_2 \dots \int_{-\infty}^{\infty} dy_p ((y_1 - \mu_1)x_1 + (y_2 - \mu_2)x_2 + \dots + (y_p - \mu_p)x_p)^2 f(y)\end{aligned}$$

Problem 4

Use rejection sampling to sample from a Gamma random variable $X \sim \mathcal{G}(\alpha, \beta)$ where

$$f(x) = \frac{\theta^{\alpha-1} e^{-\theta/\beta}}{\Gamma(\alpha) \beta^{-\alpha}} \quad \alpha, \beta > 0$$

Let $\alpha = 1$ and $\beta = 0.5$. From rejection sampling with a $N = 10^4$, compute a rejection rate for the sampling procedure. Now draw a triangle around the function and do rejection sampling. Compare the rejection rate from the triangle versus the rectangle. You may consider that the PDF is zero if $f(x) < 10^{-6}$.

Python script for rejection sampling.

Listing 1: Python Script for problem

```
#!/usr/bin/env python3

"""
Chem Calculations
"""
5 """

__author__      = "Paul Mendoza"
__copyright__    = "Copyright 2016, Planet Earth"
__credits__     = ["Sunil Chirayath",
10         "Charles Folden",
        "Jeremy Conlin"]

__license__     = "GPL"
__version__     = "1.0.1"
__maintainer__  = "Paul Mendoza"
15 __email__     = "paul.m.mendoza@gmail.com"
__status__      = "Production"

#####
##### Import packages #####
20 #####

import os.path
import pandas as pd
import numpy as np
25 import matplotlib.pyplot as plt
import datetime
from uncertainties import ufloat
from uncertainties.umath import *
from uncertainties import unumpy as unp
30 import re
import time
start_time = time.time()

import Functions as fun
35

#####
##### Examples of Calculations #####
40 #####
```

```

#####
##### Atom Fraction to Mass Fraction #####
##### and vice versa #####
#####

45 string='92235 0.285714286 0 92238 0.714285714 0'
MasstoAtom=True
Mass,Zaid=fun.StringToMass(string)
stringCalculated=fun.ConvertFractions(string,Mass,MasstoAtom,Zaid)

50 # if MasstoAtom:
#     print("Mass Fractions:")
#     print(string)
#     print("Atom Fractions:")
55 #     print(stringCalculated)
# else:
#     print("Mass Fractions:")
#     print(stringCalculated)
#     print("Atom Fractions:")
60 #     print(string)

#####
##### Calculate grams per mol of #####
##### a chemical formula #####
65 #####

#Make sure your chemical form has no repeats
#And no parentheses
ChemicalFormula='HNO_3'
70 ChemicalFormulaError=[0,0,0,0,0,0,0] #+/- error in integers of
                                     #chemical formula
ChemicalFormula=ChemicalFormula+" "
List=fun.ChemList(ChemicalFormula)

75 #Enter Modifications:
#1. Each element should be a single item in the list
#2. Format: zaid atomfraction+/-error zaid atomfraction+/-error
# or : zaid atomfraction error zaid atomfraction
Modifications=['92235 0.2883155436+/-0.0000000024 92238 0.7116844564+/-0.0000000024',stringCalcu

80 df = pd.read_csv('../Data/AtomicWeights.csv')

ModMass,ModSymbols,AtomFractions=fun.FormatMods(Modifications,df)
85 MolarMass=fun.DetermineMolarMass(List,df,
                                   ModSymbols,ModMass,
                                   AtomFractions,ChemicalFormulaError)

# print(MolarMass)

90 #####
##### Calculate Molality from #####
##### Wt % #####

```



```

#####
95  gramsOmol=MolarMass
    WtConcentration=ufloat(69,0.1)
    Molality=1000/(gramsOmol*(100/WtConcentration-1))

100 #####
    ##### Convert molality/molarity #####
    #####

    MolarityToMolality=True

105  gramsOmol=MolarMass

    #Density in grams per cc or grams per ml
    dfDen=pd.read_csv('../Data/Nitric_Acid.csv')

110  Temperature=ufloat(20,3) #Same degrees as dfDen!!!

    Molality=Molality
    Molarity=ufloat(15.43,0.06)

115  if MolarityToMolality:
        Molality=fun.ConvertMol(MolarityToMolality,Molarity,
                                gramsOmol,dfDen,Temperature)
    else:
120     Molarity=fun.ConvertMol(MolarityToMolality,Molality,
                                gramsOmol,dfDen,Temperature)

    #print("Molarity = "+str(Molarity))
125  #print("Molality = "+str(Molality))

    #####
    ##### Calculate New Concentration #####
130  #####

    Vol1=1
    Vol2=1

135  gramsOmol=gramsOmol

    m1=Molality
    m2=Molality*.25

140  Temperature=ufloat(20,3)

    dfDen=pd.read_csv('../Data/Nitric_Acid.csv')

    m3,p3,Vol3,Wt,M3=fun.NewConcentration(m1,m2,gramsOmol,
145                                     Temperature,dfDen,
                                     Vol1,Vol2)

```

150

```
M1=fun.ConvertMol(False,m1,gramsOmol,dfDen,Temperature)
M2=fun.ConvertMol(False,m2,gramsOmol,dfDen,Temperature)

##### Time To execute #####

print("--- %s seconds ---" % (time.time() - start_time))
```

This is an example citation [1].

References

- [1] E. T. Tatro, S. Heffler, S. Shumaker-Armstrong, B. Soontornniyomkij, M. Yang, A. Yermanos, N. Wren, D. J. Moore, and C. L. Achim. Modulation of bk channel by microrna-9 in neurons after exposure to hiv and methamphetamine. *J Neuroimmune Pharmacol*, 2013. Tatro, Erick T Heffler, Shannon Shumaker-Armstrong, Stephanie Soontornniyomkij, Benchawanna Yang, Michael Yermanos, Alex Wren, Nina Moore, David J Achim, Cristian L R03 DA031591/DA/NIDA NIH HHS/United States U19 AI096113/AI/NIAID NIH HHS/United States Journal article Journal of neuroimmune pharmacology : the official journal of the Society on NeuroImmune Pharmacology J Neuroimmune Pharmacol. 2013 Mar 19.