Statistical Methods for Population Health

Week 1: Introduction to Statistics

Ruoqing Zhu, Ph.D. <rqzhu@illinois.edu> July 3, 2021

Department of Statistics University Illinois Urbana-Champaign

Welcome!

· Welcome to the Statistics section!

Welcome!

- · Welcome to the Statistics section!
- This is a sequence of six lectures (including two TBL sessions)

Welcome!

- · Welcome to the Statistics section!
- This is a sequence of six lectures (including two TBL sessions)
- · Core skills
 - Statistical principles
 - · Modeling techniques
 - · Data analysis using R
 - · Result interpretation

• Week 1: The philosophy of statistics

- · Week 1: The philosophy of statistics
- · Week 2: R introduction and final project overview

- · Week 1: The philosophy of statistics
- Week 2: R introduction and final project overview
- Week 3: Testing Mean Differences and Associations

- · Week 1: The philosophy of statistics
- Week 2: R introduction and final project overview
- Week 3: Testing Mean Differences and Associations
- Week 4: TBL Correlation and Linear Regression

- · Week 1: The philosophy of statistics
- Week 2: R introduction and final project overview
- Week 3: Testing Mean Differences and Associations
- Week 4: TBL Correlation and Linear Regression
- Week 5: TBL Classification and Logistic Regression

- · Week 1: The philosophy of statistics
- Week 2: R introduction and final project overview
- Week 3: Testing Mean Differences and Associations
- Week 4: TBL Correlation and Linear Regression
- Week 5: TBL Classification and Logistic Regression
- Week 6: Survival Analysis and Beyond

• Week 1: The philosophy of statistics

The Lady Tasting Tea

The Lady Tasting Tea Problem

In 1920s Cambridge, England, a
 Lady, named Muriel Bristol, claimed
 to be able to tell whether the tea or
 the milk was added first by the taste
 of it!



"The Lady Tasting Tea: How Statistics Revolutionized Science in the Twentieth Century" (2001) by David Salsburg

The Lady Tasting Tea Problem

- In 1920s Cambridge, England, a
 Lady, named Muriel Bristol, claimed to be able to tell whether the tea or the milk was added first by the taste of it!
- A statistician Ronald Fisher what to test if thats true or not using probability principles



"The Lady Tasting Tea: How Statistics Revolutionized Science in the Twentieth Century" (2001) by David Salsburg

 Suppose Lady Bristol does not have that ability, then she will be randomly guessing

- Suppose Lady Bristol does not have that ability, then she will be randomly guessing
- Let's prepare many cups of tea for her to identify, then we would expect her to identify, on average, half of them correctly.

- Suppose Lady Bristol does not have that ability, then she will be randomly guessing
- Let's prepare many cups of tea for her to identify, then we would expect her to identify, on average, half of them correctly.
- However, if she can identify many of them correctly, then we may have to reject the assumption of random guessing

- Suppose Lady Bristol does not have that ability, then she will be randomly guessing
- Let's prepare many cups of tea for her to identify, then we would expect her to identify, on average, half of them correctly.
- However, if she can identify many of them correctly, then we may have to reject the assumption of random guessing
- The question is, how many is too many?

- · Two important concepts:
 - 1. Experimental design
 - 2. Hypothesis testing



Sir Ronald A. Fisher (1890 - 1962)

• Fisher prepared 8 cups of tea, 4 with milk added first and 4 with tea added first.

- Fisher prepared 8 cups of tea, 4 with milk added first and 4 with tea added first.
- · Lady Bristol was asked to identify them
- There are totally $\frac{8!}{4!(8-4)!}=70$ possible results. If she is randomly guessing, then each result has equal chance:

- Fisher prepared 8 cups of tea, 4 with milk added first and 4 with tea added first.
- · Lady Bristol was asked to identify them
- There are totally $\frac{8!}{4!(8-4)!} = 70$ possible results. If she is randomly guessing, then each result has equal chance:
 - The chance of identifying all 4 correctly is 1/70
 - The chance of 3 is 16/70
 - The chance of 2 is 36/70
 - The chance of 1 is 16/70
 - The chance of 0 is 1/70

- Fisher prepared 8 cups of tea, 4 with milk added first and 4 with tea added first.
- · Lady Bristol was asked to identify them
- There are totally $\frac{8!}{4!(8-4)!} = 70$ possible results. If she is randomly guessing, then each result has equal chance:
 - The chance of identifying all 4 correctly is 1/70
 - The chance of 3 is 16/70
 - The chance of 2 is 36/70
 - The chance of 1 is 16/70
 - The chance of 0 is 1/70
- What can be considered as "surprising" evidence given the assumption that she is randomly guessing?

Fisher used 0.05 as the cut-off:
 If the actual result falls into the top 5% of the most extreme cases, we consider this as a surprising evidence and claim that she is not randomly guessing.

- Fisher used 0.05 as the cut-off:
 If the actual result falls into the top 5% of the most extreme cases, we consider this as a surprising evidence and claim that she is not randomly guessing.
- How many cups Lady Bristol identified correctly?

Some **key steps** in hypothesis testing:

1). Form Null and Alternative hypotheses:

Null H_0 : Random Guessing vs. Alt. H_1 : Not Random Guessing

Some **key steps** in hypothesis testing:

1). Form Null and Alternative hypotheses:

Null H_0 : Random Guessing vs. Alt. H_1 : Not Random Guessing

2). Perform an experiment and observe that the lady identified the 4 correctly.

Some **key steps** in hypothesis testing:

1). Form Null and Alternative hypotheses:

Null H_0 : Random Guessing vs. Alt. H_1 : Not Random Guessing

- Perform an experiment and observe that the lady identified the 4 correctly.
- 3). If the Null hypothesis is correct, there is only 1.4% chance that one can guess 4 correctly

Some **key steps** in hypothesis testing:

1). Form Null and Alternative hypotheses:

Null H_0 : Random Guessing vs. Alt. H_1 : Not Random Guessing

- Perform an experiment and observe that the lady identified the 4 correctly.
- 3). If the Null hypothesis is correct, there is only 1.4% chance that one can guess 4 correctly
- 4). This is a "small probability event" (smaller than a pre-determined significance level, $\alpha=0.05$), so we will make a conclusion to reject the Null.

• If we reject the Null hypothesis, does it mean that Lady Bristol actually has the ability to identify them?

Correct or Wrong decision?

 We could still make a wrong decision. In fact, there are four situations:

	Accept H_0	Reject H_0
H_0 true	✓	Type I Error
H_0 false	Type II Error	\checkmark

- Type I error: H_0 true but we reject it.
- Type II error: H_0 false but we accept it.

Correct or Wrong decision?

- Type I error can be controlled using the α level we choose.
- 1 − Type I error is called the confidence level

Correct or Wrong decision?

- Type I error can be controlled using the α level we choose.
- 1 − Type I error is called the confidence level
- Type II error is difficult to analyze because we don't know what
 the alternative may look like. For example, the lady may have 0.7
 probability to identify a correct one, or 0.9, 0.51, etc. They all can
 have different Type II errors.
- 1 − Type II error is called the power.

Summary

- Statistics is a tool to analyze data and find patterns
- · However, statistics cannot provide a definitive answer
- · Definitive answers come from understanding the science

Homework

- Further reading (textbook): Sections 11.3.3 and 11.3.4
 - "Quantitative methods for health research: a practical interactive guide to epidemiology and statistics" by Nigel Bruce, Daniel Pope, Debbi Stanistreet. Hoboken, NJ:Wiley, 2018 2nd edition. Wiley Online Library [Download Link]
- Install RStudio and R