

Variations of atmospheric $^{12}\text{CO}_2$ and $^{13}\text{CO}_2$ from the year 1000 to 2024

W. A. van Wijngaarden¹ and W. Happer²

¹*Department of Physics and Astronomy, York University, Canada*

²*Department of Physics, Princeton University, USA*

September 30, 2026

Abstract

A simple model gives good agreement between observed emissions of CO_2 into the atmosphere from the combustion of fossil fuels and observed increases of the atmospheric concentrations C of CO_2 and decreases of the isotope delta value $\delta^{13}\text{C}$. If fossil fuel emissions were to cease the model predicts that: (1) the difference between the CO_2 concentration C and the preindustrial value, $C_0 \approx 280$ ppm, would decay exponentially with time constant $\tau_1 \approx 60$ y; (2) the difference between the concentration-delta-value product $C \times \delta^{13}\text{C}$ and the preindustrial value $C_0 \times \delta^{13}\text{C}_0 = (280 \text{ ppm}) \times (-6.5 \text{ ‰})$, would decay exponentially with a time constant $\tau_2 \approx 8$ y. The decay times are different because of isotopic exchange, largely driven by photosynthesis and respiration on the land.

1 Introduction

As is widely known, and summarized in a recent paper by Engelbeen, Hannon and Burton [1], the increasing concentration C of atmospheric CO_2 and the decrease of the isotope delta value $\delta^{13}\text{C}$ (the Suess effect), can be qualitatively explained by the addition of ^{13}C -depleted CO_2 from the combustion of fossil fuels to the atmosphere. In this note, we show that both facts are quantitatively consistent with a simple, 2-parameter model. If there were no further emissions of fossil fuels, the first parameter is the exponential decay time τ_1 of any difference of the CO_2 concentration C from the preindustrial value, $C_0 = 280$ ppm. The second parameter is the exponential decay time τ_2 of any difference of the concentration-delta-value product $C \times \delta^{13}\text{C}$ from the preindustrial value, $C_0 \times \delta^{13}\text{C}_0 = 280 \text{ ppm} \times (-6.5 \text{ ‰})$.

2 A Simple Model

2.1 Total carbon

Let the atmosphere contain N_1 atoms of the stable carbon isotope ^{12}C and N_2 atoms of the stable isotope ^{13}C . Most of these atoms will be in CO_2 molecules. To facilitate subsequent




Figure 1: Yearly fossil fuel emissions of CO_2 to the atmosphere in petagrams (Pg) or 10^{15} g calendar year. The observational data is from reference [2, 3].

discussions, we will describe the numbers of carbon atoms as the elements of a (2×1) dimensional *isotope number vector*

$$|N\rangle = \begin{bmatrix} N_1 \\ N_2 \end{bmatrix} = N \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}. \quad (1)$$

Here

$$N = N_1 + N_2, \quad n_1 = N_1/N, \quad n_2 = N_2/N \quad \text{and} \quad n_1 + n_2 = 1. \quad (2)$$

Large scale additions of CO_2 to the atmosphere began around the year 1850, as shown in Fig. 1. From Fig. 2 we see that the total number N of carbon atoms, or the concentration $C = N/N_{\text{at}}$, was nearly constant before the year 1800. The total number of atmospheric molecules [11] is

$$N_{\text{at}} = 1.065 \times 10^{44}. \quad (3)$$

The pre-industrial concentration of CO_2 was about

$$C_0 = 280 \text{ ppm}. \quad (4)$$

We denote the preindustrial numbers P_1 of ^{12}C atoms and P_2 of ^{13}C atoms with an isotope number vector analogous to (1),

$$|P\rangle = \begin{bmatrix} P_1 \\ P_2 \end{bmatrix} = P \begin{bmatrix} p_1 \\ p_2 \end{bmatrix}, \quad (5)$$

where

$$P = P_1 + P_2, \quad p_1 = P_1/P, \quad p_2 = P_2/P \quad \text{and} \quad p_1 + p_2 = 1. \quad (6)$$




Figure 2: The continuous blue line shows observed yearly averages of CO_2 concentrations $C = N/N_{\text{at}}$ in ppm (parts per million by volume) obtained from references [4, 5, 6, 7]. The continuous red line is observed isotope delta values $\delta^{13}\text{C}$ found from references [8, 9, 10]. The corresponding dotted lines are model predictions from (73) and (74). The preindustrial concentrations of CO_2 were close to $C_0 = 280$ ppm and the preindustrial isotope delta values were close to $\delta^{13}\text{C}_0 = -6.5$ ‰. The e-folding time constant $\tau_1 = 1/\gamma_1$ of (73) for the decay of concentration perturbations, $C - C_0$, was $\tau_1 = 60$ y. The e-folding time constant $\tau_2 = 1/\gamma_2$ of (74) for the decay of perturbations, $C \times \delta^{13}\text{C} - C_0 \times \delta^{13}\text{C}_0$, of concentration-delta-value products was $\tau_2 = 8$ y.

The total number of preindustrial carbon atoms in the atmosphere was

$$P = P_1 + P_2 = N_{\text{at}} C_0 = 2.98 \times 10^{40}. \quad (7)$$

We used the preindustrial CO_2 concentration C_0 from (4) and the total number of atmospheric molecules N_{at} of (3) to write the numerical value on the right of (7)

Let F_1 denote the addition rate of ^{12}C atoms from the combustion of fossil fuels, and let F_2 the addition rate of ^{13}C atoms. In analogy to (1) and (5), we will write these as the fossil-fuel *isotope source vector*

$$|F\rangle = \begin{bmatrix} F_1 \\ F_2 \end{bmatrix} = F \begin{bmatrix} f_1 \\ f_2 \end{bmatrix}, \quad (8)$$

where

$$F = F_1 + F_2, \quad f_1 = F_1/F, \quad f_2 = F_2/F \quad \text{and} \quad f_1 + f_2 = 1. \quad (9)$$

The elements of $|F\rangle$ have the units of atoms per unit time. For the elements of $|N\rangle$ and $|P\rangle$ the units are simply atoms.

Reference [12] showed that the observed number, $N = N(t)$, of atmospheric CO_2 molecules at time t is given accurately in terms of emissions $F' = F(t')$ at earlier times t' by the simple

mathematical expression

$$N = \int_{-\infty}^t dt' e^{-\gamma_1(t-t')} F' + P. \quad (10)$$

Differentiating (10) with respect to time, we see that it is a solution of the differential equation

$$\begin{aligned} \frac{d}{dt}N &= -\gamma_1 \int_{-\infty}^t dt' e^{-\gamma_1(t-t')} F' + F \\ &= -\gamma_1(N - P) + F. \end{aligned} \quad (11)$$

For no fossil fuel emissions, $F = 0$, the differential equation (11) is an analog of *Newton's law of cooling* [13]. If the number N of CO_2 differs from the preindustrial value P , and if there are no additions F of CO_2 from fossil fuel combustion (if $F = 0$), then N will decay exponentially at the rate γ_1 to the preindustrial value P .

For time-independent conditions, when $dN/dt = 0$ and $dF/dt = 0$ or $F = \text{constant}$, the solution to (11) is

$$N = \frac{P\gamma_1 + F}{\gamma_1} = \frac{D + F}{\gamma_1}. \quad (12)$$

Here we have introduced a *deep source* of CO_2 that is injected into the atmosphere,

$$D = P\gamma_1 = 4.97 \times 10^{38} \text{ y}^{-1}. \quad (13)$$

The numerical value on the right of (13) came from assuming the decay rate

$$\gamma_1 = 1/60 \text{ y}. \quad (14)$$

The numerical value (14) of γ_1 was used to draw the model curves of Fig. 2, as we will describe below. The mean mass of a terrestrial carbon atom is

$$m = 1.994 \times 10^{-23} \text{ g}. \quad (15)$$

Using the numerical value (15) with (13), we find that the pre-industrial addition rate of carbon to the atmosphere, in mass per unit time, is

$$\frac{dM}{dt} = Dm = 9.91 \text{ Pg y}^{-1}. \quad (16)$$

Here Pg is the unit petagram, or 10^{15} g of carbon atoms. From inspection of the top panel of Fig. 1, we see that the current emission rate of fossil fuel carbon, around 10.5 Pg y^{-1} in the year 2024, is slightly larger than the model preindustrial rate (16).

To facilitate subsequent analysis, we introduce the row vector

$$\langle U_1 | = [1 \quad 1]. \quad (17)$$

We note that $N = N_1 + N_2$, the sum of the number N_1 of ^{12}C atoms and N_2 of ^{13}C atoms, can be written as the product of the row vector (17) and the column vector (1),

$$\langle U_1 | N \rangle = [1 \quad 1] \begin{bmatrix} N_1 \\ N_2 \end{bmatrix} = N_1 + N_2 = N. \quad (18)$$

In like manner we can use (17) with (5) and (8) to write

$$P = \langle U_1|P \rangle \quad \text{and} \quad F = \langle U_1|F \rangle. \quad (19)$$

Letting

$$\gamma = \gamma_1, \quad (20)$$

we use (18) and (19) to write the last line of (11) as

$$\frac{d}{dt} \langle U_1|N \rangle = -\gamma_1 \langle U_1| \left(|N \rangle - |P \rangle \right) + \langle U_1|F \rangle. \quad (21)$$

2.2 Isotope fractions

The stable isotope ^{12}C makes up about 98.9% of Earth's carbon. The heavier isotope, ^{13}C , makes up almost all of the remaining 1.1%. Although the ratio N_2/N_1 of the number N_2 of ^{13}C atoms to the number N_1 of ^{12}C atoms is very nearly 0.011, *isotopic fractionation* [14] can cause slight differences in the isotope ratios of different carbon samples. For example, the organic carbon in trees has a lesser fraction of ^{13}C than the air from which the carbon was extracted and fixed by photosynthesis. This is because the lighter $^{12}\text{CO}_2$ molecules diffuse slightly faster through leaf stomata and through the leaf tissue to the chloroplasts. And the carbon-fixing enzyme rubisco works slightly more efficiently with the lighter carbon isotope.

It is customary to quantify the ratio, $[N_2/N_1]_x$, in a sample X of carbon atoms by the *isotope delta value*

$$\delta^{13}\text{C} = \delta_{xv} = \frac{[N_2/N_1]_x}{[N_2/N_1]_v} - 1. \quad (22)$$

Mass spectrometers are used to precisely measure the ratio $[N_2/N_1]_x$ in the sample X , with respect to $[N_2/N_1]_v$ in a calibrating standard V . The standard is usually *Vienna Pee Dee Belemnite* or VPDB. This standard traces back to fossil belemnite shells from a late Cretaceous formation along the Pee Dee river in South Carolina [15]. Contemporary VPDB standards are artificially prepared to have isotopic ratios nearly identical to the original belemnite standards. A recent measurement [16] found that the absolute ratio of ^{13}C to ^{12}C atoms in VPDB standards is

$$[N_2/N_1]_v = 0.0111105 \pm 0.0000042. \quad (23)$$

The pre-industrial value of $\delta^{13}\text{C}$ for the atmosphere, determined from the CO_2 of air trapped in ice core bubbles, is around

$$\delta_{pv} = -6.5 \text{‰}. \quad (24)$$

where the symbol ‰ denotes parts per thousand, or the number 10^{-3} . The value of $\delta^{13}\text{C}$ for CO_2 from fossil fuel emissions averages about

$$\delta_{fv} = -28 \text{‰}. \quad (25)$$

For the year 2026, the observed $\delta^{13}\text{C}$ for the atmosphere was

$$\delta_{nv} = -8.7 \text{‰}. \quad (26)$$

The fact that (26), the contemporary value of $\delta^{13}\text{C}$, is less than the preindustrial value (24) is what one would qualitatively expect from the addition ^{13}C -depleted carbon from fossil fuels to the atmosphere. This is called the Suess effect [17]. More discussion of the isotope delta value $\delta^{13}\text{C}$ can be found in the Appendix.

For further analysis, it will be convenient to introduce a row vector $\langle U_2|$, analogous to $\langle U_1|$ of (17), and defined in terms of the fractions p_1 of ^{12}C and p_2 of ^{13}C in the preindustrial atmosphere by

$$\langle U_2| = \begin{bmatrix} -p_2 & p_1 \end{bmatrix}. \quad (27)$$

Then we see that

$$\begin{aligned} \langle U_2|N \rangle &= \begin{bmatrix} -p_2 & p_1 \end{bmatrix} \begin{bmatrix} N_1 \\ N_2 \end{bmatrix} \\ &= -p_2 N_1 + p_1 N_2 \\ &= p_2 N_1 \left(-1 + \frac{p_1 N_2}{p_2 N_1} \right) \\ &= N p_2 n_1 \delta_{np} \\ &= N p_1 p_2 \delta_{np} + \mathcal{O}(\delta^2). \end{aligned} \quad (28)$$

To write the last line of (28) we noted from the Appendix that the fraction n_1 of ^{12}C atoms in the current atmosphere is related to the fraction p_1 in the preindustrial atmosphere by $n_1 = p_1 + \mathcal{O}(\delta)$. Therefore, to order δ we can use (28) to write

$$\delta_{np} = \frac{\langle U_2|N \rangle}{p_1 p_2 N}. \quad (29)$$

From inspection of Fig. 2 one can see that the isotope delta value, $\delta^{13}\text{C}$, remained close to the preindustrial value $\delta^{13}\text{C} = -6.5\text{‰}$ from the year 1000 to the year 1800, the beginning of the industrial age, when fossil fuel combustion began to add significant amounts of ^{13}C -depleted CO_2 to the atmosphere. It is therefore natural to assume that the variable $\langle U_2|N \rangle$ of (29), which is proportional to $N\delta_{np}$, obeys an analog of Newton's law of cooling [13], or an analog of (21) for the number N of CO_2 molecules in the atmosphere,

$$\frac{d}{dt} \langle U_2|N \rangle = -\gamma_2 \langle U_2| \left(|N \rangle - |P \rangle \right) + \langle U_2|F \rangle. \quad (30)$$

2.3 Reciprocal vectors

To facilitate further discussion we define column vectors,

$$|U_1 \rangle = \begin{bmatrix} p_1 \\ p_2 \end{bmatrix} \quad \text{and} \quad |U_2 \rangle = \begin{bmatrix} -1 \\ 1 \end{bmatrix}, \quad (31)$$

that are the reciprocals of the row vectors $\langle U_1|$ of (17) and $\langle U_2|$ of (27),

$$\langle U_1| = \begin{bmatrix} 1 & 1 \end{bmatrix} \quad \text{and} \quad \langle U_2| = \begin{bmatrix} -p_2 & p_1 \end{bmatrix}. \quad (32)$$

The reciprocal row and column vectors are readily seen to be orthonormal,

$$\langle U_j | U_k \rangle = \delta_{jk}, \quad (33)$$

where the Kronecker delta symbol is $\delta_{jk} = 1$ if $j = k$ and $\delta_{jk} = 0$ if $j \neq k$. The vectors are also complete in the sense that

$$\sum_j |U_j\rangle \langle U_j| = \hat{1} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}. \quad (34)$$

Here we have used the symbol $\hat{1}$ to denote the (2×2) identity matrix. One can verify that the product of the identity matrix $\hat{1}$ with an arbitrary column vector $|X\rangle$ on the right or an arbitrary row vector $\langle X|$ on the left simply reproduces the vectors

$$\hat{1}|X\rangle = |X\rangle \quad \text{and} \quad \langle X|\hat{1} = \langle X|. \quad (35)$$

To prove the completeness (34) of the reciprocal vectors we use (31) and (32) to write

$$\begin{aligned} \sum_j |U_j\rangle \langle U_j| &= \begin{bmatrix} p_1 \\ p_2 \end{bmatrix} \begin{bmatrix} 1 & 1 \end{bmatrix} + \begin{bmatrix} -1 \\ 1 \end{bmatrix} \begin{bmatrix} -p_2 & p_1 \end{bmatrix} \\ &= \begin{bmatrix} p_1 & p_1 \\ p_2 & p_2 \end{bmatrix} + \begin{bmatrix} p_2 & -p_1 \\ -p_2 & p_1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \hat{1}. \end{aligned} \quad (36)$$

To write the last line of (36) we recalled from (6) that $p_1 + p_2 = 1$.

Multiplying both sides of (21) on the left by $|U_1\rangle$, multiplying both sides of (30) on the left by $|U_2\rangle$, and adding the resulting equations we find

$$\begin{aligned} &|U_1\rangle \frac{d}{dt} \langle U_1 | N \rangle + |U_2\rangle \frac{d}{dt} \langle U_2 | N \rangle = \\ &- |U_1\rangle \gamma_1 \langle U_1 | \left(|N\rangle - |P\rangle \right) - |U_2\rangle \gamma_2 \langle U_2 | \left(|N\rangle - |P\rangle \right) + |U_1\rangle \langle U_1 | F \rangle + |U_2\rangle \langle U_2 | F \rangle. \end{aligned} \quad (37)$$

Using the completeness property (34) we can write (37) as

$$\frac{d}{dt} |N\rangle = -\Gamma \left(|N\rangle - |P\rangle \right) + |F\rangle. \quad (38)$$

In (38) the damping matrix Γ is

$$\begin{aligned} \Gamma &= |U_1\rangle \gamma_1 \langle U_1| + |U_2\rangle \gamma_2 \langle U_2| \\ &= \begin{bmatrix} p_1 \gamma_1 + p_2 \gamma_2 & p_1 (\gamma_1 - \gamma_2) \\ p_2 (\gamma_1 - \gamma_2) & p_2 \gamma_1 + p_1 \gamma_2 \end{bmatrix} \\ &= \begin{bmatrix} \gamma_1 + \gamma_{12} p_2 & -\gamma_{21} p_1 \\ -\gamma_{12} p_2 & \gamma_1 + \gamma_{21} p_1 \end{bmatrix}. \end{aligned} \quad (39)$$

Here we have introduced the characteristic *isotope exchange rate*

$$\gamma_{12} = \gamma_{21} = |\gamma_2 - \gamma_1|. \quad (40)$$

From inspection of the first line of (39) and the orthonormality relation (33) we see that the row vectors $\langle U_q|$ of (32) are left eigenvectors of Γ with eigenvalues γ_q , and the column vectors $|U_q\rangle$ of (31) are right eigenvectors with the same eigenvalues γ_q ,

$$\langle U_q|\Gamma = \gamma_q\langle U_q| \quad \text{and} \quad \Gamma|U_q\rangle = |U_q\rangle\gamma_q. \quad (41)$$

From inspection of (5) and (31) we see that the preindustrial isotope number vector $|P\rangle$ is proportional to the right eigenvector $|U_1\rangle$ of Γ

$$|P\rangle = P|U_1\rangle. \quad (42)$$

Using (42) and (41), we can write (38) as

$$\frac{d}{dt}|N\rangle = -\Gamma|N\rangle + \gamma_1|P\rangle + |F\rangle. \quad (43)$$

A general solution of (43) is

$$|N\rangle = \int_{-\infty}^t dt' e^{-\Gamma(t-t')} |F'\rangle + |P\rangle + e^{-\Gamma t} |K\rangle. \quad (44)$$

Here the time-independent, (2×1) vector $|K\rangle$ must be determined by boundary conditions. For $|K\rangle \neq 0$ the solution (44) is unphysical, since it goes to infinity as $t \rightarrow -\infty$. We therefore set $|K\rangle = 0$ and we write (44), in analogy to (10), as

$$|N\rangle = \int_{-\infty}^t dt' e^{-\Gamma(t-t')} |F'\rangle + |P\rangle. \quad (45)$$

We recall that e^x , the exponentiated value of any (2×2) matrix x of the form

$$x = |U_1\rangle x_1 \langle U_1| + |U_2\rangle x_2 \langle U_2| \quad (46)$$

can be evaluated with the Taylor series,

$$\begin{aligned} e^x &= \hat{1} + \frac{x}{1!} + \frac{x^2}{2!} + \dots \\ &= |U_1\rangle \langle U_1| + |U_2\rangle \langle U_2| + \frac{1}{1!} \left(|U_1\rangle x_1 \langle U_1| + |U_2\rangle x_2 \langle U_2| \right) + \frac{1}{2!} \left(|U_1\rangle x_1^2 \langle U_1| + |U_2\rangle x_2^2 \langle U_2| \right) + \dots \\ &= |U_1\rangle \left(1 + \frac{x_1}{1!} + \frac{x_1^2}{2!} + \dots \right) \langle U_1| + |U_2\rangle \left(1 + \frac{x_2}{1!} + \frac{x_2^2}{2!} + \dots \right) \langle U_2| \\ &= |U_1\rangle e^{x_1} \langle U_1| + |U_2\rangle e^{x_2} \langle U_2|. \end{aligned} \quad (47)$$

Evaluating (47) with $x = -\Gamma(t - t')$, $x_1 = -\gamma_1(t - t')$ and $x_2 = -\gamma_2(t - t')$, in accordance with (39), we see that the exponentiated matrix of (44) can be written as

$$e^{-\Gamma(t-t')} = |U_1\rangle e^{-\gamma_1(t-t')} \langle U_1| + |U_2\rangle e^{-\gamma_2(t-t')} \langle U_2|. \quad (48)$$

Eq. (48) implies that

$$\langle U_q | e^{-\Gamma(t-t')} = e^{-\gamma_q(t-t')} \langle U_q | \quad \text{and} \quad e^{-\Gamma(t-t')} | U_q \rangle = e^{-\gamma_q(t-t')} | U_q \rangle. \quad (49)$$

In component form, we can use (39) to write (43) as

$$\frac{d}{dt} \begin{bmatrix} N_1 \\ N_2 \end{bmatrix} = \begin{bmatrix} -\gamma_1 N_1 - \gamma_{12} p_2 N_1 + \gamma_{21} p_1 N_2 + \gamma_1 P_1 + F_1 \\ -\gamma_1 N_2 + \gamma_{12} p_2 N_1 - \gamma_{21} p_1 N_2 + \gamma_1 P_2 + F_2 \end{bmatrix}. \quad (50)$$

The physical significance of the terms on the right of (50) is illustrated by Fig. 3. Carbon




Figure 3: The rates of change of carbon isotopes in the atmosphere. The most abundant isotope, ^{12}C is shown in blue and the rare stable isotope ^{13}C is shown in red. Negative terms on the right of (50), or *sinks*, represent losses of carbon atoms from the atmosphere, and are denoted by downward arrows. Positive terms on the right of (50), *sources*, represent carbon atoms added to the atmosphere and are denoted with upward arrows.

atoms are transferred from the atmosphere to the deep ocean and deep terrestrial soil at the same rate, γ_1 , for both isotopes, ^{12}C and ^{13}C . The isotopes are exchanged at rates proportional to the characteristic exchange rate γ_{12} of (40), and this drives isotope delta values $\delta^{13}\text{C}$ toward preindustrial values. The rapid drawdown of atmospheric CO_2 by photosynthesis of land plants during the summer, and the steady return of carbon due to the respiration of soil and water microorganisms throughout the year makes a major contribution to the exchange rates. Upwelling deep ocean water and oxidation of deeply buried reduced carbon in the soil both inject CO_2 with preindustrial isotope delta values $\delta^{13}\text{C}$ into the atmosphere.

2.4 The Suess effect

Multiplying both sides of (44) on the left by $\langle U_1|$, and using (18) and (19) we find

$$\begin{aligned}
\langle U_1|N\rangle &= \int_{-\infty}^t dt' \langle U_1|e^{-\Gamma(t-t')}|F'\rangle + \langle U_1|P\rangle \\
&= \int_{-\infty}^t dt' e^{-\gamma_1(t-t')} \langle U_1|F'\rangle + P \\
&= \int_{-\infty}^t dt' e^{-\gamma_1(t-t')} F' + P
\end{aligned} \tag{51}$$

or

$$N = \Delta N_1 + P. \tag{52}$$

We used (49) to write the second line of (51). The *fossil fuel increments* $\Delta N_q = \Delta N_q(t)$, with $q = 1$ or $q = 2$, are defined by

$$\Delta N_q = \int_{-\infty}^t dt' e^{-\gamma_q(t-t')} F'. \tag{53}$$

Multiplying both sides of (44) on the left by $\langle U_2|$ we find,

$$\begin{aligned}
\langle U_2|N\rangle &= \int_{-\infty}^t dt' \langle U_2|e^{-\Gamma(t-t')}|F'\rangle + \langle U_2|P\rangle \\
&= \int_{-\infty}^t dt' e^{-\gamma_2(t-t')} \langle U_2|F'\rangle + \langle U_2|P\rangle.
\end{aligned} \tag{54}$$

Noting that $\langle U_2|P\rangle = P(-p_2p_1 + p_1p_2) = 0$, we use (28) to write (54), to order δ , as

$$Np_1p_2\delta_{np} = \int_{-\infty}^t dt' e^{-\gamma_2(t-t')} F' p_1p_2\delta_{fp}. \tag{55}$$

Dividing both sides of (55) by Np_1p_2 and using (53) we find

$$\delta_{np} = \frac{\Delta N_2}{N} \delta_{fp}. \tag{56}$$

As shown by (87), the isotope delta values are nearly additive so we can use (56) to write

$$\begin{aligned}
\delta_{nv} &= \delta_{np} + \delta_{pv} \\
&= \frac{\Delta N_2}{N} \delta_{fp} + \delta_{pv}.
\end{aligned} \tag{57}$$

2.4.1 No exchange

From inspection of (40) we see that in the limit of no isotopic exchange, $\gamma_{12} \rightarrow 0$

$$\gamma_2 \rightarrow \gamma_1. \tag{58}$$

Then we see from (53) and (52) that

$$\Delta N_2 \rightarrow \Delta N_1 = N - P. \quad (59)$$

Substituting (59) into (57), we find

$$\begin{aligned} \delta_{nv} &= \frac{(N - P)\delta_{fp}}{N} + \delta_{pv} \\ &= \frac{(N - P)(\delta_{fv} - \delta_{pv}) + N\delta_{pv}}{N} \\ &= \frac{(N - P)\delta_{fv} + P\delta_{pv}}{N}. \end{aligned} \quad (60)$$

To write the second line of (60) we recalled from (87) that the shift parameters δ are nearly additive, so

$$\delta_{fv} = \delta_{fp} + \delta_{pv}. \quad (61)$$

It is convenient to divide the numerator and denominator of the last line of (60) by the total number of atmospheric molecules N_{at} from (3) to find

$$\delta_{nv} = \frac{(C - C_0)\delta_{fv} + C_0\delta_{pv}}{C}. \quad (62)$$

The contemporary concentration C of atmospheric CO_2 and the preindustrial concentration C_0 are

$$C = \frac{N}{N_{\text{at}}} \quad \text{and} \quad C_0 = \frac{P}{N_{\text{at}}}. \quad (63)$$

The no-exchange limit of (60) or (62) is simply the average of $\delta^{13}\text{C} = \delta_{pv} = -6.5\text{‰}$, the $\delta^{13}\text{C}$ value for the P atoms of atmospheric carbon from natural sources, and $\delta^{13}\text{C} = \delta_{fv} = -28\text{‰}$, the $\delta^{13}\text{C}$ value for the $N - P$ atoms of atmospheric carbon from fossil fuels. For the year 2026, the concentrations needed to evaluate (62) are $C_0 = 280$ ppm and $C = 430$ ppm

$$\delta^{13}\text{C} = \frac{(430 - 280)(-28) + 280(-6.5)}{430} \text{‰} = -14.0 \text{‰}. \quad (64)$$

This is much smaller than the observed value, $\delta^{13}\text{C} = -8.7\text{‰}$, of (26).

2.4.2 Infinitely fast exchange

From inspection of (40) we see that for a fixed decay rate γ_1 of the concentration C to the preindustrial value, $C_0 = 280$ ppm, but for an infinitely fast exchange rate,

$$\gamma_2 \rightarrow \infty \quad \text{when} \quad \gamma_{12} \rightarrow \infty. \quad (65)$$

Then we see from (53) that

$$\begin{aligned}
\Delta N_2 &= \int_{-\infty}^t dt' e^{-\gamma_2(t-t')} F(t+t'-t) \\
&= \int_{-\infty}^t dt' e^{-\gamma_2(t-t')} \left[F(t) + \frac{(t'-t)}{1!} \frac{dF}{dt}(t) + \frac{(t'-t)^2}{2!} \frac{d^2F}{dt^2}(t) + \dots \right] \\
&= \frac{1}{\gamma_2} F(t) - \frac{1}{\gamma_2^2} \frac{dF}{dt}(t) + \frac{1}{\gamma_2^3} \frac{d^2F}{dt^2}(t) - \dots \\
&\rightarrow 0 \quad \text{as} \quad \gamma_2 \rightarrow \infty.
\end{aligned} \tag{66}$$

Using (66) with (57) we see that for the limit of infinitely fast isotope exchange

$$\delta^{13}\text{C} \rightarrow \delta_{pv} = -6.5\text{‰} \quad \text{as} \quad \gamma_{12} \rightarrow \infty. \tag{67}$$

This is the preindustrial value, no matter how large a fraction of fossil-fuel carbon is added to the atmosphere.

According to (67), an infinitely fast exchange rate with $\gamma_2 \rightarrow \infty$ gives no Suess effect at all. But from (64) we see that a vanishing exchange rate, $\gamma_{12} = 0$, with $\gamma_2 \rightarrow \gamma_1$, gives much too large a Suess effect. The real exchange rate γ_{12} of (40) has a value between zero and infinity.

2.5 Discretization

For model calculations with readily available observational data, it is useful to introduce a series of discrete times

$$[t_0, t_1, t_2, \dots, t_n] = [1850, 1851, 1853, \dots, 2024], \tag{68}$$

The numbers on the right side of (68) are the ends of calendar years in the industrial era. We assume that fossil fuel emissions, summarized by the data of references [2, 3], are small enough to neglect before the year $t_0 = 1850$, that is, we assume $F(t') = 0$ for $t' < t_0$. Dividing both sides of (53) by the total number of atmospheric molecules N_{at} of (3) gives the concentration increments at the end of the year t_j , with $j \geq 1$,

$$\begin{aligned}
\Delta C_{qj} = \frac{\Delta N_q(t_j)}{N_{\text{at}}} &= \int_{-\infty}^{t_j} dt' e^{-\gamma_q(t_j-t')} \frac{F(t')}{N_{\text{at}}} \\
&= \sum_{k=1}^j \int_{t_{k-1}}^{t_k} dt' e^{-\gamma_q(t_j-t')} \frac{F(t')}{N_{\text{at}}} \\
&\approx \sum_{k=1}^j e^{-\gamma_q(t_j-t_k)} \int_{t_{k-1}}^{t_k} dt' \frac{F(t')}{N_{\text{at}}},
\end{aligned} \tag{69}$$

or

$$\Delta C_{qj} = \sum_{k=1}^j e^{-\gamma_q(t_j-t_k)} E_k. \tag{70}$$

The concentration increments due to the release of fossil-fuel carbon atoms to the atmosphere in the year t_k are

$$E_k = \int_{t_{k-1}}^{t_k} dt' \frac{F(t')}{N_{\text{at}}}. \quad (71)$$

Dividing both sides of (52) by the total number of atmospheric molecules, N_{at} of (3), setting

$$C_j = \frac{N(t_j)}{N_{\text{at}}}, \quad (72)$$

using the definition (7) of C_0 and (69) of ΔC_{qj} we find for $j \geq 1$,

$$\begin{aligned} C_j &= \Delta C_{1j} + C_0 \\ &= \sum_{k=1}^j e^{-\gamma_1(t_j-t_k)} E_k + C_0, \end{aligned} \quad (73)$$

an expression given in reference [12].

In like manner, we can use (57) to write the isotope delta values for $j \geq 1$ as

$$\begin{aligned} \delta^{13}\text{C}_j &= \frac{\Delta C_{2j}}{C_j} + \delta_{pv} \\ &= \frac{1}{C_j} \sum_{k=1}^j e^{-\gamma_2(t_j-t_k)} E_k + \delta_{pv}. \end{aligned} \quad (74)$$

Here C_j was given by (73) and $\delta_{pv} = -6.5\text{‰}$ was given by (24). For the special case $j = 0$, which labels the beginning year of the industrial era, both the atmospheric concentrations of CO_2 and the isotope delta value have their preindustrial values,

$$C_0 = 280 \text{ ppm} \quad \text{and} \quad \delta^{13}\text{C}_0 = \delta_{pv} = -6.5\text{‰}. \quad (75)$$

The model curve for CO_2 concentration C of Fig. 2 was plotted from (73) with $C_0 = 280$ ppm, in accordance with (4), and with $\gamma_1 = 1/\tau_1 = 1/60$ y, which gave a good fit to observed values of C . A slightly different time constant, $\tau_1 = 50$ y was found in reference [12], where a value of $C_0 = 290$ ppm was used. The model curve for $\delta^{13}\text{C}$ in Fig. 2 was plotted from (74) and with $\gamma_2 = 1/\tau_2 = 1/8$ y. This gives a good fit to observed values of $\delta^{13}\text{C}$.

Using $\tau_1 = 60$ y and $\tau_2 = 8$ y, we can use the exchange rate γ_{12} of (40) to write the exchange time τ_{12} as

$$\tau_{12} = \frac{1}{\gamma_{12}} = \frac{\tau_1 \tau_2}{\tau_1 - \tau_2} = 9.2 \text{ y}. \quad (76)$$

3 Complete halt of fossil fuel emissions

To better understand the model, consider the hypothetical situation that all fossil fuel emissions stopped after the year 2024. Formally, we assume that the annual emissions (71) are

$$E_k = \begin{cases} \text{the values of Fig. 1 if } t_k \leq 2024, \\ 0 \text{ if } t_k > 2024. \end{cases} \quad (77)$$




Figure 4: Modelled time dependence of atmospheric CO_2 concentration C and isotopic delta value $\delta^{13}\text{C}$ if all fossil fuel emissions stopped after the year 2024, marked by the black dotted line. Because of isotope exchange, $\delta^{13}\text{C}$, shown as the smooth red line, evolves more quickly to its preindustrial value $\delta^{13}\text{C}_0 = -6.5\text{‰}$, shown as the red dotted line, than does the atmospheric concentration C , shown as the smooth blue line, to its preindustrial value, $C_0 = 280$ ppm, shown as the blue dotted line.

Substituting (77) into (73) and (74) we find the modelled CO_2 concentrations C and isotope delta values $\delta^{13}\text{C}$ of Fig. 4. Because of isotope exchange, after fossil fuel emissions stop the isotope delta value $\delta^{13}\text{C}$ relaxes more quickly to its preindustrial value $\delta^{13}\text{C}_0 = -6.5\text{‰}$, than does the atmospheric concentration C to its preindustrial value, $C_0 = 280$ ppm.

4 Conclusions

The simple model discussed here gives good agreement with observations. It could be elaborated to include additional details, for example, the changes of $\delta^{13}\text{C}$ from the beginning of the industrial era, when coal was the main fossil fuel to the present time when larger fractions of oil and natural gas are used [18]. The fraction of radioactive ^{14}C , both that steadily produced by cosmic rays and the pulse introduced by above ground nuclear tests [19] could be analyzed in a similar way.

A Appendix

A.1 Some properties of $\delta^{13}C$

In analogy to (1), we define an isotope number vector for the VPDB (Vienna Pee Dee Belemnite) standard by

$$|V\rangle = \begin{bmatrix} V_1 \\ V_2 \end{bmatrix} = V \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}, \quad (78)$$

where

$$V = V_1 + V_2, \quad v_1 = V_1/V, \quad v_2 = V_2/V \quad \text{and} \quad v_1 + v_2 = 1. \quad (79)$$

For an arbitrary sample X of carbon, with potentially different isotope ratios than that of the VPDB standard, the isotope number vector can be written as

$$|X\rangle = \begin{bmatrix} X_1 \\ X_2 \end{bmatrix} = X \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}, \quad (80)$$

where

$$X = X_1 + X_2, \quad x_1 = X_1/X, \quad x_2 = X_2/X \quad \text{and} \quad x_1 + x_2 = 1. \quad (81)$$

Then we can write the parameter $\delta^{13}C$ of (22) as

$$\begin{aligned} \delta_{xv} &= \frac{x_2/x_1}{v_2/v_1} - 1 \\ &= \frac{x_2(1 - v_2)}{v_2(1 - x_2)} - 1. \end{aligned} \quad (82)$$

Solving the second line of (82) for x_2 in terms of v_2 and δ_{xv} , and setting $x_1 = 1 - x_2$ we find

$$x_2 = v_2 + \frac{\delta_{xv}v_1v_2}{1 + \delta_{xv}v_2} \quad \text{and} \quad x_1 = v_1 - \frac{\delta_{xv}v_1v_2}{1 + \delta_{xv}v_2} \quad (83)$$

From (82) we see that the isotope delta values of a carbon samples, X compared to itself is zero,

$$\delta_{xx} = 0. \quad (84)$$

From (82) we find the *reciprocal law* for isotope delta values for two carbon samples, X and Y ,

$$(1 + \delta_{xy})(1 + \delta_{yx}) = 1. \quad (85)$$

From (82) we also find the *addition law* for isotope delta values for three carbon samples, X , Y and V ,

$$(1 + \delta_{yx})(1 + \delta_{xv}) = 1 + \delta_{yv}. \quad (86)$$

or

$$\begin{aligned} \delta_{yv} &= \delta_{yx} + \delta_{xv} + \delta_{yx}\delta_{xv} \\ &\approx \delta_{yx} + \delta_{xv}. \end{aligned} \quad (87)$$

Since the isotope delta values of natural samples of carbon are on the order of a per cent or less, they are approximately additive, as shown by the second line of (87).

Acknowledgements

The authors would like to thank their respective universities, York and Princeton, for support.

References

- [1] F. Engelbeen, R. Hannon and D. Burton, *The Human Contribution to Atmospheric Carbon Dioxide*, (2024). <https://co2coalition.org/wp-content/uploads/2024/12/Human-Contribution-to-Atmospheric-CO2-digital-compressed.pdf>
- [2] Global Carbon Emissions, <https://globalcarbonbudget.org/download/2371/?tmstv=176281725>
- [3] P. Friedlingstein et al, *Global Carbon Budget 2025*, Earth System Sci. Data, **18**, 3211–3288 (2026).
- [4] X. Lan and R. Keeling, *Trends in CO₂, CH₄, N₂O, SF₆: Trends in Atmospheric Carbon Dioxide: Mauna Loa CO₂ Annual Mean Data*. NOAA Global Monitoring Laboratory, Boulder, CO, USA, (2026). https://gml.noaa.gov/webdata/ccgg/trends/co2/co2_annmean_mlo.txt
- [5] D. M. Etheridge et al, *Natural and Anthropogenic Changes in Atmospheric CO₂ Over the Last 1000 Years from Air in Antarctic Ice and Firn*, J. Geophys. Res. **101** (D2), 4115–4128 (1996).
- [6] R. F. Keeling, S. C. Piper, A. F. Bollenbacher and S. J. Walker, *Ice-Core Merged Products: Yearly Spline*. Scripps CO₂ Program, Scripps Institution of Oceanography, University of California, La Jolla, CA, USA. https://keelinglabsites.ucsd.edu/websitedataco2/spline_merged_ice_core_yearly.csv
- [7] C. MacFarling Meure et al, *Law Dome CO₂, CH₄ and N₂O Ice Core Records Extended to 2000 Years BP*, Geophys. Res. Lett. **33**, L14810 (2006).
- [8] M. Rubino et al, *Law Dome, Antarctica 2000 Year Ice Core CO₂, CH₄, N₂O and $\delta^{13}C$ -CO₂ Data*, World Data Service for Paleoclimatology, Boulder, CO, USA, and NOAA Paleoclimatology Program, National Centers for Environmental Information (NCEI), (2024). <https://www.ncei.noaa.gov/pub/data/paleo/icecore/antarctica/law/law2018d13c-co2-noaa.txt>
- [9] M. Rubino et al, *Revised Records of Atmospheric Trace Gases CO₂, CH₄, N₂O, and $\delta^{13}C$ CO₂ over the last 2000 years from Law Dome, Antarctica*, Earth System Science Data, **11** (3), 473–492 (2019). <https://doi.org/10.5194/essd-11-473-2019>
- [10] R. F. Keeling, E. J. Morgan and C. D. Keeling, *Atmospheric Flask CO₂ and Isotopic Data Sets – Multiple Sampling Stations (Archive 2024-04-19)*, Scripps CO₂ Program Data, UC San Diego Library Digital Collections. <https://doi.org/10.6075/J0HQ3X30>

- [11] D. Alexander, J. D. Ferguson, A. Glatzle, W. Happer and W. A. van Wijngaarden, *Livestock, Methane and Climate*, Atmos. and Oceanic Phys. arXiv: 2601.18522v1 (2026). <https://co2coalition.org/publications/livestock-methane-and-climate/>
- [12] W. Happer, G. Wrightstone, and F. B. Soepyan, *Human Emissions and Atmospheric Concentrations of Carbon Dioxide*, CO₂ Coalition, (2026). <https://co2coalition.org/wp-content/uploads/2026/08/Emission-Concentration-2026-08-21-1.pdf>
- [13] Newton's law of cooling. <https://www.quadco.engineering/en/know-how/newtons-law-of-cooling.htm>
- [14] Isotope fractionation. https://en.wikipedia.org/wiki/Isotope_fractionation
- [15] Pee Dee Belemnite. <https://pubmed.ncbi.nlm.nih.gov/35312289/>
- [16] P. J. H. Dunn et al, *Redetermination of $R(^{13}\text{C}/^{12}\text{C})$ for Vienna Peedee Belemnite*, Rapid Comm. in Mass Spectrometry, John Wiley, (2024). <https://analyticalsciencejournals.onlinelibrary.wiley.com/doi/10.1002/rcm.9773>
- [17] The Suess Effect. https://en.wikipedia.org/wiki/Suess_effect
- [18] P. Wang et al, *Stable carbon isotopic characteristics of fossil fuels in China*, Science of Total Environ., **805** (2022). <https://doi.org/10.1016/j.scitotenv.2021.150240>
- [19] H. Graven, R. E. Keeling and J. Rogelj, *Changes to Carbon Isotopes in Atmospheric CO₂ Over the Industrial Era and Into the Future*, Global Biochemical Cycles, **34**, (2019). <https://doi.org/10.1029/2019GB006170>.