GAMSA · Gender-Affirming Medicines South Australia

Testosterone injection log: short-acting

Testosterone enantate (Primoteston Depot) or Sustanon · gamsa.au/using

Key points Usually every 2 to 4 weeks. Write down each injection and where it went.

Use one row for each injection, whether you give it yourself or a clinic gives it. Bring this log to each review, so your prescriber can match your blood test to your last dose.

My prescriber

Clinic _____

Phone _____


My pharmacy

Name _____

Phone _____

Good to know

- **How often:** usually every 2 to 4 weeks, as your prescriber sets. The Primoteston leaflet gives every 2 to 3 weeks at first, then every 3 to 4 weeks. Ask your prescriber before you change the gap.
- **Injecting yourself:** many people learn to. Be taught in person first, and ask for a written plan: product, dose, route, site, how often, and who to call.
- **Into the muscle at home:** the thigh is the usual site. Inject under the skin (belly) only if your prescriber has told you to. Use the next spot each time.
- **Blood test:** midway between injections, or as your prescriber asks.
- **Not for the long-acting injection** (Reandron 1000 and other brands). Use the long-acting log for that one.



Front

Spots 5 and 6 (thigh): the usual spots for an injection into the muscle at home. Spots 1 to 4 (belly): only for injections under the skin, if your prescriber has told you to. Use the sites the person who trains you shows you. Numbers match gamsa.au/using.

Get the site looked at **the same day** if redness spreads or streaks away from it, the pain gets worse, or there is pus, new blisters or a fever.

Based on gamsa.au: Using your medicine, Medicines, doses and formulations and Blood tests and monitoring.

Prepared by Theodore South, BPharm (Hons), registered pharmacist
gamsa.au · Questions or corrections: gamsaenquiries@gmail.com
A record for you to keep. Background information, not a prescription.
© 2026 GAMSA. All rights reserved.

Version 1.1 · 7 November 2026
Current version:
gamsa.au/pharmacy

GAMSA · Gender-Affirming Medicines South Australia

Testosterone injection log: short-acting

Print this page again when it is full. Page 2 of 2.

Date	Medicine and dose e.g. Primoteston 250 mg	Site map number	Given by me, or clinic	Notes

Notes: soreness, redness or a lump at the site; a late or missed dose; when the next one is due.

Based on gamsa.au: Using your medicine, Medicines, doses and formulations and Blood tests and monitoring.

Prepared by Theodore South, BPharm (Hons), registered pharmacist
gamsa.au · Questions or corrections: gamsaenquiries@gmail.com
A record for you to keep. Background information, not a prescription.
© 2026 GAMSA. All rights reserved.

Version 1.1 · 7 November 2026
Current version:
gamsa.au/pharmacy