GAMSA · Gender-Affirming Medicines South Australia

After your long-acting testosterone injection

Testosterone undecanoate 1000 mg/4 mL (Reandron 1000 and other brands), given into the side of the hip

Key points Stay at the clinic for at least 30 minutes. After you leave, ring 000 if you have trouble breathing or feel faint.

My clinic

Name _____

Phone _____

My next injection

Date _____

Time _____

Before you leave the clinic

- Stay for at least 30 minutes after the injection. Staff watch for a rare reaction to the oil, called POME. It usually starts within 30 minutes.
- Tell staff straight away about a sudden cough or urge to cough, breathlessness, chest tightness, dizziness or feeling faint.

In the next few days

- The site can ache for a day or two. This is normal.
- Walking about helps. A warm pack and simple pain relief are fine if your pharmacist says they suit you.
- Do not rub or massage the site.

Your next injection

- Book it before you leave the clinic. It is usually about 6 weeks after your first dose, then every 10 to 14 weeks.
- If you cannot make it, ring the clinic as soon as you know. Going past 14 weeks lets your testosterone level fall.

Ring 000 straight away if you

- have trouble breathing, or a lot of coughing
- have chest pain or tightness
- feel dizzy or faint, or collapse
- have swelling of the face, lips, tongue or throat.

Then tell your clinic.

Contact your clinic or doctor if

- the site is red, hot, swollen or more painful after the first few days, or leaks fluid
- you have a fever
- a lump at the site grows or does not go away
- you are not sure what to do. You can also ring 1800MEDICARE on 1800 633 422 to talk to a nurse, any time.

Report a side effect

- Tell your clinic. You or the clinic can also report it to the TGA at aems.tga.gov.au.

Need an interpreter? Ring TIS National on 131 450.

Sources: Reandron 1000 Australian Product Information and Consumer Medicine Information (Aspen, revised December 2025); ASCIA acute management of anaphylaxis (October 2026); GAMSA testosterone undecanoate SOP and training guide, version 2.6. Plain-language background: gamsa.au/using.

Document GAMSA-TU-AFTER · Next review 24 October 2027 · Scan the QR for the current version.

© 2026 GAMSA. All rights reserved.

Prepared by Theodore South, BPharm (Hons), registered pharmacist
gamsa.au · Questions or corrections: gamsaenquiries@gmail.com

General information to keep. Not a substitute for advice from your clinic.

Version 1.0 · 7 November 2026
Current version: gamsa.au/medicines/testosterone-undecanoate