

Adverse event reporting and escalation form — ventrogluteal intramuscular administration of testosterone undecanoate 1000 mg/4 mL

Version: 1.1 **Effective date:** 5 October 2025 **Revised:** 2 October 2026 **Next review:** October 2026, or on any change to the Product Information

Prepared by Theodore South, BPharm (Hons), registered pharmacist, Adelaide. Clinics remain responsible for their own procedures, credentialling and emergency arrangements. This document does not replace the Product Information [1], which prevails where they differ.

Complete immediately after any suspected adverse reaction to a testosterone undecanoate 1000 mg/4 mL injection, including pulmonary oil microembolism (POME), anaphylaxis or any other unexpected event [1][6][7][8].

Patient and event details

Patient name: _____ Date of birth: _____

Date and time of event: _____ Administering clinician: _____

Injection site: ☐ Left ventrogluteal ☐ Right ventrogluteal Needle: _____ G, _____ mm

Product batch number: _____ Expiry: _____

Time injection started: _____ Time injection finished: _____ Onset of symptoms: _____

Description of the event (symptoms, onset, duration, observations):

Immediate management and treatment given (including oxygen, adrenaline dose and time, positioning):

Escalation and notifications

1. Notify the supervising clinician immediately.
2. If severe — anaphylaxis, collapse, acute respiratory symptoms — call 000 [8].
3. Inform the prescriber and the patient's GP.
4. Notify the clinic manager and complete the internal incident report.
5. Report the event to the Therapeutic Goods Administration (TGA) at tga.gov.au/reporting-problems [9].
6. Document all actions and communication.

Local escalation contacts

Clinical manager: _____ Phone: _____

Prescriber / GP: _____ Phone: _____

Nearest emergency department: _____ Phone: _____

Follow-up

- ☐ Patient reviewed after the event; outcome recorded
- ☐ Adverse event documentation completed in the clinical record
- ☐ Report submitted to the TGA; reference number: _____
- ☐ Prescriber informed of the outcome and of any decision about future injections

Clinician signature: _____ Date: _____

References

- [1] Aspen Pharmacare Australia. Reandron 1000 (testosterone undecanoate 1000 mg/4 mL) solution for injection — Australian Product Information and Consumer Medicine Information. Accessed via the Medicine Finder, Australian Commission on Safety and Quality in Health Care, 1 October 2026. <https://www.safetyandquality.gov.au/medicine-finder/reandron-1000-solution-for-injection>
- [2] Therapeutic Goods Administration. REANDRON 1000 testosterone undecanoate 1000 mg/4 mL injection vial — ARTG 213942. Canberra: TGA. <https://www.tga.gov.au/resources/artg/213942>
- [3] Bayer New Zealand. Reandron 1000 (1000 mg/4 mL solution for injection) — New Zealand Data Sheet. Wellington: Medsafe. <https://medsafe.govt.nz/profs/Datasheet/r/Reandron1000inj.pdf>
- [4] SA Health. Changes to your long acting injection — patient brochure on the ventrogluteal site (includes the landmarking figure). Adelaide: Department for Health and Wellbeing, Government of South Australia, 2022. [https://www.sahealth.sa.gov.au/wps/wcm/connect/e185e690-bc60-402f-b264-5ed189fd0390/Long+Acting+Intramuscular+Injections+Brochure+\(1\).pdf?MOD=AJPERES](https://www.sahealth.sa.gov.au/wps/wcm/connect/e185e690-bc60-402f-b264-5ed189fd0390/Long+Acting+Intramuscular+Injections+Brochure+(1).pdf?MOD=AJPERES)
- [5] South Eastern Sydney Local Health District. Intramuscular injection in mental health — guideline SESLHDGL/076, version 2.6, 20 January 2025. Sydney: NSW Health. <https://www.seslhd.health.nsw.gov.au/sites/default/files/documents/SESLHDGL%20076%20-%20Intramuscular%20Injection%20in%20Mental%20Health.pdf>
- [6] Pastuszak AW, Hu Y, Freid JD. Occurrence of pulmonary oil microembolism after testosterone undecanoate injection: a postmarketing safety analysis. *Sex Med.* 2020;8(2):237–242. doi:10.1016/j.esxm.2020.01.009 <https://doi.org/10.1016/j.esxm.2020.01.009>

- [7] Adam JA, Pastuszak AW, Christensen MB, Spencer R, Sandberg A, Hotaling JM, Lipshultz LI. Occurrence of pulmonary oil microembolism (POME) with intramuscular testosterone undecanoate injection: literature review. *Int J Impot Res*. 2023;35(5):439–446. doi:10.1038/s41443-022-00585-1 <https://doi.org/10.1038/s41443-022-00585-1>
- [8] Australasian Society of Clinical Immunology and Allergy. ASCIA guidelines: acute management of anaphylaxis. July 2026.
https://www.allergy.org.au/images/ASCIA_HP_Guidelines_Acute_Management_Anaphylaxis_July_2026.pdf
- [9] Therapeutic Goods Administration. Reporting adverse events — medicines and vaccines. Canberra: TGA.
<https://www.tga.gov.au/reporting-problems>
- [10] Laboratorios Liconsa. Guidance on the administration of testosterone undecanoate 1000 mg/4 mL — risk minimisation material approved by the MHRA, version 2.0, July 2024. (Same formulation as Reandron 1000.) <https://www.medicines.org.uk/emc/rmm/100105/Document>