

Chapter 1: Geometrical Optics

Daniel Rice and Thomas Lockwood

September 14, 2014

Abstract

Hello this is our abstract.

1.1 The Stenopaic Spectacles

In this experiment we looked out of a lens with pinholes. We were able to observe, after looking into the glasses, a slight dim in brightness and increase of depth perception. It was easy to see small print writing. I could see the writing on my journal in a zoomed in way. The writing was tiny before the glasses, however after the width of the writing's lines was noticeably bigger. A hypothesis of why this happens is because the pinholes in the glasses make it easier for our eyes to focus on objects that are as equally small.

1.2 The Laws of Reflection and Refraction

Here we will put our graphs and calculate things.

1.3.2 Apparent Thickness of a Block of Lucite

Formula for S and S' :

$$S = P_3 - P_1, \quad S' = P_3 - P_2$$

Our results gives us

$$P_1 = 1.4mm \pm .1mm, \quad P_2 = 6.4mm \pm .1mm, \quad P_3 = 18.9mm \pm .1mm$$

$$S = 18.9 - 1.4 = 17.5mm, \quad S' = 18.9 - 6.4 = 12.5mm$$

Forumula for Index of Refraction:

$$n = \frac{S}{S'} = \frac{17.5}{12.5} = 1.4mm$$

1.5 The Prism Spectrometer

Ask our profess/classmates to calculate the minimum deviation of the yellow line.

1.6.1 Power of a Thin Prism

Formula for Power:

$$P = 100 \frac{x}{b}$$

Our results gives us

$$x = 1cm \pm .1cm, \quad b = 39.2cm \pm .1cm, \quad P = 100(1)/(39.2) = 2.55^\Delta$$

Formula for Apex:

$$\delta = (n - 1)\alpha$$

Our results gives us

$$n = 1.5, \quad \alpha = .57^\circ, \quad \delta = (1 - 1.5).57 = .285^\circ$$

1.6.2 The Risley Prism

Forumula for Lateral Displacement:

$$x = P \frac{b}{100}$$

$$P = 6^\Delta, \quad b = 44.8cm, \quad P = 100(44.8)/100 = 2.688cm$$

Our calculations were $-0.102cm$ off from the manual's expected calculated value.

1.7.1 Internal Reflection in a Bar

In this experiment we viewed the path of a laser through two different plastics bars. One was a straight rectangular bar, while the other was curved. As the laser traveled through the straight rectangular bar the laser reflected internally several times at approximately the same angle for each reflection before exiting. In the curved bar the laser also reflected, but the angle of incidence changed to follow the path of the curve before escaping from the end.

1.7.2 Internal Reflection in a Coil

In this experiment we viewed the laser as it traveled through a spiral coil. The coil seemed to illuminate as the laser entered through one end. The reason could be that the angle of the laser was reflected off of the spiral shape. It was more difficult to see the laser inside the plastic, however just like the other clear plastic objects it exited through the opposite end.

1.7.3 A Bundle of Fibers

Unfortunately, we were unable to gather fibers for this portion of the lab. Omit this section.

1.7.4 Coherent Fiber Optics

In this experiment we used a tool created of jacketed fiber bundles to view a well illuminated word in the lab manual. By placing the objective lens of the fiber close to the word, the light was reflected through the fiber optics cable to the eyepiece allowing us to view the image. The image was still visible despite the curved nature of the tool. This occurs due to the fiber jackets having different lower index material to prevent cross-feed of light that would distort the material.

1.7.5 An Illuminator

In this experiment we did not attempt to bend the light guide into an arc of radius of curvature less than about 2 inches. Instead we observed the commercial illuminator that had a white-light source. The light that came from the illuminator reflected off of the fibers through the curve of the object and exited through the end.