

# EQUAL CONSECUTIVE DIVISOR FUNCTIONS

THOMAS LOCKWOOD

## 1. THE INTEGERS

Are there infinitely many equal consecutive Divisor Functions? This problem is also known as the Erdos-Sierpinski problem. The Divisor Function,  $\sigma(n)$ , is the sum of all the positive divisors of an integer  $n$  with the prime factorization  $n = (p_1^{m_1})(p_2^{m_2})...(p_N^{m_N})$

$$\sigma(n) = \sum_{d|n} d = \left(\frac{p_1^{m_1+1} - 1}{p_1 - 1}\right)\left(\frac{p_2^{m_2+1} - 1}{p_2 - 1}\right)...\left(\frac{p_N^{m_N+1} - 1}{p_N - 1}\right)$$

For example,

$$\sigma(14) = 1 + 2 + 7 + 14 = 24$$

We want the solutions to  $\sigma(n) = \sigma(n + 1)$ . Some solutions (up to 10,000):

$$\begin{aligned}\sigma(206) &= \sigma(207) = 312 \\ \sigma(957) &= \sigma(958) = 1440 \\ \sigma(1334) &= \sigma(1335) = 2160 \\ \sigma(1364) &= \sigma(1365) = 2688 \\ \sigma(1634) &= \sigma(1635) = 2688 \\ \sigma(2685) &= \sigma(2686) = 4320 \\ \sigma(2974) &= \sigma(2975) = 4464 \\ \sigma(4364) &= \sigma(4365) = 7644\end{aligned}$$

Guy and Shanks found a pattern in the form

$$n = 2p, \quad n + 1 = 3^m q$$

when  $q$  and  $p$  are both prime for

$$q = 3^{m+1} - 4, \quad p = \frac{3^m q - 1}{2}$$

Solutions for  $m = 1, 2, 4$  yields the results  $n = 14, 206, 19358$  And another in the form

$$n = 3^m q, \quad n + 1 = 2p$$

when  $q$  and  $p$  are both prime for

$$q = 3^{m+1} - 10, \quad p = \frac{3^m q + 1}{2}$$

solutions when  $m = 4$  and  $5$ . However this does not solve the problem since it has only produced 3 and 2 solutions respectively.

## 2. THE GAUSSIAN INTEGERS

Gaussian Integers (complex integers), also known as the Integer Ring  $\mathbb{Z}[\sqrt{-1}]$ , can also be defined for the Divisor Function. Spira defined function the Divisor Function for Gaussian Integers very well, denoted  $g\sigma$ . In his definition he demonstrated that we need to use the first quadrant of the form of the complex numbers, excluding the imaginary axis.

Furthermore, using Guy and Shanks method with Gaussian Integers we have  $g\sigma$  each in the form of two distinct primes for both  $n$  and  $n + 1$ . Which gives us

$$g\sigma(n) = g\sigma(n + 1)$$

and let

$$n = (a + bi)(c + di), \quad n + 1 = (e + fi)(g + hi)$$

so

$$g\sigma(n) = (ac - bd + 1) + i(cb + da), \quad g\sigma(n + 1) = (eg - fh + 1) + i(fg + eh)$$

and we have the four functions

$$ac - bd = eg - fh + 1$$

$$bc + ad = fg + eh$$

$$ac - bd + 1 = eg - fh + 1$$

$$cb + da = fg + eh$$

Therefore our solutions are:

$$c = \frac{ea^2 - 2e^2a - ea - a + eb^2 - bf - 2ebf + ef^2 + f^2 + e^3 + e^2 + e}{a - e^2 + b - f^2}$$

$$d = \frac{a^2b - 2eab - af + b^3 - 2b^2f + bf^2 + e^2b + eb + b - f}{a - e^2 + b - f^2} - b + f$$

$$g = \frac{a^3 - 2ea^2 - a^2 + ab^2 - 2abf + af^2 + e^2a + ea - a - b^2 + bf + e}{a - e^2 + b - f^2}$$

$$h = \frac{a^2b - 2eab - af + b^3 - 2b^2f + e^2b + eb + b - f}{a - e^2 + b - f^2}$$

Some of the solutions for consecutive for  $g\sigma(n) = g\sigma(n+1)$  case

$$g\sigma(3+7i) = 10+10i = g\sigma(4+7i)$$

$$g\sigma(19+25i) = -20+60i = g\sigma(20+25i)$$

$$g\sigma(19+75i) = -100+100i = g\sigma(20+75i)$$

$$g\sigma(40+85i) = -100+140i = g\sigma(41+85i)$$

$$g\sigma(90+245i) = -320+480i = g\sigma(91+245i)$$

and for  $g\sigma(n) = g\sigma(n+i)$  case

$$g\sigma(15+9i) = 40i = g\sigma(15+10i)$$

$$g\sigma(25+39i) = -40+80i = g\sigma(25+40i)$$

$$g\sigma(25+90i) = -120+120i = g\sigma(25+91i)$$

$$g\sigma(65+60i) = -40+160i = g\sigma(65+61i)$$

Lastly for  $g\sigma(n) = g\sigma(n+1+i)$  case

$$g\sigma(8+16i) = 10+10i = g\sigma(3+12i)$$

$$g\sigma(5+5i) = 20i = g\sigma(6+6i)$$

$$g\sigma(13+4i) = 20i = g\sigma(14+5i)$$

$$g\sigma(84+5i) = 8+96i = g\sigma(85+6i)$$

$$g\sigma(90+6i) = -160+160i = g\sigma(91+7i))$$

$$g\sigma(138+102i) = -400+80i = g\sigma(139+103i)$$

$$g\sigma(149+53i) = -340+180i = g\sigma(150+54i)$$

## 3. EQUAL SIGMA FUNCTIONS

For  $g\sigma(n) = g\sigma(n+k)$  can also be solved using the previous formula. It is easy to show using the previous equations.

$$g\sigma(n) = g\sigma(n+k)$$

and let

$$n = (a + bi)(c + di), \quad n + k = (e + fi)(g + hi)$$

so

$$g\sigma(n) = (ac - bd + 1) + i(cb + da), \quad g\sigma(n + k + 1) = (eg - fh + 1) + i(fg + eh)$$

and we have the four functions

$$ac - bd = eg - fh + 1$$

$$bc + ad = fg + eh$$

$$ac - bd + 1 = eg - fh + 1$$

$$cb + da = fg + eh$$