

Welcome to the start of your EBGLYSS journey

A guide to your atopic dermatitis (also known as atopic eczema) treatment

Your doctor has prescribed EBGLYSS to help you better manage your atopic dermatitis. This booklet is designed to give you information about EBGLYSS, including how it works, how it is used and common side effects that you may experience during treatment. It should not replace the advice given by your doctor. If you have any questions or concerns, please contact your doctor, who should always be your primary source of information about atopic dermatitis.

What is EBGLYSS?

EBGLYSS is a biologic medication used to treat adults and adolescents aged 12 years and over (weighing over 40 kg) with moderate-to-severe atopic dermatitis. It contains the active ingredient lebrikizumab.

How does EBGLYSS work?

EBGLYSS belongs to a group of medicines called monoclonal antibodies, which are a type of protein that block a specific protein called interleukin-13. Interleukin-13 contributes to inflammation and plays a role in atopic dermatitis.

How is EBGLYSS administered?


EBGLYSS comes as a pre-filled pen and is given by injection under the skin. It is important that you are trained by your doctor, nurse or pharmacist before injecting yourself. A caregiver or family member may also administer your EBGLYSS injection after proper training. You can also refer to the 'Instructions for Use' that come with your EBGLYSS pre-filled pen. Please read them carefully before using EBGLYSS.



Pen image not to scale.

How often do I administer EBGLYSS?

When you are first prescribed EBGLYSS, you will receive two injections as a starting dose at Week 0 and Week 2. Following this, you will administer one injection of EBGLYSS every two weeks up to Week 16. After 16 weeks, you will administer one injection of EBGLYSS every 4 weeks. It is important to inject the whole contents of the single-use pre-filled pen each time you administer your EBGLYSS injection.



*Loading dose of 500 mg, administered as two 250 mg injections, at Weeks 0 and 2. †After completing the starting dose, and if you have had an adequate response, your doctor may prescribe the maintenance dose. If you did not have an adequate response to the starting dose, your doctor may recommend that you either: continue with one injection every 2 weeks until Week 24 and then decide if you can continue with the maintenance dose OR stop taking EBGLYSS.

What should I expect from EBGLYSS?

When you start taking EBGLYSS, you might not see immediate improvement in your signs and symptoms. Some atopic dermatitis treatments take time to show results, and this can vary from person to person.

What should I know before I use EBGLYSS?

Do not use EBGLYSS if:

- You are allergic to lebrikizumab.
- The product appears cloudy, discoloured or contains particles.
- The medicine has passed its expiration date.
- The medicine appears damaged.

Check with your doctor if you:

- Have a parasitic (helminth) infection.
- Have any other medical conditions, including new or worsening eye problems (e.g., eye pain, vision changes).
- Take or use any medicines, including topical or inhaled corticosteroid medicines.
- Have recently had, or are due to have, a vaccination. Check with your doctor to make sure that you are up to date with all vaccinations before starting EBGLYSS.
- Are pregnant or intend to become pregnant.
- Are breastfeeding or intend to breastfeed.

What should I know while using EBGLYSS?

Do:

- Keep the medicine in the original carton.
- Store the pre-filled pen in the refrigerator between 2°C to 8°C until you are ready to use it. If needed, it can be left out for up to 7 days at a temperature not above 30°C.
- Wait at least 45 minutes to let the pre-filled pen warm to room temperature before you use it.
- Make sure you place your used medicine into a sharp's container.
- Remind any doctor, dentist, or pharmacist you visit that you are using EBGLYSS.

Do not:

- Freeze the pre-filled pen. If the pre-filled pen has been frozen, do not use it.
- Expose the pre-filled pen to excessive heat greater than 30°C.
- Shake the pre-filled pen.
- Use the medicine if the pre-filled pen is damaged or you are not sure that it is working properly.
- Stop using this medicine suddenly without first checking with your doctor.
- Give EBGLYSS to anyone else, even if they have the same condition as you.

Are there any side effects?

As with all medicines, you may experience some unwanted side effects with EBGLYSS. Most of the time, these side effects are minor and temporary; however, there are some that may need medical attention.

Less serious side effects:

Eye-related:

- Eye infection
- Redness of eyes
- Eye dryness
- Itchy and watery eye(s)
- Eye pain
- Blurry vision

- Sensitivity to light
- Eye and eyelid inflammation (including redness, swelling and itching)

Injection site reactions:

- Redness, swelling or itching at the injection site

WHAT TO DO

Speak to your doctor if you have any of these less serious side effects and they worry you or are persistent and do not go away.

Serious side effects:

Hypersensitivity reactions:

- Breathing problems or wheezing
- Swelling of the face, lips, mouth, tongue, or throat
- Hives
- Itching
- Fainting, dizziness, feeling lightheaded
- Skin rash
- Cramps in your stomach area (abdomen)

Symptoms of shingles, such as:

- Headache
- Sensitivity to light
- Tingling
- Itching or painful skin rash with blisters

Blood related:

- Increase in white blood cell count, which may be found during blood tests

WHAT TO DO

Tell your doctor if you have any of these serious side effects as they may require medical attention.



Eli Lilly is committed to environmental sustainability and has decided to print on environmentally responsible paper that is carbon neutral, FSC Certified Recycled, and manufactured from 100% post-consumer recycled fiber. It is processed in a chlorine-free environment under the ISO 14001 environmental management system.

This booklet is for patients prescribed EBGLYSS for the treatment of atopic dermatitis. If you have any questions or concerns, please contact your doctor. This booklet should not replace the advice of your doctor. Before use, please review the Consumer Medicine Information leaflet. This leaflet can be found packaged with your device, or by scanning the QR code.



Ebglyss® is a registered trademark of Eli Lilly and Company.
©2024 Copyright Eli Lilly and Company. Eli Lilly Australia Pty Limited. ABN 39 000 233 992.
Telephone: 1800 454 599. Level 9, 60 Margaret Street, Sydney, NSW 2000.
Date of preparation: November 2024. PP-LK-AU-0032.

Lilly A MEDICINE COMPANY