

GeeksforGeeks

A computer science portal for geeks

Placements

Practice

GATE CS

IDE

Q&A

GeeksQuiz

Login/Register

Divide and Conquer | Set 6 (Tiling Problem)

Given a n by n board where n is of form 2^k where $k \geq 1$ (Basically n is a power of 2 with minimum value as 2). The board has one missing cell (of size 1×1). Fill the board using L shaped tiles. A L shaped tile is a 2×2 square with one cell of size 1×1 missing.




Figure 1: An example input

This problem can be solved using Divide and Conquer. Below is the recursive algorithm.

```
// n is size of given square, p is location of missing cell
```

```
Tile(int n, Point p)
```

- 1) Base case: $n = 2$, A 2×2 square with one cell missing is nothing but a tile and can be filled with a single tile.
- 2) Place a L shaped tile at the center such that it does not cover the $n/2 \times n/2$ subsquare that has a missing square. **Now all four subsquares of size $n/2 \times n/2$ have a missing cell** (a cell that doesn't need to be filled). See figure 2 below.
- 3) Solve the problem recursively for following four. Let p_1 , p_2 , p_3 and p_4 be positions of the 4 missing cells in 4 squares.
 - a) `Tile($n/2$, p_1)`
 - b) `Tile($n/2$, p_2)`
 - c) `Tile($n/2$, p_3)`
 - d) `Tile($n/2$, p_4)`

The below diagrams show working of above algorithm




Figure 2: After placing first tile




Figure 3: Recurring for first subsquare.




Figure 4: Shows first step in all four subsquares.

Time Complexity:

Recurrence relation for above recursive algorithm can be written as below. C is a constant.

$$T(n) = 4T(n/2) + C$$

The above recursion can be solved using **Master Method** and time complexity is $O(n^2)$

How does this work?

The working of Divide and Conquer algorithm can be proved using Mathematical Induction. Let the input square be of size $2^k \times 2^k$ where $k \geq 1$.

Base Case: We know that the problem can be solved for $k = 1$. We have a 2×2 square with one cell missing.

Induction Hypothesis: Let the problem can be solved for $k-1$.

Now we need to prove to prove that the problem can be solved for k if it can be solved for $k-1$. For k , we put a L shaped tile in middle and we have four subsquares with dimension $2^{k-1} \times 2^{k-1}$ as shown in figure 2 above. So if we can solve 4 subsquares, we can solve the complete square.

References:

<http://www.comp.nus.edu.sg/~sanjay/cs3230/dandc.pdf>

This article is contributed by **Abhay Rathi**. Please write comments if you find anything incorrect, or you want to share more information about the topic discussed above.

Company Wise Coding Practice Topic Wise Coding Practice

10 Comments Category: [Divide and Conquer](#) Tags: [Divide and Conquer](#)