



MCMII-IIITM



Hand-Scoring User's Guide

Dept. Clinical Psychology
Man College of Special Education
and Psychological Studies
Guna (M.P.)

MILLONTM CLINICAL MULTIAXIAL INVENTORY-III

Theodore Millon, PhD, DSc
Carrie Millon, PhD, Roger Davis, PhD, and Seth Grossman, PsyD



MCMI-III™

Hand-Scoring User's Guide

To hand score the Millon™ Clinical Multiaxial Inventory-III (MCMI-III), you will need the following materials:

- A completed MCMI-III answer sheet
- A set of plastic overlay scoring templates ("keys")
- A blank MCMI-III Hand-Scoring Worksheet
- A blank MCMI-III Hand-Scoring Profile
- The MCMI-III manual
- A calculator
- Scratch paper

Assemble these materials and then follow the instructions in this guide. It will take you approximately 40–60 minutes to hand score the MCMI-III. The major steps that you will complete are:

- Perform validity checks.
- Use the scoring templates to determine raw scale scores and record them on the worksheet.
- Look up initial BR scores and record them on the worksheet.
- Calculate and apply adjustments to BR scores and record them on the worksheet.
- Plot final BR scores on the profile.

1. Age

This is a check to ensure that the respondent's age is indicated and is in the appropriate range.

1-1: Determine if age is indicated and is 18 or greater:

- if less than 18 or not indicated, stop. This test is invalid and cannot be scored.
- if greater than or equal to 18, continue with section 2, "Omits/Double Marks."

2. Omits/Double Marks

This is a check to ensure that a sufficient number of questions were answered to provide a valid set of scores.

2-1: Count the number of omitted and double-marked responses on the answer sheet.

- if greater than 11, stop. The test results are invalid.
- if less than or equal to 11, cross out all of the double-marked responses. These responses will not be used in any score calculations. Continue with section 3, "Scale V (Invalidity) Score."

3. Scale V (Invalidity) Score

This is a check of the Invalidity score. If the score on this scale is too high, the respondent's results are invalid.

- 3-1: Count the number of True responses to Items 65, 110, and 157 on the answer sheet. The total is the Invalidity score.
- 3-2: The Invalidity score must be 0, 1, 2, or 3. Enter the score on side two of the worksheet in the upper left-hand corner in the box labeled "V Invalidity."
- 3-3: Check the Invalidity score:
- if the **Invalidity score is 2 or 3**, the test is considered invalid, and further scoring and/or interpretation is not recommended.
 - if the **Invalidity score is 1**, write "Questionable Validity based on Scale V score" at the top of the Hand-Scoring Profile and proceed to section 4, "Scale W (Inconsistency) Score."
 - if the **Invalidity score is 0**, proceed to section 4, "Scale W (Inconsistency) Score."

4. Scale W (Inconsistency) Score

This is a check of the Inconsistency score. If the score on this scale is too high, the respondent's results are invalid.

- 4-1: The Inconsistency scale consists of 44 pairs of item responses. Each blackened pair of responses adds 1 point to the scale score. The pairs are scored using the set of five plastic overlay keys for Scale W. Follow the instructions on each key to determine the score for that key, and record each score in the "Scale W Score Calculation" section on side one of the worksheet. Then sum the scores for the five keys to determine the Scale W score.
- 4-2: Record the sum on the Sum line and on side two of the worksheet in the box labeled "W Inconsistency" in the upper left-hand corner.
- 4-3: Check the Inconsistency score:
- if the **Inconsistency score is 10 or greater**, the test is considered invalid, and further scoring and/or interpretation is not recommended.

- if the **Inconsistency score is 8 or 9**, write "Questionable Validity based on Scale W score" at the top of the Hand-Scoring Profile and proceed to section 5, "Raw Scores."
- if the **Inconsistency score is less than 8**, proceed to section 5, "Raw Scores."

5. Raw Scores

In this section, raw scores are calculated for all scales (except V, W, and X) and recorded on the worksheet.

- 5-1: Find the plastic overlay key for Scale Y. The key is used to determine the total raw score based on the 1- and 2-point responses to items on the scale.
- 5-2: Place the key on the answer sheet so that the marks at the bottom of the key align with the corresponding marks at the bottom of the answer sheet. Begin scoring with the top right response circle and continue by following the lines that connect the response circles.
- 5-3: Record the numeric value (weight) of each **blackened** response circle for the scale on a separate sheet of paper. (A numeric value of 1 or 2 is printed next to each response circle on the key.) Add up the numeric values of the **blackened** response circles for the scale. The sum is the total raw score for that scale.
- 5-4: Enter this raw score in the box on the worksheet under "Raw Score" for the corresponding scale.

NOTE: Follow steps 5-1 through 5-4 for Scales Z and 1 through PP. **Be sure to use the correct key for each of the 26 scales.**

6. Scale X Raw Score

This section calculates the Scale X raw score, which is used to determine the Disclosure adjustment. It is based on the raw scores for Scales 1 through 8B.

- 6-1: Use the "Raw X Score Calculation" section on side one of the Hand-Scoring Worksheet to perform the calculations for the Scale X raw score. Record the results in the appropriate blanks.

- 6-2: Fill in the raw scores for Scales 1, 2A, 2B, 3, 4, 5, 6A, 6B, 7, 8A, and 8B in the appropriate blanks on the worksheet.
- 6-3: Multiply the raw score for Scale 5 by 2/3 and record the result on the appropriate line on the worksheet.
- 6-4: Add up the numbers in the column and write the total on the line labeled "Sum."
- 6-5: If the sum is a decimal number, round it to a whole number. Round down if the number after the decimal is less than 5 and up if it is greater than or equal to 5. Write this number on the line labeled "Raw X."
- 6-6: Enter the final value on side two of the worksheet in the Raw Score column in the box next to "X Disclosure."

7. Scale X Raw Score Check

This is a check of the raw score for Scale X. If it is too high or too low, the respondent's scores are invalid.

7-1: Check the Scale X raw score:

- if the Scale X raw score is less than 34 or greater than 178, the test is considered invalid and further scoring and/or interpretation is not recommended.
- if the Scale X raw score is between 34 and 178, proceed to section 8, "Initial Base Rate (BR) Scores."

8. Initial Base Rate (BR) Scores

In this section, initial base rate (BR) scores are determined for all scales except V and W.

- 8-1: Open the MCMI-III manual to the tables in Appendix C. The BR scores for all scales except Scale X are contained in Appendix C-1, and the BR scores for Scale X are contained in Appendix C-2. These two tables are used for both males and females.
- 8-2: Turn to the first page of Appendix C-1 and find the first column on the left. It is labeled "Raw Score" and contains a list of scores. Locate the raw score that matches the respondent's Scale 1 raw score. Then find the BR score in the Scale 1 column that

corresponds to that score. On side two of the worksheet, enter the Scale 1 BR score in the "1 Schizoid" row of the column labeled "BR From Table." Use the same procedure to determine the BR score for Scale X, but be sure to use Appendix C-2.

- 8-3: For the remaining scales (Y, Z, and 2A through PP), locate the scale's raw score in the leftmost column of the table. Determine the corresponding BR score by locating the column to the right with that scale name. Enter the BR score for the scale on the worksheet under the column labeled "BR From Table."

NOTE: Be sure to complete step 8-3 for all 27 scales (X, Y, Z, and 1-PP).

IMPORTANT: The instructions below are **cumulative**. Always use the **latest** adjusted BR scores.

9. Disclosure Adjustment

This section computes the Disclosure adjustment, which is used to adjust scores for the Clinical Personality Patterns scales (1, 2A, 2B, 3, 4, 5, 6A, 6B, 7, 8A, 8B), the Severe Personality Pathology scales (S, C, P), the Clinical Syndromes scales (A, H, N, D, B, T, R), and the Severe Clinical Syndromes scales (SS, CC, PP).

NOTE: This adjustment is based on the Scale X **RAW** score (**not** the BR score).

- 9-1: In Table 1 on the next page, locate the raw score range that contains the respondent's Scale X raw score.
- 9-2: Find the 1-8B and the S-PP adjustments that correspond to the respondent's raw score on Scale X.
- 9-3: Write these adjustment factors in the boxes labeled "Disclosure 1-8B Adjustment" and "Disclosure S-PP Adjustment" on the worksheet. These boxes are in the shaded area labeled "Adjustment Values" at the top of the worksheet.
- 9-4: Apply the 1-8B adjustment to the BR scores for Scales 1-8B. Apply the S-PP adjustment to the BR scores for Scales S-PP. After the

Table 1. Disclosure Adjustment Values

Scale X Raw Score	1-8B Adjustment	S-PP Adjustment	Scale X Raw Score	1-8B Adjustment	S-PP Adjustment
0-33*	+20	+10	138	-6	-4
34	+20	+10	139	-7	-4
35	+20	+10	140	-7	-4
36	+20	+10	141	-7	-4
37	+19	+10	142	-8	-5
38	+18	+10	143	-8	-5
39	+17	+9	144	-9	-5
40	+17	+9	145	-9	-5
41	+16	+9	146	-9	-5
42	+15	+8	147	-10	-6
43	+14	+8	148	-10	-6
44	+13	+7	149	-11	-6
45	+13	+7	150	-11	-6
46	+12	+7	151	-11	-6
47	+11	+6	152	-12	-7
48	+10	+6	153	-12	-7
49	+9	+5	154	-13	-7
50	+9	+5	155	-13	-7
51	+8	+5	156	-13	-7
52	+7	+4	157	-14	-8
53	+6	+4	158	-14	-8
54	+5	+3	159	-15	-8
55	+5	+3	160	-15	-8
56	+4	+3	161	-15	-8
57	+3	+2	162	-16	-9
58	+2	+2	163	-16	-9
59	+1	+1	164	-17	-9
60	+1	+1	165	-17	-9
61-123	None	None	166	-17	-9
124	-1	-1	167	-18	-10
125	-1	-1	168	-18	-10
126	-1	-1	169	-19	-10
127	-2	-2	170	-19	-10
128	-2	-2	171	-19	-10
129	-3	-2	172	-20	-11
130	-3	-2	173	-20	-11
131	-3	-2	174	-20	-11
132	-4	-3	175	-20	-11
133	-4	-3	176	-20	-11
134	-5	-3	177	-20	-11
135	-5	-3	178	-20	-11
136	-5	-3	179+	-20	-11
137	-6	-4			

*An individual's MCMI-III results are considered invalid if the raw score on Scale X is less than 34 or greater than 178. Scoring and/or interpretation of invalid results is not recommended.

adjustments are applied, change any BR values that are less than 0 to 0 and any BR values that are greater than 115 to 115.

Enter the adjusted BR score for each of these scales in the column of boxes labeled "Adjusted BR" under the "Disclosure 1-8B Adjustment" and "Disclosure S-PP Adjustment" boxes on the worksheet. (Note that boxes have been provided on the worksheet only for the scales that are affected by each of these adjustments.)

10. Anxiety/Depression (A/D) Adjustment

The Anxiety/Depression (A/D) adjustment is based on the respondent's Scale A (Anxiety) and Scale D (Dysthymia) scores. The A/D adjustment is applied to Clinical Personality Patterns Scales 2A, 2B, and 8B and to Severe Personality Pathology Scales S and C.

10-1: Compute the Anxiety/Depression value, which is used to select the appropriate A/D adjustment based on the BR scores for Scales A and D.

- if **A and D are both less than 75:**
No A/D adjustment is necessary. Proceed to section 11, "Inpatient Adjustment."
- if **A is greater than or equal to 75 and D is less than 75:**
Calculate $(A - 75) = A/D \text{ value}$
Proceed to step 10-2.
- if **D is greater than or equal to 75 and A is less than 75:**
Calculate $(D - 75) = A/D \text{ value}$
Proceed to step 10-2.
- if **A and D are both greater than or equal to 75:**
Calculate $[(A - 75) + (D - 75)] = A/D \text{ value}$
Proceed to step 10-2.

10-2: Using the result of the preceding calculation (the A/D value), find the A/D adjustments in Table 2, 3, or 4. Use Table 2 if the client is not an inpatient or if the client is an inpatient and the duration of his or her Axis I disorder is more than four weeks. Use Table 3 if the client is an inpatient and his or her Axis I disorder has been present for less than one week. Use Table 4 if the client is an inpatient and his or her Axis I disorder has been present for one to four weeks. Write the appropriate values

from the table in the boxes labeled "A/D 2B, 8B, C Adjustment" and "A/D 2A, S Adjustment" on the worksheet. These are the third and fourth boxes in the shaded area labeled "Adjustment Values" at the top of the worksheet.

10-3: Apply the 2B, 8B, C adjustment factor to the BR scores for Scales 2B, 8B, and C. Apply the 2A, S adjustment factor to the BR scores for Scales 2A and S. After the adjustments are applied, change any BR values that are less than 0 to 0 and any BR values that are greater than 115 to 115.

Enter the adjusted BR score for each of these scales in the column of boxes labeled "Adjusted BR" under the "A/D 2B, 8B, C Adjustment" and "A/D 2A, S Adjustment" boxes on the worksheet.

Table 2. Non-Inpatient, or Inpatient With Axis I Duration of More Than Four Weeks

2B, 8B, C		2A, S	
A/D Value	Adjustment	A/D Value	Adjustment
0	-1	0	-1
1	-1	1	-1
2	-1	2	-1
3	-1	3	-1
4	-1	4	-1
5	-1	5	-1
6	-1	6	-1
7	-1	7	-1
8	-1	8-15	-1
9	-1	16-23	-2
10	-2	24-31	-3
11	-2	32-39	-4
12	-2	40-47	-5
13	-2	48-55	-6
14	-2	56-63	-7
15-19	-3	64-71	-8
20-24	-4	72-79	-9
25-29	-5	80	-10
30-34	-6		
35-39	-7		
40-44	-8		
45-49	-9		
50-54	-10		
55-59	-11		
60-64	-12		
65-69	-13		
70-74	-14		
75-80	-15		

Table 3. Inpatient With Axis I Duration of Less Than One Week

2B, 8B, C		2A, S	
A/D Value	Adjustment	A/D Value	Adjustment
0	-1	0	-1
1	-1	1	-1
2	-1	2	-1
3	-1	3	-1
4	-1	4	-1
5	-2	5	-1
6	-2	6	-1
7	-2	7	-1
8	-3	8-11	-2
9	-3	12-15	-3
10-12	-4	16-19	-4
13-14	-5	20-23	-5
15-17	-6	24-27	-6
18-19	-7	28-31	-7
20-22	-8	32-35	-8
23-24	-9	36-39	-9
25-27	-10	40-43	-10
28-29	-11	44-47	-11
30-32	-12	48-51	-12
33-34	-13	52-55	-13
35-37	-14	56-59	-14
38-39	-15	60-63	-15
40-80	-15	64-80	-15

11. Inpatient Adjustment

The Inpatient adjustment is computed using the client's setting and Axis I duration. It is used to adjust the BR scores for the Severe Clinical Syndromes scales (SS, CC, PP).

11-1: Determine the setting and Axis I duration for the client.

11-2: Determine whether an adjustment should be applied.

If the client:

- is being evaluated using the correctional norms in the *MCMI-III Corrections Report User's Guide*, or
- is not an inpatient, or
- is an inpatient and his or her Axis I symptoms have been present for more than four weeks:

No Inpatient adjustment is necessary.

Proceed to section 12, "Denial/Complaint Adjustment."

Table 4. Inpatient With Axis I Duration of One to Four Weeks

2B, 8B, C		2A, S	
A/D Value	Adjustment	A/D Value	Adjustment
0-3	-1	0-5	-1
4-6	-1	6-10	-1
7-9	-2	11-15	-2
10-13	-3	16-21	-3
14-16	-4	22-26	-4
17-19	-5	27-31	-5
20-23	-6	32-37	-6
24-26	-7	38-42	-7
27-29	-8	43-47	-8
30-33	-9	48-53	-9
34-36	-10	54-58	-10
37-39	-11	59-63	-11
40-43	-12	64-69	-12
44-46	-13	70-74	-13
47-49	-14	75-79	-14
50-53	-15	80	-15
54-80	-15		

If the client:

- is an inpatient and his or her Axis I symptoms have been present less than one week or one to four weeks:
Proceed to step 11-3.

11-3: In Table 5, select the appropriate row to find the Inpatient adjustment.

11-4: Write the adjustment factors from Table 5 in the boxes on the worksheet labeled "SS Inpatient Adjustment," "CC Inpatient Adjustment," and "PP Inpatient Adjustment." These are the fifth, sixth, and seventh boxes in the shaded area labeled "Adjustment Values" at the top of the worksheet.

Apply the Inpatient adjustment to the BR scores for Scales SS, CC, and PP. After the adjustment is applied, change any BR values that are less than 0 to 0 and any BR values that are greater than 115 to 115.

Enter the adjusted BR score for each of these scales in the column of boxes labeled "Adjusted BR" under the "Inpatient Adjustment" boxes on the worksheet. (Note that boxes have been provided on the worksheet only for the three scales that are affected by these adjustments.)

Table 5. Inpatient/Axis I Duration Status

Inpatient/Axis I Duration Status	SS Adjustment	CC Adjustment	PP Adjustment
If setting = Inpatient and Axis I duration = Less than one week	+6	+10	+4
If setting = Inpatient and Axis I duration = One to four weeks	+4	+8	+2

12. Denial/Complaint Adjustment

The Denial/Complaint adjustment is computed using a rank ordering of the current BR scores for the Clinical Personality Patterns scales. It is used to adjust the latest BR scores for Scales 4, 5, and 7.

12-1: Determine which of the Clinical Personality Patterns scales (1, 2A, 2B, 3, 4, 5, 6A, 6B, 7, 8A, 8B) is highest based on their current **adjusted BR** scores. Any ties between scales should be broken by giving a higher ranking to the scale that appears first in the following list: 2B, 6A, 8B, 6B, 2A, 8A, 7, 4, 5, 3, 1.

12-2: The Denial/Complaint adjustment is applied as follows: if the highest ranked Clinical Personality Patterns scale is 4, 5, or 7, the adjustment value is 8. If the highest ranked Clinical Personality Patterns scale is anything else, no adjustment is made. If no adjustment is made, write "No" in the Denial/Complaint Adjustment box at the top of the worksheet and proceed to section 13, "Final BR Scale Scores."

Otherwise, write the adjustment factor (8) in the "Denial/Complaint Adjustment" box on the worksheet.

Add the 8-point adjustment factor from step 12-2 **only** to the highest ranking scale of Scales 4, 5, and 7. After the adjustment is applied, if the BR value is less than 0, change it to 0. If the BR value is greater than 115, change it to 115.

Enter the adjusted BR score for the scale in the appropriate box in the "Denial/Complaint Adjustment" column on the worksheet. Note that boxes have been provided on the worksheet only for the three scales that may be affected by this

adjustment. Copy the previous unchanged values into these boxes for the other two scales.

13. Final BR Scale Scores

The final BR scale scores for Scales X through Z and 1 through PP, which have been adjusted, are recorded on the worksheet.

13-1: Find the latest BR score for Scale X.

13-2: Transfer this final BR score for Scale X to the appropriate box in the column on the worksheet labeled "Final BR Score."

Repeat steps 13-1 and 13-2 for Scales Y, Z, and 1 through PP.

14. Score Profile

The final BR score for each scale is transferred to the Hand-Scoring Profile and plotted.

14-1: Check the final BR scores for the Clinical Personality Patterns scales (1-8B):

- if the **FINAL BR scores for Scales 1-8B are all less than 60**, stop. The result is a FLAT PROFILE, which is considered uninterpretable.
- if **at least one FINAL BR score for Scales 1-8B is greater than or equal to 60**, proceed to step 14-2.

14-2: Copy the final BR scores from the worksheet to the "BR Score" line at the bottom of the MCMI-III Hand-Scoring Profile.

Copy the Invalidity and Inconsistency scores from the worksheet to the profile.

14-3: Plot the BR score for each scale in the proper column of the profile by placing a mark at the appropriate score level. (If a BR score does not correspond exactly to any of the

numbers on the profile, approximate the location of that score between the next lowest and next highest numbers listed.)

NOTE: After you have completed the hand-scoring process, it is strongly recommended that you double-check every step.