

Worksheet 1: Understanding My Thoughts and Reactions

For: Pratham S | Age 16 | OPD No. 260703010 | Clinician: Mr. Tejas Dangodra

This worksheet helps us map out exactly what the unwanted thoughts are, what sets them off, and what you do when they show up. There are no right or wrong answers. We are just gathering information together so we can build a plan. Nothing here means you are a bad person - having a thought is not the same as wanting it or acting on it.

Safety first: If at any point you have thoughts of harming yourself, tell me, a parent, or use your safety plan / crisis contact right away. Filling this out can bring up distress - we do it together, in the room.

A. The unwanted (intrusive) thoughts

List the thoughts that keep coming even though you do not want them. Write them in your own words.

Example from our talk: "A thought that I might harm my father / a family member."

How often do they come? (circle) many times a day / a few times a day / most days / now and then

When are they strongest? e.g. at bedtime, when alone, when I see a knife/cutter/rope, after news/TV ...

B. What sets the thoughts off (triggers)

Triggers are situations, objects, places, or feelings that make the thought more likely. Rate how strong the trigger feels from 0 (no distress) to 100 (worst distress). Some are pre-filled from what we discussed - change or add anything.

Trigger (situation / object / feeling)	SUDS (0-100)	Notes
Seeing a knife or cutter		
Seeing a rope / near the fan		
Being alone in a room		
Being close to my father		
Violent scene on news / TV		
Trying to fall asleep		

C. What the thought makes me believe about myself

When the thought comes, what does it seem to say about you? (e.g. "it means I am dangerous", "it means I am a bad son", "it means I might lose control"). This is the part we will work on together.

How much do I believe that, 0-100%? _____

D. What I do to cope when the thought comes

These are the reactions we most want to find - even the ones nobody can see. They feel like they help, but they usually keep the thoughts going. Tick the ones you do and add your own.

Reaction I do when the thought hits	Do I do this? (Y/N)	How often
Try to push the thought away / block it		
Ask others (or myself) for reassurance that I won't act		
Avoid knives, cutters, rope, the kitchen		
Avoid being near family / being alone		
Hold myself tightly to stay in control		
Scratch myself when the thought is strong		
Mentally check that I haven't done anything		
Replace it with a 'safe' thought / count / pray		

Note for us: holding tight and scratching may be acting as safety behaviours. We will not just remove them - we will replace them with a safe coping skill before we drop them.

E. My OCD cycle (we fill this in together)

1. Trigger: _____

2. Intrusive thought: _____

3. Feeling + body (fear, sweating, breathing): _____

4. What I do (compulsion / avoidance): _____

5. Short-term relief, then: the thought comes back stronger

The goal of therapy is to break this cycle at step 4 - learning to let the thought be there without doing the reaction, so your brain learns the thought is not a real danger.