A LOWER BOUND FOR THE BOHR RADIUS OF THE BIDISC

TEJASVI SINGH TOMAR

Abstract. The Bohr radius K_2 of the bidisc is the largest r such that every holomorphic $f(z) = \sum c_\alpha z^\alpha$ on $\mathbb{D}^2$ with $|f| \le 1$ satisfies $\sum |c_\alpha| r^{|\alpha|} \le 1$, and its exact value is unknown. In an unrefereed solution posted on MathDB, Patel proved $K_2 > 0.30158758$. His argument combines Knese's bound in the higher degrees with three inequalities for the coefficients of degrees two and three, one of which comes from the unitary realization of Schur functions on the bidisc. We present his argument in full detail, confirm that it is correct, and check its exact computation. Knese's bound in degree n carries the factor $\sqrt{n}$, and this factor can be replaced by the Sidon constant of $\{0, 1, \dots, n\}$, which is smaller than $\sqrt{n}$ for every $n \ge 5$ by a theorem of Shapiro. We prove the computer-assisted bounds $S(\{0, 1, \dots, 5\}) \le 2.187$, $S(\{0, 1, \dots, 6\}) \le 2.363$ and $S(\{0, 1, \dots, 7\}) \le 2.552$; the first is consistent with the exact value found by Ye et al. With these bounds we obtain $K_2 > 0.301665$. The numerical steps use exact rational arithmetic and ball arithmetic. In the Lean proof assistant we also prove $K_2 \ge 0.301657$ from the first two Sidon bounds, assuming the inequality that comes from the unitary realization.

1. Introduction

For $d \ge 1$ let $\mathbb{D}^d = \{z \in \mathbb{C}^d : |z_1|, \dots, |z_d| < 1\}$ be the unit polydisc and let the Schur class $\mathcal{S}(\mathbb{D}^d)$ be the set of holomorphic maps $f \colon \mathbb{D}^d \to \overline{\mathbb{D}}$. For $f(z) = \sum_\alpha c_\alpha z^\alpha$ and $r > 0$ write $\|f_r\|_1 = \sum_\alpha |c_\alpha| r^{|\alpha|}$, where $|\alpha| = \alpha_1 + \cdots + \alpha_d$. The Bohr radius of $\mathbb{D}^d$ is

$$K_d = \sup\{r > 0 : \|f_r\|_1 \le 1 \text{ for all } f \in \mathcal{S}(\mathbb{D}^d)\}.$$

Bohr [9] proved that $K_1 \ge 1/6$. The sharp value $K_1 = 1/3$ was found independently by Riesz, Schur and Wiener, and Bohr's paper includes Wiener's proof (see [8]). Boas and Khavinson [8] introduced K_d for $d \ge 2$, proved $1/(3\sqrt{d}) < K_d < 2\sqrt{(\log d)/d}$ and asked for the exact value of K_d when $d > 1$. Defant, Frerick, Ortega-Cerdà, Ounaïes and Seip [11] showed that K_d has order $\sqrt{(\log d)/d}$, and Bayart, Pellegrino and Seoane-Sepúlveda [6] proved $K_d \sim \sqrt{(\log d)/d}$ as $d \to \infty$. No exact value of K_d is known for any $d \ge 2$.

For the bidisc, functions of one variable show $K_2 \le 1/3$. Knese [14, Corollary 1.2] proved $K_2 \ge 0.3006$: by his Theorem 1.1 and Andô's inequality, K_2 is at least the root $0.3006282\ldots$ of $\sum_{k \ge 1} \sqrt{k}\, r^k = 1/2$ (his corrigendum [15] corrects Theorem 1.4 of [14]). Baran, Pikul, Woerdeman and Wojtylak [5, Theorem 6.4] proved $K_2 < 0.3177$, and Griego [12] lowered this to $K_2 < 0.3174541$.

Patel [21] posted two bounds in an unrefereed solution to a problem on MathDB. For the upper bound he exhibits a rational inner function, and the post describes an exact integer computation that gives $K_2 < 0.302825279492$ and states its outcome. His reproducibility package [22] contains two programs that carry it out and a Lean proof. For the lower bound, a rotation and the substitution $H = (1-f)/(1-f(0))$ reduce the problem to the coefficients of holomorphic functions H on $\mathbb{D}^2$ with $\operatorname{Re} H > 0$ and $H(0) = 1$. Patel proves three inequalities for the coefficients of H of degrees two and three. The first two follow from the one-variable Carathéodory inequality on slices, the second with the help of a positive semidefinite Toeplitz form. The third

2020 *Mathematics Subject Classification.* 32A05; 30B10, 42A05, 47A48.

Key words and phrases. Bohr radius, bidisc, Carathéodory class, Sidon constant, transfer-function realization.

couples the two degrees and comes from the unitary realization of Schur functions on the bidisc [1, 4], which rests on Andô's inequality [3]. Using Knese's estimate [14, Lemma 2.1] in degrees $n \geq 4$, he then obtains $K_2 > 0.30158758$ by an exact rational computation.

A second route uses Sidon constants. For a finite set $\Lambda \subset \mathbb{Z}$ the Sidon constant $S(\Lambda)$ is the least C with

$$\sum_{\lambda \in \Lambda} |a_\lambda| \leq C \max_{|w|=1} \Big| \sum_{\lambda \in \Lambda} a_\lambda w^\lambda \Big| \qquad (a_\lambda \in \mathbb{C}).$$

Bernal-González, Cabana, García, Maestre, Muñoz-Fernández and Seoane-Sepúlveda [7, Theorem 3.3] showed that K_d is at least the root of $\sum_{k \geq 1} S(k,d)\, r^k = 1/2$, where $S(k,d)$ is the Sidon constant of the k-homogeneous polynomials in d variables. We write

$$S_k = S(\{0, 1, \ldots, k\}) \qquad (k \geq 1).$$

For $d = 2$, $S(k,2) = S_k$ (Remark 2.4). Newman's bound $S_k \leq \sqrt{k}$ [24, Chapter 3] (see also [20, (1)]) turns this equation into Knese's, and Shapiro [24] showed that $S_k = \sqrt{k}$ exactly when $k \in \{1, 2, 4\}$. Neuwirth conjectured $S_3 = 5/3$ [19] and later proved $S_3 \geq 5/3$ [20]. Yu et al. proved $S_3 \leq 5/3$, and hence $S_3 = 5/3$, with a formalized proof [26], and Ye et al. determined $S_5 = 2.186529771892569\ldots$ in a computer-assisted preprint [25]. With these values and $S_k \leq \sqrt{k}$ for $k \geq 6$, the equation of [7] gives $K_2 \geq 0.301339$ (Corollary 2.5). If the value of Ye et al. and Newman's bounds for S_6 and S_7 are replaced by the bounds of Proposition 4.2 for S_5, S_6 and S_7, the equation gives $K_2 \geq 0.301370$. Patel's coupling of degrees two and three goes beyond this.

In this paper we present Patel's argument with full details and improve its result. His post contains all the steps, and we have also checked his exact computation. Our improvement starts from the following observation. The slice argument behind Patel's inequalities also shows that the coefficients of degree n of H have absolute sum at most $2S_n$ (Lemma 2.3), which is the Carathéodory form of [7, Lemma 3.1]. Patel uses Knese's estimate $2\sqrt{n}$ in the degrees $n \geq 4$, and there $2S_n$ is smaller wherever $S_n < \sqrt{n}$. Since $S_4 = 2 = \sqrt{4}$, the first such degree is $n = 5$, and by Shapiro's theorem every larger degree qualifies. We prove $S_5 \leq 2.187$, $S_6 \leq 2.363$ and $S_7 \leq 2.552$ with computer-assisted certificates (Proposition 4.2). The exact value of S_5 found by Ye et al. [25] is consistent with the first bound and would serve equally well, but we use our own bound to keep the proof independent of that computation. With these bounds, an exact computation gives the following.

Theorem 1.1. $K_2 > 0.301665$.

The gain over Patel's bound, about $8 \cdot 10^{-5}$, comes from the terms of degrees five, six and seven. Together with Patel's upper bound, which is likewise unrefereed, Theorem 1.1 gives $0.301665 < K_2 < 0.302825279492$.

In Section 2 we set up the normalization and prove the slice estimate, including Carathéodory's inequality on slices. In Section 3 we write out Patel's treatment of the coefficients of degrees two and three, including the Gram argument behind the cubic inequality, the unitary extension and the identities of the realization. Section 4 treats the Sidon constants and proves the bounds for S_5, S_6 and S_7, Section 5 proves Theorem 1.1, and Section 6 lists the computer checks. Section 7 describes two formal proofs in Lean 4. Throughout, lower bounds are rounded down and upper bounds up.

2. The Carathéodory normalization and slices

Let $\mathcal{P}_2$ be the set of holomorphic functions H on $\mathbb{D}^2$ with $\operatorname{Re} H > 0$ and $H(0) = 1$. We write

$$H(z) = 1 + \sum_{n \geq 1} H_n(z), \qquad H_n(z) = \sum_{j+k=n} h_{jk} z_1^j z_2^k,$$

and set $\|H_n\|_1 = \sum_{j+k=n} |h_{jk}|$ and $M_r(H) = \sum_{n\geq 1} \|H_n\|_1 \, r^n$.

Patel [21] uses the following reduction.

Lemma 2.1. *Let $0 < r < 1$. If $M_r(H) \leq 1$ for every $H \in \mathcal{P}_2$, then $K_2 \geq r$.*

Proof. Constant functions satisfy $\|f_r\|_1 \leq 1$. Let $f \in \mathcal{S}(\mathbb{D}^2)$ be nonconstant. By the maximum principle $|f| < 1$ on $\mathbb{D}^2$. Multiplying f by a unimodular constant does not change $\|f_r\|_1$, so we may assume that $t = f(0)$ lies in $[0, 1)$. Then $H = (1 - f)/(1 - t)$ lies in $\mathcal{P}_2$. Its coefficients of positive degree are those of f multiplied by $-1/(1-t)$. Hence $\|f_r\|_1 = t + (1-t)M_r(H) \leq 1$. □

Remark 2.2. The normalization loses nothing: if $r < K_2$, then $M_r(H) \leq 1$ for every $H \in \mathcal{P}_2$. Indeed, for $0 < s < 1$ the function $z \mapsto H(sz)$ maps $\overline{\mathbb{D}}^2$ into a compact subset of the right half-plane, hence into a disc $\{w : |w - \rho| < \rho\}$ with $\rho \geq 1$. Then $f(z) = 1 - H(sz)/\rho$ lies in $\mathcal{S}(\mathbb{D}^2)$ with $f(0) = 1 - 1/\rho$, and $\|f_r\|_1 \leq 1$ becomes $\sum_n \|H_n\|_1 \, s^n r^n \leq 1$. Let $s \to 1$. Thus $K_2 = \sup\{r : M_r(H) \leq 1 \text{ for all } H \in \mathcal{P}_2\}$.

Lemma 2.3. *Let $H \in \mathcal{P}_2$ and $n \geq 1$. Then $|H_n(\zeta)| \leq 2$ for $\zeta \in \mathbb{T}^2$, and $\|H_n\|_1 \leq 2S_n$.*

Proof. Fix $\zeta \in \mathbb{T}^2$. The function $\varphi(\lambda) = H(\lambda\zeta) = 1 + \sum_{m\geq 1} H_m(\zeta)\lambda^m$ is holomorphic on $\mathbb{D}$ with $\operatorname{Re}\varphi > 0$. For $0 < \rho < 1$,

$$H_n(\zeta)\rho^n = \frac{1}{\pi} \int_0^{2\pi} \operatorname{Re}\varphi(\rho e^{i\theta}) \, e^{-in\theta} \, d\theta,$$

so $|H_n(\zeta)|\rho^n \leq 2 \operatorname{Re}\varphi(0) = 2$. Letting $\rho \to 1$, we obtain Carathéodory's inequality $|H_n(\zeta)| \leq 2$. Now write $H_n(\zeta_1, \zeta_2) = \zeta_2^n \, q_n(\zeta_1/\zeta_2)$ with $q_n(w) = \sum_{j=0}^n h_{j,n-j} w^j$. As ζ runs over $\mathbb{T}^2$, the quotient ζ_1/ζ_2 runs over $\mathbb{T}$ and $|H_n(\zeta)| = |q_n(\zeta_1/\zeta_2)|$. Hence $\max_{\mathbb{T}} |q_n| \leq 2$, and by the definition of S_n,

$$\|H_n\|_1 = \sum_{j=0}^n |h_{j,n-j}| \leq S_n \max_{\mathbb{T}} |q_n| \leq 2S_n. \qquad \square$$

Remark 2.4. The same identification shows that $S(k, 2) = S_k$: on $\mathbb{T}^2$ a k-homogeneous polynomial in two variables is, up to the factor ζ_2^k, a polynomial in ζ_1/ζ_2 of degree at most k. With this identification, the next corollary is the case $d = 2$ of [7, Theorem 3.3].

Corollary 2.5. *If $0 < r < 1$ and $\sum_{k\geq 1} S_k r^k \leq 1/2$, then $K_2 \geq r$.*

Proof. By Lemma 2.3, $M_r(H) \leq 2 \sum_{k\geq 1} S_k r^k \leq 1$ for every $H \in \mathcal{P}_2$. Apply Lemma 2.1. □

In [7] the corresponding estimate is proved for f itself. There Wiener's inequality takes the place of Carathéodory's: the Taylor coefficients $\hat{g}(k)$ of a holomorphic $g\colon \mathbb{D} \to \overline{\mathbb{D}}$ satisfy $|\hat{g}(k)| \leq 1 - |g(0)|^2$ for $k \geq 1$. Patel derives the bound $\|H_n\|_1 \leq 2\sqrt{n}$ for all $H \in \mathcal{P}_2$ from Knese's estimate [14, Lemma 2.1], by a limiting argument with $(1 - \varepsilon H)/(1 + \varepsilon H)$ [21]. Lemma 2.3 with Newman's bound gives it directly.

3. The coefficients of degrees two and three

Throughout this section $H \in \mathcal{P}_2$, and

$$A_2 = |h_{20}| + |h_{02}|, \quad B_2 = |h_{11}|, \qquad A_3 = |h_{30}| + |h_{03}|, \quad B_3 = |h_{21}| + |h_{12}|,$$

so that $\|H_2\|_1 = A_2 + B_2$ and $\|H_3\|_1 = A_3 + B_3$. The three inequalities and the way they are combined are due to Patel [21]. Inner products are linear in the first argument, $\langle x, y\rangle = y^* x$.

Lemma 3.1. $A_2^2 + B_2^2 \leq 4$.

Proof. Choose $\zeta_0 \in \mathbb{T}$ such that $h_{20}\zeta_0^2$ and h_{02} have the same argument, and put $a = h_{20}\zeta_0^2 + h_{02}$ and $b = h_{11}\zeta_0$. Then $|a| = A_2$, $|b| = B_2$ and $H_2(\pm\zeta_0, 1) = a \pm b$. By Lemma 2.3, $|a \pm b| \le 2$, and therefore $|a|^2 + |b|^2 = \frac{1}{2}(|a+b|^2 + |a-b|^2) \le 4$. □

Lemma 3.2. *Let R be a positive semidefinite Hermitian Toeplitz matrix of size $k+1$, and let $q(\zeta) = \sum_{j=0}^{k} c_j \zeta^j$. Then $c^* R c \le R_{00} \max_{\mathbb{T}} |q|^2$, where $c = (c_0, \dots, c_k)^{\mathsf{T}}$.*

Proof. Let $e_0, \dots, e_k$ be vectors in $\mathbb{C}^{k+1}$ with $\langle e_l, e_j \rangle = R_{jl}$, for example the columns of $R^{1/2}$. Since R is Toeplitz, the tuples $(e_0, \dots, e_{k-1})$ and $(e_1, \dots, e_k)$ have the same Gram matrix. Hence $e_j \mapsto e_{j+1}$ $(0 \le j < k)$ extends to a well-defined linear isometry from the span of $e_0, \dots, e_{k-1}$ onto the span of $e_1, \dots, e_k$, and then to a unitary operator Ω on the span of all the e_j. Then $e_j = \Omega^j e_0$, so $\sum_j c_j e_j = q(\Omega) e_0$. As Ω is unitary, $\|q(\Omega)\| \le \max_{\mathbb{T}} |q|$, and

$$c^* R c = \Big\| \sum_j c_j e_j \Big\|^2 = \|q(\Omega) e_0\|^2 \le \max_{\mathbb{T}} |q|^2 \, \|e_0\|^2 = R_{00} \max_{\mathbb{T}} |q|^2. \qquad \square$$

Lemma 3.2 is a consequence of the Carathéodory–Toeplitz theorem. Patel's post contains this argument for $k = 3$, and Ye et al. prove it in the same way [25, Lemma 3.1].

Lemma 3.3. *Let $p(\zeta) = \sum_{j=0}^{3} c_j \zeta^j$ satisfy $\max_{\mathbb{T}} |p| \le 1$. Then*

$$35\big(|c_0| + |c_3|\big)^2 + 28\big(|c_1| + |c_2|\big)^2 \le 44.$$

Consequently $35A_3^2 + 28B_3^2 \le 176$.

Proof. Replacing $p(\zeta)$ by $p(e^{i\theta}\zeta)$ for a suitable θ, we may assume that c_1 and c_2 have the same argument. Choose $\xi \in \mathbb{T}$ with $|\bar{\xi} c_0 + c_3| = |c_0| + |c_3|$, and let

$$u = (3, 4, -4, -3)^{\mathsf{T}}, \qquad v = (0, 1, 1, 0)^{\mathsf{T}}, \qquad w = (\xi, 0, 0, 1)^{\mathsf{T}}, \qquad R = uu^* + 28vv^* + 35ww^*.$$

The matrix R is positive semidefinite. Since $|\xi| = 1$, it is also Toeplitz, with first row $(44, 12, -12, -9 + 35\xi)$. By Lemma 3.2, with $c = (c_0, \dots, c_3)^{\mathsf{T}}$,

$$|u^* c|^2 + 28|v^* c|^2 + 35|w^* c|^2 = c^* R c \le 44.$$

Here $|v^* c| = |c_1 + c_2| = |c_1| + |c_2|$ and $|w^* c| = |\bar{\xi} c_0 + c_3| = |c_0| + |c_3|$, which proves the first claim. For the second, apply it to $p = q_3/2$ from the proof of Lemma 2.3. Then $|c_0| + |c_3| = A_3/2$ and $|c_1| + |c_2| = B_3/2$. □

Patel describes this argument as a weighted variant of the identity in the proof of $S_3 = 5/3$ by Yu et al. [26]. The proof above does not depend on [26], and it extends to a one-parameter family. For $-1 \le t \le 2$ and $u_t = (t, 1, -1, -t)^{\mathsf{T}}$, the matrix $u_t u_t^* + (t+1)vv^* + (2 + t - t^2)ww^*$ is positive semidefinite and Toeplitz, with first row $(2 + t, t, -t, -t^2 + (2 + t - t^2)\xi)$, so the same argument shows

$$(2 + t - t^2)\big(|c_0| + |c_3|\big)^2 + (t+1)\big(|c_1| + |c_2|\big)^2 \le 2 + t.$$

Lemma 3.3 is the case $t = 3/4$. The case $t = 1/2$ is the inequality $9(|c_0| + |c_3|)^2 + 6(|c_1| + |c_2|)^2 \le 10$, from which Yu et al. obtain $S_3 \le 5/3$ by the Cauchy–Schwarz inequality [26, (6)]. On its own, Lemma 3.3 gives only $\|H_3\|_1 \le \sqrt{396/35} \approx 3.364$, more than $2S_3 = 10/3$. It is used together with the next inequality, which bounds A_3 in terms of B_2.

Lemma 3.4. $A_3 + \frac{1}{2}B_2^2 \le 2$.

Proof. We use the transfer-function realization of Schur functions on the bidisc. The Schur–Agler class consists of the $g \in \mathcal{S}(\mathbb{D}^2)$ with $\|g(X_1, X_2)\| \le 1$ for every pair of commuting strict contractions X_1, X_2 on a Hilbert space. By Andô's inequality [3], every $g \in \mathcal{S}(\mathbb{D}^2)$ belongs to this class. By Agler's theorem [1] (see also [4, 2]) there are a Hilbert space $\mathcal{K}$, complementary

orthogonal projections P and Q on $\mathcal{K}$, and an isometry $V = \left(\begin{smallmatrix} \kappa & x^* \\ y & W \end{smallmatrix}\right)$ of $\mathbb{C} \oplus \mathcal{K}$, the colligation, such that

$$g(z) = \kappa + x^* Z (I - WZ)^{-1} y, \qquad Z = z_1 P + z_2 Q \qquad (z \in \mathbb{D}^2).$$

For $z \in \mathbb{D}^2$ we have $\|Z\| \le \max_j |z_j| < 1$, so $I - WZ$ and the other operators $I - ZW$, $I - ZU$ inverted below are invertible. We may take V unitary, which matters if the realization is given with an isometric colligation, as in [14, Section 2]. Indeed, the operator $\left(\begin{smallmatrix} V & I - VV^* \\ 0 & V^* \end{smallmatrix}\right)$ on $(\mathbb{C} \oplus \mathcal{K}) \oplus (\mathbb{C} \oplus \mathcal{K})$ is unitary and agrees with V on the first summand. We use it as the new colligation, with state space $\mathcal{K} \oplus (\mathbb{C} \oplus \mathcal{K})$ and projections $P \oplus I$ and $Q \oplus 0$. The new y is the old one, the new W and x^* agree with the old ones on $\mathcal{K}$, and the new projections map $\mathcal{K}$ into itself, so each term $x^* Z (WZ)^m y$ of the expansion of g is unchanged. From now on $\mathcal{K}$, V, x, y, W, P and Q denote the new data.

Apply this to $g = (H - 1)/(H + 1)$, which lies in $\mathcal{S}(\mathbb{D}^2)$ and vanishes at 0. Then $\kappa = 0$, and since V is unitary,

$$\|x\| = \|y\| = 1, \qquad Wx = 0, \qquad W^* y = 0, \qquad xx^* + W^* W = I, \qquad yy^* + WW^* = I.$$

Put $U = W + yx^*$. These identities imply $U^*U = UU^* = I$. Let $G = I - ZW$. Then $Gx = x$ and $x^* G^{-1} Z y = x^* Z (I - WZ)^{-1} y = g$. Since $I - ZU = G - Zyx^*$ is a rank-one perturbation of G and $1 - g \neq 0$,

$$x^* (I - ZU)^{-1} x = x^* G^{-1} x + \frac{(x^* G^{-1} Z y)(x^* G^{-1} x)}{1 - x^* G^{-1} Z y} = 1 + \frac{g}{1 - g} = \frac{1}{1 - g}.$$

Hence $x^*(I + ZU)(I - ZU)^{-1} x = 2x^*(I - ZU)^{-1} x - 1 = (1 + g)/(1 - g) = H$, and expanding the resolvent,

$$H_n(z) = 2x^* (ZU)^n x \qquad (n \ge 1).$$

Relative to $\mathcal{K} = P\mathcal{K} \oplus Q\mathcal{K}$ write

$$U = \begin{pmatrix} U_{11} & U_{12} \\ U_{21} & U_{22} \end{pmatrix}, \qquad x = \begin{pmatrix} a \\ b \end{pmatrix}, \qquad Ux = \begin{pmatrix} \tilde{a} \\ \tilde{b} \end{pmatrix},$$

so that $\|a\|^2 + \|b\|^2 = \|\tilde{a}\|^2 + \|\tilde{b}\|^2 = 1$. Collecting the monomials $z_1 z_2$, z_1^3 and z_2^3 in $2x^*(ZU)^n x$, we find

$$\tfrac{1}{2} h_{11} = a^* U_{12} \tilde{b} + b^* U_{21} \tilde{a}, \qquad \tfrac{1}{2} h_{30} = a^* U_{11}^2 \tilde{a}, \qquad \tfrac{1}{2} h_{03} = b^* U_{22}^2 \tilde{b}.$$

Put $\sigma = \|U_{11}^* a\|^2 + \|U_{22}^* b\|^2$ and $\tau = \|U_{11} \tilde{a}\|^2 + \|U_{22} \tilde{b}\|^2$. Since U^* is an isometry, $\|U_{11}^* a\|^2 + \|U_{12}^* a\|^2 = \|a\|^2$ and $\|U_{21}^* b\|^2 + \|U_{22}^* b\|^2 = \|b\|^2$, so $\|U_{12}^* a\|^2 + \|U_{21}^* b\|^2 = 1 - \sigma$. Since U is an isometry, $\|U_{12} \tilde{b}\|^2 + \|U_{21} \tilde{a}\|^2 = 1 - \tau$ in the same way. Applying the Cauchy–Schwarz inequality to $\frac{1}{2} h_{11} = \langle \tilde{b}, U_{12}^* a \rangle + \langle \tilde{a}, U_{21}^* b \rangle = \langle U_{12} \tilde{b}, a \rangle + \langle U_{21} \tilde{a}, b \rangle$ in the two groupings, we obtain

$$\tfrac{1}{4} B_2^2 \le 1 - \sigma \qquad \text{and} \qquad \tfrac{1}{4} B_2^2 \le 1 - \tau.$$

Finally $\frac{1}{2}|h_{30}| = |\langle U_{11} \tilde{a}, U_{11}^* a \rangle| \le \frac{1}{2}(\|U_{11}^* a\|^2 + \|U_{11} \tilde{a}\|^2)$, and likewise for h_{03}, so $A_3 \le \sigma + \tau \le 2 - \frac{1}{2} B_2^2$. □

Remark 3.5. Lemmas 3.1 and 3.4 are sharp. Equality holds in both for $H = (1 + z_1)/(1 - z_1)$, where $A_2 = A_3 = 2$ and $B_2 = 0$, and for $H = (1 + z_1 z_2)/(1 - z_1 z_2)$, where $A_2 = A_3 = 0$ and $B_2 = 2$.

Lemma 3.6. *Let $\alpha, \beta > 0$, $\gamma \ge 0$ and $r > 0$, and put*

$$\mathcal{B}(r) = 4\alpha + 176\beta + 2\gamma + \frac{1}{4\alpha} + \frac{1}{4\alpha + 2\gamma} + \frac{(r - \gamma)^2}{140\beta} + \frac{r^2}{112\beta}.$$

Then $\|H_2\|_1 + r\|H_3\|_1 \le \mathcal{B}(r)$.

Proof. Expanding the squares verifies the identity

$$\begin{aligned}\mathcal{B}(r)-(A_2+B_2+r(A_3+B_3)) &= \alpha(4-A_2^2-B_2^2)+\beta(176-35A_3^2-28B_3^2)+\gamma(2-A_3-\tfrac{1}{2}B_2^2)\\ &\quad+\alpha\Big(A_2-\frac{1}{2\alpha}\Big)^2+\Big(\alpha+\frac{\gamma}{2}\Big)\Big(B_2-\frac{1}{2\alpha+\gamma}\Big)^2\\ &\quad+35\beta\Big(A_3-\frac{r-\gamma}{70\beta}\Big)^2+28\beta\Big(B_3-\frac{r}{56\beta}\Big)^2.\end{aligned}$$

By Lemmas 3.1, 3.3 and 3.4, each term on the right is nonnegative. □

Thus $\mathcal{B}(r)$ is the Lagrangian bound, with multipliers α, β and γ, for the maximum of $A_2+B_2+r(A_3+B_3)$ under the three inequalities of Lemmas 3.1, 3.3 and 3.4.

4. Sidon constants and certificates for S_5, S_6 and S_7

Table 1 lists what is known about S_k for small k. The proof of Theorem 1.1 uses upper bounds for S_1 and for S_k with $k \geq 4$. Degrees two and three are covered by Lemma 3.6.

Table 1. Sidon constants $S_k = S(\{0,1,\dots,k\})$.

k	S_k	source
1	1	elementary
2	$\sqrt{2}$	Newman (see [19])
3	5/3	Yu et al. [26]; lower bound Neuwirth [20]
4	2	Newman's bound; equality by Shapiro [24]
5	$2.186529771892569\ldots$; ≤ 2.187	Ye et al. [25]; Proposition 4.2
6	≤ 2.363	Proposition 4.2
7	≤ 2.552	Proposition 4.2
$k \geq 8$	$\leq \sqrt{k}$	Newman [24, Chapter 3]

Ye et al. [25, Theorem 1.1] show that S_5 is the root in $(2.18, 2.19)$ of $675X^4 - 1026X^3 - 936X^2 - 414X + 677$. Proposition 4.2 below gives the weaker bound $S_5 \leq 2.187$, which suffices for Theorem 1.1, together with $S_6 \leq 2.363$ and $S_7 \leq 2.552$. Its proof rests on the following lemma.

Lemma 4.1. *Let $k \geq 1$. For $w \in \mathbb{C}^{k+1}$ put*

$$\nu(w) = \max\Big\{\operatorname{Re}\sum_{j=0}^{k}\bar{w}_j c_j :\ \max_{\mathbb{T}}\Big|\sum_{j=0}^{k} c_j z^j\Big| \leq 1\Big\}.$$

(a) $S_k = \max\{\nu(w) : w_0 = w_1 = 1,\ |w_j| = 1\ (2 \leq j \leq k)\}$.
(b) *ν is convex.*
(c) *Let $y \in \mathbb{C}^k$, and let $T(y)$ be the Hermitian Toeplitz matrix with diagonal entries 1 and entries $T(y)_{jl} = y_{l-j}$ for $l > j$. If $T(y)$ is positive definite, then $\nu(w)^2 \leq w^* T(y)^{-1} w$.*
(d) *For $0 < \delta \leq \pi/4$, the arc $\{e^{i\varphi} : |\varphi - \varphi_0| \leq \delta\}$ lies in the triangle with vertices $e^{i(\varphi_0-\delta)}$, $e^{i(\varphi_0+\delta)}$ and $e^{i\varphi_0}/\cos\delta$.*
(e) *$\nu(\bar{w}) = \nu(w)$, where $\bar{w} = (\bar{w}_0, \dots, \bar{w}_k)$.*

Proof. (a) Write $q(z) = \sum_j c_j z^j$. If $|w_j| = 1$ for all j, then $\nu(w) \leq \max\{\sum_j |c_j| : \max_{\mathbb{T}} |q| \leq 1\} = S_k$. Equality holds when c attains this maximum and $w_j = c_j/|c_j|$ (any unimodular number if $c_j = 0$). The substitution $c_j \mapsto c_j e^{i(\psi+j\theta)}$ replaces $q(z)$ by $e^{i\psi} q(e^{i\theta} z)$ and w_j by $w_j e^{-i(\psi+j\theta)}$ without changing ν, and a choice of ψ and θ makes $w_0 = w_1 = 1$. (b) ν is a supremum of

real-linear functions of w. (c) By Lemma 3.2, $c^*T(y)c \le \max_{\mathbb{T}} |q|^2$ for $c = (c_0, \dots, c_k)^{\mathsf{T}}$. Hence, by the Cauchy–Schwarz inequality,

$$\operatorname{Re} w^*c \le \bigl(w^*T(y)^{-1}w\bigr)^{1/2}\bigl(c^*T(y)c\bigr)^{1/2} \le \bigl(w^*T(y)^{-1}w\bigr)^{1/2} \max_{\mathbb{T}} |q|.$$

(d) After a rotation, $\varphi_0 = 0$. The triangle is $\{u \in \mathbb{C} : \operatorname{Re} u \ge \cos\delta,\ \operatorname{Re}(ue^{-i\delta}) \le 1,\ \operatorname{Re}(ue^{i\delta}) \le 1\}$, and $u = e^{i\varphi}$ with $|\varphi| \le \delta$ satisfies $\cos\varphi \ge \cos\delta$ and $\cos(\varphi \mp \delta) \le 1$. (e) The map $c \mapsto \bar{c}$ maps the set of admissible c onto itself, since $\sum_j \bar{c}_j z^j = \overline{q(\bar{z})}$, and $\operatorname{Re}\sum_j w_j\bar{c}_j = \operatorname{Re}\sum_j \bar{w}_j c_j$. Hence $\nu(\bar{w}) = \max \operatorname{Re}\sum_j w_j c_j = \max \operatorname{Re}\sum_j w_j\bar{c}_j = \nu(w)$. □

Parts (a) and (c) are variants of the reduction to phase functionals and of the Toeplitz domination bound used by Ye et al. [25, (14) and Proposition 3.2], who normalize the phases differently [25, (17)]. Proposition 4.2 also differs from their proof in the covering of the phase space.

Proposition 4.2. *$S_5 \le 2.187$, $S_6 \le 2.363$ and $S_7 \le 2.552$.*

Proof (computer-assisted). Let (k, b_k) be $(5, 2.187)$, $(6, 2.363)$ or $(7, 2.552)$. By Lemma 4.1(a) it suffices to show that $\nu(1, 1, e^{i\varphi_2}, \dots, e^{i\varphi_k}) \le b_k$ for $(\varphi_2, \dots, \varphi_k) \in [-\pi, \pi]^{k-1}$. By Lemma 4.1(e), which replaces $(\varphi_2, \dots, \varphi_k)$ by $(-\varphi_2, \dots, -\varphi_k)$, we may also assume $\varphi_2 \ge 0$. For $k = 5$ and $k = 6$ the certificate covers $[-\pi, \pi]^{k-1}$ by boxes, obtained from a uniform grid of 8^{k-1} boxes with half-widths $\pi/8$ by repeated bisection. For $k = 7$ it covers $[0, \pi] \times [-\pi, \pi]^5$ in the same way, starting from a uniform grid of $2 \cdot 4^5$ boxes with half-widths $\pi/4$. By Lemma 4.1(d), the vectors $(1, 1, e^{i\varphi_2}, \dots, e^{i\varphi_k})$ with φ in the box lie in a product of $k - 1$ triangles. By Lemma 4.1(b), the maximum of ν over this product is attained at one of its 3^{k-1} vertices. For each vertex v that occurs, the certificate gives $y \in \mathbb{C}^k$ with $T(y)$ positive definite and $v^*T(y)^{-1}v < b_k^2$, so that $\nu(v) < b_k$ by Lemma 4.1(c). A vertex with a coordinate -1 occurs with two labels, phase $-\pi$ and phase π, and has a certificate for each. Table 2 gives the sizes of the three certificates, and Section 6 describes the check. □

Table 2. The certificates of Proposition 4.2.

	$k = 5$	$k = 6$	$k = 7$
bound b_k	2.187	2.363	2.552
initial grid	8^4	8^5	$2 \cdot 4^5$
boxes	8,912	61,118	210,960
vertex certificates	269,769	9,893,956	65,041,100
smallest margin $b_k^2 - v^*T(y)^{-1}v$	$1.49 \cdot 10^{-5}$	$8.48 \cdot 10^{-8}$	$8.88 \cdot 10^{-8}$

For $k = 6$ the certificate is split into 48 files, and a vertex that occurs in several files has a certificate in each, so the count 9,893,956 includes repetitions. There are 4,333,585 distinct vertex labels. For $k = 7$ it is split into 576 files in the same way. Numerical searches, which are not part of the proof, give $S_6 \approx 2.36256$ and $S_7 \approx 2.55166$, so the bounds 2.363 and 2.552 are close to the true values.

5. Proof of Theorem 1.1

Let $N = 40$. Let $s_4 = 2$, $s_5 = 2.187$, $s_6 = 2.363$, $s_7 = 2.552$, and $s_n = \lceil 10^8\sqrt{n}\,\rceil/10^8$ for $8 \le n \le N$. By Table 1 and Proposition 4.2, $S_n \le s_n$ for $4 \le n \le N$, and $S_n \le \sqrt{n} \le n$ for all n. Let $H \in \mathcal{P}_2$, let $0 < r < 1$, and fix $\alpha, \beta > 0$ and $\gamma \ge 0$. Lemma 2.3 bounds $\|H_n\|_1$ by 2 for $n = 1$, by $2s_n$ for $4 \le n \le N$ and by $2n$ for $n > N$, and Lemma 3.6 bounds $\|H_2\|_1 + r\|H_3\|_1$.

Summing the series $\sum_{n>N} nr^n = r^{N+1}(N+1-Nr)/(1-r)^2$, we obtain a majorant of the form that Patel uses [21]:

$$M_r(H) \le E(r) := 2r + r^2\mathcal{B}(r) + 2\sum_{n=4}^{N} s_n r^n + \frac{2r^{N+1}(N+1-Nr)}{(1-r)^2}.$$

We take

$$r_* = \frac{60333}{200000}, \qquad \alpha = \frac{20674053}{62500000}, \qquad \beta = \frac{3901457}{1562500000}, \qquad \gamma = \frac{101990031}{10^9}.$$

All quantities in $E(r_*)$ are then rational, and an exact computation shows $E(r_*) < 0.99999556 < 1$. Since α, β and γ are fixed, E is continuous on $(0,1)$, so there is $r' > r_*$ with $E(r') < 1$. By Lemma 2.1, as in Patel's argument, $K_2 \ge r' > r_*$, which proves Theorem 1.1.

With Patel's parameters $\alpha = 264631/800000$, $\beta = 249627/10^8$ and $\gamma = 10197021/10^8$, with $N = 20$ and with $s_n = \lceil 10^8\sqrt{n}\,\rceil/10^8$ for all $4 \le n \le 20$, the function E is the rational majorant that Patel evaluates in [21]. Its value at his radius $r_P = 15079379/(5\cdot 10^7)$ is less than 0.99999997857, below his bound $1 - 1/(5\cdot 10^7)$ (Table 3).

6. Verification

Four programs check the computations, and a fifth checks Patel's computation. All of them are in the archive listed under Data availability, together with the certificates, the programs that produced them and the output of all runs.

The first program checks the certificate for S_5 in ball arithmetic, using Arb [13] through python-flint [23] at a precision of 128 bits. The certificate stores the entries of each y as decimal numerals, which Python parses into the nearest binary64 numbers. These binary64 numbers are the certificate, and the program reads them exactly into balls. It rebuilds the boxes from the uniform grid and the recorded bisections, so the boxes cover $[-\pi,\pi]^4$ by construction. For each box it forms the 81 vertices of Lemma 4.1(d) from rational multiples of π. For each vertex label it checks that the leading principal minors of $T(y)$ are positive and that $v^*T(y)^{-1}v < 2.187^2$, solving the linear system in ball arithmetic. The run takes about 30 seconds and 0.6 GB of memory on one core of an x86-64 server.

The certificate for S_6 occupies about 3.4 GB and is split into 48 files, and the certificate for S_7 occupies about 26.8 GB and is split into 576 files. Each file contains some cells of the initial grid, their subtrees and the vertex certificates they need. The second program carries out the checks of the first program on one such file and lists the cells it has covered, together with the name of the file and its SHA-256 checksum. The third reads these lists and checks that the 48 files together cover each of the 8^5 cells exactly once, and likewise for the 576 files and the $2\cdot 4^5$ cells of the certificate for S_7. It recomputes the checksum of each file named in the lists and compares it with the listed one, but it trusts that the lists come from runs of the second program. The logs in the archive record such a run. Each of the 48 runs of the second program takes under a minute and about 0.7 GB of memory on one core, and each of the 576 runs for S_7 takes between a few seconds and 15 minutes, with at most 11.1 GB of memory.

The fourth program evaluates $E(r_*)$ of Section 5 in exact rational arithmetic and compares it with 1. The fifth checks, in exact rational arithmetic, that $\mathcal{B}(r_P)$ equals the fraction displayed in [21], that his table of rational upper bounds for $\sqrt{n}$ consists of the integers $\lceil 10^8\sqrt{n}\,\rceil$, that the two parts of his majorant lie below the two rational bounds he displays, and that the sum of these bounds is below $1 - 1/(5\cdot 10^7)$. Table 3 lists the results, with the exact values rounded up.

With the same α, β, γ and s_n, the fourth program finds $E(r_* + 10^{-6}) \ge 1$, and it finds $E(r_*) \ge 1$ when $s_7 = 2.552$ is replaced by $\lceil 10^8\sqrt{7}\,\rceil/10^8$. The first program rejects the certificate

TABLE 3. Results of the programs.

run	result
certificate for S_5, bound 2.187	accepted
certificate for S_6, bound 2.363 (48 files and their coverage)	accepted
certificate for S_7, bound 2.552 (576 files and their coverage)	accepted
$E(r_*)$ with $s_5 = 2.187$, $s_6 = 2.363$ and $s_7 = 2.552$	< 0.99999556
$E(r_P)$ in Patel's computation [21]	< 0.99999997857

for S_5 with the bound 2.186, which is less than S_5 [25], and the second rejects a file of the certificate for S_6 with the bound 2.362 and a file of the certificate for S_7 with the bound 2.551, below the numerical value 2.55166 of S_7. The programs also reject deliberately altered inputs, and the archive contains the log of these tests.

7. Formal verification

We also prove the bound $K_2 \geq 0.301657$ in the Lean 4 proof assistant [16], with the Mathlib library [17]. This bound uses the bounds of Proposition 4.2 for S_5 and S_6 but not the one for S_7.The formal proof has one hypothesis, the inequality of Lemma 3.4. Its proof rests on Andô's inequality and Agler's realization, which are not in Mathlib. The formal statements use Mathlib's definitions. A family $c\colon \mathbb{N}^2 \to \mathbb{C}$ is admissible if the series $\sum_\alpha c_\alpha z^\alpha$ converges absolutely at every point of $\mathbb{D}^2$ and its value there has modulus less than 1. Bohr's inequality holds at r if $\sum_{\alpha\in F} |c_\alpha| r^{|\alpha|} \leq 1$ for every admissible family c and every finite set $F \subset \mathbb{N}^2$, and K_2 is the supremum of the radii $r > 0$ at which it holds. It agrees with the definition in the introduction: a nonconstant $f \in \mathcal{S}(\mathbb{D}^2)$ maps $\mathbb{D}^2$ into $\mathbb{D}$, and a constant of modulus 1 satisfies Bohr's inequality with equality. The hypothesis states $A_3 + \frac{1}{2}B_2^2 \leq 2$ for every family h with $h_{00} = 1$ whose series converges absolutely on $\mathbb{D}^2$ to values with positive real part, that is, for the coefficients of every $H \in \mathcal{P}_2$. The theorem `Bohr59aN.K2_ge_N` states that this hypothesis implies $K_2 \geq 0.301657$.

The formal proof follows Sections 2 to 5. It proves Carathéodory's inequality on slices, Lemmas 2.3, 3.1, 3.2, 3.3 and 3.6, Newman's bound $S_n \leq \sqrt{n}$, the reduction of Section 5 and the exact evaluation of the majorant. The bounds $S_5 \leq 2.187$ and $S_6 \leq 2.363$ come from the certificates of Proposition 4.2. A checking program written in Lean validates each part of a certificate in exact rational arithmetic and ball arithmetic, and a Lean proof shows that whenever it accepts a part, the bound of Lemma 4.1(c) holds on the boxes that the part covers. Our scripts convert the certificates into Lean source files. None of these files needs to be trusted, since the correctness proof holds for every input that the program accepts.

The runs of the checking program on the 49 parts of the certificates, one for S_5 and 48 for S_6, are evaluated by `native_decide`, which executes compiled code. The trusted base therefore includes the Lean compiler, the C compiler that it uses, the Lean runtime and the compiled core-library functions that the checks call. Besides the standard axioms `propext`, `Classical.choice` and `Quot.sound`, the theorem `K2_ge_N` depends on 49 auxiliary axioms, one for each part, each stating that the compiled check returns true. The other parts of the proof use only the standard axioms. Among them are the theorems `Bohr59a.K2_ge_patel`, which proves Patel's bound $K_2 \geq 0.30158758$ under the same hypothesis, and `Bohr59a.K2_ge_of_sidon`, which derives $K_2 \geq 0.301657$ from the hypothesis and the two Sidon bounds. The checking program contains no definitions marked `implemented_by`, `extern`, `unsafe`, `opaque` or `partial`. Rebuilt from a clean copy of the sources on other machines, the proof gives the same axioms, and regenerating the Lean files of the certificates reproduces them byte for byte. The exported proof was accepted

by `leanchecker` and by two further kernel implementations, nanoda [18] and lean4lean [10], with the 49 auxiliary axioms admitted. The Lean files, for Lean 4 version `v4.35.0-rc3` and Mathlib commit `c55e6e786f49`, are in the directory `lean/` of the archive.

A second formal proof does without compiled code. Its checking program uses only integers and structural recursion, so that the Lean kernel can evaluate it, and it is again proved correct in Lean. For each vertex of the certificates for S_5 and S_6, one of its vertex certificates becomes a theorem checked by the kernel, and further theorems check the boxes of each cell. This gives $K_2 \geq 0.301657$ under the same hypothesis (theorem `Bohr59aK.K2_ge_K`), with only the standard axioms. Its Lean files are in the directory `lean_kernel/` of the archive.

Data availability

The certificates of Proposition 4.2, the programs of Section 6, the programs that produced the certificates, the controls and the output of all runs, and the Lean files of Section 7, are archived at doi:10.5281/zenodo.23553495. The repository `https://github.com/tejasvi/bidisc-bohr-radius-v3` holds the same files except the 48 files of the certificate for S_6 and the 576 files of the certificate for S_7. The certificate for S_5 has the SHA-256 checksum

`ce3e9dd41a0dbd9c3a8ea49b04aa7aaf21f0fe557291a2084c7739461a460a14`,

and the archive lists the checksums of the other files.

Use of AI tools

Generative AI tools were used to find and prove the results, to write the programs, and to draft the text. The author reviewed the arguments and the programs and takes responsibility for the content.

References

[1] J. Agler, *On the representation of certain holomorphic functions defined on a polydisc*, in: Topics in Operator Theory: Ernst D. Hellinger Memorial Volume, Oper. Theory Adv. Appl. **48**, Birkhäuser, Basel, 1990, 47–66.

[2] J. Agler and J. E. McCarthy, *Pick interpolation and Hilbert function spaces*, Grad. Stud. Math. **44**, Amer. Math. Soc., Providence, RI, 2002.

[3] T. Andô, *On a pair of commutative contractions*, Acta Sci. Math. (Szeged) **24** (1963), 88–90.

[4] J. A. Ball and T. T. Trent, *Unitary colligations, reproducing kernel Hilbert spaces, and Nevanlinna–Pick interpolation in several variables*, J. Funct. Anal. **157** (1998), 1–61.

[5] R. Baran, P. Pikul, H. J. Woerdeman and M. Wojtylak, *Contractive realization theory for the annulus and other intersections of disks on the Riemann sphere*, J. Funct. Anal. **290** (2026), no. 8, 111346; arXiv:2504.03236.

[6] F. Bayart, D. Pellegrino and J. B. Seoane-Sepúlveda, *The Bohr radius of the n-dimensional polydisk is equivalent to* $\sqrt{(\log n)/n}$, Adv. Math. **264** (2014), 726–746.

[7] L. Bernal-González, H. J. Cabana, D. García, M. Maestre, G. A. Muñoz-Fernández and J. B. Seoane-Sepúlveda, *A new approach towards estimating the n-dimensional Bohr radius*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **115** (2021), no. 2, Paper No. 44.

[8] H. P. Boas and D. Khavinson, *Bohr's power series theorem in several variables*, Proc. Amer. Math. Soc. **125** (1997), 2975–2979.

[9] H. Bohr, *A theorem concerning power series*, Proc. London Math. Soc. (2) **13** (1914), 1–5.

[10] M. Carneiro, *Lean4Lean: Verifying a typechecker for Lean, in Lean*, arXiv:2403.14064, 2024.

[11] A. Defant, L. Frerick, J. Ortega-Cerdà, M. Ounaïes and K. Seip, *The Bohnenblust–Hille inequality for homogeneous polynomials is hypercontractive*, Ann. of Math. (2) **174** (2011), 485–497.

[12] S. Griego, *Improve* C_{59} *upper bound*, pull request no. 75 to the repository teorth/optimizationproblems, GitHub, May 2026, `https://github.com/teorth/optimizationproblems/pull/75`.

[13] F. Johansson, *Arb: efficient arbitrary-precision midpoint-radius interval arithmetic*, IEEE Trans. Comput. **66** (2017), 1281–1292.

[14] G. Knese, *Three radii associated to Schur functions on the polydisk*, Proc. Amer. Math. Soc. Ser. B **12** (2025), 48–63; arXiv:2410.21693v3.

[15] G. Knese, *Corrigendum to "Three radii associated to Schur functions on the polydisk"*, Proc. Amer. Math. Soc. Ser. B **13** (2026), no. 4, 35–36; doi:10.1090/bproc/280.
[16] L. de Moura and S. Ullrich, *The Lean 4 theorem prover and programming language*, in: Automated Deduction – CADE 28, Lecture Notes in Comput. Sci. **12699**, Springer, 2021, 625–635.
[17] The mathlib Community, *The Lean mathematical library*, in: Proc. 9th ACM SIGPLAN Int. Conf. Certified Programs and Proofs (CPP 2020), 2020, 367–381.
[18] `nanoda_lib`, a type checker for Lean 4, `https://github.com/ammkrn/nanoda_lib`.
[19] S. Neuwirth, *The Sidon constant of sets with three elements*, arXiv:math/0102145, 2001.
[20] S. Neuwirth, *On the (Fourier analytic) Sidon constant of* $\{0,1,2,3\}$, arXiv:2603.28229, 2026 (note dated 2013).
[21] S. Patel, *Improved lower and upper bounds for the bidisc Bohr radius*, solution to the problem "Exact value of the Bohr radius of the bidisc", MathDB, second solution on the page, posted 27 August 2026 (UTC) and edited the same day, `https://mathdb.com/p/392062/exact-value-of-the-bohr-radius-of-the-bidisc#s2` (consulted 21 October 2026).
[22] S. Patel, *Complex phases and a certified upper bound for the bidisc Bohr radius: reproducibility package*, version 1.0.0, Zenodo, 5 September 2026, doi:10.5281/zenodo.22341928.
[23] python-flint: Python bindings for FLINT, version 0.9.0 (FLINT 3.6.0), `https://github.com/flintlib/python-flint`.
[24] H. S. Shapiro, *Extremal problems for polynomials and power series*, Master's thesis, Massachusetts Institute of Technology, 1951.
[25] Z. Ye et al., *The exact Sidon constant of six consecutive frequencies*, preprint, October 2026, `https://github.com/ABCDEFGCoder/The-Sidon-Constant-of-n-Consecutive-Frequencies`.
[26] H. Yu et al., *The exact Sidon constant of* $\{0,1,2,3\}$ *and its formalization*, preprint, ai.viXra.org:2606.0009, 2026 (dated 10 May 2026); Lean formalization at `https://github.com/yhx-12243/Sidon3`.

ORCID: `https://orcid.org/0000-0003-4668-3639`