

[illegible]

SUMMARY OUTPUT

Regression Statistics

Multiple R	0.9787069
R Square	0.9578673
Adjusted R Square	0.9536540
Standard Error	6.6565414
Observations	12


ANOVA


	df	SS	MS	F	Significance F
Regression	1	10073.571	10073.571	227.34540	0.0000000
Residual	10	443.09544	44.309544		
Total	11	10516.666			

	Coefficients	Standard Error	t Stat	P-value	Lower 95%	Upper 95%	Lower 95.0%	Upper 95.0%
Intercept	-9.140912	6.8290303	-1.338537	0.2103568	-24.35694	6.0751153	-24.35694	6.0751153
X Variable 1	1.1081223	0.0734927	15.077977	0.0000000	0.9443702	1.2718744	0.9443702	1.2718744

RESIDUAL OUTPUT

Observation	Predicted Y	Residuals
1	42.940839	-4.940839
2	59.562675	2.4373247
3	62.887047	-9.887042
4	68.427654	-1.427654
5	68.427654	15.572345
6	77.292633	1.7073666
7	96.130711	-3.130713
8	101.67132	4.3286741
9	117.18503	-0.185039
10	121.61752	-5.617528





		11	128.2662	-1.266263				
		12	131.5906	2.409369				

	Normal Distribution						
Marks	f(x)						
6	0.00417739717		Mean	50.727272			
12	0.005631893802		SD	28.948515			
25	0.009284827905		Variance	838.01652			
30	0.01066499062						
45	0.01351400868						
51	0.01378048492						
62	0.01277486148						
65	0.01220388619						
76	0.009414141944						
88	0.006015878378						
98	0.003632657595						
	Fitting	of a	Normal	Distribution			
X	f=freq	x=mid-val	x*f	x-mean		f*(x-mean)^2	
0-10	6	5	30	-22		2904	
10-20	8	15	120	-12		1152	
20-30	14	25	350	-2		56	
30-40	16	35	560	8		1024	
40-50	4	45	180	18		1296	
50-60	2	55	110	28		1568	
>60	0		0	-27		0	
TOTAL	50		1350			8000	
			Mean=	27			

(x-mean)^2
484
144
4
64
324
784
729