

(1) EM Spectrum

Ranges	γ	X	UV	vis	IR	micro	radio
Energy (E)	↑						↓
Frequency (ν)	↑						↓
Wavelength (λ)	↓						↑
Wavelength (λ)	<0.01	0.01–10	10–400	400–700	700–1e6	1e6–9	>1e9

• violet • indigo • blue • green • yellow • orange • red

(2) $c = \lambda \nu$ (Speed of light = wavelength × frequency)

c	ν (Hz)	kHz	MHz	GHz
λ (m)	2.9979e8 (m/s)	2.9979e5 (m·kHz)	2.9979e2 (m·MHz)	2.9979e-1 (m·GHz)
nm	2.9979e17 (nm·Hz)	2.9979e14 (nm·kHz)	2.9979e11 (nm·MHz)	2.9979e8 (nm·GHz)

(3) $E = h\nu = h(c/\lambda)$ (Energy = Planck × frequency)

hc , h	λ (m)	λ (nm)	ν (Hz)
E (J / photon)	1.9864e-25 (J·m)	1.9864e-16 (J·nm)	6.626e-34 (J·s)
J / mol	1.1962e-1 (J·m/mol)	1.1962e8 (J·nm/mol)	3.9902e-10 (J·s/mol)
kJ / mol	1.1962e-4 (kJ·m/mol)	1.1962e5 (kJ·nm/mol)	3.9902e-13 (kJ·s/mol)

• Grams → mol → ×6.022e23 for photon count

(4) Quantum Principles

Maxwell	Light is electromagnetic radiation
Planck	Energy emitted only in fixed amounts (quanta) (black-body radiation)
Einstein	Different light has different amounts of fixed energy (photons) (photoelectric effect)
Bohr	Electron quanta (H atom line spectra); Rydberg
De Broglie	Electrons are particle and wave (5)
Schrodinger	Wave equation (energy and spatial prob. of e-) (n, l, m_l, m_s)
Heisenberg	Position and velocity not both knowable

$$E = h\nu = h(c/\lambda) = R(1/n_f^2 - 1/n_i^2)$$

$$n_i = \sqrt{1/(1/n_f^2 - hc/(R\lambda))}; n_f = \sqrt{1/(1/n_i^2 + hc/(R\lambda))}$$

Variable	Value	Variable	Value
h	6.626e-34 J·s	h/R	3.0394e-16 s
c	2.9979e8 m/s	hc	1.9864e-25 J·m
R	2.18e-18 J	hc/R	9.1120e-8 m

(5) $\lambda = h/(m \cdot u)$ (De Broglie)

- 1 mm = 1e-3 m, 1 nm = 1e-9 m
- e- (9.11e-31 kg), p+ (1.6726e-27 kg), n (1.6749e-27 kg)

h	u (m/s)	km/s
m (kg)	6.626e-34 (J·s)	6.626e-37 (kJ·s)
g	6.626e-31 (mJ·s)	6.626e-34 (J·s)
amu	3.9892e-7 (J·s·amu/kg)	3.9892e-10 (kJ·s·amu/kg)

(6) Electron Configuration (Aufbau, Pauli, Hund)

- 1s 2s 2p 3s 3p 4s 3d 4p 5s 4d 5p 6s 4f 5d 6p 7s 5f 6d 7p
- Round-robin sublevel before double occupaton (Hund)
- Two e- per orbital (Pauli's exclusion)
- n^2 orbitals, $2n^2$ e- per shell

Symbol	Name	Value
n	Principal energy level (shell)	(1, 2, 3, ...)
l	Sublevel = subshell (shape)	(0, ..., $n - 1$)
m_l	Orbital (orientations)	(-l, ..., l)
m_s	Spin (2 per orbital)	$\pm 1/2$

Sublevel	l	m_l	# orbitals	# e-
s	0	0	1	2
p	1	± 1	3	6
d	2	± 2	5	10
f	3	± 3	7	14

Element	EC	Element	EC
Cr	[Ar]4s ¹ 3d ⁵	Ag	[Kr]5s ¹ 4d ¹⁰
Mo	[Kr]5s ¹ 4d ⁵	Au	[Xe]6s ¹ 4f ¹⁴ 5d ¹⁰
Cu	[Ar]4s ¹ 3d ¹⁰	Pd	[Kr]4d ¹⁰ (no 5s)

(7-8) Reading EC, Anion, Cation

EC → element	add every superscript to get Z
Paramagnetic	at least 1 unpaired val. e- (else dia.)
Ground state	follows Aufbau (else excited)
Valence electrons	highest / outer shell (else core)
Anion	count val. e- needed until noble gas
Cation	remove val. e- until noble gas
Tx metal	remove s before d
Isoelectric	same EC (e.g. Na ⁺ , Al ³⁺ , F ⁻ , N ³⁻ , Ne)

(9-10) Penetration, Shielding, Z_{eff} , Size

$$Z_{eff} \approx Z - (\text{core } e^-)$$

- ↑ penetration (core electrons) = ↓ shielding = ↓ energy
- Energy s < p < d < f (same shell)
- Core e- shield valence e-
- Atomic size (cation < neutral < anion)

(11-13) Trends

	Trend	Top	Bot	Left	Right
(11) Atomic size (radius)	↙ (Cs) (f > p)	↓	↑	↑	↓
(12) Ionization energy (IE)	↗ (He)	↑	↓	↓	↑
(13) Electron affinity (EA)	↗ (Cl)	↑	↓	↓	↑
Z_{eff} (+ ch. toward nucl.)	↘	↓	↑	↓	↑
Metallic	↙ (Fr)	↓	↑	↑	↓
Nonmetallic	↗ (F)	↑	↓	↓	↑
Electronegativity (EN)	↗ (F)	↑	↓	↓	↑

• EA equation: X + e- = X-

(14) Families & Diagonal Rule

Family	Name	Reactivity	e-	Charge
1A	alkali metals	(downward)	lose 1	1+
2A	alkaline earth	less than 1A	lose 2	2+
7A	halogens	(upward)	gain 1	1-
8A	noble gases	inert	—	0

- Diagonal rule (Li~Mg, Be~Al, B~Si)
- Lanthanide (58–71), Actinide (90–103)
- Diatomic (H₂, N₂, O₂, F₂, Cl₂, Br₂ (l), I₂ (s))
- Oxides (basic = metal, acidic = nonmetal)
- Family reactions
 - 2K(s) + 2H₂O(l) → 2KOH(aq) + H₂(g)
 - H₂(g) + Cl₂(g) → 2HCl(g)

(15) Lattice Energy

$$U \propto \frac{q_1 q_2}{r_1 + r_2}$$

- Energy released (exo.) forming ionic crystal (from gas)
 - (Lattice dissociation energy = endothermic)
- max U: 1. ↑ Q+/Q- (ionic charge), 2. ↓ AS (rad.)

Trend	Bond	Enthalpy	Exception
EA	Form	Exothermic	Endo: Be, Mg, noble, EA2 (usually)
IE	Break	Endothermic	--

(16-17) Electronegativity, Bonds

$$\Delta EN = |EN_A - EN_B|$$

- EN = e- attraction in cov. bond.
- ↑ ΔEN EN = ↑ polar = ↑ bond (not always)
- Stronger = shorter (triple > double > single)

ΔEN	Label
0	pure non cov.
0.1 – 0.5	slightly pol. cov.
0.5 – 1.6	pol. cov.
> 1.6	ionic

(18) Formal Charge & Resonance

$$FC = (\text{valence } e^-) - [\text{nonbonding } e^- + \# \text{ bonds}]$$

- Overall 0 charge is better
- Else -1 on the **more electronegative** atom
- Resonance (e.g. NO₂·: two equivalent N-O at 1½)
- Hybrid = ONE averaged structure, all bonds identical — "1 single + 1 double" describes a *form*, not the real ion
- Bond order = total bonds / positions (NO₂· 3/2 = 1½ · NO₃·, CO₃· 4/3 = 1⅓)
- Hybrid bond length/strength sits **between** single and double
- Between forms only e- move (lone pairs ⇌ double bonds), never atoms; same total e-

(19) Lewis Structures (Kelter)

- Sum valence e⁻ (including +/- for ions)
- Sum "wants" (octet)
 - H (2), He (2), - Be (4), Al (6), B (6)
 - 3p or below (>8)
- Sum bonds = (2. - 1.) / 2 (round down)
- Central atom = min EN (never H)
- Remaining e⁻ = 1. outside, central,
 - Consider double/trip. bond for central octet / nonzero FC
 - Ions go in brackets with the charge outside
- FC: overall 0, or 1- on max EN element

(20) Photoelectric Effect ($h\nu$ - binding / work)

$$KE = h\nu - \Phi = h(c/\lambda) - \Phi = \frac{1}{2}m_e u^2$$

Variable	Value
h	6.626e-34 J·s
m_e	9.11e-31 kg
hc	1.9864e-25 J·m
$2hc/m_e$	4.3609e5 m ² /s ²

Have	Want	Use
λ, λ_{thr}	u	$u = \sqrt{(2hc/m_e)(1/\lambda - 1/\lambda_{thr})}$
Φ	λ_{thr} (max ej. λ)	$\lambda_{thr} = hc/\Phi$
ν_{thr} or λ_{thr}	$\Phi, \Phi / \text{mol}$	$\Phi = h\nu_{thr} = h(c/\lambda_{thr})$, n.b. $\Phi \times N_A$ (/1e3 for kJ/mol)
Φ, λ	u	1. $KE = h(c/\lambda) - \Phi$ 2. $u = \sqrt{2KE/m_e}$


(21) Heisenberg Uncertainty

$$\Delta x \cdot \Delta p \geq \frac{h}{4\pi} = 5.2728e-35 \text{ J·s}, \quad \Delta p = m\Delta u$$

- $\Delta p = m \cdot \Delta u$ (kg, m/s)
- $\Delta x \geq h/(4\pi \cdot \Delta p)$ (minimum uncertainty)

(22) Born-Haber Cycle For Lattice Energy (U)

$$\Delta H^\circ_f = \Delta H_{sub} + \sum IE + \frac{1}{2}BE + EA + U$$



- List every step with its sign: sublimation (+), IE (+), bond dissociation (+), EA (usually -), U (-)
- Halve** the bond dissociation if you need only one atom
- Chain: elements → gaseous atoms → gaseous ions → solid
- Set the chain equal to ΔH°_f
- Solve for the single unknown
 - Use IE_2 as well when the cation is 2+
 - U runs ions(g) → solid, so it comes out **negative**
 - Report |U| if asked for a positive lattice energy

(23) Bond Enthalpy

$$\Delta H_f = \sum BE(\text{reactants}) - \sum BE(\text{products})$$

$$\Delta H_f = \sum \Delta H(\text{broken}) + \sum \Delta H(\text{formed})$$

- n.b. **the sign**: BE table values are positive → 1st form **subtracts**; in the 2nd form each ΔH (formed) is already **negative** (energy released) → **add**
- List bonds broken (reactants) and formed (products), using the balanced coefficients
 - $\sum BE(\text{broken})$, always positive
 - $\sum BE(\text{formed})$, subtracted
 - $\Delta H = \sum \text{broken} - \sum \text{formed}$
 - Bonds unchanged on both sides cancel — skip them
 - Negative ΔH = exothermic
 - For "heat released by g of X," convert g → mol and scale

Average Bond Energies (kJ/mol)

Bond	kJ/mol	Bond	kJ/mol	Bond	kJ/mol
H-H	436	N-N	163	Br-F	237
H-C	414	N=N	418	Br-Cl	218
H-N	389	N≡N	946	Br-Br	193
H-O	464	N-O	222	I-Cl	208
H-S	368	N=O	590	I-Br	175
H-F	565	N-F	272	I-I	151
H-Cl	431	N-Cl	200	Si-H	323
H-Br	364	N-Br	243	Si-Si	226
H-I	297	N-I	159	Si-C	301
C-C	347	O-O	142	S-O	265
C=C	611	O=O	498	Si=O	368
C≡C	837	O-F	190	S=O	523
C-N	305	O-Cl	203	Si-Cl	464
C=N	615	O-I	234	S=S	418
C≡N	891	F-F	159	S-F	327
C-O	360	Cl-F	253	S-Cl	253
C=O	736 (799 in CO ₂)	Cl-Cl	243	S-Br	218
C≡O	1072			S-S	266
C-Cl	339				

Lewis Gallery

