

**(1) EM Spectrum**

| Ranges                   | $\gamma$     | X       | UV     | vis     | IR      | micro | radio        |
|--------------------------|--------------|---------|--------|---------|---------|-------|--------------|
| Energy ( $E$ )           | $\uparrow$   |         |        |         |         |       | $\downarrow$ |
| Frequency ( $\nu$ )      | $\uparrow$   |         |        |         |         |       | $\downarrow$ |
| Wavelength ( $\lambda$ ) | $\downarrow$ |         |        |         |         |       | $\uparrow$   |
| Wavelength ( $\lambda$ ) | <0.01        | 0.01-10 | 10-400 | 400-700 | 700-1e6 | 1e6-9 | >1e9         |

• violet • indigo • blue • green • yellow • orange • red

**(2)  $c = \lambda \nu$  (Speed of light = wavelength  $\times$  frequency)**

| $c$           | $\nu$ (Hz)        | kHz                | MHz                | GHz               |
|---------------|-------------------|--------------------|--------------------|-------------------|
| $\lambda$ (m) | 2.9979e8 (m/s)    | 2.9979e5 (m·kHz)   | 2.9979e2 (m·MHz)   | 2.9979e-1 (m·GHz) |
| nm            | 2.9979e17 (nm·Hz) | 2.9979e14 (nm·kHz) | 2.9979e11 (nm·MHz) | 2.9979e8 (nm·GHz) |

**(3)  $E = h\nu = h(c/\lambda)$  (Energy = Planck  $\times$  frequency)**

| $hc$ , $h$       | $\lambda$ (m)        | $\lambda$ (nm)       | $\nu$ (Hz)            |
|------------------|----------------------|----------------------|-----------------------|
| $E$ (J / photon) | 1.9864e-25 (J·m)     | 1.9864e-16 (J·nm)    | 6.626e-34 (J·s)       |
| J / mol          | 1.1962e-1 (J·m/mol)  | 1.1962e8 (J·nm/mol)  | 3.9902e-10 (J·s/mol)  |
| kJ / mol         | 1.1962e-4 (kJ·m/mol) | 1.1962e5 (kJ·nm/mol) | 3.9902e-13 (kJ·s/mol) |

• Grams  $\rightarrow$  mol  $\rightarrow \times 6.022e23$  for photon count

**(4) Quantum Principles**

|                    |                                                                                        |
|--------------------|----------------------------------------------------------------------------------------|
| <b>Maxwell</b>     | Light is electromagnetic radiation                                                     |
| <b>Planck</b>      | Energy emitted only in fixed amounts (quanta) (black-body radiation)                   |
| <b>Einstein</b>    | Different light has different amounts of fixed energy (photons) (photoelectric effect) |
| <b>Bohr</b>        | Electrons are quantum (line spectra); Rydberg eq.                                      |
| <b>De Broglie</b>  | Electrons are particle and wave (5)                                                    |
| <b>Schrodinger</b> | Wave equation (energy and spatial prob. of e-) ( $n, l, m_l, m_s$ )                    |
| <b>Heisenberg</b>  | Position and velocity not both knowable                                                |

$$E = h\nu = h(c/\lambda) = R(1/n_f^2 - 1/n_i^2)$$

$$n_i = \sqrt{1/(1/n_f^2 - hc/(R\lambda))}; n_f = \sqrt{1/(1/n_i^2 + hc/(R\lambda))}$$

| Variable | Value         | Variable | Value          |
|----------|---------------|----------|----------------|
| $h$      | 6.626e-34 J·s | $h/R$    | 3.0394e-16 s   |
| $c$      | 2.9979e8 m/s  | $hc$     | 1.9864e-25 J·m |
| $R$      | 2.18e-18 J    | $hc/R$   | 9.1120e-8 m    |

**(5)  $\lambda = h/(m \cdot u)$  (De Broglie)**

- 1 nm = 1e-9 m
- 1 e- = 9.11e-31 kg

| $h$        | $u$ (m/s)              | km/s                     |
|------------|------------------------|--------------------------|
| $m$ (kg)   | 6.626e-34 (J·s)        | 6.626e-37 (kJ·s)         |
| <b>g</b>   | 6.626e-31 (mJ·s)       | 6.626e-34 (J·s)          |
| <b>amu</b> | 3.9892e-7 (J·s·amu/kg) | 3.9892e-10 (kJ·s·amu/kg) |

**(6) Electron Configuration (Aufbau, Pauli, Hund)**

- 1s 2s 2p 3s 3p 4s 3d 4p 5s 4d 5p 6s 4f 5d 6p 7s 5f 6d 7p
- Round-robin sublevel before double occupaton (Hund)
- Two e- per orbital (Pauli's exclusion)
- $n^2$  orbitals,  $2n^2$  e- per shell

| Symbol | Name                           | Value           |
|--------|--------------------------------|-----------------|
| $n$    | Principal energy level (shell) | (1, 2, 3, ...)  |
| $l$    | Sublevel = subshell (shape)    | (0, ..., n - 1) |
| $m_l$  | Orbital (orientations)         | (-l, ..., l)    |
| $m_s$  | Spin (2 per orbital)           | $\pm 1/2$       |

| Sublevel | $l$ | $m_l$   | # orbitals | # e- |
|----------|-----|---------|------------|------|
| <b>s</b> | 0   | 0       | 1          | 2    |
| <b>p</b> | 1   | $\pm 1$ | 3          | 6    |
| <b>d</b> | 2   | $\pm 2$ | 5          | 10   |
| <b>f</b> | 3   | $\pm 3$ | 7          | 14   |

| Element   | EC                                   | Element   | EC                                                    |
|-----------|--------------------------------------|-----------|-------------------------------------------------------|
| <b>Cr</b> | [Ar]4s <sup>1</sup> 3d <sup>5</sup>  | <b>Ag</b> | [Kr]5s <sup>1</sup> 4d <sup>10</sup>                  |
| <b>Mo</b> | [Kr]5s <sup>1</sup> 4d <sup>5</sup>  | <b>Au</b> | [Xe]6s <sup>1</sup> 4f <sup>14</sup> 5d <sup>10</sup> |
| <b>Cu</b> | [Ar]4s <sup>1</sup> 3d <sup>10</sup> | <b>Pd</b> | [Kr]4d <sup>10</sup> (no 5s)                          |

**(7-8) Reading EC, Anion, Cation**

|                                            |                                                                                           |
|--------------------------------------------|-------------------------------------------------------------------------------------------|
| <b>EC <math>\rightarrow</math> element</b> | add every superscript to get $Z$                                                          |
| <b>Paramagnetic</b>                        | at least 1 unpaired val. e- (else dia.)                                                   |
| <b>Ground state</b>                        | follows Aufbau (else excited)                                                             |
| <b>Valence electrons</b>                   | highest / outer shell (else core)                                                         |
| <b>Anion</b>                               | count val. e- needed until noble gas                                                      |
| <b>Cation</b>                              | remove val. e- until noble gas                                                            |
| <b>Tx metal</b>                            | remove s before d                                                                         |
| <b>Isoelectric</b>                         | same EC (e.g. Na <sup>+</sup> , Al <sup>3+</sup> , F <sup>-</sup> , N <sup>3-</sup> , Ne) |

**(9-10) Penetration, Shielding,  $Z_{eff}$ , Size**

$$Z_{eff} \approx Z - (\text{core } e^-)$$

- $\uparrow$  penetration (core electrons) =  $\downarrow$  shielding =  $\downarrow$  energy
- **Energy** s < p < d < f (same shell)
- **Core** e- shield valence e-
- **Atomic size** (cation < neutral < anion)

**(11-13) Trends**

|                                    | Trend                   | Top          | Bot          | Left         | Right        |
|------------------------------------|-------------------------|--------------|--------------|--------------|--------------|
| <b>(11) Atomic size</b> (radius)   | $\swarrow$ (Cs) (f > p) | $\downarrow$ | $\uparrow$   | $\uparrow$   | $\downarrow$ |
| <b>(12) Ionization energy</b> (IE) | $\nearrow$ (He)         | $\uparrow$   | $\downarrow$ | $\downarrow$ | $\uparrow$   |
| <b>(13) Electron affinity</b> (EA) | $\nearrow$ (Cl)         | $\uparrow$   | $\downarrow$ | $\downarrow$ | $\uparrow$   |
| $Z_{eff}$ (+ ch. toward nucl.)     | $\searrow$              | $\downarrow$ | $\uparrow$   | $\downarrow$ | $\uparrow$   |
| <b>Metallic</b>                    | $\swarrow$ (Fr)         | $\downarrow$ | $\uparrow$   | $\uparrow$   | $\downarrow$ |
| <b>Nonmetallic</b>                 | $\nearrow$ (F)          | $\uparrow$   | $\downarrow$ | $\downarrow$ | $\uparrow$   |
| <b>Electronegativity</b> (EN)      | $\nearrow$ (F)          | $\uparrow$   | $\downarrow$ | $\downarrow$ | $\uparrow$   |

**(14) Families & Diagonal Rule**

| Family    | Name           | Reactivity                      | e-     | Charge |
|-----------|----------------|---------------------------------|--------|--------|
| <b>1A</b> | alkali metals  | most reactive metals (downward) | lose 1 | 1+     |
| <b>2A</b> | alkaline earth | reactive, less than 1A          | lose 2 | 2+     |
| <b>7A</b> | halogens       | reactive nonmetals (upward)     | gain 1 | 1-     |
| <b>8A</b> | noble gases    | <b>inert</b>                    | —      | 0      |

- **Diagonal rule** (Li~Mg, Be~Al, B~Si)
- **Lanthanide** (58-71), **Actinide** (90-103)
- **Diatomic** (H<sub>2</sub>, N<sub>2</sub>, O<sub>2</sub>, F<sub>2</sub>, Cl<sub>2</sub>, Br<sub>2</sub> (l), I<sub>2</sub> (s))
- **Oxides** (basic = metal, acidic = nonmetal)
- **Family reactions**
  - 2K(s) + 2H<sub>2</sub>O(l)  $\rightarrow$  2KOH(aq) + H<sub>2</sub>(g)
  - H<sub>2</sub>(g) + Cl<sub>2</sub>(g)  $\rightarrow$  2HCl(g)

### (15) Lattice Energy

$$U \propto \frac{q_1 q_2}{r_1 + r_2}$$

- Energy **released** (exo.) forming ionic crystal (from gas)
  - (Lattice *dissociation* energy = endothermic)
- max U: 1. ↑ Q+/Q- (ionic charge), 2. ↓ AS (rad.)

| Trend     | Bond  | Enthalpy    | Exception                          |
|-----------|-------|-------------|------------------------------------|
| <b>EA</b> | Form  | Exothermic  | Endo: Be, Mg, noble, EA2 (usually) |
| <b>IE</b> | Break | Endothermic | --                                 |

### (16-17) Electronegativity, Bond Polarity

$$\Delta EN = |EN_A - EN_B|$$

- $EN$  = e- attraction in cov. bond.
- ↑  $EN$  = ↑ bond = ↑ polar = shorter (not always)
- triple > double > single (ranking)

| $\Delta EN$ | Label              |
|-------------|--------------------|
| 0           | pure non cov.      |
| 0.1 – 0.5   | slightly pol. cov. |
| 0.5 – 1.6   | pol. cov.          |
| > 1.6       | ionic              |

### (18) Formal Charge & Resonance

$$FC = (\text{valence } e^-) - [\text{nonbonding } e^- + \# \text{ bonds}]$$

1. Overall 0 charge is better
2. Else –1 on the **more electronegative** atom
- Equivalent resonance → real bonds are **average** (e.g. 1½)

### (19) Lewis Structures (Kelter)

1. Sum valence e- (including +/- for ions)
2. Sum "wants" (octet)
  - H (2), He (2), - Be (4), Al (6), B (6)
  - 3p or below (>8)
3. Sum bonds = (2. - 1.) / 2 (round down)
4. Central atom = min EN (never H)
5. Remaining e- = 1. outside, central,
  - Consider double/trip. bond for central octet / nonzero FC
  - Ions go in brackets with the charge outside
6. FC: overall 0, or 1- on max EN element

### (20) Photoelectric Effect ( $h\nu$ - binding / work)

$$KE = h\nu - \Phi = h(c/\lambda) - \Phi = \frac{1}{2} m_e u^2$$

| Variable  | Value          |
|-----------|----------------|
| $h$       | 6.626e-34 J·s  |
| $m_e$     | 9.11e-31 kg    |
| $hc$      | 1.9864e-25 J·m |
| $2hc/m_e$ | 4.3609e5 m³/s² |

| Have                           | Want                                 | Use                                                                                   |
|--------------------------------|--------------------------------------|---------------------------------------------------------------------------------------|
| $\lambda, \lambda_{thr}$       | $u$                                  | $u = \sqrt{(2hc/m_e)(1/\lambda - 1/\lambda_{thr})}$                                   |
| $\Phi$                         | $\lambda_{thr}$ (max ej. $\lambda$ ) | $\lambda_{thr} = hc/\Phi$                                                             |
| $\nu_{thr}$ or $\lambda_{thr}$ | $\Phi, \Phi / \text{mol}$            | $\Phi = h\nu_{thr} = h(c/\lambda_{thr}),$<br>n.b. $\Phi \times N_A$ (/1e3 for kJ/mol) |
| $\Phi, \lambda$                | $u$                                  | 1. $KE = h(c/\lambda) - \Phi$<br>2. $u = \sqrt{2 KE/m_e}$                             |

### (21) Heisenberg Uncertainty

$$\Delta x \cdot \Delta p \geq \frac{h}{4\pi}, \quad \Delta p = m\Delta u$$

1.  $\Delta p = m \cdot \Delta u$  (kg, m/s)
2.  $\Delta x \geq h/(4\pi \cdot \Delta p)$ 
  - Result is the **minimum** uncertainty
  - Rearrange the same way to solve for  $\Delta u$
  - Keep everything in kg, m, s

### (22) Born-Haber Cycle

$$\Delta H^\circ_f = \Delta H_{sub} + \Sigma IE + \frac{1}{2} BE + EA + U$$

1. List every step with its sign: sublimation (+),  $IE$  (+), bond dissociation (+),  $EA$  (usually –),  $U$  (–)
2. **Halve** the bond dissociation if you need only one atom
3. Chain: elements → gaseous atoms → gaseous ions → solid
4. Set the chain equal to  $\Delta H^\circ_f$
5. Solve for the single unknown
  - Use  $IE_2$  as well when the cation is 2+
  - U runs ions(g) → solid, so it comes out **negative**; report |U| if asked for a positive lattice energy

### (23) Bond Enthalpy

$$\Delta H_{rxn} = \Sigma \Delta H(\text{bonds broken}) - \Sigma \Delta H(\text{bonds formed})$$

1. List bonds broken (reactants) and formed (products), using the balanced coefficients
2.  $\Sigma BE(\text{broken})$ , always positive
3.  $\Sigma BE(\text{formed})$ , subtracted
4.  $\Delta H = \Sigma \text{broken} - \Sigma \text{formed}$ 
  - Bonds unchanged on both sides cancel — skip them
  - Negative  $\Delta H$  = exothermic
  - For "heat released by g of X," convert g → mol and scale