

Lewis Structures (Kelter) — step 0 for (1–8)

- Sum valence e⁻ (= **group #**; add 1 per -, remove 1 per + for ions)
- Sum "wants" (table; period 3+ central may exceed 8)
- Bonds = (wants - valence) ÷ 2 (round down)
- Central atom = lowest EN (**never H**); ions go in brackets, charge outside
- Remaining e⁻ = valence - 2×bonds: outer atoms first, then central; double/triple bond if central lacks octet or nonzero FC
- FC = valence - (lone e⁻ + # bonds): want overall 0, else - on most EN atom

	H	Be	B, Al	C, Si	N, P	O, S	F-I	Noble
Val e	1	2	3	4	5	6	7	8
Wants	2	4	6	8	8	8	8	8

(1–2) Molecular Geometry (VSEPR)

- Central atom BG + LP (double/triple bond or radical e⁻ = 1 group)

e ⁻	Hyb	BG	LP	e ⁻ geo	Mol. geo	Angle	Pol
2	sp	2	0	linear	linear	180°	X
3	sp ²	3	0	tri. planar	tri. planar	120°	X
3	sp ²	2	1	tri. planar	bent	<120°	✓
4	sp ³	4	0	tetra.	tetra.	109.5°	X
4	sp ³	3	1	tetra.	tri. pyr.	<109.5°	✓
4	sp ³	2	2	tetra.	bent	<109.5°	✓
5	sp ³ d	5	0	tri. bipy.	tri. bipy.	90/120°	X
5	sp ³ d	4	1	tri. bipy.	seesaw	90/120°	✓
5	sp ³ d	3	2	tri. bipy.	T-shaped	90, 180°	✓
5	sp ³ d	2	3	tri. bipy.	linear	180°	X
6	sp ³ d ²	6	0	octa.	octa.	90°	X
6	sp ³ d ²	5	1	octa.	sq. pyr.	90°	✓
6	sp ³ d ²	4	2	octa.	sq. planar	90°	X

- Lone pairs squeeze angles (SCl₂ a little < 109.5°) · SO₂, BF₃ tri. planar · PCl₃, PH₃, NH₃ tri. pyr. · CS₂, CO₂, N₂, I₂ linear · SiH₄, CH₄, PO₄³⁻ tetra. · SCl₂, H₂S, H₂O, O₃, ClNO, NH₃ bent · ClF₃ T-shaped · XeF₄ sq. planar

(3–4) Polarity & Dipole Moment

- Dipole** = separation of + / - charge; net dipole moment (μ ≠ 0) = **polar** molecule
- Polar = polar bonds (ΔEN) whose vectors **don't cancel**. Pol column above assumes identical outer atoms — **mixed outer atoms** → **polar** (CH₃Cl, CH₂Cl₂, CHCl₃)
- Nonpolar: same-element diatomic (Br₂), all hydrocarbons, X-row shapes: XeF₄, XeF₆, CCl₄, PBr₅, PCl₅, BeCl₂, CO₂, BF₃, BH₃, SO₂
- Polar: different-element diatomic (CO, IBr, HCl) + lone-pair shapes: BrF₃ (sq. pyr.), NH₃, OF₂, H₂O, SO₂ (bent), NCl₃, NF₃
- ↑ΔEN = ↑μ: CH₃F > CH₃Cl > CH₃Br > CH₃I · NH₃ > PH₃ > AsH₃ (EN falls down group)
- Polar dissolves polar: NH₃ in H₂O ✓, H₂S in NH₃ ✓, CCl₄/SiO₂/O₂/CO₂ in polar X

(5–6) Hybridization of an Atom in a Structure

- Count e⁻ groups **on that atom** (bonded atoms + lone pairs) → table (1): 2 sp, 3 sp², 4 sp³, 5 sp³d, 6 sp³d²
- sp² geometry is trigonal planar (**not** square planar)
- C, 4 single bonds → **sp³** · C with 1 double bond (C=O, C=C, benzene ring) → **sp²** · C with triple or 2 doubles → **sp**
- O in -O-H or ether (2 atoms + 2 LP) → **sp³** · carbonyl =O → **sp²**
- N with 3 atoms + 1 LP (amine) → **sp³**
- Each atom separately: CH₃-CHO → left C sp³, carbonyl C sp²

(7–8) σ and π Bonds: Counting & Concepts

- single** = 1σ · **double** = 1σ + 1π · **triple** = 1σ + 2π
- Count **every** bond, incl. each C-H! σ = # of atom-atom connections; π = extras on doubles/triples
- CH₃-CH=CH-COOH: 11 σ, 2 π · benzene C₆H₆: 12 σ (6 C-C + 6 C-H) + 3 π
- σ: end-to-end overlap, e⁻ density **on** internuclear axis, rotation free, stronger
- π: **side-by-side p-p** overlap, density above & below axis → **restricted rotation** (energy needed to break the p-p interaction) → cis/trans in C=C (1,2-dichloroethene), none around C-C single (1,2-dichloroethane)
- Two 1s (or head-on p+p along bond axis) → σ and σ* · sideways p+p (perpendicular axis) → π and π*
- Benzene delocalized π MOs: e⁻ **free to move around the ring**; all C-C equal

(9–10) Intermolecular Forces

Force	Who has it
Dispersion (London, van der Waals, induced dipole)	ALL molecules; ↑ w/ size & molar mass; straight-chain > branched
Dipole-dipole	polar molecules
H-bond	H bonded directly to N, O, F
Ion-dipole	ion + polar solvent (NaCl in H ₂ O)

- Dispersion only (nonpolar): Br₂, Cl₂, F₂, H₂, CO₂, BCl₃, CF₄, hydrocarbons, noble gases
- Disp + dip-dip: HCl, HBr, SO₂, CHCl₃, CH₃Br, ethers (CH₃OCH₃), CH₃CHO, CH₃CN
- + H-bond: H₂O, NH₃, HF, CH₃OH, CH₃CH₂OH, CH₃NH₂, CH₃COOH, H₂O₂, HOCH₂CH₂OH
- Traps: H-bond **with water** just needs N/O/F (CH₃COOH ✓, CH₃NH₂ ✓; C₂H₆ X; Na⁺ = ion-dipole) · S never H-bonds (H₂S, CH₃SCH₃) · ether O has no H → none with itself · polar **bonds** ≠ polar molecule (CO₂)

(11–12) Rank Vapor Pressure & Boiling Point

- Strength: **ionic > H-bond > dipole-dipole > dispersion**, but size can trump: big nonpolar can beat small polar
- ↑IMF → ↑bp, ↑mp, ↑viscosity, ↑surf. tension, ↑ΔHvap, ↑critical T, ↓ **vapor pressure**
- VP order = reverse of bp order** (lowest bp = highest VP at room T)
- Weak IMF ⇒ low ΔHvap, high VP, low bp


VP ↓	CF ₄ > PF ₅ > BrF ₃ (polar last)
bp ↑	CH ₄ < CH ₃ Cl < CH ₃ OH < RbCl (ionic)
bp (disp. size)	I ₂ > Br ₂ > Cl ₂ > F ₂ · CH ₃ CHO > CH ₃ OCH ₃ > Rn
bp (H-bond)	HF > HI > HBr > HCl
bp (# of OH)	HOCH ₂ CH ₂ OH > ClCH ₂ CH ₂ OH > CH ₃ CH ₂ OH > CH ₃ CH ₂ Cl

(13) Viscosity, Capillary Action, Surface Tension

Surface tension	energy required to stretch/increase surface area by unit amount
Viscosity	resistance of a liquid to flow ; ↑ with IMF, ↓ with ↑T; spherical/compact → lower
Capillary action	liquid flows up a thin tube against gravity = cohesion (like molecules) + adhesion (liquid-glass)

- Concave meniscus (H₂O): adhesion > cohesion; convex (Hg): cohesion > adhesion. Highest viscosity = most H-bonding (HOCH₂CH₂OH)

(14) Phase Diagrams



- Lines = 2 phases coexist · **triple point** = all 3 · **critical point**: above T_c no pressure can liquefy gas → **supercritical fluid**
- ↑T at const P (→): solid → liquid → gas. ↑P at const T (↑): gas → liquid → solid. Normal bp/mp read at 1 atm
- H₂O**: fusion line leans **left** (ice less dense) → ↑P melts ice · **CO₂**: triple pt at 5.1 atm → at 1 atm solid **sublimes** s→g (dry ice)

(15–16) Type of Solid

Type	Held by	Properties	Examples
Ionic	± attraction	hard, brittle, high mp, poor cond.	NaCl, AgCl, CaF ₂ , MgO, CaCO ₃
Molecular	IMFs	soft, low mp, poor cond.	ice H ₂ O, CO ₂ , I ₂ , sucrose
Metallic (atomic)	e ⁻ sea	low-high mp, good cond.	Au, Cu, Fe, Na
Network covalent (atomic)	covalent bonds	v. hard, v. high mp, poor cond.	diamond, quartz SiO ₂ , graphite, SiC
Nonbonding (atomic)	dispersion	v. low mp	noble gases: Ar(s)
Amorphous	no long-range order	melts over a range	glass, rubber, plastic, butter, wax
Ceramic	inorganic nonmetal; powder + H ₂ O + heat	brittle, heat-resistant	clay/kaolinite, porcelain, Al ₂ O ₃

- Graphite sp² sheets: conducts along sheets, slippery · diamond sp³: insulator · fcc packing = most space-efficient (12 neighbors)

(17) Band Theory & Doping

- Bands = MOs delocalized over the whole crystal: **valence band** = **bonding MOs**, **conduction band** = **antibonding MOs**, **band gap** between
- Conductor** (metal): bands continuous/overlap · **Semiconductor** (metalloids Si, Ge): small gap, e⁻ promotable · **Insulator**: **LARGE gap**
- Doping** = add impurity to ↑ conductivity:

	Dopant	Carrier
n-type	Group V/5A (P, As)	extra e ⁻ in conduction band (- charges)
p-type	Group III/3A (B, Al, Ga, In)	holes in valence band (e ⁻ hop between holes)
p-n junction	—	diode : current flows one way

(18) Heat of Solution & Hydration

$$\Delta H_{\text{solution}} = \Delta H_{\text{hydration}} - \Delta H_{\text{lattice energy}}$$

- ΔHsolution**: heat absorbed/released on dissolving = ΔH(solvent-solvent) + ΔH(solute-solute) + ΔH(solvent-solute)
- ΔHhydration**: heat **released** when 1 mol **gaseous ions** dissolves in water (forms ion-dipole; always exo). ΔHsolute = -ΔHlattice
- Highest ΔHhydr**: smallest ion + highest charge: Al³⁺ > Mg²⁺ > Na⁺ > K⁺ · F⁻ > Cl⁻
- ΔHsoln **exo (-)**: ΔHhydration > ΔHsolute · **endo (+)**: solute > hydration ≈ 0: comparable

(19) Saturation; Like Dissolves Like

Unsaturated	less than equilibrium amount — more can dissolve
Saturated	max amount at that T; dissolved $\rightleftharpoons$ solid dynamic equilibrium
Supersaturated	more than equilibrium amount; unstable — seed crystal/scratch $\rightarrow$ excess crystallizes out

- Solution forms if solute-solvent attraction **comparable** to solute-solute + solvent-solvent · polar/ionic $\leftrightarrow$ polar (H₂O, ethanol, NH₃); nonpolar $\leftrightarrow$ nonpolar (hexane, CCl₄, benzene)
- Most soluble in ethanol = most H-bonds (ethylene glycol); in hexane = longest chain (1-pentanol); all **nitrates** soluble

(20) Solubility vs P & T; When Bubbles/Boiling

$\uparrow$ gives	Gas	Solid / liquid
$\uparrow$ Pressure	$\uparrow$ solubility (Henry)	no effect
$\uparrow$ Temperature	$\downarrow$ solubility	$\uparrow$ solubility

- Dissolve the most gas: **low T + high P** (cold soda holds CO₂)
- Liquid **boils** when its vapor pressure = **external (atm) pressure**; gas **bubbles out** of solution when its partial (vapor) pressure > atm pressure

(21) Molarity (and Molality)

$M = \frac{\text{mol solute}}{\text{L solution}}$	$m = \frac{\text{mol solute}}{\text{kg solvent}}$
---	---

- mol = M $\times$ V(L) · V(L) = mol $\div$ M · mL $\div$ 1000 = L · g $\leftrightarrow$ mol via molar mass · dilution: M₁V₁ = M₂V₂
- 0.250 L of 0.500 M $\rightarrow$ 0.125 mol · need 0.300 mol of 0.450 M $\rightarrow$ 0.667 L = 667 mL
- Molality m uses **kg of solvent**, not solution — for $\Delta T_f/\Delta T_b$. Only molarity changes with T (volume swells); m, mass %, χ don't

(22) Percent Concentration (% w/v)

$\% w/v = \frac{\text{g solute}}{100 \text{ mL soln}} \Rightarrow g = \frac{\% \times \text{mL}}{100}$
--

- 1.5% (w/v), 250 mL $\rightarrow$ 1.5 $\times$ 250/100 = **3.75 g** (n.b. the summary's "31.25 g" contradicts its own formula — trust g = % $\times$ mL / 100)
- mL from % w/v: 16.6 g $\div$ 0.0450 g/mL = 369 mL
- % m/m** = g solute / (g solute + g solvent) $\times$ 100 · g water = g soln $\times$ (1 - %/100): 300 g of 9.71% KBr $\rightarrow$ 271 g water
- % v/v** = vol component / total vol $\times$ 100

(23) Henry's Law (C_{gas} = k_H·P_{gas})

$C_{\text{gas}} = k_H P_{\text{gas}}$	$k_H = C_{\text{gas}} / P_{\text{gas}}$
---------------------------------------	---

- k_H = Henry's constant = "solubility constant of a gas" (M/atm); C = molar solubility
- mmHg $\rightarrow$ atm: $\div$ **760**
- k_H = C/P: 1.3 M at 0.022 atm $\rightarrow$ **59 M/atm**
- 4.7e-4 M at 522 mmHg $\rightarrow$ k_H = 4.7e-4 $\div$ (522/760) = **6.8e-4 M/atm**

(24) Osmosis

- Osmosis**: flow of **solvent** (water) across a semipermeable membrane from **low solute conc.** $\rightarrow$ **high solute conc.** (dilute $\rightarrow$ concentrated)
- Semipermeable membrane**: passes solvent, **blocks solute** (ions) — it does *not* stop water
- Hyperosmotic** = more solute: water flows **into** it (cell in it shrinks/crenates)
- Hypotonic** = less solute: water flows **out** into the stronger side (cell in it swells/bursts)
- Isosmotic** = equal: no net flow (isotonic saline = 0.9% w/v NaCl) · n = pressure needed to **stop** osmotic flow

(25) WORD: MO Theory — Bond Order & Magnetism


B ₂	C ₂	N ₂	O ₂ filled (F ₂ , Ne ₂ , NO, OF ⁺ , Cl ₂ ...)
σ_{2p}^+ _____			σ_{2p}^+ _____
π_{2p}^+ _____			π_{2p}^+ _____
σ_{2p} _____			π_{2p} _____
π_{2p} _____			σ_{2p} _____
σ_{2s}^+ _____			σ_{2s}^+ _____
σ_{2s} _____			σ_{2s} _____

- Valence e⁻ total: **add** 1 per - charge, **remove** 1 per +
- Fill low $\rightarrow$ high, 2 per orbital, Hund across the n pair
- BO = (bonding e⁻ - antibonding e⁻) / 2**
- Any unpaired e⁻ $\rightarrow$ **paramagnetic**; all paired $\rightarrow$ **diamagnetic**
- BO > 0 $\rightarrow$ stable, can form (H₂ 1 $\checkmark$; He₂ 0 $\times$ can't exist); **higher BO = stronger & shorter** bond (N₂⁺ weaker/longer than N₂). σ_{2s} + σ_{2s}^* cancel — you may count 2p only

Sp.	e ⁻	BO	Mag	Sp.	e ⁻	BO	Mag
N ₂	10	3	dia	O ₂	12	2	para (2 $\uparrow$)
N ₂ ⁺	9	2.5	para	O ₂ ⁺	11	2.5	para
N ₂ ⁻	11	2.5	para	O ₂ ⁻	13	1.5	para
F ₂ , Cl ₂	14	1	dia	Cl ₂ ⁺	13	1.5	para
B ₂	6	1	para	Cl ₂ ⁻	15	0.5	para

- Heteronuclear (NO, OF⁺) $\rightarrow$ the O₂-type (right) ordering (more EN atom sits lower) · period 3 (Cl₂, Na₂) $\rightarrow$ same shapes, n = 3

(26) WORD: Phase-Change Heat (gas $\rightleftharpoons$ solid)



- Sketch the path; skip segments you don't cross (65 °C liq $\rightarrow$ 115 °C gas: heat liq 65 $\rightarrow$ 100, vaporize, heat gas 100 $\rightarrow$ 115)
- Sloped (one phase): **q = m(g)·Cs·ΔT** — flat (phase change, T constant): **q = n(mol)·ΔH**
- Sum; convert J $\rightarrow$ kJ. Cooling path (gas $\rightarrow$ solid): every q **negative** ($\Delta H_{\text{cond}} = -\Delta H_{\text{vap}}$, $\Delta H_{\text{freeze}} = -\Delta H_{\text{fus}}$)
 - H₂O: C(ice) 2.09, C(liq) 4.184, C(steam) $\approx$ 2.0 J/g·°C; ΔH_{fus} 6.02, ΔH_{vap} 40.7 kJ/mol; MM 18.02
 - 36.0 g H₂O, 65 °C liq $\rightarrow$ 115 °C gas: 36 $\times$ 4.184 $\times$ 35 = 5.27 kJ; (36/18.02) $\times$ 40.7 = 81.3 kJ; 36 $\times$ 2.01 $\times$ 15 = 1.09 kJ $\rightarrow$ **87.7 kJ**
 - Cooling: 125 g benzene (78.11) gas 425 K $\rightarrow$ liq 335 K (bp 353 K, mp 279 K — never freezes): 125 $\times$ 1.06 $\times$ 72 = 9.5 kJ; (125/78.11) $\times$ 33.9 = 54.3 kJ; 125 $\times$ 1.73 $\times$ 18 = 3.9 kJ $\rightarrow$ **67.7 kJ removed**
 - Endo (+): melt, vaporize, sublime · Exo (-): freeze, condense, deposit · $\Delta H_{\text{sub}} = \Delta H_{\text{fus}} + \Delta H_{\text{vap}}$

(27) WORD: Raoult's Law (VP Lowering)

$$P_{\text{soln}} = \chi_{\text{solvent}} P_{\text{solvent}}^{\circ} \quad \Delta P = \chi_{\text{solute}} P_{\text{solvent}}^{\circ}$$
$$\chi_{\text{solvent}} = \frac{\text{mol}_{\text{solvent}}}{\text{mol}_{\text{solvent}} + i \cdot \text{mol}_{\text{solute}}}$$

- mol solvent = g $\div$ MM; mol solute = g $\div$ MM (**$\times$ i if ionic**)
- $\chi(\text{solvent}) = \text{mol solvent} \div \text{total mol}$ — always < 1
- P(soln) = $\chi(\text{solvent}) \times P^{\circ}$; lowering $\Delta P = P^{\circ} - P(\text{soln}) = \chi(\text{solute}) \cdot P^{\circ}$
 - 75.0 g citric acid (192.12) + 420 g H₂O, P^o 71.93 mmHg: 0.390 & 23.31 mol $\rightarrow \chi = 0.9835 \rightarrow$ **70.7 mmHg**
 - 200 g sucrose (342.3) + 350 g H₂O, P^o 17.5: 0.584 & 19.42 mol $\rightarrow \chi = 0.9708 \rightarrow$ **17.0 mmHg**
 - Both volatile: P(total) = $\chi_A \cdot P^{\circ}_A + \chi_B \cdot P^{\circ}_B$

(28) WORD: Freezing $\downarrow$ / Boiling $\uparrow$ (van't Hoff)

- $\Delta T_f = i K_f m$ $\Delta T_b = i K_b m$
- m = mol solute / **kg solvent** · H₂O: K_f 1.86, K_b 0.512 °C/m · T_f = T^o - ΔT_f , T_b = T^o + ΔT_b
 - i** = particles per formula unit: nonelectrolytes (glucose, sucrose, urea) 1 · NaCl, KCl, NaOH, NH₄Cl 2 · CaCl₂, (NH₄)₂SO₄ 3 · Al(NO₃)₃, FeCl₃ 4 (measured i slightly less — ion pairing)
 - Find i: **i = $\Delta T_b / (K_b \cdot m)$** — 1.83 m boils 102.5 °C: 2.5/(0.512 $\times$ 1.83) = **2.7**
 - Molar mass from fp: $\Delta T_f \rightarrow m = \Delta T_f / (i \cdot K_f) \rightarrow \text{mol} = m \times \text{kg} \rightarrow$ **MM = g/mol**: 20.0 g per 500 g benzene (T^o 5.444, K_f 5.12) freezes 3.77 °C: ΔT 1.674 $\rightarrow$ 0.327 m $\rightarrow$ 0.163 mol $\rightarrow$ **$\approx$ 120 g/mol**
 - Lowest fp, equal grams = most mol $\times$ i (small MM wins: NaOH > KCl, CaCl₂)

(29) WORD: Osmotic Pressure $\rightarrow$ Molar Mass

- $\pi = i M R T \Rightarrow M = \frac{\pi}{i R T}$
- R = **0.08206** L·atm/mol·K · T in **K** (+273.15) · n in **atm** (torr $\div$ 760) · i = 1 nonelectrolyte
 - 1. M = n / (iRT)
 - 2. mol = M $\times$ V(L)
 - 3. **MM = g solute $\div$ mol**
 - 15.2 **mg** in 150.0 mL, n 8.44 torr, 25 °C: M = (8.44/760)/(0.08206 $\times$ 298) = 4.54e-4 $\rightarrow$ 6.81e-5 mol $\rightarrow$ 0.0152/6.81e-5 = **223 g/mol**
 - n $\rightarrow$ freezing pt: M from n $\rightarrow$ per 1 L: g solute = M $\times$ MM; g soln = 1000 $\times$ density; kg solvent = (g soln - g solute)/1000 $\rightarrow$ molality $\rightarrow$ $\Delta T_f = i \cdot K_f \cdot m$