

(1) EM Spectrum							
Ranges	γ	X	UV	vis	IR	micro	radio
Energy (E)	↑						↓
Frequency (ν)	↑						↓
Wavelength (λ)	↓						↑
Wavelength (λ)	<0.01	0.01-10	10-400	400-700	700-1e6	1e6-9	>1e9

• violet • indigo • blue • green • yellow • orange • red

(2) $c = \lambda\nu$ (Speed of light = wavelength × frequency)

c	ν (Hz)	kHz	MHz	GHz
λ (m)	2.9979e8 (m/s)	2.9979e5 (m·kHz)	2.9979e2 (m·MHz)	2.9979e-1 (m·GHz)
nm	2.9979e17 (nm·Hz)	2.9979e14 (nm·kHz)	2.9979e11 (nm·MHz)	2.9979e8 (nm·GHz)

(3) $E = h\nu = h(c/\lambda)$ (Energy = Planck × frequency)

hc , h	λ (m)	λ (nm)	ν (Hz)
E (J / photon)	1.9864e-25 (J·m)	1.9864e-16 (J·nm)	6.626e-34 (J·s)
J / mol	1.1962e-1 (J·m/mol)	1.1962e8 (J·nm/mol)	3.9902e-10 (J·s/mol)
kJ / mol	1.1962e-4 (kJ·m/mol)	1.1962e5 (kJ·nm/mol)	3.9902e-13 (kJ·s/mol)

• Grams → mol → ×6.022e23 for photon count

(4) Quantum Principles

Maxwell	Light is electromagnetic radiation
Planck	Energy emitted only in fixed amounts (quanta) (black-body radiation)
Einstein	Different light has different amounts of fixed energy (photons) (photoelectric effect)
Bohr	Electrons are quantum (line spectra); Rydberg eq.
De Broglie	Electrons are particle and wave (5)
Schrodinger	Wave equation (energy and spatial prob. of e-) (n, l, m_l, m_s)
Heisenberg	Position and velocity not both knowable

$$E = h\nu = h(c/\lambda) = R(1/n_f^2 - 1/n_i^2)$$

$$n_i = \sqrt{1/(1/n_f^2 - hc/(R\lambda))}; \quad n_f = \sqrt{1/(1/n_i^2 + hc/(R\lambda))}$$

Variable	Value	Variable	Value
h	6.626e-34 J·s	h/R	3.0394e-16 s
c	2.9979e8 m/s	hc	1.9864e-25 J·m
R	2.18e-18 J	hc/R	9.1120e-8 m

(5) $\lambda = h/(m \cdot v)$ (De Broglie)

- 1 nm = 1e-9 m
- 1 e- = 9.11e-31 kg

h	u (m/s)	km/s
m (kg)	6.626e-34 (J·s)	6.626e-37 (kJ·s)
g	6.626e-31 (mJ·s)	6.626e-34 (J·s)
amu	3.9892e-7 (J·s·amu/kg)	3.9892e-10 (kJ·s·amu/kg)

(6) Electron Configuration (Aufbau, Pauli, Hund)

- 1s 2s 2p 3s 3p 4s 3d 4p 5s 4d 5p 6s 4f 5d 6p 7s 5f 6d 7p
- Round-robin sublevel before double occupaton (Hund)
- Two e- per orbital (Pauli's exclusion)
- n^2 orbitals, $2n^2$ e- per shell

Symbol	Name	Value
n	Principal energy level (shell)	(1, 2, 3, ...)
l	Sublevel = subshell (shape)	(0, ..., n - 1)
m_l	Orbital (orientations)	(-l, ..., l)
m_s	Spin (2 per orbital)	$\pm 1/2$

Sublevel	l	m_l	# orbitals	# e-
s	0	0	1	2
p	1	± 1	3	6
d	2	± 2	5	10
f	3	± 3	7	14

Element	EC	Element	EC
Cr	[Ar]4s ¹ 3d ⁵	Ag	[Kr]5s ¹ 4d ¹⁰
Mo	[Kr]5s ¹ 4d ⁵	Au	[Xe]6s ¹ 4f ¹⁴ 5d ¹⁰
Cu	[Ar]4s ¹ 3d ¹⁰	Pd	[Kr]4d ¹⁰ (no 5s)

(7-8) Reading EC, Anion, Cation

EC → element	add every superscript to get Z
Paramagnetic	at least 1 unpaired val. e- (else dia.)
Ground state	follows Aufbau (else excited)
Valence electrons	highest / outer shell (else core)
Anion	count val. e- needed until noble gas
Cation	remove val. e- until noble gas
Tx metal	remove s before d
Isoelectric	same EC (e.g. Na+, Al ³⁺ , F-, N ³⁻ , Ne)

(9-10) Penetration, Shielding, Z_{eff}

- ↑ penetration (core electrons) = ↓ shielding = ↓ energy
- **Energy** s < p < d < f (same shell)
- **Core** e- shield valence e-
 $Z_{eff} \approx Z - (\text{core } e^-)$
- Positive charge felt by e- toward nucleus
- Core electrons = non-valence

(11-13) Trends

	Trend	Top	Bot	Left	Right
(11) Atomic size (radius)	↘ (Cs; f > p)	↓	↑	↑	↓
(12) Ionization energy (IE)	↗ (He)	↑	↓	↓	↑
(13) Electron affinity (EA)	↗ (Cl)	↑	↓	↓	↑
Z_{eff}	↘	↓	↑	↓	↑
Metallic	↗ (Fr)	↓	↑	↑	↓
Nonmetallic	↗ (F)	↑	↓	↓	↑
Electronegativity (EN)	↗ (F)	↑	↓	↓	↑

- **Atomic size** (cation < neutral < anion)

Trend	Bond	Enthalpy	Exception
IE	break	endothermic	--
EA	form	exothermic	Endo: Be, Mg, noble, EA2 (usually)

(14) Families & Diagonal Rule

Family	Name	Reactivity	e-	Charge
1A	alkali metals	most reactive metals (downward)	lose 1	1+
2A	alkaline earth	reactive, less than 1A	lose 2	2+
7A	halogens	reactive nonmetals (upward)	gain 1	1-
8A	noble gases	inert	—	0

- **Diagonal rule** (Li~Mg, Be~Al, B~Si)
- **Lanthanide** (58-71), **Actinide** (90-103)
- **Diatomic** (H₂, N₂, O₂, F₂, Cl₂, Br₂ (l), I₂ (s))
- **Oxides** (basic = metal, acidic = nonmetal)

HW-only gaps — off the exam list

Formula from charges — 1A 1+ · 2A 2+ · 3A 3+ · 5A 3- · 6A 2- · 7A 1-; criss-cross and reduce

- Same-group swap keeps the pattern: BaS → **CaO**
- 2A + 3F₂ → 2AF₃ ⇒ A is 3+ (**Al**) · 3Li + Z → Li₃Z ⇒ Z is 3-, so Mg gives **Mg₃Z₂**

Bond-type ID — ionic **and** covalent together = a salt of a **polyatomic ion** (NH₄NO₃, K₂SO₄); all-nonmetal = covalent only

Family reactions

- 2K(s) + 2H₂O(l) → 2KOH(aq) + H₂(g)
- H₂(g) + Cl₂(g) → 2HCl(g)

(15) Lattice Energy

Energy **released** when gaseous ions form the solid crystal.

$$U \propto \frac{q_1 q_2}{r_1 + r_2}$$

1. Compare charges first
 2. If charges tie, smaller ions win
- $\uparrow$ charge $\rightarrow \uparrow$ **U** (dominant) • $\uparrow$ ionic radius $\rightarrow \downarrow$ U

(16) Electronegativity

EN = an atom's ability to attract the electrons shared in a bond.

$$\Delta EN = |EN_A - EN_B|$$

- Read EN off the chart — $EN \uparrow$ up and to the right, F highest, noble gases excluded

(17) Bond Polarity

1. Take the absolute difference (ΔEN)
2. Classify

~ 0	nonpolar covalent
0.4 – 1.9	polar covalent
≥ 2.0	ionic

- **"Most polar"** $\rightarrow \Delta EN$ **alone** (largest difference wins)
- **"Strongest / shortest bond"** $\rightarrow$ **bond order FIRST:** triple > double > single. ΔEN only breaks ties between equal orders — a C $\equiv$ N beats a high- ΔEN Si=O
- Larger $\Delta EN \rightarrow$ stronger, shorter bond, but the guide flags this as *not always true*

(18) Formal Charge & Resonance

$$FC = (\text{valence } e^-) - [\text{nonbonding } e^- + \# \text{ bonds}]$$

1. FC on every atom
 2. Check ΣFC = the overall charge
 3. Pick the structure with FC s **nearest 0**
 4. Put any -1 on the **more electronegative** atom
- Equivalent resonance forms $\rightarrow$ real bonds are the **average** (e.g. $1\frac{1}{2}$)

(19) Lewis Structures x4

1. Total valence e^- — $+1$ per negative charge, -1 per positive
2. Central atom = least electronegative, never H
3. Single bonds out to every atom
4. Fill outer octets with lone pairs (H takes 2)
5. Leftover e^- onto the central atom
6. Central below 8 $\rightarrow$ pull an outer lone pair into a double/triple bond, **unless** Al/B/Be
7. Check formal charges — see (18)

Four structures — one of each type:

(a) Normal octet	8 valence e^- per atom (H wants 2) — H_2O , CH_4 , NH_3 , CO_2 , CH_2Cl_2 , CH_3CH_2Cl
(b) Incomplete octet	Al, B, Be only — content below 8, do not add double bonds to fix. $BF_3/BCl_3/AlI_3$ leave 6 e^- on the center; $BeCl_2$ leaves 4
(c) Expanded octet	Central atom period 3 or below (S, P, Cl, Xe, As, I) uses empty d orbitals for >8 — $AsCl_5$, XeI_2 , SO_4^{2-}
(d) Organic	$-OH$ alcohol • $C=O$ aldehyde/ketone • $-COOH$ carboxylic acid • $-NH_2$ amine

- Every C makes 4 bonds; a $-COOH$ carbon carries one **$=O$** and one **$-O-H$** ; every O gets 2 lone pairs
- Ions go in brackets with the charge outside
- If the all-single-bond form leaves the center with a nonzero FC , add double bonds to zero it (e.g. two S=O in SO_4^{2-})

(20) Photoelectric Effect ($h\nu$ - binding / work)

$$KE = h\nu - \Phi = h(c/\lambda) - \Phi = \frac{1}{2}m_e u^2$$

Variable	Value
h	6.626e-34 J·s
m_e	9.11e-31 kg
hc	1.9864e-25 J·m
$2hc/m_e$	4.3609e5 m ³ /s ²

Have	Want	Use
λ, λ_{thr}	u	$u = \sqrt{(2hc/m_e)(1/\lambda - 1/\lambda_{thr})}$
Φ	λ_{thr} (max ej. λ)	$\lambda_{thr} = hc/\Phi$
ν_{thr} or λ_{thr}	$\Phi, \Phi / \text{mol}$	$\Phi = h\nu_{thr} = h(c/\lambda_{thr})$, n.b. $\Phi \times N_A$ (/1e3 for kJ/mol)
Φ, λ	u	1. $KE = h(c/\lambda) - \Phi$ 2. $u = \sqrt{2 KE/m_e}$

(21) Heisenberg Uncertainty

$$\Delta x \cdot \Delta p \geq \frac{h}{4\pi}, \quad \Delta p = m\Delta u$$

1. $\Delta p = m \cdot \Delta u$ (kg, m/s)
 2. $\Delta x \geq h/(4\pi \cdot \Delta p)$
- Result is the **minimum** uncertainty
 - Rearrange the same way to solve for Δu
 - Keep everything in kg, m, s

(22) Born-Haber Cycle

$$\Delta H^\circ_f = \Delta H_{sub} + \Sigma IE + \frac{1}{2}BE + EA + U$$

1. List every step with its sign: sublimation (+), IE (+), bond dissociation (+), EA (usually $-$), U ($-$)
 2. **Halve** the bond dissociation if you need only one atom
 3. Chain: elements $\rightarrow$ gaseous atoms $\rightarrow$ gaseous ions $\rightarrow$ solid
 4. Set the chain equal to ΔH°_f
 5. Solve for the single unknown
- Use IE_2 as well when the cation is $2+$
 - U runs ions(g) $\rightarrow$ solid, so it comes out **negative**; report |U| if asked for a positive lattice energy

(23) Bond Enthalpy

$$\Delta H_{rxn} = \Sigma \Delta H(\text{bonds broken}) - \Sigma \Delta H(\text{bonds formed})$$

1. List bonds broken (reactants) and formed (products), using the balanced coefficients
 2. $\Sigma BE(\text{broken})$, always positive
 3. $\Sigma BE(\text{formed})$, subtracted
 4. $\Delta H = \Sigma \text{broken} - \Sigma \text{formed}$
- Bonds unchanged on both sides cancel — skip them
 - Negative ΔH = exothermic
 - For "heat released by g of X ," convert $g \rightarrow$ mol and scale