

Tautologies and Conditional Proofs

Announcements

- Quiz 1: today (20-min; noon - 11:59 PM)
- HW 1: next Wed (probs assigned this week)
- No class next Mon (MLK)
- OH next week: TW 4:30 PM - 6:00 PM (Zoom)

Contents

① Some Remarks on Logical Connectives

② Conditional Proofs

Some Remarks on Logical Connectives

Parentheses

$$a + (b \times c) + d$$

The order of priority of the logical connectives (from highest to lowest):

$$\neg, \wedge, \vee, \Rightarrow, \Leftrightarrow$$

Examples.

$$\neg P \wedge \neg Q \quad \text{means} \quad (\neg P) \wedge (\neg Q)$$

$$P \wedge Q \vee R \quad \text{means} \quad (P \wedge Q) \vee R$$

$$P \wedge Q \Rightarrow P \vee Q \quad \text{means} \quad (P \wedge Q) \Rightarrow (P \vee Q)$$

$$P \Rightarrow (Q \Rightarrow R) \Leftrightarrow P \wedge Q \Rightarrow R \quad \text{means} \quad [P \Rightarrow (Q \Rightarrow R)] \Leftrightarrow [(P \wedge Q) \Rightarrow R]$$

-
- Economical writing by omitting some parentheses
 - Judicious inclusion of some dispensable parenthesis may enhance readability

Associativity of Conditionals

The logical connective \Rightarrow is not associative, that is,

$$P \Rightarrow (Q \Rightarrow R) \quad \text{and} \quad (P \Rightarrow Q) \Rightarrow R$$

are logically inequivalent. Furthermore, neither of the two is logically equivalent to

$$(P \Rightarrow Q) \wedge (Q \Rightarrow R)$$

For further discussions, see Exercise 11 and Remarks 2.18 and 2.19.

However, people commonly use

$$P \Rightarrow Q \Rightarrow R$$

to mean

$$(P \Rightarrow Q) \wedge (Q \Rightarrow R)$$

Associativity of Biconditionals

The logical connective \Leftrightarrow is associative, that is,

$$P \Leftrightarrow (Q \Leftrightarrow R) \quad \text{and} \quad (P \Leftrightarrow Q) \Leftrightarrow R$$

are logically equivalent. However, neither of the two is logically equivalent to

$$(P \Leftrightarrow Q) \wedge (Q \Leftrightarrow R)$$

For further discussions, see Exercise 12 and Remark 2.20.

However, people commonly use

$$P \Leftrightarrow Q \Leftrightarrow R$$

to mean


$$(P \Leftrightarrow Q) \wedge (Q \Leftrightarrow R)$$

Tautologies

In logic, a *tautology* is a sentence which is true under any possible truth values of its propositional variables.

Examples.

- $P \vee \neg P$
- $[\neg(P \wedge Q)] \Leftrightarrow [\neg P \vee \neg Q]$
- $(P \wedge Q) \Rightarrow P$
- $P \Rightarrow (P \vee Q)$
- If $R \equiv S$, then the sentence $R \Leftrightarrow S$ is a tautology. (Why?)



P	$\neg P$	$P \vee \neg P$
T	F	T
F	T	T

Exercise. Construct a tautology using three sentences P , Q , and R .

Conditional Proofs

Conditional Proofs

always true, i.e., is a tautology.

Conditional Proofs

To show that $A \Rightarrow B$ is true, it suffices to consider the case where A is true and to show that in this case, B must also be true. This approach is known as the method of *conditional proof*.

Template of Conditional Proof. To show $A \Rightarrow B$ is a tautology:

A1: Suppose that A is true.

Work to show that B is true under A1.

Discharging A1, we see that $A \Rightarrow B$ is true under no assumptions.
Therefore $A \Rightarrow B$ is a tautology.

Conditional Proofs (cont')

$Q \Rightarrow (P \vee Q)$ is also
a tautology.

Example

Use the method of conditional proof to explain in words why

$$\underbrace{P}_{\text{Antecedent}} \Rightarrow \underbrace{(P \vee Q)}_{\text{Consequent}}$$

is a tautology. (Do not use cases. Be explicit about discharging assumptions.)

Solution.

A1: Suppose P is true.

Then $P \vee Q$ is true.

We have shown that $P \vee Q$ is true under the assumption A1 that P is true.

Discharging A1, we see that $P \Rightarrow (P \vee Q)$ is true under no assumptions.

Therefore $P \Rightarrow (P \vee Q)$ is a tautology, because we have shown that it is true under no assumptions on the truth values of P and Q . \square

Modus Ponens

The following is useful in proofs.

Modus Ponens

If $P \Rightarrow Q$ is true and P is also true, then Q must be true. This rule of inference is called *modus ponendo ponens* or, more commonly, *modus ponens*.

Exercise 1

Use the method of conditional proof to explain in words why the sentence

$$[(P \Rightarrow Q) \wedge (Q \Rightarrow R)] \Rightarrow (P \Rightarrow R)$$

is a tautology. Be explicit about discharging assumptions.

Solution

A1: Suppose $\overbrace{(P \Rightarrow Q) \wedge (Q \Rightarrow R)}^{A_1}$ is true.

We wish to show that $\underbrace{P \Rightarrow R}_{C_1}$ is true.

A2: Suppose P is true.

We wish to show that R is true.

From A1, it follows that $P \Rightarrow Q$ is true.

From this and A2, we see that Q is true, by modus ponens.

From A1 again, it follows that $Q \Rightarrow R$ is true.

From this and the fact that Q is true, we see that R is true, by modus ^{ponens.}

We have shown that $P \Rightarrow R$ is true under A1 and A2.

Discharging A2, we see that $P \Rightarrow R$ is true under A1 alone.

Discharging A1, we see that $\overbrace{A_1 \Rightarrow C_1}^{A_1 \Rightarrow C_1}$ is true under no assumptions. So it is a tautology.

Exercise 2

Use the method of conditional proof to explain in words why the sentence

$$[P \Rightarrow (Q \Rightarrow R)] \Rightarrow [(P \Rightarrow Q) \Rightarrow (P \Rightarrow R)]$$

is a tautology. Be explicit about discharging assumptions.

Homework (1/14; due Wed 1/19)

Section 2: # 8, 9, 10, 15, 17