

Multidimensional Rootfinding

Root finding (Scalar)

- Given: $f: \mathbb{R} \rightarrow \mathbb{R}$
- Want: $r \in \mathbb{R}$ s.t. $f(r) = 0$

\uparrow
a root or zero
of f

Iteration methods

① Fixed point iteration


$$|g(r) = r \quad (g(x) = x - f(x))$$

- $\begin{cases} \text{do initial guess} \\ x_{k+1} = g(x_k), \quad k=0, 1, 2, \dots \end{cases}$
- converges if $|g'(r)| < 1$

- linear convergence ($\epsilon_{k+1} \approx g'(r) \epsilon_k$)

② Newton's method

- Idea: Replace f by a simpler function and find its root.




- $\begin{cases} \text{do initial guess} \\ x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)}, \quad k=0, 1, 2, \dots \end{cases}$
- form. for x -intercept of t -line.

- quad. convergence:

$$\epsilon_{k+1} \approx \frac{f''(r)}{2f'(r)} \epsilon_k^2$$

③ Secant method

- Idea: Replace f by a simpler function and look for its root.



- x_{-1}, x_0 initial guesses

$$\begin{cases} x_{k+1} = x_k - \frac{f(x_k)}{(\text{slope})} = x_k - \frac{x_k - x_{k-1}}{f(x_k) - f(x_{k-1})} f(x_k), \quad k=0, 1, 2, \dots \end{cases}$$

- Superlinear conv. (but slower than quad.)

$$\epsilon_{k+1} \approx \left(\frac{f''(r)}{2f'(r)} \right)^{\alpha-1} \epsilon_k^\alpha \quad \text{where}$$

$$\alpha = \frac{1+\sqrt{5}}{2} = 1.618 \dots$$

(the golden ratio)

Example (Computation of \sqrt{a} using Newton) Assume $a > 0$.

• Note that \sqrt{a} is a root of $\underline{f(x) = x^2 - a}$.

• Newton's iter. formula:

$$x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)}$$

$$= x_k - \frac{x_k^2 - a}{2x_k}$$

$$= x_k - \frac{1}{2}x_k + \frac{a}{2x_k} = \frac{1}{2}x_k + \frac{a}{2x_k}.$$

$$f(x) = 0$$

$$x^2 - a = 0$$

$$x^2 = a$$

$$x = \pm\sqrt{a}$$

$$\downarrow$$
$$f'(x) = 2x$$

Newton's Method for Nonlinear Systems

Multidimensional Rootfinding Problem

Rootfinding Problem: Vector Version

Given a continuous vector-valued function $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$, find a vector $\mathbf{r} \in \mathbb{R}^n$ such that $\mathbf{f}(\mathbf{r}) = \mathbf{0}$.

The rootfinding problem $\mathbf{f}(\mathbf{x}) = \mathbf{0}$ is equivalent to solving the *nonlinear* system of n scalar equations in n unknowns:

$$\vec{f} = \begin{bmatrix} f_1 \\ f_2 \\ \vdots \\ f_n \end{bmatrix} \quad \left\{ \begin{array}{l} f_1(x_1, \dots, x_n) = 0, \\ f_2(x_1, \dots, x_n) = 0, \\ \vdots \\ f_n(x_1, \dots, x_n) = 0. \end{array} \right.$$

f_1, \dots, f_n are scalar functions.

Multidimensional Taylor Series

If f is differentiable, we can write

$$f(\underbrace{\mathbf{x}}_{\text{center}}) + \underbrace{(\mathbf{h})}_{\text{perturbation}} = f(\mathbf{x}) + \boxed{\mathbf{J}(\mathbf{x})}\mathbf{h} + O(\|\mathbf{h}\|^2),$$

where \mathbf{J} is the **Jacobian matrix** of f

$$\mathbf{J}(\mathbf{x}) = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} & \cdots & \frac{\partial f_1}{\partial x_n} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} & \cdots & \frac{\partial f_2}{\partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial x_1} & \frac{\partial f_n}{\partial x_2} & \cdots & \frac{\partial f_n}{\partial x_n} \end{bmatrix} = \left[\frac{\partial f_i}{\partial x_j} \right]_{i,j=1,\dots,n}.$$

$$\text{cf) } f(x+h) = f(x) + \boxed{f'(x)}h + O(h^2)$$

“multidimensional analog of derivative.”

$$\vec{x} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}, \quad \vec{h} = \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix}$$

- The first two terms $f(\mathbf{x}) + \mathbf{J}(\mathbf{x})\mathbf{h}$ is the “linear approximation” of f near \mathbf{x} .
- If f is actually linear, i.e., $f(\mathbf{x}) = A\mathbf{x} - \mathbf{b}$, then the Jacobian matrix is the coefficient matrix A and the rootfinding problem $f(\mathbf{x}) = \mathbf{0}$ is simply $A\mathbf{x} = \mathbf{b}$.

Example

Let

$$f_1(x_1, x_2, x_3) = -x_1 \cos(x_2) - 1,$$

$$f_2(x_1, x_2, x_3) = x_1 x_2 + x_3,$$

$$f_3(x_1, x_2, x_3) = e^{-x_3} \sin(x_1 + x_2) + x_1^2 - x_2^2.$$

Then

$$\mathbf{J}(\mathbf{x}) = \begin{bmatrix} -\cos(x_2) & x_1 \sin(x_2) & 0 \\ x_2 & x_1 & 1 \\ e^{-x_3} \cos(x_1 + x_2) + 2x_1 & e^{-x_3} \cos(x_1 + x_2) - 2x_2 & -e^{-x_3} \sin(x_1 + x_2) \end{bmatrix}.$$

Exercise. Write out the linear part of the Taylor expansion of

$$f_1(x_1 + h_1, x_2 + h_2, x_3 + h_3), \quad \text{near } (x_1, x_2, x_3).$$

The Multidimensional Newton's Method

Recall the idea of Newton's method:

If finding a zero of a function is difficult, replace the function with a simpler approximation (linear) whose zeros are easier to find.

Applying the principle:

- Linearize f at the k th iterate \mathbf{x}_k :

$$\underline{f(\mathbf{x})} \approx L(\mathbf{x}) = \underline{f(\mathbf{x}_k) + \mathbf{J}(\mathbf{x}_k)(\mathbf{x} - \mathbf{x}_k)}.$$

- Define the next iterate \mathbf{x}_{k+1} by solving $L(\mathbf{x}_{k+1}) = \mathbf{0}$:

$$\mathbf{0} = f(\mathbf{x}_k) + \mathbf{J}(\mathbf{x}_k)(\mathbf{x} - \mathbf{x}_k) \implies \mathbf{x}_{k+1} = \mathbf{x}_k - [\mathbf{J}(\mathbf{x}_k)]^{-1} f(\mathbf{x}_k).$$

Note that $\mathbf{J}^{-1}f$ plays the same role as f/f' in the scalar Newton.

cf) $x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)}$

The Multidimensional Newton's Method (cont')

- In practice, we do not compute \mathbf{J}^{-1} . Rather, the k th Newton step $\mathbf{s}_k = \mathbf{x}_{k+1} - \mathbf{x}_k$ is found by solving the square linear system

$$\mathbf{J}(\mathbf{x}_k)\mathbf{s}_k = -\mathbf{f}(\mathbf{x}_k),$$

which is solved using the backslash in MATLAB.

- Suppose `f` and `J` are MATLAB functions calculating \mathbf{f} and \mathbf{J} , respectively. Then the Newton iteration is done simply by

```
% x is a Newton iterate (a column vector).  
% The following is the key fragment  
% inside Newton iteration loop.  
fx = f(x)  
s = -J(x)\fx;  
x = x + s;
```

- Since $\mathbf{f}(x_k)$ is the residual and \mathbf{s}_k is the gap between two consecutive iterates at the k th step, monitor their norms to determine when to stop iteration.

Computer Illustration

- 1 Define f and J , either as anonymous functions or as function m-files.

```
f = @(x) [exp(x(2)-x(1)) - 2;  
          x(1)*x(2) + x(3);  
          x(2)*x(3) + x(1)^2 - x(2)];  
J = @(x) [-exp(x(2)-x(1)), exp(x(2)-x(1)), 0;  
          x(2), x(1), 1;  
          2*x(1), x(3)-1, x(2)];
```

- 2 Define an initial iterate x , say $x_0 = (0, 0, 0)^T$.

```
x = [0 0 0]';
```

- 3 Iterate.

```
for k = 1:7  
    s = -J(x) \ f(x);  
    x = x + s;  
end
```

Implementation

```
function x = newtonsys(f,x1)
% NEWTONSYS    Newton's method for a system of equations.
% Input:
%   f          function that computes residual and Jacobian matrix
%   x1         initial root approximation (n-vector)
% Output
%   x          array of approximations (one per column, last is best)

% Operating parameters.
funtol = 1000*eps;  xtol = 1000*eps;  maxiter = 40;

x = x1(:);
[y,J] = f(x1);
dx = Inf;
k = 1;

while (norm(dx) > xtol) && (norm(y) > funtol) && (k < maxiter)
    dx = -(J\y);    % Newton step
    x(:,k+1) = x(:,k) + dx;

    k = k+1;
    [y,J] = f(x(:,k));
end

if k==maxiter, warning('Maximum number of iterations reached. '), end
end
```