

# **Zonal momentum budget of the Madden-Julian Oscillation: The source and strength of mechanical damping**

**Jialin Lin<sup>1</sup>, Minghua Zhang<sup>2</sup> and Brian Mapes<sup>1</sup>**

<sup>1</sup> NOAA-CIRES Climate Diagnostics Center, Boulder, CO 80305

<sup>2</sup> State University of New York, Stony Brook, NY 11794

Submitted to *J. Atmos. Sci.*, January 2004

*Corresponding author address:* Dr. Jialin Lin,  
NOAA-CIRES Climate Diagnostics Center,  
325 Broadway, R/CDC1, Boulder, CO 80305  
Email: jialin.lin@noaa.gov

## ABSTRACT

Linear, dissipative models with resting base states are sometimes used in theoretical studies of the Madden-Julian Oscillation (MJO). Mechanical damping in such models ranges from nonexistent to strong, since an observational basis for its source and strength has been lacking. This study examines the zonal momentum budget of a composite MJO over the equatorial western Pacific region, constructed using filtering and regression techniques from 15 years (1979-1993) of daily global reanalysis data. Two different reanalyses (NCEP/NCAR and ERA-15) give qualitatively similar results for all major terms, including the budget residual, whose structure is consistent with its interpretation as eddy momentum flux convergence (EMFC) in convection.

The results show that the MJO is a highly viscous oscillation, with a 3-5 day damping time scale, in the upper as well as lower troposphere. Upper-level damping is mainly in the form of large-scale advection terms which are linear in MJO amplitude but involve horizontal and vertical background flow. Specifically, the leading terms are the advection of time-mean zonal shear by MJO vertical motion anomalies, and advection of MJO wind anomalies by time-mean ascent. This upper-level damping in the western Pacific is mostly confined between 10N and 10S. In contrast, zonal wind damping in the lower troposphere involves EMFC (budget residual) and zonal mean linear meridional advection.

Stated another way, the strong upper-level damping necessitates upper-level geopotential height gradients to maintain the observed zonal wind anomalies over the time scales implied by the MJO's low frequency. The existence of the background flow thus

tends to shift MJO temperature perturbations westward, so that the warm anomaly ahead (east) of the convective center is shifted back into the convection. This shifting effect is fully realized only for anomalies with a period much longer than the 3-5 day damping time.

## 1. Introduction

Discovered by Madden and Julian (1971, 1972), the Madden-Julian Oscillation (MJO) is the dominant intraseasonal mode of variability in tropical convection and circulation (e.g. Weickmann et al. 1985, Lau and Chan 1985, Salby and Hendon 1994, Wheeler and Kiladis 1999). It affects a wide range of tropical weather such as the onset and breaks of the Indian and Australian summer monsoons (e.g. Yasunari 1979, Hendon and Liebmann 1990), and the formation of tropical cyclones (e.g. Nakazawa 1986, Liebmann et al. 1994). It also drives teleconnections to the extratropics (e.g., Lau and Phillips 1986) and impacts some important extratropical weather (e.g. Higgins and Mo 1997, Higgins et al. 2000). On a longer timescale, the MJO is observed to trigger or terminate some El Nino events (e.g. Kessler et al. 1995, Takayabu et al. 1999, Bergman et al. 2001). Therefore, the MJO is important for both extended-range weather prediction and long-term climate prediction.

Tropical intraseasonal variability is poorly simulated in general circulation models (GCMs). Typically, simulated phenomena are too weak and propagate too fast (e.g. Hayashi and Sumi 1986, Hayashi and Golder 1986, 1988, 1993, Lau et al. 1988, Slingo et al. 1996). To solve this problem, many theoretical studies have examined the feedback mechanisms which may affect the MJO's amplitude and phase speed, especially the wave-heating feedback mechanisms. Different types of heating parameterizations have been studied, such as wave-CISK (Convective Instability of the Second Kind; e.g. Lau and Peng 1987; Chang and Lim 1988), frictional wave-CISK (e.g. Wang 1988; Salby et al. 1994), WISHE (Wave Induced Surface Heat Exchange; (e.g. Emanuel

1987; Neelin et al. 1987), charge-discharge (e.g. Blade and Hartmann 1993; Hayashi and Golder 1997), and cloud-radiation interaction (Raymond 2001).

In these theoretical studies, linear, dissipative models were often used, with the momentum equation linearized about a state at rest:

$$\frac{\partial u'}{\partial t} = -\frac{\partial \phi'}{\partial x} + f v' - \epsilon u' \quad (1)$$

$$\frac{\partial v'}{\partial t} = -\frac{\partial \phi'}{\partial y} - f u' - \epsilon v' \quad (2)$$

where  $u$  is the zonal wind,  $v$  is the meridional wind,  $\phi$  the geopotential height, and  $f$  the Coriolis parameter. The mechanical damping is usually represented as a Rayleigh friction with a time-scale of  $\epsilon^{-1}$ . Such mechanical damping has been shown to affect the growth rate and phase speed of simulated intraseasonal oscillations. For example, in the case of a free Kelvin wave, Chang (1977) found that damping can significantly increase the intrinsic vertical wavelength and decrease the intrinsic phase speed. In a wave-CISK type model, Chao (1987) found that the phase speed of an intraseasonal oscillation decreases with increasing damping. In a WISHE model of the MJO, Neelin et al. (1987) found that the growth rate of the unstable Kelvin modes significantly decreases with increasing mechanical damping, and this was later confirmed by Neelin and Yu (1994) and Goswami and Rao (1994). However, the choices of damping magnitude in these and other theoretical models vary over a wide range, from highly viscous (e.g. 1 day in Chao 1987), to weakly viscid (e.g. 25 days in Salby et al. 1994), to totally inviscid (e.g. Xie 1994).

The budget terms which are not explicitly expressed in the linearized Eqs. (1)

and (2) include the advective tendency and the effect of convective eddy momentum transport (CMT, also called "cumulus friction" in some previous studies), and therefore they are the possible sources of mechanical damping. In MJO models that did use a strong damping, the source of the damping was usually assumed to be the CMT. The importance of CMT for synoptic-scale waves has been shown in both the momentum budget (Stevens 1979) and the vorticity budget (e.g. Reed and Johnson 1974; Shapiro 1978; Stevens 1979; Esbenson et al. 1982; see review by Sui and Yanai 1986).

For the MJO, the importance of CMT was also shown by two recent observational studies (Houze et al. 2000; Tung and Yanai 2002a,b). Using Doppler radar data collected by aircraft and ship radars during the TOGA COARE field program, Houze et al. (2000) inferred the sense of mesoscale momentum transport in the "super convective systems" in different phases of the MJO event. They found that this mesoscale momentum transport provides damping to the low-level zonal wind in the westerly onset phase, but acceleration during the mature westerly wind burst phase. Using TOGA COARE sounding array momentum budgets, Tung and Yanai (2002a,b) studied the area-averaged CMT, which includes both the convective-scale momentum transport and the mesoscale momentum transport. They found that the area-averaged CMT is strongly modulated by the MJO event, and provides acceleration in the westerly onset phase, but damping in the westerly wind burst phase: opposite in sign to the effect of mesoscale momentum transport. In these two studies, the magnitude of CMT effect was not compared with those of the other momentum

budget terms at the MJO time-scale, and the equivalent damping strength was not estimated. Weickmann et al. (1997) calculated the vertically-integrated angular momentum budget for the MJO, and found that the advective tendency is a dominant budget term. This suggests that the advective tendency may be another important source of mechanical damping.

The purpose of this study is to evaluate the sources and strength of mechanical damping in the observed MJO. To do this, we calculate the MJO momentum budget using 15 years of daily reanalysis data, and then identify the damping term in (1) with the terms proportional to  $u'$  in the full zonal momentum equation. Because the MJO circulation near the equator is dominated by the zonal wind, we focus on the zonal momentum budget. The main budget uncertainties associated with pressure gradient forces and advection terms are examined by (1) comparing between NCEP and ECMWF reanalyses, whose consistencies gave us more confidence, and (2) constructing long-term (15 years) MJO composite and considering only the statistically significant signals, which are likely real signals unless the reanalyses have errors coherent with the MJO.

The datasets used in this study are described in section 2. The time filtering and intraseasonal composite (regression) methods are described in section 3. The zonal momentum budget results are reported in section 4. Summary and discussions are given in section 5.

## 2. Data

The datasets used include 15 years (1979-1999) of daily reanalyses data from two different centers: NCEP (Kalnay et al. 1996) and ECMWF (the ERA-15 reanalysis, Gibson et al. 1999) The variables used include upper air wind, geopotential height, and vertical pressure velocity on pressure surfaces. The horizontal resolution is 2.5 degree longitude by 2.5 degree latitude. The zonal momentum budget is calculated for both reanalyses following Carr and Bretherton (2001), based on the zonal momentum equation:

$$\frac{\partial u}{\partial t} = -u \frac{\partial u}{\partial x} - v \frac{\partial u}{\partial y} - \omega \frac{\partial u}{\partial p} + f(y)v - \frac{\partial \phi}{\partial x} + X \quad (3)$$

where  $u$  is the grid-resolved zonal wind,  $v$  the meridional wind,  $\omega$  the vertical pressure velocity;  $x$  and  $y$  are E-W and N-S distance,  $f$  the Coriolis parameter, and  $\phi$  the geopotential. Here  $X$  represents accelerations due to all subgrid-scale processes. Each budget term except  $X$  was calculated using daily average data at each 2.5 degree grid point.  $X$  is computed as the residual, meaning that all errors in the other terms are included in its observational estimate, requiring caution in interpretation. Derivatives were evaluated using 3-point central differencing. The results were then averaged to pentad data along the equator (between 5N and 5S) with a zonal resolution of 10 degree longitude.

For constructing the MJO composite, we also used 15 years (1979-1993) of pentad CMAP precipitation data (Xie and Arkin 1997) averaged along the equator (between 5N and 5S).



### 3. Method

The MJO is a broad-band phenomenon, with an averaged period of 45 days but a fairly wide spread from 20 to 80 days (see review by Madden and Julian 1994). Its deep convection signal is dominated by wavenumber 1-6, while its circulation signal is dominated by wavenumber 1 (e.g. Salby and Hendon 1994, Wheeler and Kiladis 1999). The variance of its deep convection signal has two centers: western Pacific and eastern Indian Ocean. Over these two centers, the MJO propagates eastward with a slow phase speed of about 5 m/s (e.g. Wheeler and Kiladis 1999, Lin et al. 2004).

These characteristics were used to isolate the MJO signal. Lin et al. (2004) discussed the different methods used in previous observational and modeling studies. The different methods often give qualitatively similar results in terms of propagation characteristics and phase difference among different variables. This consistency is because intraseasonally filtered deep convection along the equator is dominated, both in number and strength, by coherent eastward propagating events (Wang and Rui 1990). Linear composite methods used in most of the previous studies (e.g. phase sum, correlation, regression) are apparently dominated by these strong eastward propagating events. The method used in this study is similar to that used in Lin et al. (2004) and Lin and Mapes (2004). The procedure is as follows:

- (1) The zonal means were first removed from all datasets.
- (2) The datasets were then filtered using a 30-70 day Murakami (1976) filter, whose response function was shown in Lin et al. (2004). The central frequency corresponds to a period of 45 days, with half amplitude at periods of 30 and 70 days. We also

tested the Lanczos filter (Duchan 1979), and the results were not sensitive.

(3) For the master MJO index, we filtered the CMAP precipitation in time as above, and also in space, retaining wavenumbers 1-6.

(4) To study the MJO phenomena in its two centers of variance, we averaged the datasets in two boxes: a western Pacific box between 150-160E and 5N-5S, and an eastern Indian Ocean box between 80-90E and 5N-5S. For each box, an MJO composite was constructed using linear regression with respect to the filtered CMAP precipitation at the same box, and the regression coefficient was multiplied by one standard deviation of the filtered CMAP precipitation. The confidence level of linear correlation was estimated following Oort and Yienger (1996).

(5) Regressions were done for all seasons of the year, as well as for individual seasons (December-February, March-May, June-August, and September-November).

## 4. Results

### *4.1 Vertical structure of the MJO at the equator*

Before examining the MJO zonal momentum budget, we first look at the vertical structure of the MJO, which is important for understanding later the zonal momentum budget. The composite vertical structure of MJO in the western Pacific box is shown in Figs. 1 and 2. The time lag is with respect to the time of maximum precipitation. The time evolution is from the right to the left, showing the local evolution of measured variables as the eastward-moving MJO passes the measurement longitude. The zonal wind  $u$  (Fig. 1a) shows a simple two-layer structure with the upper layer

out of phase with the lower layer. This two-layer structure is well-known from many previous observations (e.g. Madden and Julian 1971, 1972, Weickmann et al. 1985, Knutson and Weickmann 1987).

One interesting subtlety is that the maxima of both the two layers are at higher altitudes than the 850 mb and 200 mb levels used in many previous MJO studies. The maximum of the lower layer is around 600 mb (just below the 0 C level), while that of the upper layer is around 150 mb (just below the tropopause). Thus at the time of maximum precipitation, the zonal convergence is just below the 0 C level, and the zonal divergence is just below the tropopause. The zonal divergence pattern is shifted to the east relative to vertical motion and rainfall, indicative of the meridional component of divergence associated with the Rossby wave component of the flow. In the vertical, however, there is a consistency between the zonal divergence profile and the vertical motion (Fig. 1b) which concentrates high in the upper troposphere. This middle troposphere inflow and near tropopause outflow structure are consistent with the top-heavy heating profile in the MJO shown in Lin et al. (2004).

The vertical motion shows a westward phase tilt with height. It develops first in the lower troposphere, and then shifts upward as it intensifies. This westward phase tilt of vertical motion is consistent with the westward phase tilt of the associated heating anomaly shown in Lin et al. (2004), which is caused by vertical phase tilts of convective heating in the earlier stages, and stratiform heating in the later stages.

The geopotential height  $Z$  (contours in Fig. 1c) has a large amplitude in the near-tropopause outflow layer but a small amplitude in the middle-troposphere inflow layer.

For clarity, the two-dimensional flow is also plotted using arrows in Fig. 1c.  $Z$  lags  $u$  at both the inflow and outflow layers, by about 5 days in the outflow layer and 10 days in the inflow layer. This phase shift is one of the features most different between the NCEP and ECMWF reanalyses (cf. Fig. 1c, 2c). The phase difference in the upper layer is smaller than those found by Kiladis and Weickmann (1992) and Hsu (1996). The reason is that we removed the zonal mean component from all datasets while they didn't, and our test using data retaining zonal mean showed that the zonal mean component in geopotential height increases the  $u$  and  $Z$  phase difference in western Pacific (not shown).

Consistent with the geopotential height structure, the temperature (Fig. 1d) has its largest anomalies in three layers: the lower troposphere below 800 mb, the upper troposphere between 200 and 600 mb, and near the tropopause above 150 mb. The temperature anomaly in the upper troposphere is likely associated with the top-heavy heating profile shown in Lin et al. (2004), because modeling studies showed that the vertical structure of temperature perturbation follows that of heating perturbation (e.g. Haertel and Kiladis 2004). The temperature anomaly near the tropopause is out of phase with that in the upper troposphere, and has a larger amplitude than the later. This is similar to the observations in convectively-coupled Kelvin waves (Wheeler et al. 2000, Straub and Kiladis 2002).

Composites constructed using 15 years (1979-1993) of ECMWF reanalysis data show similar wave structure as the NCEP data (Fig. 2). We also checked the composites for each individual seasons (DJF, MAM, JJA and SON). The wave structures

are generally similar to the all-seasons composites.

In summary, the equatorial wave structure of the MJO has the following three characteristics:

- (1) It is associated with a vertical mode concentrated in the upper half of the troposphere, with middle-troposphere inflow and near-tropopause outflow at the time of maximum precipitation.
- (2) The vertical motion has a westward phase tilt with height.
- (3) Geopotential height lags zonal wind by about 5-10 days at both the inflow and outflow layers.

The phase difference between  $u$  and  $Z$  has implications for the zonal momentum budget. Eq. (1) suggests that when the damping is zero, the balance near the equator is simply between the local tendency and the pressure gradient force. In a neutral wave, this balance leads to an in-phase relationship between  $u$  and  $Z$ . Since zonal wind extrema in the MJO exist within pressure gradients, we can already anticipate that some other (damping) terms must contribute importantly to the zonal momentum budget.

#### *4.2 Zonal momentum budget of the MJO*

From Eq. (3) we can write the zonal momentum equation for intraseasonal anomalies of deviations from the zonal mean:

$$\frac{\partial \tilde{u}'}{\partial t} = -\frac{\partial \tilde{\phi}'}{\partial x} + \tilde{f}v' + \left(-u\frac{\partial \tilde{u}'}{\partial x} - v\frac{\partial \tilde{u}'}{\partial y} - \omega\frac{\partial \tilde{u}'}{\partial p}\right) + \tilde{X}' \quad (4)$$

where the tilde denotes deviation from zonal mean, and primes denotes 30-70 day temporally filtered anomaly.

Major MJO zonal momentum budget terms at the equator (averaged between 150E-160E and 5N-5S) are shown in Figs. 3 and 4 for NCEP and ECMWF reanalyses, respectively. The Coriolis force is small near the equator and is not shown. The two different reanalyses show similar patterns for each of the three budget terms, giving us more confidence in the results. Moreover, only signals with correlation above the 95% confidence level are plotted in Figs. 3 and 4. These signals are likely real unless there are errors in the reanalyses which are systematically coherent with the MJO.

In the upper troposphere, the dominant balance is between the pressure gradient force and the advective tendency. The local tendency is much smaller, as is the budget residual. In the lower troposphere, the advective tendency and the budget residual are of similar amplitude as the local tendency and pressure gradient force. The MJO zonal momentum budgets for each individual seasons are similar to the all-season results (not shown).

The budget residual (Figs. 3d, 4d) shows a two-layer structure, with the residual near the tropopause out of phase with that in the lower troposphere. This is consistent with the vertical structure of budget residual in the TOGA COARE MJO events observed by Tung and Yanai (2002b, their Figs. 8b, 9b). During the westerly wind burst, the budget residual provides a damping to the zonal wind, which is also consistent with the TOGA COARE results. These results suggest that the budget residual from both reanalyses represent the effect of convective momentum transport

(CMT). It is worth noting that the ERA-15 reanalysis model had a parameterization of CMT, while the NCEP reanalysis model did not.

The large contribution of advective tendency to the MJO zonal momentum budget is subdivided into zonal, meridional and vertical components in Fig. 5. In the upper troposphere, the advective tendency is mainly contributed by zonal (Fig. 5a) and vertical components (Fig. 5c), while that in the lower troposphere is mainly contributed by the meridional component (Fig. 5b). Each of these components contain both linear and nonlinear effects, which can be decomposed into (see Appendix):

$$-\widetilde{u} \frac{\partial u'}{\partial x} = -[\widetilde{u}] \frac{\partial \widetilde{u}'}{\partial x} - \widetilde{u}' \frac{\partial [\widetilde{u}]}{\partial x} + \text{other terms} \quad (5)$$

$$-\widetilde{v} \frac{\partial v'}{\partial y} = -[\widetilde{v}] \frac{\partial \widetilde{v}'}{\partial y} - \widetilde{v}' \frac{\partial [\widetilde{v}]}{\partial y} + \text{other terms} \quad (6)$$

$$-\widetilde{\omega} \frac{\partial u'}{\partial p} = -[\widetilde{\omega}] \frac{\partial \widetilde{u}'}{\partial p} - \widetilde{\omega}' \frac{\partial [\widetilde{u}]}{\partial p} + \text{other terms} \quad (7)$$

where  $[\ ]$  denotes time-mean. On the right side of each equation, the first two terms are linear advection terms associated with time-mean zonal asymmetric flow, while the other terms include the linear terms associated with zonal-mean and the nonlinear terms (see Appendix). The first two linear terms were calculated from the corresponding time-mean and the MJO anomalies, with the sum of all remaining terms left together as a (small) residual. For the zonal advection and the vertical advection, the time-mean was taken to be the 15-year annual mean. For the meridional advection, because there is a strong seasonal variation in meridional wind, the time-mean was taken to be 15-year seasonal mean for one season, using DJF here as an example.

First we look at the decomposition of the zonal advection (Fig. 6) and the vertical advection (Fig. 7), which are important in the upper troposphere. The zonal advection (Fig. 6) is dominated by the first two linear terms, both of which contribute substantially to the budget. The vertical advection (Fig. 7) is also dominated by the first two linear terms, especially  $-\tilde{\omega} \frac{\partial \tilde{u}'}{\partial p}$  (Fig. 7a). Therefore, the zonal momentum budget of the MJO is basically linear in the upper troposphere, but with an importantly non-resting basic state. This is consistent with the finding of Weickmann et al. (1997) that the vertically-integrated MJO angular momentum budget is basically linear, since the vertically-integrated momentum budget is mainly contributed by the upper troposphere.

The term  $-\tilde{u} \frac{\partial \tilde{u}'}{\partial x}$ , which describes the advection of the MJO anomaly by the time-mean zonal flow, cannot contribute to damping since it is in quadrature with  $\tilde{u}'$ . In contrast,  $-\tilde{u}' \frac{\partial \tilde{u}}{\partial x}$ , the advection of time-mean zonal flow by the MJO zonal wind anomaly, has an opposite phase with  $\tilde{u}'$  in the upper troposphere, and therefore provides a damping. Meanwhile,  $-\tilde{\omega} \frac{\partial \tilde{u}'}{\partial p}$ , the advection of the MJO zonal wind anomaly by the time-mean vertical motion, tends to reduce the vertical windshear anomaly in the MJO, and therefore also provides a damping. The meridional advection (Fig. 8), which is important in the lower troposphere, is dominated by the terms other than the first two. Further analysis shows that it is actually dominated by the zonal mean linear term  $-\tilde{v}' \frac{\partial \tilde{v}}{\partial y}$  (not shown).

In summary, we find that the MJO zonal momentum budget near the equator can



be simplified to:

$$\frac{\partial \tilde{u}'}{\partial t} \approx -\frac{\partial \tilde{\phi}'}{\partial x} - [\tilde{u}] \frac{\partial \tilde{u}'}{\partial x} - \tilde{u}' \frac{\partial [\tilde{u}]}{\partial x} - [\tilde{\omega}] \frac{\partial \tilde{u}'}{\partial p} - \tilde{v}' \frac{\partial [\tilde{u}]}{\partial y} + \tilde{X}' \quad (8)$$

#### 4.3 Understanding the linear advection terms in the upper troposphere

The dominant linear advection terms in the upper troposphere owe their existence to the time-mean zonal wind  $[\tilde{u}]$ , zonal divergence  $\frac{\partial [\tilde{u}]}{\partial x}$ , and vertical velocity  $[\tilde{\omega}]$ . To help understand these terms, we plot their 15 year (1979-1993) climatological mean in Fig. 9. For consistency with the MJO anomaly, we plot here the zonal asymmetric components, but the zonal means are actually much smaller.

The warm pool region has abundant deep convection, associated with strong upward motion (Fig. 9a). The vertical velocity profile is very top-heavy, and is therefore associated with strong upper-level zonal divergence concentrating in a thin layer near the tropopause (Fig. 9b). The upper-level zonal wind over the warm pool is easterly (Fig. 9c), consistent with the Gill (1980) pattern associated with a stationary heating source. The wind is strong within a thin layer near the tropopause because the zonal divergence is concentrated in this layer. Fig. 9 only shows the annual mean, so the upper-level upward motion and zonal divergence are weaker over the eastern Indian Ocean. The long-term composite seasonal variation of upper-level vertical velocity, zonal divergence, and zonal wind (Fig. 10) show that they are large over the eastern Indian Ocean during northern summer.

When the MJO travels in such a large-scale environment, three terms become important: (1)  $-\tilde{u} \frac{\partial \tilde{u}'}{\partial x}$  (Fig. 6a) lags  $\tilde{u}'$  (Fig. 1a) by a quarter cycle because the time-mean upper-level wind is easterly over the warm pool (Fig. 9c). (2)  $-\tilde{u}' \frac{\partial \tilde{u}}{\partial x}$  (Fig. 6b) is out of phase with  $\tilde{u}'$  because the time-mean upper level zonal wind gradient (zonal divergence) is positive over the warm pool (Fig. 9b), and is advected by the zonal wind anomaly in the MJO. (3)  $-\tilde{\omega} \frac{\partial \tilde{u}'}{\partial p}$  (Fig. 7a) opposes the upper-level  $\tilde{u}'$  (Fig. 1a) because zonal wind shear  $\frac{\partial \tilde{u}'}{\partial p}$  is in phase with upper-level  $\tilde{u}'$  and the time-mean vertical motion is upward (negative  $[\tilde{\omega}]$ , Fig. 9a).

A simple scale analysis clarifies the magnitudes of these terms and of local rate of change. The basic scales for the MJO and time-mean flow are:

$T \approx 40$ days	period
$L \approx 4 \times 10^7$ m	zonal wavelength
$[\tilde{u}] \approx 10$ m/s	time-mean upper-level zonal wind
$\frac{\partial \tilde{u}}{\partial x} \approx 0.2$ day <sup>-1</sup>	time-mean upper-level zonal divergence

From these scales we can estimate the magnitude of the local tendency and the dominant linear zonal advection terms:

$$\begin{aligned}
\frac{\partial \tilde{u}'}{\partial t} &\approx 2\tilde{u}'/(0.5T) \approx \tilde{u}'/(10 \text{ days}) \\
-\tilde{u} \frac{\partial \tilde{u}'}{\partial x} &\approx [\tilde{u}] \times 2\tilde{u}'/(0.5L) \approx \tilde{u}'/(10 \text{ days}) \\
-\tilde{u}' \frac{\partial [\tilde{u}]}{\partial x} &\approx \tilde{u}'/(5 \text{ days})
\end{aligned}$$

Therefore, the linear zonal advection terms is larger than the local tendency. Because the vertical advection terms enhance the zonal advection terms, the total advective

tendency (Fig. 3c), when scaled using  $\tilde{u}'$  (Fig. 1a), is equivalent to a time-scale of 2-3 days, which is much shorter than the 10 days time-scale of local tendency. Therefore, the dominant balance between advective tendency and pressure gradient force in the upper troposphere is a physically sensible feature for the MJO.

#### 4.4 Equivalent linear damping

To get quantitative estimates of the equivalent linear damping strength in the MJO, we regressed the MJO anomalies of the budget terms to the MJO zonal wind anomaly (Fig. 11). When a budget term is in phase with (or in opposite phase to) the zonal wind, it provides a driving (or damping). The unit of the regression coefficient is 1/day, i.e., the reciprocal of the coefficient gives the damping time-scale in the unit of days. The corresponding correlation coefficient indicates how well the budget term is represented by a linear damping.

The pressure gradient force (Fig. 11a) provides a driving with a time scale of 2-3 days in the upper troposphere (150-300 mb), and 3-5 days in the lower troposphere (below 700 mb). This strong forcing cannot be balanced by the local tendency, which, as shown in section 4.3, has a time-scale of about 10 days. The Coriolis force is also small near the equator. Therefore, some strong dampings are needed to balance the pressure gradient force.

In the upper troposphere, the damping mainly comes from the advective tendency (Fig. 11c), which, as shown in section 4.2, includes  $-\tilde{u}' \frac{\partial [\tilde{u}]}{\partial x}$  and  $-\tilde{\omega} \frac{\partial \tilde{u}'}{\partial p}$ . The sub-grid term X (Fig. 11e) also contributes slightly near the tropopause. In the lower

troposphere, the damping comes equally from the advective tendency (the nonlinear meridional advection term) and  $X$ . The linear correlation coefficients (Figs. 11b,d,f) are generally large when the regression coefficient are large, suggesting the usefulness of linear damping representation.

Since the damping involves mean flow, its value varies in space. For example, Fig. 12 shows the meridional distribution at 200 mb of damping due to the advective tendency, in longitudes 150-160. The damping is large only near the equator between 10N-10S, because the strong upper-level zonal divergence and vertical motion are confined near the equator. The damping in the lower levels also has its largest value near the equator (not shown).

As mentioned in section 3, deep convection signal in the MJO has two centers of variance: western Pacific and eastern Indian Ocean. The above results are for the western Pacific. We also did similar analysis for the eastern Indian Ocean. The wave structures and zonal momentum budgets in the eastern Indian Ocean are generally similar to those in the western Pacific except during the boreal summer (not shown). Many previous studies have shown that the boreal summer intraseasonal oscillation has a prominent northward-propagating component (e.g. Yasunari 1979, Madden and Julian 1994), which is different from the eastward-propagating MJO mode originally described by Madden and Julian (1971, 1972), although there are some connections between the northward-propagating mode and the eastward-propagating MJO mode (Lawrence and Webster 2002). Therefore we restrict this study to the eastward-propagating MJO mode and will report the results for the boreal summer intrasea-

sonal oscillation in a separate study.

## 5. Summary and discussions

The observed MJO wave structure and zonal momentum budget at the equator are schematically shown in Fig. 13. The equatorial wave structure of the MJO is associated with a vertical mode concentrated in the upper half of the troposphere, with middle-troposphere inflow and near-tropopause outflow at the time of maximum precipitation. The vertical motion has a westward phase tilt with height, but zonal divergence is offset to the east of the ascent by meridional effects (Rossby wave flow components), making it slightly difficult to display both tilts with a schematic circulation in the zonal plane. The main feature to note here is that geopotential height lags zonal wind by about 5-10 days at both the inflow and outflow layers.

In the MJO zonal momentum budget, the advective tendency is a major budget term in both the upper and lower tropospheres, while subgrid transports are a major budget term only in the lower troposphere. The advective tendency in the upper troposphere comes from linear terms associated with time-mean zonal and vertical motions, while that in the lower troposphere comes from zonal mean linear meridional advection. The simplified MJO zonal momentum equation is Eq. 8.

The strong linear advection terms in the upper troposphere can be understood from the time-mean flow in the warm-pool region where abundant climatological deep convection is associated with strong upward motion ( $[\omega] < 0$ ), upper-level zonal divergence ( $\partial[u]/\partial x > 0$ ), and mean upper-level easterly wind ( $[u] < 0$ ). When the

MJO travels in such a large-scale environment, at the upper levels,  $-[u]\partial u'/\partial x$  lags  $u'$  by a quarter cycle, while both  $-u'\partial[u]/\partial x$  and  $-\omega\partial u'/\partial p$  are out of phase with  $u'$ . Given the low frequency of the MJO, the magnitudes of the time-mean flows are bound to cause an advective tendency much larger than the local tendency. The advective tendency and convective eddy momentum flux convergence together provide a 3-5 day time-scale damping to the MJO both in the upper troposphere and in the lower troposphere. Therefore, the MJO is a highly viscous oscillation. At western Pacific longitudes, the damping is large only near the equator between 10N-10S. The full 3D spatial structure of mean flow may constitute a spatial envelope that shapes the MJO importantly, but the assumptions behind linearized composite diagnostics here do not warrant production of detailed maps of damping rates.

It may be possible to anticipate the effect of the observed strong damping on theoretical models of intraseasonal oscillations. One obvious effect is to reduce growth rates, perhaps stabilizing unstable modes found under weak-damping assumptions, for example as shown in the WISHE model of Goswami and Rao (1994, their Fig. 5c). It may require several positive feedback mechanisms to generate unstable modes in the presence of damping (e.g. WISHE plus cloud-radiation feedback). Another possible effect is on the phase speed of intraseasonal modes. For example, the wave-CISK model of Chao (1987) shows that strong damping reduces phase speed. Phase speed or period will also be importantly affected by quadrature terms like  $-[\tilde{u}]\frac{\partial \tilde{u}'}{\partial x}$ , which might be more conveniently viewed as a simple Doppler shifting by the mean flow.

It is also instructive to turn these diagnostic findings about momentum damping around, and orient the results toward thermodynamic considerations. We have seen that the existence of warm-pool mean flow necessitates baroclinic  $Z'$  anomalies in the MJO, in order to sustain the zonally divergent zonal wind anomalies associated with its low-frequency convection anomalies. These  $Z'$  anomalies correspond to temperature anomalies, as indicated on Fig. 13. Specifically, the lowness of MJO frequency relative to the time-scale of mean-flow advective damping forces the warm anomaly ahead of the convection to shift rearward, into the convective heating anomaly.

In the terms of a conventional short-hand of energetics reasoning (e.g. Emanuel et al. 1994), this phase shift tends to increase the correlation between heating and temperature, thus increasing eddy energy production. It is tempting to claim that this reasoning might comprise an explanation for the enhanced power in low frequencies that we call the MJO, or an interpretation of the constant-frequency rather than dispersion-line spectral characteristics of the MJO (Wheeler and Kiladis 1999). But circumspection requires acknowledging that the spatial correlation in the equatorial zonal plane between time-filtered zonal deviations of heating and temperature is at best a very partial study of the energetics of observed variability, and falls far short of a useful predictive theory of why that observed variability exists.

Nonetheless, this study points to the fact that, for low frequency disturbances, the background flow takes on a special importance which pervades a disturbance's thermodynamic as well as momentum fields. This is in line with the finding of Slingo et al. (1996) from the Atmospheric Model Intercomparison Project (AMIP) that GCMs

which produce more realistic intraseasonal variability also tend to produce stronger and deeper time-mean vertical motion. It is clear that the effort to improve intraseasonal variability in global models must be considered in concert with the effort to simulate the mean climate well.

### *Acknowledgments*

We gratefully thank Klaus Weickmann and George Kiladis for carefully reading an early version of the manuscript and making very helpful comments. We also thank Don Hooper for assistance in reading the reanalyses data. This work was supported by the National Science Foundation Grants ATM-0097116, ATM-0073206 and ATM-0112715.

### **Appendix. Decomposition of the advection terms**

Because the MJO anomaly in this study does not include the zonal mean, decomposition of the advection terms is more complex than when zonal mean is retained. Following Peixoto and Oort (1992, pp61-64), a quantity  $A$  can be decomposed into:

$$A = [\overline{A}] + [\tilde{A}] + \overline{A'} + \tilde{A'} \quad (\text{A1})$$

where overbar represents zonal mean, tilde represents deviation from the zonal mean (zonal asymmetric component), bracket represents time mean, and prime represents deviation from the time mean. It can be derived that:

$$\widetilde{AB'} = ([\tilde{A}]\tilde{B'} + [\tilde{B}]\tilde{A'}) + ([\overline{A}]\tilde{B'} + [\overline{B}]\tilde{A'}) + ([\tilde{A}]\overline{B'} + [\tilde{B}]\overline{A'})$$



$$+(-[\widetilde{A}]\widetilde{B}' - [\widetilde{B}]\widetilde{A}' + \widetilde{A}'\widetilde{B}' - [\widetilde{A}'\widetilde{B}']) \quad (\text{A2})$$

On the right hand of the equation, the terms on the first line are the linear terms, among which the first two are associated with time-mean zonal asymmetric component, and are calculated in Eqs. 5 and 7. The third and fourth are associated with time-mean zonal-mean component. The fifth and sixth are associated with temporally filtered zonal-mean component. The terms on the second line are the nonlinear terms. When considering only the 30-70 day anomalies, the nonlinear terms will include some more terms corresponding to interaction with other time-scales.

## REFERENCES

- Bergman, J. W., H. H. Hendon, K. M. Weickmann, 2001: Intraseasonal Air-Sea Interactions at the Onset of El Nino. *Journal of Climate*, **14**, 1702-1719.
- Blade, I., and D. L. Hartmann, 1993: Tropical intraseasonal oscillations in a simple non-linear model. *J. Atmos. Sci.*, **50**, 2922-2939.
- Carr, M. T., Bretherton, C. S. 2001: Convective Momentum Transport over the Tropical Pacific: Budget Estimates. *J. Atmos. Sci.*, **58**, 1673-1693.
- Chang, C. P., 1977: Viscous internal gravity waves and low-frequency oscillations in the tropics. *J. Atmos. Sci.*, **34**, 901-910.
- Chang, C. P., and H. Lim, 1988: Kelvin wave-CISK: A possible mechanism for the 30-50 day oscillations. *J. Atmos. Sci.*, **45**, 1709-1720.
- Chao, W. C., 1987: On the origin of the tropical intraseasonal oscillation. *J. Atmos. Sci.*, **44**, 1940-1949.
- Duchan, C.E., 1979: Lanczos filtering in one and two dimensions. *J. Appl. Meteor.*, **18**, 1016-1022.
- Emanuel, K. A., 1987: An air-sea interaction model of intraseasonal oscillation in the Tropics. *J. Atmos. Sci.*, **44**, 2324-2340.
- Emanuel, K. A., J. D. Neelin, and C. S. Bretherton, 1994: On large-scale circulations in convecting atmospheres. *Quart. J. Roy. Meteor. Soc.*, **120**, 1111-1143.
- Esbensen, S. K., E. I. Tollerud, and J.-H. Chu, 1982: Cloud-cluster-scale circulations and the vorticity budget of synoptic-scale waves over the eastern Atlantic intertropical convergence zone. *Mon. Wea. Rev.*, **110**, 1677-1692.

- Gibson, J. K., P. Kallberg, S. Uppala, A. Hernandez, A. Nomura, and E. Serrano, 1999: ERA description. Reanalysis (ERA) Project Report Series, No. 1, European Centre for Medium-Range Weather Forecasts, 84 pp.
- Gill, A. E., 1980: Some simple solutions for heat-induced tropical circulation. *Quart. J. Roy. Meteor. Soc.*, **106**, 447-462.
- Goswami, P., and R. K. Rao, 1994: A dynamical mechanism for selective excitation of the Kelvin mode at timescale of 30-50 days. *J. Atmos. Sci.*, **51**, 2769-2779.
- Haertel, P. T., and G. N. Kiladis, 2004: On the dynamics of two-day equatorial disturbances. *J. Atmos. Sci.*, submitted.
- Hartmann, D. L., M. L. Michelson, and S. A. Klein, 1992: Seasonal variations of tropical intraseasonal oscillations: A 20-25-day oscillation in the western Pacific. *J. Atmos. Sci.*, **49**, 1277-1289.
- Hayashi, Y., and A. Sumi, 1986: The 30-40 day oscillation simulated in an "aqua planet" model. *J. Meteor. Soc. Japan.*, **64**, 451-466.
- Hayashi, Y., and D. G. Golder, 1986: Tropical intraseasonal oscillations appearing in a GFDL general circulation model and FGGE data. Part I: Phase propagation. *J. Atmos. Sci.*, **43**, 3058-3067.
- Hayashi, Y., and D. G. Golder, 1988: Tropical intraseasonal oscillations appearing in a GFDL general circulation model and FGGE data. Part II: Structure. *J. Atmos. Sci.*, **45**, 3017-3033.
- Hayashi, Y., and D. G. Golder, 1993: Tropical 40-50 and 25-30-day oscillations appearing in realistic and idealized GFDL climate models and the ECMWF dataset. *J. Atmos.*

*Sci.*, **50**, 464-494.

Hayashi, Y., and D. G. Golder, 1997: United mechanisms for the generation of low- and high-frequency tropical waves. Part I: Control experiments with moist convective adjustment. *J. Atmos. Sci.*, **54**, 1262-1276.

Hendon, H. H., and B. Liebmann, 1990: A composite study of onset of the Australia monsoon. *J. Atmos. Sci.*, **47**, 2227-2240.

Houze, R. A., Chen, S. S., Kingsmill, D. E., Serra, Y., Yuter, S. E. 2000: Convection over the Pacific Warm Pool in relation to the Atmospheric Kelvin-Rossby Wave. *J. Atmos. Sci.*, **57**, 3058-3089.

Hsu, H.-H., 1996: Global view of the intraseasonal oscillation during northern winter. *J. Climate*, **9**, 2386-2406.

Kessler, W. S., and M. J. McPhaden, and K. M. Weickmann, 1995: Forcing of intraseasonal Kelvin waves in the equatorial Pacific. *J. Geophys. Res.*, **100**, 10613-10631.

Kiladis, G. N., and K. M. Weickmann, 1992: Circulation anomalies associated with tropical convection during northern winter. *Mon. Wea. Rev.*, **120**, 1900-1923.

Knutson, T. R., and K. M. Weickmann, 1987: 30-60 day atmospheric oscillations: Composite life cycles of convection and circulation anomalies. *Mon. Wea. Rev.*, **115**, 1407-1436.

Lawrence, D. M., and P. J. Webster, 2002: The Boreal Summer Intraseasonal Oscillation: Relationship between Northward and Eastward Movement of Convection. *J. Atmos. Sci.*, **59**, 1593-1606.

Lau, K. M., and P. H. Chan, 1985: Aspects of the 40-50-day oscillation during the northern

- winter as inferred from outgoing longwave radiation. *Mon. Wea. Rev.*, **113**, 1889-1909.
- Lau, K. M., and L. Peng, 1987: Origin of low-frequency (intraseasonal) oscillations in the tropical atmosphere. *J. Atmos. Sci.*, **44**, 950-972.
- Lau, K.-M., and T. J. Phillips, 1986: Coherent fluctuations of extratropical geopotential height and tropical convection in intraseasonal timescales. *J. Atmos. Sci.*, **43**, 1164-1181.
- Lau, N. C., I. M. Held, and J. D. Neelin, 1988: The Madden-Julian oscillations in an idealized general circulation model. *J. Atmos. Sci.*, **45**, 3810-3831.
- Liebmann, B., H. H. Hendon, and J. D. Glick, 1994: The relationship between the tropical cyclones of the western Pacific and Indian Oceans and the Madden-Julian oscillation. *J. Meteor. Soc. Japan*, **72**, 401-411.
- Lin, J. L., B. E. Mapes, M. H. Zhang, and M. Newman, 2004: Stratiform precipitation, vertical heating profiles, and the Madden-Julian Oscillation. *J. Atmos. Sci.*, **61**, 296-309.
- Lin, J. L., and B. E. Mapes, 2004: Radiation budget of the tropical intraseasonal oscillations. *J. Atmos. Sci.*, accepted. Available at <http://www.cdc.noaa.gov/~jlin>
- Madden, R. A., and P. R. Julian, 1971: Detection of a 40-50 day oscillation in the zonal wind in the tropical Pacific. *J. Atmos. Sci.*, **28**, 702-708.
- Madden, R. A., and P. R. Julian, 1972: Description of global-scale circulation cells in the tropics with a 40-50 day period. *J. Atmos. Sci.*, **29**, 1109-1123.
- Madden, R. A., and P. R. Julian, 1994: Observations of the 40-50-day oscillation -A review.

- Mon. Wea. Rev.*, **122**, 814-837.
- Murakami, M., 1979: Large-scale aspects of deep convective activity over the GATE area. *Mon. Wea. Rev.*, **107**, 994-1013.
- Nakazawa, T., 1986: Mean features of 30-60 day variations as inferred from 8-year OLR data. *J. Meteor. Soc. Japan*, **64**, 777-786.
- Neelin, J. D., I. M. Held, and K. H. Cook, 1987: Evaporation-wind feedback and low-frequency variability in the tropical atmosphere. *J. Atmos. Sci.*, **44**, 2341-2348.
- Oort, A. H., and J. J. Yienger, 1996: Observed long-term variability in the Hadley circulation and its connection to ENSO. *J. Climate*, **9**, 2751-2767.
- Raymond, D. J., 2001: A New Model of the Madden-Julian Oscillation. *J. Atmos. Sci.*, **58**, 2807-2819.
- Reed, R. J., and R. H. Johnson, 1974: The vorticity budget of synoptic-scale wave disturbances in the tropical western Pacific. *J. Atmos. Sci.*, **31**, 1784-1790.
- Salby, M. L., and H. H. Hendon, 1994: Intraseasonal behavior of clouds, temperature, and motion in the Tropics. *J. Atmos. Sci.*, **51**, 2207-2224.
- Salby, M. L., R. B. Garcia, and H. H. Hendon, 1994: Planetary-scale circulations in the presence of climatological and wave-induced heating. *J. Atmos. Sci.*, **51**, 2344-2367.
- Shapiro, L. J., 1978: The vorticity budget of a composite African tropical wave disturbance. *Mon. Wea. Rev.*, **106**, 806-817.
- Slingo, J. M., and Coauthors, 1996: Intraseasonal oscillations in 15 atmospheric general circulation models: Results from an AMIP diagnostic subproject. *Climate Dyn.*, **12**, 325-357.

- Stevens, D. E., 1979: Vorticity, momentum, and divergence budgets of synoptic-scale wave disturbances in the tropical eastern Atlantic. *Mon. Wea. Rev.*, **107**, 535-550.
- Straub, K. H., and G. N. Kiladis, 2002: Observations of a Convectively Coupled Kelvin Wave in the Eastern Pacific ITCZ. *J. Atmos. Sci.*, **59**, 30-53.
- Sui, C.-H., and M. Yanai, 1986: Cumulus ensemble effects on the large-scale vorticity and momentum fields of GATE. Part I: Observational evidence. *J. Atmos. Sci.*, **43**, 1618-1642.
- Takayabu, Y. N., Toshio Iguchi, Misako Kachi, Akira Shibata and Hiroshi Kanzawa, 1999: Abrupt termination of the 1997-98 El Nino in response to a Madden-Julian oscillation. *Nature*, **402**, 279-282.
- Tung, W. W., Yanai, M., 2002a: Convective Momentum Transport Observed during the TOGA COARE IOP. Part I: General Features. *J. Atmos. Sci.*, **59**, 1857-1871.
- Tung, W. W., Yanai, M., 2002b: Convective Momentum Transport Observed during the TOGA COARE IOP. Part II: Case Studies. *J. Atmos. Sci.*, **59**, 2535-2549.
- Wang, B., 1988: Dynamics of tropical low-frequency waves: An analysis of the moist Kelvin wave. *J. Atmos. Sci.*, **45**, 2051-2065.
- Wang, B., and H. Rui, 1990: Synoptic climatology of transient tropical intraseasonal convection anomalies. *Meteor. Atmos. Phys.*, **44**, 43-61.
- Weickmann, K. M., G. R. Lussky, and J. E. Kutzbach, 1985: Intraseasonal (30-60 day) fluctuations of outgoing longwave radiation and 250 mb streamfunction during northern winter. *Mon. Wea. Rev.*, **113**, 941-961.
- Weickmann, K. M., G. N. Kiladis, and P. D. Sardeshmukh, 1997: The dynamics of intrasea-

- sonal atmospheric angular momentum oscillations. *J. Atmos. Sci.*, **54**, 1445-1461.
- Wheeler, M., and G. N. Kiladis, 1999: Convectively coupled equatorial waves: Analysis of clouds and temperature in the wavenumber-frequency domain. *J. Atmos. Sci.*, **56**, 374-399.
- Wheeler, M., Kiladis, G. N., Webster, P. J. 2000: Large-Scale Dynamical Fields Associated with Convectively Coupled Equatorial Waves. *J. Atmos. Sci.*, **57**, 613-640.
- Xie, P., and P. A. Arkin, 1997: Global precipitation: A 17-year monthly analysis based on gauge observations, satellite estimates, and numerical model outputs. *Bull. Amer. Meteor. Soc.*, **78**, 2539-2558.
- Xie, S.-P., 1994: On the preferred zonal scale of wave-CISK with conditional heating. *J. Meteor. Soc. Japan*, **72**, 19-30.
- Yasunari, T., 1979: Cloudiness fluctuations associated with the northern hemisphere summer monsoon. *J. Meteor. Soc. Japan*, **57**, 227-242.



## FIGURE CAPTIONS

**Fig. 1** The vertical structure of the MJO anomaly for (a) zonal wind (m/s), (b) vertical velocity (mb/day), (c) geopotential height (m), and (d) temperature (K) for the 15 years (1979-1993) of NCEP reanalysis data averaged over 150E-160E and 5N-5S. Negative values are shaded. The arrows in (c) are the wind vectors whose horizontal component is the zonal wind, and vertical component the vertical velocity.

**Fig. 2** Same as Fig. 1 except for the ECMWF reanalysis data.

**Fig. 3** Terms of the MJO zonal momentum budget: (a) local tendency  $\frac{\partial \tilde{u}'}{\partial t}$ , (b) pressure gradient force  $-\frac{\partial \tilde{\phi}'}{\partial x}$ , (c) advection terms  $(-\widetilde{u \frac{\partial u}{\partial x}}' - \widetilde{v \frac{\partial u}{\partial y}}' - \widetilde{\omega \frac{\partial u}{\partial p}}')$ , and (d) the residual term  $\widetilde{X}'$ , for 15 years (1979-1993) of NCEP reanalysis data averaged over 150E-160E and 5N-5S. Unit is  $ms^{-1}day^{-1}$ .

**Fig. 4** As in Fig. 3 except for ECMWF reanalysis data.

**Fig. 5** As in Fig. 3 except for three advection terms for NCEP reanalysis data: (a) zonal advection  $-\widetilde{u \frac{\partial u}{\partial x}}'$ , (b) meridional advection  $-\widetilde{v \frac{\partial u}{\partial y}}'$ , and (c) vertical advection  $-\widetilde{\omega \frac{\partial u}{\partial p}}'$ .

**Fig. 6** As in Fig. 3 except for three components of the zonal advection term for NCEP reanalysis data: (a)  $-\widetilde{[u] \frac{\partial u'}{\partial x}}$ , (b)  $-\widetilde{u' \frac{\partial [u]}{\partial x}}$ , and (c) other terms.

**Fig. 7** As in Fig. 3 except for three components of the vertical advection term for NCEP reanalysis data: (a)  $-\widetilde{[\omega] \frac{\partial u'}{\partial p}}$ , (b)  $-\widetilde{\omega' \frac{\partial [u]}{\partial p}}$ , and (c) other terms.

**Fig. 8** As in Fig. 3 except for three components of the meridional advection term for NCEP reanalysis data for DJF season: (a)  $-\widetilde{[\tilde{v}] \frac{\partial u'}{\partial y}}$ , (b)  $-\widetilde{\tilde{v}' \frac{\partial [\tilde{u}]}{\partial y}}$ , and (c) other terms.

**Fig. 9** Annual mean (a)  $[\tilde{\omega}]$  (mb/day), (b)  $\frac{\partial[\tilde{u}]}{\partial x}$  (1/day), and (c)  $[\tilde{u}]$  (m/s), for 15 years (1979-1993) of NCEP reanalysis data along the equator (5N-5S).

**Fig. 10** Seasonal variation of (a)  $[\tilde{\omega}]$  (mb/day) at 300 mb, (b)  $\frac{\partial[\tilde{u}]}{\partial x}$  (1/day) at 150 mb, and (c)  $[\tilde{u}]$  (m/s) at 150 mb, for NCEP reanalysis data along the equator (5N-5S).

**Fig. 11** Vertical profile of linear regression coefficient of the MJO (a) pressure gradient force anomaly, (c) advective tendency anomaly, and (e) budget residual anomaly, with respect to the zonal wind anomaly at the same level for 15 years (1979-1993) of NCEP (solid line) and ECMWF (dashed line) reanalyses data averaged over 150E-160E and 5N-5S. The corresponding correlation coefficients are shown in (b), (d), and (f), respectively.

**Fig. 12** Meridional profile of (a) linear regression coefficient, and (b) linear correlation coefficient, between the 200 mb MJO advective tendency anomaly and zonal wind anomaly averaged between 150E-160E for 15 years (1979-1993) of NCEP (solid line) and ECMWF (dashed line) reanalyses data.

**Fig. 13** Schematic depiction of the time-mean flow over the warm-pool, and the observed MJO wave structure and zonal momentum budget. Regions of enhanced large-scale convection are indicated schematically by the clouds. The dark shading inside the clouds represents the maximum of the diabatic heating. “H” and “L” represent the high and low geopotential height anomalies, respectively. “W” inside thick solid circle and “C” inside thick dashed circle represent the warm and cold temperature anomalies, respectively. Thin arrows represent the winds. Thick arrows above the MJO wave represent components of zonal momentum budget in the upper

troposphere, while those below the MJO wave represent components in the lower troposphere.




Figure 1:




Figure 2:




Figure 3:




Figure 4:




Figure 5:






Figure 6:




Figure 7:




Figure 8:




Figure 9:




Figure 10:




Figure 11:




Figure 12:

# MJO in warm-pool time-mean flow

## Time-mean flow





Figure 13: