


The dataset chosen: The forest fire dataset.

	X	Y	month	day	FFMC	DMC	DC	ISI	temp	RH	wind	rain	area	month_num
0	7	5	mar	fri	86.2	26.2	94.3	5.1	8.2	51	6.7	0.0	0.00	3
1	7	4	oct	tue	90.6	35.4	669.1	6.7	18.0	33	0.9	0.0	0.00	10
2	7	4	oct	sat	90.6	43.7	686.9	6.7	14.6	33	1.3	0.0	0.00	10
3	8	6	mar	fri	91.7	33.3	77.5	9.0	8.3	97	4.0	0.2	0.00	3
4	8	6	mar	sun	89.3	51.3	102.2	9.6	11.4	99	1.8	0.0	0.00	3
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
512	4	3	aug	sun	81.6	56.7	665.6	1.9	27.8	32	2.7	0.0	6.44	8
513	2	4	aug	sun	81.6	56.7	665.6	1.9	21.9	71	5.8	0.0	54.29	8
514	7	4	aug	sun	81.6	56.7	665.6	1.9	21.2	70	6.7	0.0	11.16	8
515	1	4	aug	sat	94.4	146.0	614.7	11.3	25.6	42	4.0	0.0	0.00	8
516	6	3	nov	tue	79.5	3.0	106.7	1.1	11.8	31	4.5	0.0	0.00	11

Distributions for each feature:




The pair plot between each two features:



One example that holds to the expressiveness principle:

The below chart is a scatter plot between two features: DMC and FPMC, with the color channel indicating month. We can observe from the color channel that the value of DMC generally tends to increase with month until Sep, and there doesn't seem to be a salient correlation between FPMC and month.



One example that violates the expressiveness principle:

Below is a line chart plotted between two features: temperature and wind. Since there's no indicated order in the dataset, the default sample order, which does not make any sense, generates highly chaotic line charts for any two features.

