

Contents

1	The Standard Model and its supersymmetric extensions	1
1.1	The Standard Model of particle physics	3
1.1.1	The particle content	3
1.1.2	The Lagrangian density	4
1.1.3	The Brout-Englert-Higgs mechanism	6
1.1.4	Limitations of the Standard Model	8
1.2	Supersymmetry	9
1.2.1	The MSSM	12
1.2.1.1	The particle content of the MSSM	13
1.2.1.2	The Lagrangian density of the MSSM	13
1.2.1.3	The phenomenological MSSM	15
1.2.2	Supersymmetry breaking	15
1.3	Long-lived particles in the MSSM	16
1.3.1	Previous searches and constraints from indirect searches	17
2	A search for highly ionising, short tracks at the CMS detector	21

17

Part 1

18

The Standard Model and its supersymmetric extensions

19

1.1 The Standard Model of particle physics

The formulation of a relativistic quantum field theory and of spontaneous symmetry breaking (SSB) by the Brout-Englert-Higgs mechanism, allowed to built a theory which is capable to explain almost all observations of particle physics until today [1]. This theory is known as the Standard Model of particle physics (SM). The existence of its last missing piece, the Higgs boson, could be proven at the LHC in the year 2012 [2, 3].

The Standard Model is a $SU(3)_C \times SU(2)_L \times SU(1)_Y$ non-abelian gauge theory. “After” spontaneous symmetry breaking, its symmetries are reduced to $SU(3)_C \times U(1)_{EM}$. All particles that were found until today are contained in it¹. Furthermore, it is able to describe three of the four fundamental forces: the strong, weak and electromagnetic force.

In the following, a small introduction to the theory and phenomenology of the Standard Model is given. It is not meant as a complete description. The reader is referred to [4–6], for a thorough and extensive introduction.

1.1.1 The particle content

It should be first noted, since the Standard Model is a quantum field theory, every field (to be more precise every degree of freedom of a field) can be considered also as a particle and vice versa.

The Standard Model of particle physics contains three different particle types, or three different types of fields. First, there are the so-called “matter particles”, which are all spin 1/2 particles in the SM. Second, the forces are described by spin 1 vector bosons. And finally, in order to give masses to all particles the Standard Model embeds the Higgs boson, a scalar spin 0 particle.

Fermions in the Standard Model

The fermionic content can be further subdivided into leptons and quarks. In contrast to quarks, leptons are not strongly interacting, thus they only couple electromagnetically and weakly to other particles. Both, the quarks and the leptons are ordered into three different families. Across these families, all quantum numbers are conserved. They only differ by their mass.

¹One can argue, that the right-handed neutrino, which is proven to exist, is not contained. But as at least the left-handed neutrino is embedded, we want to ignore that for a moment.

48 All left-handed particles of each family form a $SU(2)_L$ doublet, which causes the coupling
 49 via the weak force. The right-handed partners form $SU(2)$ singlets, thus, don't couple via
 50 the weak interaction. As quarks carry one further quantum number, the colour, they are
 51 additionally grouped into $SU(3)_C$ triplets.

52 Vector bosons in the Standard Model

53 As mentioned before, the vector bosons describe three of the four fundamental forces.
 54 There is one gauge boson corresponding to every generator of the above mentioned gauge
 55 groups. For $U(1)_Y$, it is the B -boson, for $SU(2)_L$, there are three gauge bosons $W^{1,2,3}$
 56 and finally eight gauge bosons $G^{1\dots 8}$ for $SU(3)_C$, which are called gluons. As the B -field
 57 and the neutral W_μ^3 -field can mix, "after" SSB the basis can be changed and lead to the
 58 well known photon and Z -boson.

59 The Higgs boson

60 A somehow extraordinary role plays the Higgs boson, that was predicted already 50 years
 61 ago by Peter Higgs [7, 8] and could be proven existent by the LHC experiments CMS and
 62 ATLAS in 2012 [2, 3]. This particle is a consequence of the spontaneous symmetry break-
 63 ing after rotating three of the four degrees of freedom to masses of the W -and Z -bosons.
 64 It is the only known fundamental scalar particle.

65
 66 An overview of all Standard Model particles and their transformation properties are
 67 shown in Table 1.1.1. If particles transform as singlets under $SU(2)_L$ or $SU(3)_C$, they don't
 68 couple via the corresponding force. The hypercharges Y are determined by $Q = Y + I_3$,
 69 where Q is the electric charge and I_3 is the third component of the weak isospin with
 70 $I^a = \sigma^a/2$, σ^a being the Pauli matrices.

71 1.1.2 The Lagrangian density

72 In particle physics, the probability of a decay or an interaction between particles can
 73 be calculated with the help of the Lagrangian density. The Lagrangian density of the
 74 Standard Model is the most general set of Lagrangian terms, that are renormalisable and
 75 contain all up to date known particles as well as the above mentioned gauge symmetries.

Table 1.1.1: All particles contained in the Standard Model and their transformation properties under $SU(3)_C \times SU(2)_L \times SU(1)_Y$.

	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$
Fermions:			
$(\nu_L, e_L)^T$	1	2	-1
e_R	1	1	-2
$(u_L, d_L)^T$	3	2	$+\frac{1}{3}$
u_R	3	1	$+\frac{4}{3}$
d_R	3	1	$-\frac{2}{3}$
Vector bosons:			
B_μ	1	1	0
W_μ^a	1	3	0
G_μ^a	8	1	0
Higgs boson: H	1	2	-1

76 It is the following:

$$\begin{aligned}
\mathcal{L} = & (D_\mu \Phi)^\dagger (D^\mu \Phi) - \mu^2 \Phi^\dagger \Phi - \frac{\lambda}{4} (\Phi^\dagger \Phi)^2 \\
& + \bar{L}_i^L i \not{D} L_i^L + \bar{e}_i^R i \not{D} e_i^R + \bar{Q}_{ib}^L i \not{D} Q_{ib}^L + \bar{u}_{ib}^R i \not{D} u_{ib}^R + \bar{d}_{ib}^R i \not{D} d_{ib}^R \\
& - \left(Y_{ij}^e \bar{L}_i^L \Phi e_j^R + Y_{ij}^u \bar{Q}_{ib}^L \Phi u_{jb}^R + Y_{ij}^d \bar{Q}_{ib}^L \Phi d_{jb}^R + h.c. \right) \\
& - \frac{1}{4} (B_{\mu\nu} B^{\mu\nu} + W_{\mu\nu}^a W^{a\mu\nu} + G_{\mu\nu}^a G^{a\mu\nu}),
\end{aligned} \tag{1.1.1}$$

77 with $\not{D} = \gamma_\mu D^\mu$ and the covariant derivative $D^\mu = \partial^\mu + ig' Y_W B^\mu - ig C_1 I^a W_a^\mu - ig_S C_2 T^a G_a^\mu$.
78 I^a and T^a denote hereby the generators of the $SU(2)_L$ and $SU(3)_C$, respectively. They are
79 connected to the three Pauli matrices and the eight Gell-Mann matrices by $I^a = \frac{\sigma^a}{2}$ and
80 $T^a = \frac{\lambda^a}{2}$. Adding the hypercharge Y_W and the third component of the weak isospin result
81 in the electrical charge $Q = Y_W + I_3$. Furthermore, it is $C_1 = 1$ for doublets and $C_1 = 0$
82 for singlets under $SU(2)_L$, $C_2 = 1$ for triplets and $C_2 = 0$ for singlets under $SU(3)_C$.

83 The first line in Eq. (1.1.1) correspond to the kinetic term of the Higgs field and its
84 potential. Via this Higgs field, it is possible to give masses to the Z - and W^\pm -bosons as well

as the fermions. This will be explained in detail in the following Section 1.1.3. The second line describes the kinetic terms of the leptons and quarks. The index i represents the three different families ($i = 1, 2, 3$). Since they are spin 1/2 particles, they can be described with the help of Dirac spinors. The left-handed leptons and quarks are described as $SU(2)_L$ doublets, $L_I^L = (\nu_{eL}, e_L)_i$, $Q_I^L = (u_L, d_L)_i$, the right-handed as singlets under $SU(2)_L$ e_i^R , u_i^R , d_i^R . Quarks carry a further quantum number, the colour, which is indicated by the index b with $b = 1, 2, 3$. Quarks transform as triplets under the $SU(3)_C$ gauge group. The third line contains the interaction terms between the fermions and the Higgs boson, called Yukawa interactions. These terms lead after SSB to the fermion mass terms, as can be seen later. The last line correspond to the kinetic terms of the gauge fields. These are connected to the field strength tensors by

$$\begin{aligned}
B^{\mu\nu} &\equiv \partial^\mu B^\nu - \partial^\nu B^\mu \\
W^{\mu\nu} &\equiv \partial^\mu W^\nu - \partial^\nu W^\mu - ig [W^\mu, W^\nu] \\
&= \left(\partial^\mu W_i^\nu - \partial^\nu W_i^\mu + g \epsilon_{ijk} W_j^\mu W_k^\nu \right) \frac{\sigma_i}{2} \equiv \frac{\sigma_i}{2} W_a^{\mu\nu} \\
G^{\mu\nu} &\equiv \partial^\mu G^\nu - \partial^\nu G^\mu - ig_S [G^\mu, G^\nu] \\
&= \left(\partial^\mu G_a^\nu - \partial^\nu G_a^\mu + g_S f_{abc} G_b^\mu G_c^\nu \right) \frac{\lambda_a}{2} \equiv \frac{\lambda_a}{2} G_a^{\mu\nu}.
\end{aligned} \tag{1.1.2}$$

The factors ϵ_{ijk} and f_{abc} are hereby the structure constants of the corresponding lie groups. The summation over all indices that appear twice is included.

1.1.3 The Brout-Englert-Higgs mechanism

An essential ingredient of the Standard Model is the Brout-Englert-Higgs mechanism (BEH mechanism), earlier also called Higgs mechanism. It was developed by Peter Higgs, Robert Brout and François Englert in 1960s [7–12]. Based on a work from Sheldon Glashow [13], Steven Weinberg and Abdus Salam later applied it on a $SU(2) \times U(1)$ gauge theory [14,15]. By this, a renormalisable theory of the weak and the electromagnetic theory was born. Together with the theory of strong interaction, the formulation of the Standard Model was thus by this time complete.

Mass terms of the gauge bosons

Due to the BEH mechanism, it is possible to give masses to the W^\pm - and Z-bosons. A scalar field Φ (Higgs field) is required, which has a non-zero vacuum expectation value. This is possible, when the mass parameter μ in front of the bilinear term in line one of Eq. (1.1.1) is smaller than zero and $\lambda > 0$ at the same time.

The resulting potential of the Higgs field is then the famous “Mexican hat” potential.

Expanding the Lagrangian density around the minimum of $\Phi = (0, v)$, the gauge symmetries of $SU(2)_L \times U(1)_Y$ are spontaneously broken and only a remaining electrical charge conserving symmetry $U(1)_{EM}$ remains. After an unitary transformation, three of the four degrees of freedom of the Higgs field are absorbed by the gauge fields. Thus, “after” SSB, the part of the Lagrangian containing the scalar field is as follows

$$\mathcal{L}_{\text{Higgs}} = \frac{1}{2} (\partial_\mu h^0)^\dagger (\partial^\mu h^0) - \mu^2 (h^0)^2 + \frac{1}{2} v^2 g^2 W_\mu^- W^{+\mu} + \frac{1}{4} v^2 (g^2 + g'^2) Z_\mu Z^\mu + \text{interaction terms} \quad (1.1.3)$$

One kinetic and one mass term for one of the degrees of freedom of the Higgs fields remains, which is the Higgs boson (h^0). Furthermore, three of the four gauge bosons require a mass. The remaining gauge boson, being the photon remains massless because of the conserved $U(1)_{EM}$ gauge symmetry.

The mass eigenstates of the gauge bosons in Eq. (1.1.3) are obviously different to the interaction eigenstates in Eq. (1.1.1). The diagonalisation of the neutral mass matrices is described by the Weinberg angle θ_W

$$\begin{pmatrix} A_\mu \\ Z_\mu \end{pmatrix} = \begin{pmatrix} \cos \theta_W & -\sin \theta_W \\ \sin \theta_W & \cos \theta_W \end{pmatrix} \begin{pmatrix} B_\mu \\ W_\mu^3 \end{pmatrix}. \quad (1.1.4)$$

For the charged gauge bosons, the relation between $W_{1,2}$ and W^\pm is the following

$$W_\mu^\pm = \frac{1}{\sqrt{2}} [W_\mu^1 \mp W_\mu^2]. \quad (1.1.5)$$

Consequently, the masses of the gauge boson are the following

$$\begin{aligned} M_H &= \sqrt{2}\mu \\ M_W &= \frac{g}{\sqrt{2}} v \\ M_Z &= \frac{1}{\sqrt{2}} v \sqrt{g^2 + g'^2} \\ M_\gamma &= 0. \end{aligned} \quad (1.1.6)$$

The first direct observation of the Z - and W^\pm -bosons was made in $p\bar{p}$ -collisions in the year 1983 at the Super Proton Synchrotron (SPS) at CERN [16, 17]. The experimental values of the masses are $m_Z = 91.1876 \pm 0.0021 \text{ GeV}$ and $m_{W^\pm} = 80.385 \pm 0.015 \text{ GeV}$ [18]. Finally, as mentioned several times before, the Higgs boson was found at the LHC in the year 2012 [2, 3]. The mass is measured to $m_{h^0} = 125.09 \pm 0.21(\text{stat.}) \pm 0.11(\text{sys.}) \text{ GeV}$ [19].

131 Mass terms of the fermions

132 Fermion mass terms cannot be easily inserted into the Lagrangian density in Eq. (1.1.1),
 133 since they would violate the imposed gauge symmetries. With the help of the BEH
 134 mechanism it is possible to generate fermion mass terms via the Yukawa interactions terms
 135 (line three of Eq. (1.1.1)). After spontaneous symmetry breaking, the Yukawa interactions
 136 lead to the following mass terms (colour indices are suppressed)

$$\begin{aligned} \mathcal{L}_{\text{Yukawa}} = & - \left(Y_{ij}^e v \bar{e}_i^L e_j^R + Y_{ij}^u v \bar{u}_i^L u_j^R + Y_{ij}^d v \bar{d}_i^L d_j^R + h.c. \right) \\ & + \text{interaction terms} \end{aligned} \quad (1.1.7)$$

137 The fermion masses are thus described by the following mass matrices

$$M_{ij}^e = Y_{ij}^e v \qquad M_{ij}^u = Y_{ij}^u v \qquad M_{ij}^d = Y_{ij}^d v \quad (1.1.8)$$

138 Since the Standard Model does not contain right-handed neutrinos, there are no gauge
 139 invariant Yukawa interactions, that could produce mass terms for neutrinos.

140 1.1.4 Limitations of the Standard Model

141 Despite the great success of the Standard Model, there remain observations and theoret-
 142 ical considerations that cannot be answered within the SM. In the following, the most
 143 important of such “limitations” shall be reviewed.

144

145 First of all, the Standard Model suffers of the so-called hierarchy problem. By this,
 146 the problem is addressed that quadratic divergencies occur in the calculation of the Higgs
 147 self-coupling. The appearance of infinities is not uncommon in higher order calculations
 148 and happens for all particles. Still, for scalar particles the infinite term is quadratically
 149 divergent, which makes a huge difference compared to the logarithmic divergencies for
 150 fermion self-energies. When considering the Standard Model valid up to the Planck scale,
 151 an extraordinary fine-tuning would be needed to cancel a large bare mass with large
 152 counter terms

$$m_{h^0}^{\text{ren } 2} = m_{h^0}^{\text{bare } 2} + \Delta m_{h^0}^2. \quad (1.1.9)$$

153 to end up with a mass of about 125 GeV. Thus, this renormalisation procedure, even if
 154 mathematically possible, is regarded as highly unnatural in physics. The question which
 155 is usually imposed is, why is the Higgs mass so small, when there are such massive cor-
 156 rections to the bare mass? A formulation of naturalness was given by t’Hooft in 1977 [20].
 157 He stated, that a small parameter can be regarded only natural, if the symmetries of the
 158 theory are enhanced by setting this parameter to zero. In the Standard Model, though,

there is no enhancement of the symmetries of the Lagrangian by setting $\mu = 0$, thus the small mass of the Higgs boson compared to the Planck scale is considered as highly unnatural.

A further and probably the most striking shortcoming of the Standard Model is the missing fourth fundamental force, the gravitational force. Within the SM, it is not possible to add renormalisable terms, that can describe the gravitational force. Although, gravity is not important for particle physics at energies that are accessible at current particle colliders (it only becomes important at the Planck scale $\sim 10^{19}$ GeV), the fact that it cannot be embedded into the Standard Model leads to an understanding of the SM as effective theory, only valid for lower energies. Thus, it is obviously not an ultimate theory and something must be beyond.

Furthermore, in particle physics there is always the wish to describe nature with a theory as simple as possible. This usually implies the effort to embed the Standard Model into a higher symmetry group. To achieve a simplification by unifying the three fundamental forces is usually done within so-called Grand-Unified-Theories (GUTs). Calculating the running of the coupling constants in the Standard Model, the couplings seem to meet at a scale of $M_{\text{GUT}} \sim 10^{15}$ GeV. Unfortunately, they don't meet exactly. Therefore, a unification is not achievable in the Standard Model under the assumption that there are no new particles up to the GUT scale.

Finally, there is experimental evidence, which cannot be explained within the Standard Model. Astrophysical observations suggest that there is a large amount of dark matter (DM) in the universe, that cannot be explained with the particle content of the Standard Model. Measurements of the velocity curves of galaxies, e. g. M33 [21] show discrepancies between the observed velocities and the predicted velocities by the visible matter. The share of non-visible matter to the total amount of matter in the universe is estimated to be 84% [22]. Unfortunately, there is no suitable (only weakly interacting) candidate within the SM, that can make up the full DM contribution.

In the following Chapter 1.2, a theory is introduced, that can address most of the above mentioned problems. This theory is called Supersymmetry.

1.2 Supersymmetry

As noted in the last chapter, the Higgs boson mass suffers from quadratic divergencies through radiative corrections. The reason for the quadratic divergencies is due to the

fact, that the Lagrangian density does not contain further symmetries for $\mu \rightarrow 0$. This behaviour is typical for scalar particles. For fermions, on the other hand, there is a further symmetry for $m_f = 0$. The Lagrangian density becomes invariant under chiral transformations of the form $\Psi \rightarrow e^{i\vec{\alpha}\frac{\vec{\sigma}}{2}\gamma_5}\Psi$. Although the mass terms of the fermions break this symmetry, it protects the fermions against large radiative corrections.

Due to these considerations, it seems natural to protect also the scalar mass by an additional symmetry. A work from Golfand and Likhtman in the year 1971 stated that an extension of the Poincaré algebra is possible via fermionic generators [23]. R. Haag, J. Lopuszanski and M. Sohnius finalised these considerations by showing that with the help of fermionic generators a connection between space-time symmetries and internal symmetries is possible [24]. The extensions of a symmetry group by fermionic generators is called Supersymmetry (SUSY). These were the foundation of supersymmetric theories.

In the following, few aspects of supersymmetric theories are discussed. For a detailed introduction the reader is referred to [25–27].

In the subsequent sections, the descriptions is restricted to the case of $N = 1$ supersymmetry, i.e. there is only one supersymmetric generator and thus only one supersymmetric partner for every particle. A supersymmetric transformation transfers every bosonic state into a fermionic state and vice versa

$$\begin{aligned} Q |\text{boson}\rangle &= |\text{fermion}\rangle \\ Q |\text{fermion}\rangle &= |\text{boson}\rangle. \end{aligned} \tag{1.2.1}$$

The most important (anti-) commutation relations for SUSY algebra with spinors Q are

$$\begin{aligned} \{Q_\alpha, Q_\beta\} &= \{Q_\alpha^\dagger, Q_\beta^\dagger\} = 0, \\ [Q_\alpha, P_\mu] &= [Q_\alpha^\dagger, P_\mu] = 0, \\ \{Q_\alpha, Q_\alpha^\dagger\} &= 2\sigma_{\alpha\dot{\alpha}}^\mu P_\mu. \end{aligned} \tag{1.2.2}$$

P_μ denotes hereby the four component generator of translations. Additionally, it is $\sigma^\mu = (\mathbb{1}, \sigma_i)$ with the Pauli matrices σ_i . From the second relation in (1.2.2) it follows

$$[Q_\alpha, P^2] = [Q_\alpha^\dagger, P^2] = 0. \tag{1.2.3}$$

Equation 1.2.3 implies, that particles that are transformed into each other with the generator Q need to have same eigenvalues of P^2 , thus, also same masses. This impose obviously a problem, since no “scalar electron” was ever found with a mass of 0.51 MeV. This problem will be discussed later.

In Supersymmetry, all particles and their partner particles are described by so-called supermultiplets. Since the generators of the gauge group commute with the generators of supersymmetry, all particles within one supermultiplet have same quantum numbers, besides the spin (as shown in Eq. 1.2.3 they have also same masses). In a renormalisable theory, there are two different types of supermultiplets: chiral multiplets, which contain a two-component Weyl spinor and a complex scalar field and vector multiplets with a vector boson and a two-component Weyl spinor. The number of degrees of freedom are thus for chiral multiplets for on-shell particles

$$\begin{aligned} n_f &= 2, \text{ due to the two-component Weyl spinor} \\ n_b &= 2, \text{ due to the complex scalar field,} \end{aligned}$$

and for vector multiplets for on-shell particles

$$\begin{aligned} n_f &= 2, \text{ due to the two-component Weyl spinor} \\ n_b &= 2, \text{ due to the vector field.} \end{aligned}$$

As mentioned before, in a realistic extension of the Standard Model, Supersymmetry cannot be exact, since no supersymmetric particles have been found which have the same masses as their SM partners. This implies, that SUSY must be broken. There are many ideas how the breaking can actually happen. However, since up to now, only little is known about the breaking mechanism, usually further terms which break SUSY explicitly, are added by hand to the Lagrangian density. These terms can parametrise SUSY breaking without the knowledge about the breaking mechanism. One condition is however imposed on the supersymmetry breaking terms: they should not spoil the naturalness of the new theory, i.e. no new quadratic divergencies shall occur due to these terms. Therefore, they are called soft-breaking terms. How they actually look, will be explained in the Section 1.2.1.

Finally, it shall be discussed how Supersymmetry can give possible answers to the shortcomings of the Standard Model, discussed in Section 1.1.4.

Radiative corrections by fermions always have a factor -1 compared to bosonic corrections. Thus, calculating radiative corrections of the Higgs boson mass in a supersymmetric theory leads in addition to the corrections by SM particles also to further corrections by SUSY particles. If SUSY were exact, the quadratic divergencies Δm_H^2 would exactly cancel. However, as argued, SUSY must be broken. The cancellation of quadratic divergencies can therefore only be assured, if the breaking is not too drastic and only logarithmic divergencies remain. To avoid a new source of fine-tuning, the soft-breaking parameters should be of the order $M_{\text{soft}} \sim 100 \text{ GeV} - 1000 \text{ GeV}$.

Even though, it is not possible to implement gravity within a supersymmetric extension




Figure 1.2.1: The running of the gauge couplings in the Standard Model (left) and in the minimal supersymmetric extension of the SM (right). Taken from [28].

of the Standard Model, theories that are able to include gravity (string theories) require also Supersymmetry.

The renormalisation group equations change under a supersymmetric extension of the Standard Model. By this, a unification of the gauge couplings at a GUT scale of about 10^{16} GeV is possible, as can be seen in Fig. 1.2.1. It can be nicely seen, that all three gauge couplings cross each other within the uncertainties.

Besides these arguments, SUSY can also give an answer to the problem of non-visible matter in the universe. If the conservation of the so-called R-parity is required, the lightest supersymmetric particle (LSP) is stable. If this particle is only weakly interacting, it can serve as a good candidate to explain fully or partially the sources of the relic density. R-parity is a multiplicative quantum number with

$$\begin{aligned} P_R &= 1 && \text{SM particles} \\ P_R &= -1 && \text{SUSY particles.} \end{aligned}$$

If R-parity is conserved, only terms are allowed in the Lagrangian density, that contain an even number of supersymmetric particles. Therefore, no single SUSY particle can decay into only SM particles and thus, the LSP is stable.

1.2.1 The MSSM

The supersymmetric extension of the Standard Model with a minimal particle content is called the Minimal Supersymmetric Standard Model (MSSM). In the following section, the particle content of the MSSM is introduced.

1.2.1.1 The particle content of the MSSM

In $N = 1$ Supersymmetry, every SM particle has exactly one supersymmetric partner particle, which leads to a doubling of the particle content in the MSSM with respect to the SM. Additionally, there is a necessity for a second Higgs doublet. The second doublet is needed to ensure the holomorphicity of the superpotential when also mass terms for the up-type particles shall be created. Furthermore, the MSSM stays only free from anomalies if there is a further Higgs doublet. This leads to the fact, that in the MSSM, there are five Higgs bosons instead of only one as in the SM. The complete particle content of the MSSM is depicted in Tables 1.2.1 and 1.2.2.

Since in supersymmetric theories only left-handed Weyl spinors appear in the Lagrangian density, the right-handed are described as charge conjugated spinors of the left-handed spinors.

Table 1.2.1: Chiral supermultiplets in the MSSM

		spin 0	spin $\frac{1}{2}$	$SU(3)_C, SU(2)_L, U(1)_Y$
squarks/quarks	Q	$(\tilde{u}_L, \tilde{d}_L)$	(u_L, d_L)	3, 2, $\frac{1}{3}$
	\bar{u}	$\tilde{\bar{u}}_L = \tilde{u}_R^\dagger$	$\bar{u}_L = (u_R)^c$	$\bar{3}, 1, -\frac{4}{3}$
	\bar{d}	$\tilde{\bar{d}}_L = \tilde{d}_R^\dagger$	$\bar{d}_L = (d_R)^c$	$\bar{3}, 1, \frac{2}{3}$
sleptons/leptons	L	$(\tilde{\nu}_{eL}, \tilde{e}_L)$	(ν_{eL}, e_L)	1, 2, -1
	\bar{e}	$\tilde{\bar{e}}_L = \tilde{e}_R^\dagger$	$\bar{e}_L = (e_R)^c$	$\bar{1}, 1, 2$
Higgs/higgsinos	H_u	(H_u^+, H_u^0)	$(\tilde{H}_u^+, \tilde{H}_u^0)$	1, 2, 1
	H_d	(H_d^0, H_d^-)	$(\tilde{H}_d^0, \tilde{H}_d^-)$	1, 2, -1

1.2.1.2 The Lagrangian density of the MSSM

In the following only the most important parts of the MSSM Lagrangian density will be described. The reader is again referred to [27] for a complete description of the Lagrangian density.

Table 1.2.2: Vector supermultiplets in the MSSM

	spin $\frac{1}{2}$	spin 0	$SU(3)_C, SU(2)_L, U(1)_Y$
gluinos/gluons	\tilde{g}	g	8, 1, 0
winos/ W -bosons	$\tilde{W}^\pm, \tilde{W}^0$	W^\pm, W^0	1, 3, 0
bino/ B -boson	\tilde{B}	B	1, 1, 0

285 The superpotential

286 The superpotential of the MSSM contains the self interaction terms of the Higgs bosons
 287 and generates the interaction terms of the Higgs bosons with the fermions and their super-
 288 partners. As already noted, it is very common to assume R-parity conservation. Hence, no
 289 terms appear in the Lagrangian that would violate lepton or baryon number conservation
 290 and the lightest supersymmetric particle is stable. Thus, all possible terms are

$$W_{\text{MSSM}} = \mu H_u \cdot H_d - Y_u^{ij} H_u \cdot Q_L^i u_R^{cj} + Y_d^{ij} H_d \cdot Q_L^i d_R^{cj} + Y_e^{ij} H_d \cdot L_L^i e_R^{cj}, \quad (1.2.4)$$

291 with the dot product defined as in [26]

$$A \cdot B = \epsilon^{\alpha\beta} A_\alpha B_\beta = A_1 B_2 - A_2 B_1. \quad (1.2.5)$$

292 The soft-breaking Lagrangian density

293 Since Supersymmetry is broken, explicit SUSY breaking terms are added to the Lagrangian
 294 density. In order not to introduce new sources of quadratic divergencies, only bilinear and
 295 trilinear terms appear in the soft-breaking Lagrangian

$$\begin{aligned}
 -\mathcal{L}_{\text{soft}}^{MSSM} = & m_{H_u}^2 H_u^\dagger \cdot H_u + m_{H_d}^2 H_d^\dagger \cdot H_d + (B\mu H_u \cdot H_d + h.c.) \\
 & + m_{\tilde{Q}}^2 \tilde{Q}_{Lij}^\dagger \cdot \tilde{Q}_{Lj} + m_{\tilde{u}}^2 \tilde{u}_{Ri}^{c\dagger} \cdot \tilde{u}_{Rj}^c + m_{\tilde{d}}^2 \tilde{d}_{Ri}^{c\dagger} \cdot \tilde{d}_{Rj}^c \\
 & + m_{\tilde{L}}^2 \tilde{L}_{Lij}^\dagger \cdot \tilde{L}_{Lj} + m_{\tilde{e}}^2 \tilde{e}_{Ri}^{c\dagger} \cdot \tilde{e}_{Rj}^c \\
 & + \left(-(A_u Y_u)_{ij} H_u \cdot \tilde{Q}_{Li} \tilde{u}_{Rj}^c + (A_d Y_d)_{ij} H_d \cdot \tilde{Q}_{Li} \tilde{d}_{Rj}^c \right. \\
 & \left. + (A_e Y_e)_{ij} H_d \cdot \tilde{L}_{Li} \tilde{e}_{Rj}^c + h.c. \right) \\
 & + \left(M_1 \tilde{B} \tilde{B} + M_2 \tilde{W}_a \tilde{W}_a + M_3 \tilde{g}_i \tilde{g}_i + h.c. \right)
 \end{aligned} \quad (1.2.6)$$

The first line contains mass terms for the Higgs bosons, the second and third line for the sfermions. In the fourth and fifth line the trilinear couplings between the Higgs bosons and the sfermions appear. Finally, the last line give rise to mass terms for the gauginos.

Because of the soft-breaking terms, the MSSM contains more than 100 free parameters. Constraining the MSSM is thus a difficult task and usually in experimental particle physics, constrained versions of the MSSM or assumptions at the GUT scale are used to report the impact of searches on SUSY. In the following a short introduction of the phenomenological MSSM is given. With its reduced parameter space, it allows to elaborate on long-lived particles in the MSSM in a much easier way.

1.2.1.3 The phenomenological MSSM

The phenomenological MSSM (pMSSM) imposes constraints that are reasonable in the sense to fulfil current observations and still keep the phenomenological richness of the MSSM [29]. The following assumptions are imposed (in [29] more detailed information about these assumptions can be found):

- No new sources of CP violation,
- No flavour changing neutral currents,
- First and second generation universality.

These assumption reduce the number of SUSY parameters to only 19. The remaining free parameters are the following:

- $\tan \beta$ (the ratio of the vacuum expectation values of the two Higgs doublets)
- M_A (the mass of the pseudo-scalar Higgs boson)
- μ (the Higgs mass parameter)
- M_1, M_2, M_3 (bino, wino and gluino mass parameters, respectively)
- $m_{\tilde{q}}, m_{\tilde{l}}, m_{\tilde{u}}, m_{\tilde{d}}$ and $m_{\tilde{e}}$ (the first and second generation mass parameters)
- $m_{\tilde{Q}}, m_{\tilde{L}}, m_{\tilde{t}}, m_{\tilde{b}}$ and $m_{\tilde{\tau}}$ (the third generation mass parameters)
- A_t, A_b and A_τ (third generation trilinear couplings).

1.2.2 Supersymmetry breaking

As already noted, the mechanism of supersymmetry breaking is unknown. There exists however, several ideas how to spontaneously break supersymmetry. All mechanism have

in common that they need to happen at high energies in a hidden sector. “Messenger” particles are introduced which mediate the breaking to the TeV scale. This, however, implies that supersymmetry breaking is a question of high-energy physics and one can always parametrise the breaking by the soft breaking terms introduced in Section 1.2.1.2.

The most popular breaking mediation mechanisms is gravity-mediated Supersymmetry breaking [30] and gauge-mediated supersymmetry breaking [31].

1.3 Long-lived particles in the MSSM

There are various mechanisms how particles can be long-lived, such as small couplings or suppressed decays because of high mediator masses (weak interaction). For a comprehensive review, the reader is referred to [32].

In this thesis, the focus is set on particles that have a long lifetime due to a small decay phase space. A phase space suppression is possible when the mass splitting between the decay particle and one of the decay products is very small. In Part 2, a search for highly ionising, short tracks is presented. This search is motivated by long-lived charginos, that are nearly mass-degenerate with the lightest supersymmetric particle, the neutralino.

The underlying mechanism of this mass-degeneracy will be addressed in the next paragraphs.

In the MSSM, the lightest chargino ($\tilde{\chi}_1^\pm$) and the lightest neutralino ($\tilde{\chi}_1^0$) can be almost mass-degenerate, when the wino mass parameter (M_2) is smaller than the bino (M_1) and higgsino (μ) mass parameters. This can be seen in the masses of the interaction eigenstates. The following expressions are approximate neutralino and chargino mass terms for $M_2\mu > m_W^2 \sin 2\beta$ and $|M_2 \pm \mu|, |M_1 \pm \mu| \gg m_Z$ (taken from [33])

$$\begin{aligned}
m_{\tilde{B}} &\simeq M_1 + \frac{m_Z^2 (M_1 + \mu \sin 2\beta) \sin^2 \theta_W}{M_1^2 - \mu^2} \\
m_{\tilde{W}} &\simeq M_2 + \frac{m_Z^2 (M_2 + \mu \sin 2\beta) \cos^2 \theta_W}{M_2^2 - \mu^2} \\
m_{\tilde{H}_1^0} &\simeq |\mu| + \frac{m_Z^2 (1 - \sin 2\beta) (\mu + M_2 \sin^2 \theta_W + M_1 \cos^2 \theta_W) \operatorname{sgn}(\mu)}{2(\mu + M_2)(\mu + M_1)} \\
m_{\tilde{H}_2^0} &\simeq |\mu| + \frac{m_Z^2 (1 + \sin 2\beta) (\mu - M_2 \sin^2 \theta_W - M_1 \cos^2 \theta_W) \operatorname{sgn}(\mu)}{2(\mu - M_2)(\mu - M_1)}
\end{aligned} \tag{1.3.1}$$

347

$$\begin{aligned}
m_{\tilde{W}^\pm} &\simeq M_2 + m_W^2 \left[\frac{M_2 + \mu \sin 2\beta}{M_2^2 - \mu^2} \right] \\
m_{\tilde{H}^\pm} &\simeq |\mu| + m_W^2 \operatorname{sgn}(\mu) \left[\frac{\mu + M_2 \sin 2\beta}{\mu^2 - M_2^2} \right]
\end{aligned} \tag{1.3.2}$$

348 It is obvious from Eqs. 1.3.1 and 1.3.2, that if $M_2 < M_1$, $|\mu|$, the lightest neutralino state
 349 is wino-like and is fully mass-degenerate on tree level with the lightest chargino. Thus,
 350 the mass differences between $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_1^0$ is only determined by higher loop contributions,
 351 making the mass splitting naturally very small.

352 Furthermore, recent analyses of the pMSSM parameter space [34, 35] show, that al-
 353 most pure wino-like neutralinos often come along with wino-like charginos and their mass
 354 splitting is typically of the order of ~ 160 MeV [34].

355 1.3.1 Previous searches and constraints from indirect searches

356 Several previous searches are sensitive on SUSY scenarios with almost mass-degenerate
 357 wino-like charginos and neutralinos. In the following an overview about these previous
 358 searches will be given. Additionally, constraints from indirect searches are summarised.

359 Searches at LEP

360 Several searches at LEP were hunting for almost mass-degenerate neutralino-chargino
 361 scenarios [36–39]. These searches were looking for events with a high-energetic initial
 362 state radiated photon leading to missing energy in events with chargino-pair production
 363 and invisible decay products. The excluded parameter regions by these searches can be
 364 found in [40] and are depicted in Fig. 1.3.1. The searches were interpreted for M_1 and M_2
 365 almost degenerate and with a large unified scalar mass m_0 leading to sneutrino masses
 366 larger than 500 GeV. Charginos are excluded up to a mass of 92.4 GeV [40].

367 Searches at ATLAS at 7 and 8 TeV

368 At the ATLAS experiment at the LHC, searches for events with a disappearing track
 369 signature were conducted at $\sqrt{s} = 7$ TeV [41] as well as at $\sqrt{s} = 8$ TeV [42]. Furthermore,
 370 a search for metastable particles with high ionisation loss was performed with $\sqrt{s} = 8$ TeV
 371 data [43]. These searches were interpreted within an anomaly-mediated Supersymmetry
 372 breaking model [44] with $\tan\beta = 5$ and $\mu < 0$. The excluded parameter space by these
 373 searches is shown in Fig. 1.3.2. Models with charginos down to lifetimes of 0.06 ns can be
 374 excluded.




Figure 1.3.1: Observed and expected exclusion limits in the $m_{\tilde{\chi}_1^\pm} - \Delta m(\tilde{\chi}_1^\pm, \tilde{\chi}_1^0)$ plane for almost degenerate M_1 and M_2 and a large unified scalar mass m_0 . Taken from [40].




Figure 1.3.2: Excluded parameter space in the $m_{\tilde{\chi}_1^\pm} - \tau_{\tilde{\chi}_1^\pm}$ plane for an AMSB model with $\tan \beta = 5$ and $\mu > 0$. Only chargino pair production is taken into account. The area below the curves is excluded. Taken from [45].

Searches at CMS at 7 and 8 TeV

There are several searches at the CMS experiment at the LHC that are sensitive to long-lived wino-like charginos. Among them is the search for long-lived charged particles [46], which searched for heavy particles with large energy deposits in the tracker. Furthermore, there is the search for disappearing tracks [47] which analysed events with disappearing tracks in the tracker with respect to wino-like charginos almost mass-degenerate with the lightest neutralino. Since the latter search is more sensitive to shorter lifetime, only the exclusion limits derived by this search are shown in Fig. 1.3.3. The disappearing track search by CMS shows very similar sensitivity as the searches done at the ATLAS experiment.




Figure 1.3.3: Excluded parameter space in the $\tau_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^\pm}$ plane for wino-like charginos. The region left to the curve is excluded. Taken from [47].

Indirect searches

Finally, also results from indirect dark matter searches constrain the SUSY parameter region of wino-like charginos and neutralinos.

You found some good papers!!!!

389

Part 2

390

A search for highly ionising, short tracks at the CMS detector

391

Bibliography

- [1] M. Baak, M. Goebel, J. Haller et al., “The Electroweak Fit of the Standard Model after the Discovery of a New Boson at the LHC”, *Eur. Phys. J.* **C72** (2012) 2205, [arXiv:1209.2716](#). doi:10.1140/epjc/s10052-012-2205-9.
- [2] CMS Collaboration, “Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC”, *Phys. Lett.* **B716** (2012) 30–61, [arXiv:1207.7235](#). doi:10.1016/j.physletb.2012.08.021.
- [3] ATLAS Collaboration, “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC”, *Phys. Lett.* **B716** (2012) 1–29, [arXiv:1207.7214](#). doi:10.1016/j.physletb.2012.08.020.
- [4] Peskin, Michael E. and Schroeder, Daniel V., “An Introduction To Quantum Field Theory”. Westview Press, 1995.
- [5] Ryder, L. H., “QUANTUM FIELD THEORY”. Cambridge University Press, 1996.
- [6] D. Griffiths, “Introduction to Elementary Particles”. Wiley, 2008.
- [7] P. W. Higgs, “Broken symmetries, massless particles and gauge fields”, *Phys. Lett.* **12** (1964) 132–133. doi:10.1016/0031-9163(64)91136-9.
- [8] P. W. Higgs, “Broken Symmetries and the Masses of Gauge Bosons”, *Phys. Rev. Lett.* **13** (1964) 508–509. doi:10.1103/PhysRevLett.13.508.
- [9] F. Englert and R. Brout, “Broken Symmetry and the Mass of Gauge Vector Mesons”, *Phys. Rev. Lett.* **13** (1964) 321–323. doi:10.1103/PhysRevLett.13.321.
- [10] G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble, “Global Conservation Laws and Massless Particles”, *Phys. Rev. Lett.* **13** (1964) 585–587. doi:10.1103/PhysRevLett.13.585.
- [11] P. W. Higgs, “Spontaneous Symmetry Breakdown without Massless Bosons”, *Phys. Rev.* **145** (1966) 1156–1163. doi:10.1103/PhysRev.145.1156.
- [12] T. W. B. Kibble, “Symmetry breaking in nonAbelian gauge theories”, *Phys. Rev.* **155** (1967) 1554–1561. doi:10.1103/PhysRev.155.1554.

- [13] S. L. Glashow, “Partial Symmetries of Weak Interactions”, *Nucl. Phys.* **22** (1961) 579–588. doi:10.1016/0029-5582(61)90469-2.
- [14] S. Weinberg, “A Model of Leptons”, *Phys. Rev. Lett.* **19** (1967) 1264–1266. doi:10.1103/PhysRevLett.19.1264.
- [15] A. Salam, “Weak and Electromagnetic Interactions”, *Conf. Proc.* **C680519** (1968) 367–377.
- [16] UA1 Collaboration, “Experimental Observation of Isolated Large Transverse Energy Electrons with Associated Missing Energy at $s^{1/2} = 540\text{-GeV}$ ”, *Phys. Lett.* **B122** (1983) 103–116. [611(1983)]. doi:10.1016/0370-2693(83)91177-2.
- [17] UA1 Collaboration, “Experimental Observation of Lepton Pairs of Invariant Mass Around $95\text{-GeV}/c^2$ at the CERN SPS Collider”, *Phys. Lett.* **B126** (1983) 398–410. doi:10.1016/0370-2693(83)90188-0.
- [18] Particle Data Group Collaboration, “Review of Particle Physics”, *Chin. Phys.* **C38** (2014) 090001. doi:10.1088/1674-1137/38/9/090001.
- [19] ATLAS, CMS Collaboration, “Combined Measurement of the Higgs Boson Mass in pp Collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS Experiments”, *Phys. Rev. Lett.* **114** (2015) 191803, arXiv:1503.07589. doi:10.1103/PhysRevLett.114.191803.
- [20] G. ’t Hooft, C. Itzykson, A. Jaffe et al., “Recent Developments in Gauge Theories. Proceedings, Nato Advanced Study Institute, Cargese, France, August 26 - September 8, 1979”, *NATO Sci. Ser. B* **59** (1980) pp.1–438.
- [21] J. E. Gunn and B. A. Peterson, “On the Density of Neutral Hydrogen in Intergalactic Space”, *Astrophys. J.* **142** (1965) 1633. doi:10.1086/148444.
- [22] Planck Collaboration, “Planck 2015 results. XIII. Cosmological parameters”, arXiv:1502.01589.
- [23] Yu. A. Golfand and E. P. Likhtman, “Extension of the Algebra of Poincare Group Generators and Violation of p Invariance”, *JETP Lett.* **13** (1971) 323–326. [Pisma Zh. Eksp. Teor. Fiz.13,452(1971)].
- [24] R. Haag, J. T. Lopuszanski, and M. Sohnius, “All Possible Generators of Supersymmetries of the s Matrix”, *Nucl. Phys.* **B88** (1975) 257. doi:10.1016/0550-3213(75)90279-5.

- [25] M. Drees, “An Introduction to Supersymmetry”, in *Current topics in physics. Proceedings, Inauguration Conference of the Asia-Pacific Center for Theoretical Physics (APCTP), Seoul, Korea, June 4-10, 1996. Vol. 1, 2.* 1996.
arXiv:hep-ph/9611409.
- [26] I. J. R. Aitchison, “Supersymmetry and the MSSM: An Elementary introduction”,
arXiv:hep-ph/0505105.
- [27] M. Drees, R. Godbole, and P. Roy, “Theory and phenomenology of sparticles: An account of four-dimensional N=1 supersymmetry in high energy physics”. World Scientific, 2004.
- [28] U. Amaldi, W. de Boer, and H. Furstenau, “Comparison of grand unified theories with electroweak and strong coupling constants measured at LEP”, *Phys. Lett.* **B260** (1991) 447–455. doi:10.1016/0370-2693(91)91641-8.
- [29] MSSM Working Group Collaboration, A. Djouadi et al., “The Minimal supersymmetric standard model: Group summary report”, in *GDR (Groupement De Recherche) - Supersymetrie Montpellier, France, April 15-17, 1998.* 1998.
arXiv:hep-ph/9901246.
- [30] H. P. Nilles, “Supersymmetry, Supergravity and Particle Physics”, *Phys. Rept.* **110** (1984) 1–162. doi:10.1016/0370-1573(84)90008-5.
- [31] G. F. Giudice and R. Rattazzi, “Theories with gauge mediated supersymmetry breaking”, *Phys. Rept.* **322** (1999) 419–499, arXiv:hep-ph/9801271.
doi:10.1016/S0370-1573(99)00042-3.
- [32] M. Fairbairn, A. C. Kraan, D. A. Milstead et al., “Stable massive particles at colliders”, *Phys. Rept.* **438** (2007) 1–63, arXiv:hep-ph/0611040.
doi:10.1016/j.physrep.2006.10.002.
- [33] CMS Collaboration, “Phenomenological MSSM interpretation of the CMS 2011 5 fb^{-1} results”, *CMS Physics Analysis Summary* **CMS-PAS-SUS-12-030** (2013).
- [34] M. W. Cahill-Rowley, J. L. Hewett, A. Ismail et al., “More energy, more searches, but the phenomenological MSSM lives on”, *Phys. Rev.* **D88** (2013), no. 3, 035002, arXiv:1211.1981. doi:10.1103/PhysRevD.88.035002.
- [35] M. W. Cahill-Rowley, J. L. Hewett, S. Hoeche et al., “The New Look pMSSM with Neutralino and Gravitino LSPs”, *Eur. Phys. J.* **C72** (2012) 2156,
arXiv:1206.4321. doi:10.1140/epjc/s10052-012-2156-1.

- [36] ALEPH Collaboration, “Search for charginos nearly mass degenerate with the lightest neutralino in $e^+ e^-$ collisions at center-of-mass energies up to 209-GeV”, *Phys. Lett.* **B533** (2002) 223–236, [arXiv:hep-ex/0203020](#).
[doi:10.1016/S0370-2693\(02\)01584-8](#).
- [37] OPAL Collaboration, “Search for nearly mass degenerate charginos and neutralinos at LEP”, *Eur. Phys. J.* **C29** (2003) 479–489, [arXiv:hep-ex/0210043](#).
[doi:10.1140/epjc/s2003-01237-x](#).
- [38] DELPHI Collaboration, “Searches for supersymmetric particles in $e^+ e^-$ collisions up to 208-GeV and interpretation of the results within the MSSM”, *Eur. Phys. J.* **C31** (2003) 421–479, [arXiv:hep-ex/0311019](#). [doi:10.1140/epjc/s2003-01355-5](#).
- [39] DELPHI Collaboration, “Search for SUSY in the AMSB scenario with the DELPHI detector”, *Eur. Phys. J.* **C34** (2004) 145–156, [arXiv:hep-ex/0403047](#).
[doi:10.1140/epjc/s2004-01715-7](#).
- [40] ALEPH, DELPHI, L3 and OPAL Collaborations, “Joint SUSY Working Group”.
<http://lepsusy.web.cern.ch/lepsusy>. LEPSUSYWG (June 2004).
- [41] ATLAS Collaboration, “Search for direct chargino production in anomaly-mediated supersymmetry breaking models based on a disappearing-track signature in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *JHEP* **01** (2013) 131, [arXiv:1210.2852](#). [doi:10.1007/JHEP01\(2013\)131](#).
- [42] ATLAS Collaboration, “Search for charginos nearly mass degenerate with the lightest neutralino based on a disappearing-track signature in pp collisions at $\sqrt{s}=8$ TeV with the ATLAS detector”, *Phys. Rev.* **D88** (2013), no. 11, 112006, [arXiv:1310.3675](#). [doi:10.1103/PhysRevD.88.112006](#).
- [43] ATLAS Collaboration, “Search for metastable heavy charged particles with large ionisation energy loss in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS experiment”, *Eur. Phys. J.* **C75** (2015), no. 9, 407, [arXiv:1506.05332](#).
[doi:10.1140/epjc/s10052-015-3609-0](#).
- [44] G. F. Giudice, M. A. Luty, H. Murayama et al., “Gaugino mass without singlets”, *JHEP* **12** (1998) 027, [arXiv:hep-ph/9810442](#).
[doi:10.1088/1126-6708/1998/12/027](#).
- [45] ATLAS collaboration, “Summary plots from the ATLAS Supersymmetry physics group”. <https://atlas.web.cern.ch/Atlas/GROUPS/PHYSICS/CombinedSummaryPlots/SUSY/>.
Accessed: 2016-01-17.

-
- 516 [46] CMS Collaboration, “Searches for long-lived charged particles in pp collisions at
517 $\sqrt{s}=7$ and 8 TeV”, *JHEP* **07** (2013) 122, [arXiv:1305.0491](#).
518 [doi:10.1007/JHEP07\(2013\)122](#).
- 519 [47] CMS Collaboration, “Search for disappearing tracks in proton-proton collisions at
520 $\sqrt{s} = 8$ TeV”, *JHEP* **01** (2015) 096, [arXiv:1411.6006](#).
521 [doi:10.1007/JHEP01\(2015\)096](#).

