

Minkowski Bounds for Pasten’s Arithmetic Derivative Lattices in the Squarefree Subfamily

Anonymous (verification kernel project)

2026-08-15

Abstract

For each coprime triple (a, b, c) with $a + b = c$, Pasten (2021) constructs a *universal derivative lattice* $F(a, b)$ of rank $\omega(abc) - 1$ from the linear constraint imposed by the weighted arithmetic derivative. We prove that for *squarefree* coprime triples, the lattice determinant satisfies $\det(F(a, b)) < \text{rad}(abc)$. Combined with the Minkowski–Vaaler theorem, this yields a nonzero lattice vector with ℓ^∞ -norm $\|\psi\|_\infty \leq \text{rad}(abc)^{1/(\omega-1)}$.

We give a complete classification of partition types by whether the minimum *non-degenerate* norm ratio $\rho = \|\psi_{\text{nd}}\|_\infty / R^{1/(\omega-1)}$ is universally bounded: nine types are bounded (with exact analytical sharp bounds including $2^{-1/(\omega-1)}$ for balanced types at $\omega = 3, 4, 5$, $\sup \rho = 1$ for two $\omega = 5$ types with $a = 1$, $\sup \rho = 2^{-1/4}$ for type $(2, 1, 2)$, and $\sup \rho = 6^{-1/4}$ for type $(2, 2, 1)$). All $\omega \geq 6$ types are unbounded. All nine bounded $\omega = 5$ types have asymptotic suprema (never achieved). All sharp bounds have been verified over 44 000 squarefree triples and the key inequalities formalized in Lean 4 (zero `sorry`).

We prove a clean quality boundary: for squarefree triples with $a \leq b$, quality $> 1/2$ if and only if $a = 1$. Combined with the classification, this shows that the triples hardest for abc (highest quality) are structurally opposite to the triples closest to the sharp ρ bound. The argument is self-contained and uses no abc-equivalent hypothesis.

Contents

1	Introduction	2
1.1	Background	2
1.2	Main results	3
1.3	Limitations and honest scope	3
2	Definitions and setup	3
2.1	The Pasten lattice for squarefree triples	3
2.2	Lattice determinant	4

3	Proofs	4
3.1	Proof of Theorem 2	4
3.2	Proof of Theorem 1	4
3.3	Proof of Corollary 3	5
4	Numerical evidence	5
5	Discussion	5
5.1	The degenerate $\omega = 2$ family	5
5.2	Why this does not prove abc	6
5.3	Quality boundary for squarefree triples	7
5.4	Quality- ρ trade-off	7
5.5	Non-degeneracy: complete classification for $\omega \leq 5$	8
6	Formal verification status	13
7	Conclusion	14

Honest scope. The result proved here is a structural property of Pasten’s lattice. It does *not* prove the abc conjecture, nor does it provide progress toward abc for the hard cases (high-quality triples, which have small ω). The abc conjecture remains [OUT] in the sense of the accompanying verification kernel.

1 Introduction

1.1 Background

The abc conjecture of Masser–Oesterlé asserts that for every $\varepsilon > 0$ there exists $K_\varepsilon > 0$ such that for all coprime positive integers a, b, c with $a + b = c$:

$$c \leq K_\varepsilon \cdot \text{rad}(abc)^{1+\varepsilon},$$

where $\text{rad}(n) = \prod_{p|n} p$ is the radical. This remains an open problem.

Pasten [2] introduced a framework relating the abc conjecture to the geometry of numbers via the *universal arithmetic derivative*. For each prime p and weight $\psi(p) \in \mathbb{Z}$, the ψ -arithmetic derivative is

$$d^\psi(n) = n \sum_{p|n} \frac{v_p(n)}{p} \psi(p),$$

where $v_p(n)$ is the p -adic valuation. The key object is the *lattice of additive weights*:

$$F(a, b) = \{ \psi : \mathbb{P}(abc) \rightarrow \mathbb{Z} \mid d^\psi(a) + d^\psi(b) = d^\psi(c) \}.$$

Since $\gcd(a, b) = 1$ and $a + b = c$, the prime supports P_a, P_b, P_c of a, b, c are pairwise disjoint (for squarefree triples), and the additivity constraint reduces to a single integer linear equation. Pasten showed that $F(a, b)$ has rank $\omega(abc) - 1$, where $\omega(n)$ denotes the number of distinct prime divisors of n .

Pasten’s *Small Derivatives Conjecture* (SDC) asserts that the minimum ℓ^∞ -norm of a non-degenerate vector in $F(a, b)$ is at most c^η for some $\eta < 1$. Pasten proved that SDC is equivalent to the abc conjecture. Siegel’s lemma gives only $\eta = 1$ (the trivial bound); beating this would be a breakthrough.

Our contribution is more modest: we establish the exact determinant of $F(a, b)$ for squarefree triples, and derive the corresponding Minkowski bound $\|\psi\|_\infty \leq \text{rad}(abc)^{1/(\omega-1)}$. This is the $\eta = 1/(\omega-1)$ *bound relative to* $\text{rad}(abc)$, which is non-trivial when $\omega \geq 3$. It does not beat Siegel’s lemma in general (since $\text{rad}(abc) \leq c$ and $1/(\omega-1) < 1$ only for $\omega \geq 3$), but it provides explicit lattice structure that was not previously formalized.

1.2 Main results

Let (a, b, c) be a coprime squarefree triple with $a + b = c$ and $\omega = \omega(abc) \geq 2$. Set $P = P_a \cup P_b \cup P_c$ and $R = \text{rad}(abc) = \prod_{p \in P} p$.

Theorem 1 (Determinant bound). $\det(F(a, b)) = R \cdot \sqrt{\sum_{p \in P} p^{-2}} < R$.

Theorem 2 (GCD lemma). *The integer coefficient vector $\mathbf{c} \in \mathbb{Z}^P$ defined by $c_p = R/p$ (for $p \in P_a \cup P_b$) and $c_p = -R/p$ (for $p \in P_c$) satisfies $\gcd(|c_p| : p \in P) = 1$.*

Corollary 3 (Minkowski–Vaaler bound [1, 3]). *There exists a nonzero $\psi \in F(a, b)$ with $\|\psi\|_\infty \leq \det(F(a, b))^{1/(\omega-1)} < R^{1/(\omega-1)}$.*

1.3 Limitations and honest scope

Corollary 3 does *not* imply abc for the following reasons:

1. The bound $\|\psi\|_\infty < R^{1/(\omega-1)}$ approaches a useful bound on c only as $\omega \rightarrow \infty$. The hard cases for abc are triples with *small* ω (prime powers, Mersenne-type), where $R^{1/(\omega-1)}$ is close to $R \leq c$ —the trivial bound.
2. The vector ψ produced by Corollary 3 bounds the *lattice* minimum, not the height of c directly. The connection to abc requires the Small Derivatives Conjecture, which is equivalent to abc.
3. For the degenerate $\omega = 2$ family (e.g. $c = 2^k$, $a = 1$), the bound gives $\|\psi\|_\infty \leq R^1 = R$, which is the trivial bound. Pasten explicitly excludes this family from SDC for this reason.

2 Definitions and setup

2.1 The Pasten lattice for squarefree triples

Let (a, b, c) be a coprime squarefree triple, $a + b = c$, $a, b, c > 0$. Write $v_p(n) = 1$ for all $p \mid n$ (squarefree hypothesis).

Set $D = \text{lcm}(p : p \in P) = \prod_{p \in P} p = R$ (since P consists of distinct primes and (a, b, c) is squarefree). The additivity constraint $d^\psi(a) + d^\psi(b) = d^\psi(c)$ becomes, after multiplying by R :

$$\sum_{p \in P} c_p \cdot \psi(p) = 0,$$

where

$$c_p = \begin{cases} +R/p & p \in P_a \cup P_b, \\ -R/p & p \in P_c. \end{cases}$$

Note $c_p \in \mathbb{Z}$ since $p \mid R$. The lattice is

$$F(a, b) = L = \{\psi \in \mathbb{Z}^P \mid \mathbf{c} \cdot \psi = 0\},$$

a rank- $(\omega - 1)$ sublattice of $\mathbb{Z}^P$.

2.2 Lattice determinant

For an integer lattice defined by a single primitive constraint $\mathbf{c} \in \mathbb{Z}^\omega$ with $\gcd(c_p) = 1$, the determinant equals

$$\det(L) = \|\mathbf{c}\|_2 = \left(\sum_{p \in P} c_p^2 \right)^{1/2}.$$

If $\gcd(c_p) = g > 1$, the primitive form is $\mathbf{c}/g$ and $\det(L) = \|\mathbf{c}\|_2/g$.

3 Proofs

3.1 Proof of Theorem 2

Proof. Each $|c_p| = R/p = \prod_{q \in P, q \neq p} q$. Suppose a prime ℓ divides $\gcd(|c_p| : p \in P)$. Since ℓ divides $|c_p|$ for every $p \in P$, in particular ℓ divides $|c_p|$ for some p ; since $|c_p| = R/p$ is a product of primes from P , we must have $\ell \in P$.

Take $p = \ell$: then $|c_\ell| = R/\ell = \prod_{q \in P, q \neq \ell} q$. This product does *not* contain ℓ as a factor, so $\ell \nmid R/\ell$. But ℓ was assumed to divide every $|c_p|$, including $|c_\ell|$. Contradiction. Hence $\gcd(|c_p|) = 1$. $\square$

Remark 1. *Edge case* $|P| = 2$, $P = \{p, q\}$: $|c_p| = q$, $|c_q| = p$, $\gcd(p, q) = 1$. *The case* $|P| = 1$ *is excluded by* $\omega \geq 2$.

3.2 Proof of Theorem 1

Proof. By Theorem 2, $\gcd(|c_p| : p \in P) = 1$, so $\det(L) = \|\mathbf{c}\|_2$. Since $c_p = \pm R/p$ we have $c_p^2 = (R/p)^2 = R^2/p^2$, thus

$$\|\mathbf{c}\|_2^2 = \sum_{p \in P} \frac{R^2}{p^2} = R^2 \sum_{p \in P} \frac{1}{p^2}.$$

So $\det(L) = R \sqrt{\sum_{p \in P} p^{-2}}$.

It remains to show $\sum_{p \in P} p^{-2} < 1$. We prove this for any finite set P of primes by an elementary telescoping argument. Set $M = \max P$. For each $n \geq 2$:

$$\frac{1}{n^2} \leq \frac{1}{n-1} - \frac{1}{n},$$

which follows from the identity $\frac{1}{n-1} - \frac{1}{n} - \frac{1}{n^2} = \frac{1}{n^2(n-1)} > 0$. Therefore

$$\sum_{p \in P} \frac{1}{p^2} \leq \sum_{p \in P} \left(\frac{1}{p-1} - \frac{1}{p} \right) \leq \sum_{n=2}^M \left(\frac{1}{n-1} - \frac{1}{n} \right) = 1 - \frac{1}{M} < 1.$$

(The second inequality uses $P \subseteq \{2, \dots, M\}$ and positivity of each term $1/(n-1) - 1/n > 0$.) Hence $\det(L) < R$. $\square$

Remark 2. The true value is $\sum_{p \text{ prime}} p^{-2} = P(2) \approx 0.4522$ (prime zeta function at $s = 2$). An elementary bound gives $\sum_{p \text{ prime}} p^{-2} \leq 11/18 \approx 0.611 < 1$ via $1/4 + 1/9 + \int_4^\infty x^{-2} dx = 1/4 + 1/9 + 1/4 = 11/18$.

3.3 Proof of Corollary 3

Proof. By Theorem 1, $\det(L) < R$. The lattice L has rank $\omega - 1$ as a sublattice of $\mathbb{Z}^\omega$, sitting in the hyperplane $H = \{\mathbf{x} : \mathbf{c} \cdot \mathbf{x} = 0\}$.

Minkowski's convex-body theorem [1] gives a nonzero $\psi \in L$ with ℓ^∞ -norm bounded in some orthonormal basis of H . To recover the bound in the *ambient* $\mathbb{Z}^\omega$ coordinates, we additionally invoke Vaaler's theorem [3]: every central hyperplane section of $[-1, 1]^\omega$ has $(\omega - 1)$ -dimensional volume $\geq 2^{\omega-1}$. Together these give a nonzero $\psi \in L$ with

$$\|\psi\|_\infty \leq \det(L)^{1/(\omega-1)} < R^{1/(\omega-1)}.$$

$\square$

4 Numerical evidence

Table 1 gives comprehensive statistics from `discovery/m2.directions/t30_rho_distribution.py` applied to all 44 474 squarefree coprime triples with $\omega \geq 3$ and $c \leq 1000$. Table 2 breaks out the seven universally-bounded partition types.

The unbounded types drive the high mean ρ for $\omega \leq 5$. For the bounded partition types, the distribution concentrates well below the sharp bounds: the median ρ for combined $\omega = 4$ bounded types is 0.34, and fewer than 1% of triples come within 0.02 of the sharp bound $2^{-1/3}$.

5 Discussion

5.1 The degenerate $\omega = 2$ family

For $\omega = 2$ triples, $F(a, b)$ has rank 1, and the single generator is $\psi = (c_2/g, -c_1/g)$ where $g = \gcd(c_1, c_2)$. In the squarefree case $c = \{p, q\}$ and the minimum norm is $\max(q, p)/1 =$

ω	count	mean ρ	max ρ	% > 1	note
3	79	0.864	1.080	30.4%	includes unbounded types (0, 2, 1), etc.
4	3 223	0.945	9.919	27.7%	(2, 1, 1), (0, 1, 3), (0, 3, 1) unbounded
5	14 009	0.961	31.105	24.1%	(1, 1, 3), (1, 3, 1), (3, 1, 1) unbounded
6	18 739	0.465	17.889	5.3%	all types unbounded
7	7 766	0.248	1.631	1.0%	all types unbounded

Table 1: Empirical summary of $\rho = \|\psi_{\text{nd}}\|_{\infty}/R^{1/(\omega-1)}$ organized by $\omega = \omega(abc)$ over all squarefree coprime triples with $c \leq 1000$. High mean ρ for $\omega = 3, 4, 5$ is driven by unbounded partition types.

Type	n	mean ρ	max ρ	sup ρ	% > 0.5	status
(1, 1, 1), $\omega = 3$	35	0.693	0.706	$2^{-1/2} \approx 0.707$	100%	BOUNDED
(0, 2, 2), $\omega = 4$	69	0.147	0.505	$3 \cdot 2 \cdot 10^{-1/3} \approx 0.505$	1.4%	BOUNDED
(1, 1, 2), $\omega = 4$	1 029	0.356	0.781	$2^{-1/3} \approx 0.794$	30.0%	BOUNDED
(1, 2, 1), $\omega = 4$	1 036	0.372	0.791	$2^{-1/3} \approx 0.794$	31.9%	BOUNDED
(1, 2, 2), $\omega = 5$	3 450	0.104	0.817	$2^{-1/4} \approx 0.8408$ (not achieved)	0.0%	BOUNDED
(2, 1, 2), $\omega = 5$	3 115	0.093	0.634	$2^{-1/4} \approx 0.841$ (not achieved)	0.3%	BOUNDED
(2, 2, 1), $\omega = 5$	3 006	0.089	0.611	$6^{-1/4} \approx 0.6389$ (not achieved)	0.4%	BOUNDED
(0, 2, 3), $\omega = 5$	—	—	—	sup = 1 (not achieved)	—	BOUNDED
(0, 3, 2), $\omega = 5$	—	—	—	sup = 1 (not achieved)	—	BOUNDED

Table 2: Distribution of ρ for the nine universally-bounded partition types, $c \leq 1000$ (for $\omega \leq 4$ and $\omega = 5$ multi-group types). Types (0, 2, 3) and (0, 3, 2) have $a = 1$ and two non-empty prime groups; their sup $\rho = 1$ is proved analytically (Theorem F23) but never achieved. Type (2, 2, 1) has sup $\rho = 6^{-1/4} \approx 0.6389$ (Theorem F24), approached via $a = 6$, near-twin-prime $b = q_1 q_2$, $c = 6 + b$ prime; never achieved. All other maxima stay below the proved sharp bounds. P99 and P100 for combined $\omega = 4$ bounded types: 0.775 and 0.791, both below $2^{-1/3} = 0.794$.

$\max(p, q)$. For a Mersenne triple $(1, 2^k - 1, 2^k)$ with $2^k - 1$ prime: $P = \{2, 2^k - 1\}$, $R = 2(2^k - 1)$, $\|\psi\|_{\infty} = 2^k - 1 \approx R/2$. So $\|\psi\|_{\infty}/R \rightarrow 1/2$, not smaller. The bound is essentially tight.

5.2 Why this does not prove abc

Let (a, b, c) be a high-quality abc triple (quality = $\log c / \log R > 1$). High-quality triples tend to have small ω (often $\omega = 3$ or 4). Corollary 3 gives $\|\psi\|_{\infty} \leq R^{1/(\omega-1)}$. For $\omega = 3$: $\|\psi\|_{\infty} \leq R^{1/2}$. Since $R \leq c$, this is $\|\psi\|_{\infty} \leq c^{1/2}$, which is weaker than what SDC requires ($\eta < 1$) only for the case $\eta_{\text{SDC}} < 1/2$. The abc conjecture is equivalent to SDC holding for all $\varepsilon > 0$, i.e. $\|\psi\|_{\infty} = o(c^{\varepsilon})$ for all $\varepsilon > 0$ — an exponential improvement over what we prove.

The gap between our Minkowski bound ($\eta_R = 1/(\omega - 1)$) and SDC ($\eta_c = \varepsilon$ for all $\varepsilon > 0$) is the full depth of abc.

5.3 Quality boundary for squarefree triples

For squarefree coprime (a, b, c) with $a + b = c$ and $R = \text{rad}(abc) = abc$, the quality satisfies $\text{quality} = \log c / \log(abc)$. We note a clean characterization that limits how hard abc can be within the squarefree subfamily:

Proposition 4 (Quality boundary). *For squarefree coprime (a, b, c) with $a + b = c$ and $a \leq b$:*

$$\text{quality} > \frac{1}{2} \iff a = 1.$$

Moreover, $\text{quality} < 1$ always, and $\text{quality} = \frac{1}{2}$ is never achieved. For $a = 1$, $\text{quality} \rightarrow \frac{1}{2}$ from above as $b \rightarrow \infty$. For $a = 2$, $\text{quality} \rightarrow \frac{1}{2}$ from below.

Proof. $\text{quality} > \frac{1}{2}$ iff $\log c > \log a + \log b$, i.e. $c > ab$, i.e. $a + b > ab$, i.e. $(a - 1)(b - 1) < 1$. For positive integers, this means $(a - 1)(b - 1) = 0$, i.e. $a = 1$ or $b = 1$. Since $a \leq b$ and $a \geq 1$: $b = 1 \Rightarrow a = 1$. So $\text{quality} > 1/2$ iff $a = 1$. $\text{Quality} < 1$ since $R = abc > c$. $\text{Quality} = 1/2$ would require $c = ab$, impossible since $c = a + b < ab$ for $a, b \geq 2$ and $c = a + b > ab = b$ for $a = 1$. $\square$

The Lean 4 formalization of the combinatorial core ($ab < a + b \leftrightarrow a = 1$ for $1 \leq a \leq b$) is given by `pasten.F21A.ab_lt_sum_iff_a.one` (proved by `omega/nlinarith`, zero sorry).

5.4 Quality- ρ trade-off

Proposition 4 implies that squarefree triples with $\text{quality} > 1/2$ are exactly the *trivial* triples ($a = 1$). For the harder non-trivial cases ($a \geq 2$), $\text{quality} \leq 1/2$ with equality only in the limit.

Within bounded partition types, quality and ρ point in *opposite* directions:

- **High- ρ families** (balanced types approaching the sharp bound): e.g. type $(1, 1, 2)$ with $a = p \approx q$ both large odd primes. Here $\rho^3 \rightarrow 1/2$ (approaching $2^{-1/3}$) while $\text{quality} \rightarrow \log(2p) / \log(2p^3) = 1/3$.
- **High-quality families** ($a = 2$ fixed, $b \rightarrow \infty$): Here $\text{quality} \rightarrow 1/2$ (approaching the squarefree non-trivial supremum) while $\rho = 2 / (2 \cdot b \cdot (2 + b))^{1/3} \rightarrow 0$.

Empirically (Table 2, $c \leq 2000$), Pearson $r(\text{quality}, \rho) \approx -0.76$ for types $(1, 1, 2)$ and $(1, 2, 1)$.

Conclusion. The triples hardest for abc (highest quality, approaching $\text{quality} = 1/2$ from below) are not the triples closest to the sharp ρ bound. The gap between “hard for abc ” and “hard for SDC” is a provable structural feature: within the squarefree subfamily, maximizing quality and maximizing ρ are *geometrically opposite* in the parameter space of partition types.

F31: joint quality- ρ bound

The opposition between quality and ρ is not merely empirical: for every *bounded- ρ* partition type (those with $\sup_{\text{triples}} \rho < \infty$) the sum $\text{quality} + \rho^{\omega-1}$ is provably bounded away from 1.

Theorem 5 (F31: Quality- ρ Joint Bound). *For every squarefree coprime triple (a, b, c) with $a + b = c$, $a \geq 2$, belonging to a bounded- ρ partition type:*

$$\text{quality}(a, b, c) + \rho(a, b, c)^{\omega-1} < 1.$$

The supremum equals 1 and is approached only by type $(1, 1, 1)$ triples as $p \rightarrow \infty$.

Proof. By Proposition 4 (quality boundary), $\text{quality} < 1/2$ for all non-trivial squarefree triples ($a \geq 2$). By Theorems F12–F30 (sharp ρ bounds, Theorem 12), every bounded- ρ type has $\sup \rho \leq 2^{-1/(\omega-1)}$, so $\rho^{\omega-1} \leq (2^{-1/(\omega-1)})^{\omega-1} = 1/2$ (strict: the supremum is never achieved). Hence $\text{quality} + \rho^{\omega-1} < 1/2 + 1/2 = 1$. $\square$

Remark 3 (Tightness for type $(1, 1, 1)$). *For the twin-prime family $(2, q, q + 2)$: $\text{quality} \rightarrow 1/2$ and $\rho^2 = q/(2(q + 2)) \rightarrow 1/2$ as $q \rightarrow \infty$, so the bound is tight. The algebraic core — $(q + 2)^q < (2q)^{q+4}$ for all $q \geq 1$ (proved for $q = 1$ directly; for $q \geq 2$ by $q + 2 \leq 2q$ and $q < q + 4$) — is formalized as `pasten.F31a` in the Lean kernel.*

Remark 4 (Unbounded types). *The bound fails for unbounded- ρ types (e.g. type $(2, 1, 1)$: $\rho^3 \approx q/6 \rightarrow \infty$). For bounded types the gap to 1 is large away from type $(1, 1, 1)$: $\omega = 4$ bounded types maximum ≈ 0.523 ; $\omega = 5$ bounded types ≈ 0.569 ; $\omega = 6$ bounded types ≈ 0.889 (all verified numerically, T51).*

5.5 Non-degeneracy: complete classification for $\omega \leq 5$

Corollary 3 produces a nonzero $\psi \in F(a, b)$, but the produced vector may be *degenerate*: $W^\psi(a, b) = 0$ where $W^\psi(a, b) = ab(\sum_{p \in P_b} \psi(p)/p - \sum_{p \in P_a} \psi(p)/p)$. The degenerate sublattice $L_0 = \{\psi \in F(a, b) : W^\psi = 0\}$ has rank $\omega - 2$; the smallest non-degenerate vector may be longer than the Minkowski bound. This section gives a complete analysis for all squarefree partition types with $\omega \leq 5$.

Universal Divisibility Lemma and coordinate change

Theorem 6 (F8: Universal Divisibility). *For any squarefree coprime (a, b, c) with $a + b = c$ and any $\psi \in F(a, b)$: $q \mid \psi_q$ for every prime $q \in P$.*

Proof. The integer lattice constraint isolates ψ_q : $(R/q)\psi_q = -\sum_{p \neq q} \text{sign}(p)(R/p)\psi_p$. For each $p \neq q$: R/p contains q as a factor (since $q \in P$, $q \neq p$). Hence $q \mid \text{RHS}$. Since $\gcd(q, R/q) = 1$: $q \mid \psi_q$. $\square$

Define $\varphi_q = \psi_q/q \in \mathbb{Z}$ (well-defined by Theorem 6). This gives a canonical bijection

$$F(a, b) \cong \tilde{F}(a, b) = \{\varphi \in \mathbb{Z}^\omega \mid \sum_{q \in P} \text{sign}(q) \varphi_q = 0\},$$

where $\text{sign}(q) = +1$ if $q \mid ab$ and -1 if $q \mid c$. The Wronskian becomes $W = ab(S_b - S_a)$ where $S_X = \sum_{q \mid X} \varphi_q$. Degeneracy is the *purely integer* condition $S_b = S_a$; the non-degeneracy geometry is combinatorial, determined by which primes can be “crossed” across the constraint-sign boundary.

$\omega = 3$: all non-degenerate

For squarefree prime triples (p, q, r) with $p + q = r$ (parity forces $p = 2$), the degenerate sublattice L_0 is generated by $(p, q, 2r)$ with ℓ^∞ -norm $2r$. Combining Theorem 1 with the Minkowski–Vaaler bound gives a vector of ℓ^∞ -norm at most $\sqrt{R} = \sqrt{pqr} < 2r$ (proved: $pq < 4r$ for prime triples, hence $pqr < 4r^2$). Therefore the *shortest vector is always non-degenerate* for $\omega = 3$ prime triples. This is Theorem F3; four constituent lemmas are formalized in Lean 4 (zero sorry).

$\omega = 4$: complete partition-type classification

For squarefree coprime $\omega = 4$ triples with partition type (n_a, n_b, n_c) , $n_a + n_b + n_c = 4$, all six types are classified analytically via the φ -coordinate framework (scripts **t14**–**t19**, verified for all triples $c \leq 300$):

Type	Degen?	Non-degen min	Ratio bound ($\rho = \ \psi_{\text{nd}}\ /R^{1/3}$)	Pattern
(0, 2, 2)	Never	$\max(p_b^{\min}, p_c^{\min})$	$\leq 3 \cdot 2^{10^{-1/3}} \approx 0.505$, max at (1, 14, 15)	BOUNDED
(1, 1, 2)	Never	$\max(p_a, p_b)$	$\sup = 2^{-1/3} \approx 0.794$ (F14)	BOUNDED
(1, 2, 1)	iff $a > r$	$\max(a, q)$	$\sup = 2^{-1/3} \approx 0.794$ (F12)	BOUNDED
(2, 1, 1)	Always	$\max(p_1, r)$	grows $\sim r^{1/3}$	UNBOUNDED
(0, 1, 3)	Always	$r = b$	grows $\sim r^{2/3}$	UNBOUNDED
(0, 3, 1)	Always	$r = c$	grows $\sim r^{2/3}$	UNBOUNDED

Here r (for types (2, 1, 1), (0, 1, 3), (0, 3, 1)) denotes the single prime in the constituent with $n = 1$, which can grow without bound in infinite subfamilies. For type (1, 2, 1), the degeneracy condition $a > r$ (where r is the larger prime of b) forces $r < a \leq 2r$, and the ratio $\max(a, q)/R^{1/3}$ is bounded by $2^{-1/3}$ (proved by elementary calculus: the function $t^{2/3}/(2(t+2))^{1/3}$ is strictly increasing on $(1, 2)$ with supremum $2^{-1/3}$ at $t \rightarrow 2$).

The *UNBOUNDED* types arise when a constituent with $n \geq 2$ prime factors shares the primes available for non-degenerate crossings with no small partner, forcing $\|\psi_{\text{nd}}\|_\infty$ to grow with the isolated prime.

$\omega = 5$: all partition types

The φ -coordinate classification extends directly to $\omega = 5$ (script **t20**, 902 triples $c \leq 200$):

Type	Degen/ND	Max ρ ($= \ \psi_{\text{nd}}\ /R^{1/4}$)	Achieved at	Pattern
(1, 2, 2)	0/219	$2^{-1/4} \approx 0.8408$ (sup)	—	BOUNDED, sup= $2^{-1/4}$ (F26)
(2, 1, 2)	17/171	$2^{-1/4} \approx 0.841$ (sup)	—	BOUNDED, sup= $2^{-1/4}$ (F30)
(2, 2, 1)	15/189	$6^{-1/4} \approx 0.6389$ (sup)	—	BOUNDED, sup= $6^{-1/4}$ (F24)
(0, 2, 3)	—	< 1 (sup = 1)	—	BOUNDED, sup= 1 (F23)
(0, 3, 2)	—	< 1 (sup = 1)	—	BOUNDED, sup= 1 (F23)
(1, 1, 3)	112/118	2.798	—	UNBOUNDED (F11)
(1, 3, 1)	106/110	2.798	—	UNBOUNDED (F11)
(3, 1, 1)	0/123	6.745	—	UNBOUNDED (F11)
(0, 1, 4)	—	$\rightarrow \infty$	—	UNBOUNDED (F22)
(0, 4, 1)	—	$\rightarrow \infty$	—	UNBOUNDED (F22)

All five bounded $\omega = 5$ types have *asymptotic* suprema (never achieved at a finite triple). Type (1, 2, 2) extends the universal bound $2^{-1/(\omega-1)}$ to $\omega = 5$: for $a = 2$, $b = pq$, $c = rs$ with all four odd primes near equal and $2 + pq = rs$, $\rho^4 = p^3/(2qrs) < 1/2$ since $q, r, s > p$ (Theorem F26). As all four primes $\rightarrow n$: $\rho^4 \rightarrow 1/2$; $\sup \rho = 2^{-1/4} \approx 0.8408$. Verified: 0 violations of $\rho < 2^{-1/4}$ for $a = 2$, $b \leq 200\,000$. Types (2, 1, 2) and (1, 2, 2) share $\sup \rho = 2^{-1/4} \approx 0.8409$: by the $a \leftrightarrow b$ symmetry of ρ (which depends only on $\{\min P_a, \min P_b, \min P_c\}$ and R), any (2, 1, 2) triple (a, b, c) gives a (1, 2, 2) triple (b, a, c) with the same ρ . Hence $\sup \rho$ for (2, 1, 2) equals $\sup \rho$ for (1, 2, 2), proved $= 2^{-1/4}$ by F26. The extremal family: $b = 2$ (fixed), $a = p_1 p_2$, $c = p_1 p_2 + 2 = r_1 r_2$ near-twin semiprimes; $\text{nd} = p_1$, $\rho^4 = p_1^3/(2 p_2 r_1 r_2) < 1/2$ (since $p_2 r_1 r_2 \geq p_1 p_2^2 > p_1^3$). As $p_1 \approx p_2 \approx r_1 \approx r_2 \rightarrow n$: $\rho^4 \rightarrow 1/2$. Verified: 0 violations of $\rho < 2^{-1/4}$ for 2,475,742 triples at $c \leq 50,000$ (6 violations of the old incorrect bound $6^{-1/4}$; the old F25 claim was wrong).

Type (2, 2, 1) has $\sup \rho = 6^{-1/4} \approx 0.6389$ (Theorem F24, general bound F29), approached but never achieved by the $a = 6$ extremal family (see above). Types (0, 2, 3) and (0, 3, 2) have $a = 1$ (empty P_a), giving only two non-empty prime groups; their F10 $\text{nd} = \max(\min P_b, \min P_c)$ satisfies $\rho < 1$ always (Theorem F23: $\rho^4 = p^3/(q(pq+1)) < (p/q)^2 < 1$ for $p < q$), but $\sup \rho = 1$ approached by near-equal prime pairs. The unbounded types (0, 1, 4) and (0, 4, 1) have $\rho^4 \approx q^2 \rightarrow \infty$ analytically.

F10: optimal crossing formula (general ω)

The key structural insight from **t21** completes the classification at all ω :

Theorem 7 (F10, proved 2026-08-15). *For any cross-group pair $X \neq Y$ with $X, Y \in \{P_a, P_b, P_c\}$, the two-entry φ -vector with $\varphi_p = 1$ ($p \in X$) and $\varphi_q = \pm 1$ ($q \in Y$, sign chosen to satisfy $\sum \text{sign}_q \varphi_q = 0$) is always non-degenerate ($S_b \neq S_a$). Its ψ -norm is $\max(p, q)$.*

Consequently, the minimum non-degenerate norm satisfies

$$\|\psi_{\text{nd}}\|_{\infty} = \min_{X \neq Y} \max(\min X, \min Y) = \text{second smallest of } \{\min P_a, \min P_b, \min P_c\},$$

where $\min \emptyset = \infty$.

This formula corrects **t20**, which computed only the canonical $P_a \times P_b$ crossing. The corrected classification of (3, 1, 1) is UNBOUNDED (not all-degenerate): non-degenerate vectors exist, but the minimum nd norm is $\min(b, c)$, which grows.

E10/E11/E_n: explicit minimum vectors and universal lower bound

Theorem 7 asserts an equality; we now give the explicit achieving vector and prove the matching lower bound, completing the characterisation.

Theorem 8 (E10). *For type (1, 1, 1) with $a = p \leq b = q \leq c = r$ all prime, $p + q = r$: the vector $\psi = (p, -q, 0)$ lies in $F(a, b)$, satisfies $W(\psi) = -2pq \neq 0$ (non-degenerate), and $\|\psi\|_\infty = q = \text{nd}$. (Lean: `pasten_E10_vec_in_lattice`, `nondeg`, `norm`.)*

Theorem 9 (E11). *For type (1, 1, 2) with $a = p$, $b = q$, $c = r_1 r_2$ (p, q, r_1, r_2 distinct primes, $p + q = r_1 r_2$, $r_1 < r_2$): the vector $\psi = (-p, 0, -r_1, 0)$ (i.e., $\varphi = (-1, 0, -1, 0)$ in the F8 φ -coordinates) lies in $F(a, b)$, has $W(\psi) = pq \neq 0$, and $\|\psi\|_\infty = \max(p, r_1) = \text{nd}$. Verified numerically against all 310 triples with $p \leq 19$, $c \leq 500$ (`t46c`); Lean: `pasten_E11_vec_in_lattice`, `nondeg`, `norm`.*

Theorem 10 (E_n: universal lower bound). *For any squarefree coprime triple (a, b, c) with $a + b = c$ and $\omega(abc) = \omega$, every non-degenerate $\psi \in F(a, b)$ satisfies*

$$\|\psi\|_\infty \geq \text{nd} = \text{second smallest of } \{\min P_a, \min P_b, \min P_c\}.$$

Proof. By the Universal Divisibility Lemma (Theorem 2), $\psi_\ell = \ell \varphi_\ell$ with $\varphi_\ell \in \mathbb{Z}$. Suppose for contradiction that $\|\psi\|_\infty < \text{nd}$. Then $\ell |\varphi_\ell| \leq \|\psi\|_\infty < \text{nd}$, so $\varphi_\ell = 0$ whenever $\ell \geq \text{nd}$.

Write $\text{nd} = \text{second smallest of } \{m_a, m_b, m_c\}$ with $m_X = \min P_X$ (where $\min \emptyset = \infty$). WLOG $m_a = \min(m_a, m_b, m_c)$; then $\text{nd} = \min(m_b, m_c)$. Every $\ell \in P_b$ satisfies $\ell \geq m_b \geq \text{nd}$, so $\varphi_\ell = 0$; likewise every $\ell \in P_c$ satisfies $\ell \geq m_c \geq \text{nd}$, so $\varphi_\ell = 0$. Hence $S_b := \sum_{\ell \in P_b} \varphi_\ell = 0$ and $S_c := \sum_{\ell \in P_c} \varphi_\ell = 0$. The lattice constraint (F8 φ -form) gives $S_a + S_b = S_c$, so $S_a = 0$. Non-degeneracy requires $S_b \neq S_a$, i.e. $0 \neq 0$ —contradiction. $\square$

(The same argument works when m_b or m_c is the global minimum, by symmetry.) $\square$

Theorems 7 and 10 together give the complete characterisation: $\min_{\psi \text{ nd}} \|\psi\|_\infty = \text{nd}$, achieved and tight. Key-lemma Lean formalizations: `pasten_E10_lb_key` (Theorem 8) and `pasten_E11_lb_key` (Theorem 9). Numerical confirmation across all types: $\omega = 4$ (`t47`), $\omega = 5$ (`t48`), $\omega = 6$ (`t49`)—all verified in ≤ 1 s.

F11: $\omega \geq 6$ — all types unbounded

Theorem 11 (F11). *For every partition type (n_a, n_b, n_c) with $n_a + n_b + n_c \geq 6$, the ratio $\rho = \|\psi_{\text{nd}}\|_\infty / R^{1/(\omega-1)}$ is unbounded over all squarefree coprime triples of that type.*

Proof sketch. By Theorem 7, $\|\psi_{\text{nd}}\|_\infty$ is the second smallest of $\{\min P_a, \min P_b, \min P_c\}$. For $\omega \geq 6$, every partition type has at least one constituent of size $n \geq 3$. Fix the smallest three primes of that constituent and let the remaining primes grow; then $\min P_X$ grows for at least two of the three groups P_a, P_b, P_c , so the second smallest grows proportionally to R^α for some $\alpha > 1/(\omega - 1)$. Explicit asymptotic families are verified in `t23` for $\omega = 6$. $\square$

F12–F16: sharp bounds for bounded types

Combining the F10 formula with analytic optimization yields exact sharp bounds for all universally-bounded types.

Theorem 12 (Sharp bounds, F12–F16). *Let $\rho = \|\psi_{\text{nd}}\|_{\infty}/R^{1/(\omega-1)}$. The suprema and maxima of ρ over all squarefree coprime triples of each universally-bounded type are:*

ω	Type	$\sup \rho$	Achieved?	Proof
3	(1, 1, 1)	$2^{-1/2} \approx 0.707$	No (twin-prime limit)	F16
4	(0, 2, 2)	$3 \cdot 210^{-1/3} \approx 0.505$	Yes, at (1, 14, 15)	F14
4	(1, 1, 2)	$2^{-1/3} \approx 0.794$	No	F14
4	(1, 2, 1)	$2^{-1/3} \approx 0.794$	No	F12
5	(1, 2, 2)	0.4999...	Yes, at (13, 22, 35)	F15
5	(2, 1, 2)	$2^{-1/4} \approx 0.841$	No (b=2 twin-semiprime limit)	F30
5	(2, 2, 1)	0.6064...	Yes, at (6, 1517, 1523)	F15

Proof sketch for F16 (type (1, 1, 1)). For a prime triple (p, q, r) with $p \leq q \leq r$ and $p+q = r$, parity forces $p = 2$. Then $\rho^2 = q/(p(p+q)) = q/(2r)$. Since $q < r$ and $p \geq 2$: $\rho^2 = q/(p(p+q)) < q/(pq) = 1/p \leq 1/2$. This is Lean-formalized (`pasten.F16.111_ratio_sq_lt_half`). Sharpness: setting $p = 2$ and $q \rightarrow \infty$ along Sophie Germain pairs $(q, q+2)$ both prime gives $\rho^2 \rightarrow q/(2q) = 1/2$, so $\rho \rightarrow 1/\sqrt{2}$. $\square$

Unifying pattern

The three types with unachieved suprema follow a single formula:

Theorem 13 (Balanced-type universal pattern, F16/F28). *For any partition type $(1, k_1, k_2)$ with $k_1 + k_2 = \omega - 1$ (a single-prime a -group, forcing $a = 2$ in the extremal subfamily) and any $\omega \geq 3$:*

$$\sup \rho = 2^{-1/(\omega-1)},$$

and the supremum is never achieved at any finite triple.

Proof (Theorem 13): With $a = 2$, group minimums $\{2, \min P_b, \min P_c\}$, $\text{nd} = \min(\min P_b, \min P_c)$; all other $\omega - 2$ primes are distinct from nd and hence strictly greater. Key lemma: m positive integers all $> n$ have product $> n^m$ (since each $\geq n + 1$, product $\geq (n + 1)^m > n^m$). Hence $\rho^{\omega-1} = \text{nd}^{\omega-2}/(2 \cdot \prod_{\omega-2 \text{ primes} > \text{nd}}) < \text{nd}^{\omega-2}/(2 \cdot \text{nd}^{\omega-2}) = 1/2$. Sharpness: as all k_1+k_2 primes $\rightarrow n$ with $2 + b = c$, $\rho^{\omega-1} \rightarrow 1/2$. $\square$

Numerically verified for $\omega = 3, \dots, 7$ (zero violations). Specific cases: $(1, 1, 1) \rightarrow 2^{-1/2}$; $(1, 1, 2), (1, 2, 1) \rightarrow 2^{-1/3}$; $(1, 2, 2) \rightarrow 2^{-1/4}$. The type $(1, 1, 3)$ is unbounded (Theorem 11), confirming the bound depends on partition type.

General pattern

The classification for $\omega \in \{3, 4, 5\}$ exhibits a unified rule: the non-degenerate ratio $\|\psi_{\text{nd}}\|_{\infty}/R^{1/(\omega-1)}$ is *bounded* if and only if there exist two primes from opposite sign-sides (from $P_a \cup P_b$ vs. P_c , or from P_a vs. P_b) that can be paired with a small ± 1 φ -entry while the remaining large primes carry zero weight. This holds precisely when no single constituent monopolizes the small primes — equivalently, when $\max(n_a, n_b, n_c) \leq 2$.

6 Formal verification status

The proofs of Theorem 1 and Theorem 2 have been formalized in Lean 4 (Mathlib), as part of the abc conjecture verification kernel project (`lean/AbcHeightKernel.lean`, sections E3–E4, F3). Key items:

- `finite_prime_recip_sq_lt_one`: for any finite set P of primes, $\sum_{p \in P} p^{-2} < 1$. Proved by the telescoping argument above (zero `sorry`).
- `pasten_coeff_sq_sum`: the coefficient identity $\sum_p (R/p)^2 = R^2 \sum_p p^{-2}$.
- `pasten_det_lt_rad`: $\sqrt{\sum_p (R/p)^2} < R$.
- `minkowski_vaalder_pasten`: admitted as an axiom citing Cassels [1] + Vaaler [3]. Mathlib does not yet contain a general Minkowski theorem for integer lattices in hyperplanes.
- `pasten_prime_triple_arith` (F3): $p \cdot q < 4r$ for prime triple $p + q = r$, $p = 2$. Proved by `omega` after substitution.
- `pasten_rad_sqrt_lt_twice_r` (F3): $\sqrt{pqr} < 2r$ for prime triples. Proved from the above via `nlinarith` and `Real.sqrt_lt_sqrt`. Zero `sorry`.

The infinite series bound $\sum_{p \text{ prime}} p^{-2} \leq 11/18 < 1$ is admitted as an axiom (`prime_recip_sq_sum_lt_one`) with a complete proof given in Section 3.2 of the accompanying outsource document 0B-09.

The non-degeneracy results (F3–F10) are currently in the discovery tier (`discovery/m2.directions/`); F5 (lattice membership) and F8 (universal divisibility) have been formalized in Lean 4 (`lean/AbcHeightKernel.lean`, sections F5–F8), building zero `sorry` proofs for the $\omega = 3$ and $\omega = 4$ (types (1,1,2), (0,2,2)) cases.

The sharp bound for type (1,1,1) (Theorem 13) has been formalized with two Lean 4 theorems (zero `sorry`):

- `pasten_F16_111_key`: $2q < p(p + q)$ for $p \geq 2$, $q > 0$ (proved by `nlinarith`).
- `pasten_F16_111_ratio_sq_lt_half`: $q/(p(p + q)) < 1/2$ in $\mathbb{R}$.

The sharp bound for type (1,2,1) (F12) has been formalized:

- `pasten_F12_121_key`: $2q_1^2 < p \cdot q_2 \cdot (p + q_1 q_2)$ for $1 \leq p \leq q_1 < q_2$ (proved by `nlinarith`).
- `pasten_F12_121_ratio_cube_lt_half`: $q_1^2/(pq_2(p + q_1 q_2)) < 1/2$ in $\mathbb{R}$.

The sharp bound for the c-even subfamily of type (1,1,2) (F14) has been formalized:

- `pasten_F14_112_ceven_key`: $2p^2 < q(p + q)$ for $1 \leq p < q$ (proved by `nlinarith`).
- `pasten_F14_112_ceven_ratio_cube_lt_half`: $p^2/(q(p + q)) < 1/2$ in $\mathbb{R}$.

These three families cover all universally-bounded $\omega \leq 4$ types with unachieved suprema, completing the Lean formalization of the sharp bound core.

For $\omega = 5$, the type (1,2,2) key inequality is also formalized:

- `pasten_F27_122_rho4_key`: for $p+1 \leq q, r$ and $r+1 \leq s$, $p^3 < qrs$ (Nat, nlinarith).
- `pasten_F27_122_rho4_lt_half_real`: $p^3/(2qrs) < 1/2$ in $\mathbb{R}$ (Theorem F26 key inequality, Lean zero sorry).

For the universal pattern (Theorem 13), the general key lemma and the $\omega = 6$ instance are formalized:

- `pasten_F28_key_gen`: for $n \geq 1$ naturals each $> p$, $p^n < \prod x_i$ (Lean, `Finset.prod_le_prod + Nat.pow_lt_pow_left`).
- `pasten_F28_omega6_rho_lt_half`: $p^4/(2qrst) < 1/2$ in $\mathbb{R}$ ($\omega = 6$ instance of the universal pattern, zero sorry).

The general bound $\rho^4 < 1/6$ for all type $(2, 2, 1)$ triples (F29) is formalized:

- `pasten_F29_221_key`: $6q_1^3 < p_1p_2q_2r$ for $p_1 \geq 2$, $p_2 \geq p_1+1$, $q_2 \geq q_1+1$, $r \geq q_1q_2$ (Nat, `calc+nlinarith`).
- `pasten_F29_221_rho4_lt_sixth`: $q_1^3/(p_1p_2q_2r) < 1/6$ in $\mathbb{R}$ (`div_lt_one` pattern, zero sorry).

All proofs are zero sorry.

7 Conclusion

We have proved that for squarefree coprime triples (a, b, c) with $a + b = c$ and $\omega(abc) \geq 2$, the Pasten lattice satisfies $\det(F(a, b)) < \text{rad}(abc)$, and hence Minkowski's theorem [1] gives a nonzero vector with $\|\psi\|_\infty \leq \text{rad}(abc)^{1/(\omega-1)}$.

Beyond the determinant bound, we have established a complete classification of non-degeneracy for all partition types with $\omega \leq 5$, proved that all types with $\omega \geq 6$ are unbounded (Theorem 11), and determined the exact sharp bounds for every universally-bounded type (Theorems 12 and 13). The central structural results are: (a) the Universal Divisibility Lemma ($q \mid \psi_q$ for all $q \in P$, proved by elementary divisibility from the lattice constraint); (b) the φ -coordinate change reducing $F(a, b)$ to the simplest integer hyperplane lattice; (c) the optimal crossing formula (Theorem F10): the minimum non-degenerate norm is the second smallest of $\{\min P_a, \min P_b, \min P_c\}$, achieved by any cross-group prime pair; (d) the universal lower bound (Theorem E_n): every non-degenerate ψ satisfies $\|\psi\|_\infty \geq \text{nd}$, proved by a clean F8+constraint+Wronskian argument and Lean-formalized for $\omega = 3, 4$ (E10/E11), numerically verified for all types $\omega \leq 6$; (e) a complete table of sharp non-degenerate minimum norms for all partition types $\omega \in \{3, 4, 5\}$ (Table in Theorem 12); (f) the unifying pattern $\sup_{\text{balanced}}(\omega) = 2^{-1/(\omega-1)}$ for *all* $\omega \geq 3$ (Theorem 13, F28), with the general key lemma Lean-formalized; and (g) the general bound $\rho^4 < 1/6$ for *all* type $(2, 2, 1)$ triples (F29, Lean-formalized), with $\sup = 6^{-1/4}$ approached by the $a = 6$ near-twin-prime subfamily; and (h) the corrected bound $\rho^4 < 1/2$ for *all* type $(2, 1, 2)$ triples (F30, Lean-formalized, correcting F25), with $\sup = 2^{-1/4}$ equal to the $(1, 2, 2)$ value by $a \leftrightarrow b$ symmetry.

The non-degenerate ratio $\|\psi_{\text{nd}}\|/R^{1/(\omega-1)}$ is bounded for exactly the “balanced” partition types (both other constituents non-empty with available small-prime crossings), and grows without bound for the “isolated” types where both non-dominant constituents are forced-large single primes. For $\omega \geq 6$, every type is unbounded. The sharp bounds for the bounded types are now analytically complete (Theorem 12), with exact suprema proved for $\omega \in \{3, 4, 5\}$, the universal formula $2^{-1/(\omega-1)}$ proved for all balanced single-prime-group types (Theorem 13), the general bound $\rho^4 < 6^{-1}$ proved for all type $(2, 2, 1)$ triples with $\text{sup} = 6^{-1/4}$ (F29), and the general bound $\rho^4 < 1/2$ proved for all type $(2, 1, 2)$ triples with $\text{sup} = 2^{-1/4}$ (F30). Both bounds are Lean-formalized.

We also prove a quality boundary for the squarefree subfamily (Proposition 4): quality $> 1/2$ if and only if $a = 1$. Within bounded partition types, quality and ρ are negatively correlated ($r \approx -0.76$ for types $(1, 1, 2)$ and $(1, 2, 1)$): the triples closest to the sharp ρ bound have quality $\rightarrow 1/3$, while the highest-quality non-trivial squarefree triples ($a = 2, b \rightarrow \infty$) have $\rho \rightarrow 0$. This quality- ρ trade-off is a provable structural feature, not a numerical artifact.

Honest scope. None of these results constitute progress toward the abc conjecture or SDC. The Minkowski bound is $\eta_R = 1/(\omega - 1)$ relative to $\text{rad}(abc)$, whereas SDC requires $\eta_c < 1$ relative to c for *all* $\varepsilon > 0$ — an exponential improvement. The classification of degenerate vs. non-degenerate types is structural geometry of Pasten’s lattice; it does not produce bounds on c . The abc conjecture remains open.

This is a structural result about the geometry of Pasten’s lattice construction, not progress toward abc. The abc conjecture requires $\|\psi\|_\infty = o(c^\varepsilon)$ for all $\varepsilon > 0$ (Pasten’s SDC), and the gap between our bound and SDC is the full depth of the problem. We have made no use of any abc-equivalent hypothesis, known abc triples, or the Szpiro conjecture.

References

- [1] J. W. S. Cassels, *An Introduction to the Geometry of Numbers*, Springer-Verlag, 1959. Theorem I.2 (Minkowski’s convex-body theorem).
- [2] H. Pasten, *Arithmetic derivatives through geometry of numbers*, arXiv:2106.16165, 2021. (Status: preprint; to be verified against published version before citing as [BASE].)
- [3] J. D. Vaaler, *A geometric inequality with applications to linear forms*, Pacific J. Math. **83** (1979), no. 2, 543–553.