

Matroid Theory

Héctor Manuel Téllez Gómez

January 27, 2014 - April 16, 2014

Contents

1	Union Of Closed Sets That Is Not Closed	2
2	First Test	3
2.1	Oxley, Section 1.1, Problem 2	3
2.2	Oxley, Section 1.1, Problem 7	7
2.3	Oxley, Section 1.1, Problem 9	8
2.4	Oxley, Section 1.2, Problem 1	9
2.5	Oxley, Section 1.2, Problem 6	10
2.6	Oxley, Section 1.3, Problem 4	11
2.7	Oxley, Section 1.3, Problem 5	12
2.8	Oxley, Section 1.4, Problem 2	13
2.9	Oxley, Section 1.4, Problem 6	14
2.10	Oxley, Section 1.6, Problem 1	16
2.11	Oxley, Section 1.6, Problem 3	17
2.12	Oxley, Section 1.6, Problem 4	18
2.13	Oxley, Section 1.7, Problem 5	19
2.14	Oxley, Section 1.8, Problem 1	20
2.15	Oxley, Section 1.8, Problem 4	23
3	Second Test	24
3.1	Oxley, Section 2.1, Problem 1	24
3.2	Oxley, Section 2.1, Problem 2	28
3.3	Oxley, Section 2.1, Problem 6	29
3.4	Oxley, Section 2.1, Problem 10	30
3.5	Oxley, Section 2.2, Problem 2	31
3.6	Oxley, Section 2.2, Problem 4	32
3.7	Oxley, Section 2.2, Problem 8	33
3.8	Oxley, Section 2.3, Problem 1	36
3.9	Oxley, Section 2.3, Problem 2	38
3.10	Oxley, Section 2.3, Problem 6	41
3.11	Oxley, Section 2.3, Problem 10	43
3.12	Oxley, Section 2.4, Problem 3	44
3.13	Oxley, Section 2.4, Problem 4	46
3.14	Oxley, Section 3.1, Problem 1	47
3.15	Oxley, Section 3.1, Problem 4	48
3.16	Oxley, Section 3.2, Problem 3	49
3.17	Oxley, Section 3.2, Problem 5	50

Note for the readers:

The goal on writing this in english instead of writing it in my first language (spanish) is because this way I can start building a portfolio that could not be as important as a research project, but it is work after all. So it would be better to have something to show that nothing to show at all, and much better if the audience capable of read it is as wide as possible.

As my first language is not english, this text could have a lot of misspellings and bad use of english. I apologize in advance and I'll try to be as careful as possible.

Here is a link to this project's version controller repository:

<https://github.com/tellezhector/matroids>

There you can see what are the changes that have occurred along all this project's lifetime.

If you want to leave comments you can do it there, where anyone else is able to notice it. Or you could write directly to my personal e-mail address, which is: `tellez.hector@gmail.com` where it will be read only by me.

1

Union Of Closed Sets That Is Not Closed

Definition 1 (Closure). *Given a matroid $M = (E, \mathcal{I})$ with rank function f . The **closure** of a set $S \subset E$ is*

$$Cl(S) = \{x \in E \mid r(S \cup \{x\}) = r(S)\}.$$

Definition 2 (Closed set). *Given a matroid $M = (E, \mathcal{I})$ with rank function f . We say that a set $S \subset E$ is **closed** if*

$$Cl(S) = S.$$

Problem 3. *Give an example of two closed sets whose union is not closed.*

— . —

Proof. We are going to use the linear matroid of \mathbb{Z}^2 , where the independent sets are

$$\emptyset, \{(1, 0)\}, \{(0, 1)\}, \{(1, 1)\}, \{(1, 0), (0, 1)\}$$

and the rank function is the size of the biggest independent set contained.

It is clear that $A = \{(1, 0), (0, 0)\}$ and $B = \{(0, 1), (0, 0)\}$ are both closed. But $C = A \cup B = \{(1, 0), (0, 1), (0, 0)\}$ is not, because $r(C) = 2 = r(C \cup \{(1, 1)\})$ and $(1, 1) \notin C$. \square

2

First Test

February 17, 2014

Test for Oxley's book first chapter (Basic definitions and examples).

2.1 Oxley, Section 1.1, Problem 2

Problem 4. Let A be the matrix

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 \end{pmatrix}$$

For q in $\{2, 3\}$, let $M_q[A]$ be the vector matroid of A when A is viewed over $GF(q)$, the field of q elements. Show that:

- (i) The sets of circuits of $M_2[A]$ and $M_3[A]$ are different.
- (ii) $M_2[A]$ is graphic but $M_3[A]$ is not.
- (iii) $M_2[A]$ is representable over $GF(3)$, but $M_3[A]$ is not representable over $GF(2)$.

— . —

Proof. (i) Lets look for the circuits over $GF(2)$. As 0 and 1 are the only scalars over $GF(2)$, we can only pay attention to sums of the vector columns and forget about full linear combinations.

There are no loops, the only way to get a loop is having a zero column and it is not the case.

There are no circuits of length two (or parallel columns), the only way to get them is having two exactly equal columns and it is not the case.

Lets call the columns 1, 2, and 3 the “simple columns”, and the columns 4, 5 and 6 the “double columns”.

For any pair of the columns from the simple columns there is a column from the double columns such that the sum is zero. Such column is the one you can get from adding the chosen pair. That is, $\{1, 2, 4\}$, $\{1, 3, 5\}$, $\{2, 3, 6\}$ are three dependent sets.

If you try to add all the simple columns, you can't add up to zero.

If you add two of the the double columns, you get the third one, so $\{4, 5, 6\}$ is dependent.

These four sets are circuits given that there are no smaller dependent sets.

These four are all the circuits of size three. Lets think about the circuits of size four.

Again, if you add two of the double columns, you get the third one, and there are two simple columns that can add up this third one. So, $\{1, 2, 5, 6\}$, $\{1, 3, 4, 6\}$, $\{2, 3, 4, 5\}$ are dependent sets. And no one of the four already given circuits are contained in them, so these are circuits themselves.

At any time you choose the three simple columns, you will get no way to add up to zero just adding double columns. So, any set containing all of the simple columns will not be a circuit.

If you choose all the double columns, you have already chosen a circuit, so if you add any other vector, you will get a dependent set that is not a circuit.

So these seven are all the circuits of $M_2[A]$.

Now lets look for the circuits over $GF(3)$. Again and for the same reason, there are not loops or parallel columns.

If you choose two simple colums and you add them, you get one of the double columns, so if you add to them the negative of this double column, you get the zero vector. So $\{1, 2, 4\}$, $\{1, 3, 5\}$, $\{2, 3, 6\}$ are circuits.

$\{4, 5, 6\}$ is not dependent, you can check that the determinant of their submatrix is 2 (or -2 if you switched something).

If you choose two doble columns and add to them a simple one, again the determinant is not zero (it is always 1 or -1). So $\{1, 2, 4\}$, $\{1, 3, 5\}$, $\{2, 3, 6\}$ are all the circuits of size three.

Now lets look for circuits of size four.

If you choose two double columns and add one to the negative of the second one, you can find two simple columns that can make them add up the zero vector (using the correct signs), these two vectors are exactly the ones that add up to the double column that you didn't choose. So none of these selections contains the circuits of size three and therefore they are circuits. So this give us the circuits $\{1, 2, 5, 6\}$, $\{1, 3, 4, 6\}$, $\{2, 3, 4, 5\}$.

If choose all of the double columns, you can choose any of the simple one so the four of them add up the zero vector using an appropriate selection for the signs. So this give us the dependent sets $\{1, 4, 5, 6\}$, $\{2, 4, 5, 6\}$, $\{3, 4, 5, 6\}$ and they are circuits because any of the smaller circuits contain at least two simple columns.

The only other chance is choosing all of the simple columns, but wathever you add, you will get any of the three sized circuits as a subset. So these six are all of the circuits of size four.


There are no circuits of size five, because there are only two possibilities, having three simple columns and two double columns, or having three double columns and two simple columns. The first one, again contains as a subset a circuit of size three. The second one always contains as a subset a circuit of size four.

So in total, there are nine circuits for $M_3[A]$.

Only by their cardinality, you can tell that the set of circuits of $M_2[A]$ and $M_3[A]$ are different.

- (ii) We are going to give a graphic representation of a graph that has $M_2[A]$ as an isomorphic matroid. You can verify that it has all the seven circuits that we gave previously and only those.

Figure 2.1:



Now let's think what could happen if $M_3[A]$. For $M_3[A]$, we have the circuits $\{1, 4, 5, 6\}$, $\{2, 4, 5, 6\}$, $\{3, 4, 5, 6\}$. Take any two of them, let's say $\{1, 4, 5, 6\}$, $\{2, 4, 5, 6\}$, they have three edges in common $\{4, 5, 6\}$.

If there is a graphic matroid isomorphic to $M_3[A]$, $\{4, 5, 6\}$ defines three edges of a 4-cycle, that is, a 3-trajectory. In any 3-trajectory we will find 4 vertices, so the fourth edge is already determined by two of those vertices (the start vertex and the last vertex), but in this case we have that $\{1, 4, 5, 6\}$, $\{2, 4, 5, 6\}$ are both 4-cycles over the same set of vertices, and so 1 and 2 have exactly the same two vertices, that is, they are parallel. But in [4.(i)] we saw that $M_3[A]$ has no parallel columns. So this is a contradiction.

- (iii) We are going to give a matrix B over $GF(3)$ such that $M_3[B]$ is isomorphic to $M_2[A]$.

We are going to take the incidence matrix for the graphic representation [2.1], and choose one "1" from each column and change it by "-1". What we are doing is giving an orientation to each of the edges. So, if a cycle is such that each edge has exactly outdegree 1 and indegree 1, it is easy to see that the sum of the respective columns will be the zero vector (each 1 will have exactly one -1 in its same row).

But let's remember that we are taking linear combination over $GF(3)$, that is, we can take a column, its negative, or take the zero vector instead. So, if we have a cycle, not necessarily "well-oriented", we can give it a "good-orientation" by multiplying its edges by -1 or -2 such way that each vertex has outdegree 1 and indegree 1. If we sum them, we will get the zero vector, and giving a "good-orientation" is no other thing than finding a linear combination that sums up 0.

So, let B be the matrix

$$\begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 & 5 & 6 \end{matrix} \\ \begin{matrix} a \\ b \\ c \\ d \end{matrix} & \begin{pmatrix} -1 & 0 & 0 & 1 & -1 & 0 \\ 0 & -1 & 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & 0 & 1 & -1 \\ 1 & 1 & 1 & 0 & 0 & 0 \end{pmatrix} \end{matrix}$$

By the explanation just given, each circuit in $M_2[A]$ is a circuit in $M_3[B]$ (note that we used the same labels in both matrices so the subsets of labels can have the same representation).

Also, note that if you choose a subset of edges such that you have leaves, the row for any leaf cannot be zero under any linear combination unless you multiply it by 0. Having said these, it is easy to see that the subsets of columns representing forests are independent.

Now, let's suppose there is a matrix C such that $M_2[C]$ is isomorphic to $M_3[A]$. Again, let's suppose that the labels are the same in both matrices. Then, you must have that the subsets of columns $\{1, 4, 5, 6\}$ and

$\{2, 4, 5, 6\}$ are circuits. So, the sum of the columns 4, 5, 6 and 1 must be the zero vector, that is the column 1 is equal to the sum of the columns 4, 5 and 6. But the same occurs with the other circuit, the sum of columns 4, 5, 6 and 2 must be zero, so the column 2 is equal to the sum of the columns 4, 5 and 6. Then we have that the columns 1 and 2 must be the equal. That is, the columns 1 and 2 are parallel. Which is a contradiction to what we proved in [4.(i)].

□

2.2 Oxley, Section 1.1, Problem 7

Problem 5. Let M_1 and M_2 be matroids on disjoint sets E_1 and E_2 and with independent sets \mathcal{I}_1 and \mathcal{I}_2 respectively. Let $E = E_1 \cup E_2$ and $\mathcal{I} = \{I_1 \cup I_2 : I_1 \in \mathcal{I}_1, I_2 \in \mathcal{I}_2\}$. Prove that (E, \mathcal{I}) is a matroid.

— . —

Proof. Lets see that (E, \mathcal{I}) satisfies I1, I2 and I3.

(I1) $\emptyset \in \mathcal{I}$. This is clear, $\emptyset \in \mathcal{I}(M_1)$ and $\emptyset \in \mathcal{I}(M_2)$, so $\emptyset = \emptyset \cup \emptyset \in \mathcal{I}$.

(I2) If $I \in \mathcal{I}$, $J \subset I$ then $J \in \mathcal{I}$. $I \in \mathcal{I}$ means that there are $I_1 \in \mathcal{I}_1$ and $I_2 \in \mathcal{I}_2$ such that $I = I_1 \cup I_2$. Given that $J \subset I$ it is clear that $J = J \cap I = J \cap (I_1 \cup I_2) = (J \cap I_1) \cup (J \cap I_2)$. We have that $(J \cap I_1) \subset I_1$ and $(J \cap I_2) \subset I_2$, given that M_1 and M_2 are matroids, then $(J \cap I_1) \in \mathcal{I}_1$ and $(J \cap I_2) \in \mathcal{I}_2$. Then $J = (J \cap I_1) \cup (J \cap I_2) \in \mathcal{I}$.

(I3) If $I, J \in \mathcal{I}$, $|J| < |I|$ then there exists $x \in I \setminus J$ such that $J \cup \{x\} \in \mathcal{I}$. By hypotesis, there are $I_1, J_1 \in \mathcal{I}_1$ and $I_2, J_2 \in \mathcal{I}_2$ such that $I = I_1 \cup I_2$ and $J = J_1 \cup J_2$. Then $|J| = |J_1| + |J_2| < |I_1| + |I_2| = |I|$. Then there must be that $|J_1| < |I_1|$ or that $|J_2| < |I_2|$. Withtout loss of generality, lets say that $|J_1| < |I_1|$, as M_1 is a matroid, then, there exists $x \in I_1 \setminus J_1$ such that $J_1 \cup \{x\} \in \mathcal{I}_1$. Then, $x \in I$ and $J \cup \{x\} = (J_1 \cup \{x\}) \cup J_2 \in \mathcal{I}$. And we are done.

□

2.3 Oxley, Section 1.1, Problem 9

Problem 6. Let M_1 and M_2 be matroids on a set E . Give an example to show that $(E, \mathcal{I}(M_1) \cap \mathcal{I}(M_2))$ need not be a matroid.

— . —

Proof. Let $E = \{a, b, c\}$.

Let $\mathcal{I}(M_1) = \{\{a, b\}, \{a, c\}, \{a\}, \{b\}, \{c\}, \emptyset\}$. Let's check that this is a matroid. It has the empty set, given any set, any of its subsets is independent. To see that the independence augmentation axiom (if $I, J \in \mathcal{I}$ and $|J| < |I|$ then there exists $x \in I \setminus J$ such that $J \cup \{x\} \in \mathcal{I}$), I will give you a table, where the first row is for the set I , the second for J and the third one for x .

I	J	x
$\{a, b\}$	$\{a\}$	b
$\{a, b\}$	$\{b\}$	a
$\{a, b\}$	$\{c\}$	a
$\{a, b\}$	\emptyset	a
$\{a, c\}$	$\{a\}$	c
$\{a, c\}$	$\{b\}$	a
$\{a, c\}$	$\{c\}$	a
$\{a, c\}$	\emptyset	a
$\{a\}$	\emptyset	a
$\{b\}$	\emptyset	b
$\{c\}$	\emptyset	c

So $(E, \mathcal{I}(M_1))$ is a matroid.

Now, let $\mathcal{I}(M_2) = \{\{a, b\}, \{b, c\}, \{a\}, \{b\}, \{c\}, \emptyset\}$, to see that $(E, \mathcal{I}(M_2))$ the first two axioms can be justified just as above, and just below is the table to check the augmentation axiom.

I	J	x
$\{a, b\}$	$\{a\}$	b
$\{a, b\}$	$\{b\}$	a
$\{a, b\}$	$\{c\}$	b
$\{a, b\}$	\emptyset	a
$\{b, c\}$	$\{a\}$	b
$\{b, c\}$	$\{b\}$	c
$\{b, c\}$	$\{c\}$	b
$\{b, c\}$	\emptyset	b
$\{a\}$	\emptyset	a
$\{b\}$	\emptyset	b
$\{c\}$	\emptyset	c

So $(E, \mathcal{I}(M_2))$ is also a matroid.

Then $\mathcal{I}' = \mathcal{I}(M_1) \cap \mathcal{I}(M_2) = \{\{a, b\}, \{a\}, \{b\}, \{c\}, \emptyset\}$ cannot be an independent set for E . It satisfies (I1) and (I2), but not I3 since $\{a, b\}, \{c\} \in \mathcal{I}'$, and $|\{c\}| < |\{a, b\}|$ but $\{c\} \cup \{a\} \notin \mathcal{I}'$ and $\{c\} \cup \{b\} \notin \mathcal{I}'$. \square

2.4 Oxley, Section 1.2, Problem 1

Problem 7. Prove that \mathfrak{B} is the collection of bases of a matroid on E if and only if \mathfrak{B} satisfies (B1) and the following two conditions:

(B2)' If $B_1, B_2 \in \mathfrak{B}$ and $e \in B_1$, then there is an element f of B_2 such that $(B_1 \setminus \{e\}) \cup \{f\} \in \mathfrak{B}$.

(B3) If $B_1, B_2 \in \mathfrak{B}$ and $B_1 \subset B_2$, then $B_1 = B_2$.

— . —

Proof.

Sufficiency

Suppose that \mathfrak{B} is the collection of bases of a matroid.

Then, it satisfies (B1) and (B2). As \mathfrak{B} is a clutter, if $B_1, B_2 \in \mathfrak{B}$, then $B_1 = B_2$. Proving that it satisfies (B3).

Now, let's suppose that $B_1, B_2 \in \mathfrak{B}$ and $e \in B_1$.

If $e \in B_1 \cap B_2$ then we could choose $f = e \in B_2$ and then $(B_1 \setminus \{e\}) \cup \{f\} = B_1 \in \mathfrak{B}$.

If $e \in B_1 \setminus B_2$ then there exists an element $f \in B_2 \setminus B_1$ such that $(B_1 \setminus \{e\}) \cup \{f\} \in \mathfrak{B}$.

Necessity

Now let's suppose that \mathfrak{B} satisfies (B1), (B2)' and (B3).

Let be B_1, B_2 in \mathfrak{B} and $x \in B_1 \setminus B_2$. By (B2)' we have that there is an element y in B_2 such that $(B_1 \setminus \{x\}) \cup \{y\} \in \mathfrak{B}$. If $y \in B_1$, then $(B_1 \setminus \{x\}) \cup \{y\} = B_1 \setminus \{x\} \in \mathfrak{B}$, but as $B_1 \setminus \{x\} \subset B_1$, (B3) implies that $B_1 \setminus \{x\} = B_1$, which is a contradiction. So $y \in B_1 \setminus B_2$, proving that \mathfrak{B} satisfies (B2) and therefore, is the collection of bases of a matroid. \square

2.5 Oxley, Section 1.2, Problem 6

Problem 8. Suppose B is a basis of a matroid M , $f \in E(M)$ and $e \in E(M) \setminus B$. Prove that $(B \cup \{e\}) \setminus \{f\}$ is a basis of M if and only if $f \in C(e, B)$.

— . —

Proof.

Sufficiency

Suppose that $f \in C(e, B)$. $C(e, B)$ is the only circuit contained in $B \cup \{e\}$ by definition. Then $(B \cup \{e\}) \setminus \{f\}$ does not contain any circuit, that means, it is independent. On the other hand, $e \notin B$ and $f \in B$, then $|(B \cup \{e\}) \setminus \{f\}| = (|B| + 1) - 1 = |B|$, therefore, $(B \cup \{e\}) \setminus \{f\}$ is a basis, as it is an independent set with the size of a basis.

Necessity

We know that $(B \cup \{e\})$ contains only one circuit (and it is $C(e, B)$). If $(B \cup \{e\}) \setminus \{f\}$ is a basis of M , then it can not contain $C(e, B)$. That is, $C(e, B) \subseteq (B \cup \{e\})$, but $C(e, B) \not\subseteq (B \cup \{e\}) \setminus \{f\}$, then f must be in $C(e, B)$.

□

2.6 Oxley, Section 1.3, Problem 4

Problem 9. *Prove that a matroid M is uniform if and only if it has no circuits of size less than $r(M) + 1$.*

— . —

Proof.

Sufficiency

Suppose that M is a uniform matroid $U_{m,n}$, then $r(M) = m$ by definition. Moreover, any $A \subset E(M)$ such that $|A| < m$ will be subset of some basis and therefore independent. Then if there is any circuit C , it must have size greater than m , that is $r(C) \geq m + 1 = r(M) + 1$.

Necessity

Suppose now that M is such that it has no circuits of size less than $r(M) + 1$. Then, for any $A \subset E(M)$ with $|A| \leq r(M)$ it must be independent, if not, then A should contain a circuit, but any circuit has size at least $r(M) + 1$.

Then M must be isomorphic to an uniform matroid $U_{r(M), |E(M)|}$. □

2.7 Oxley, Section 1.3, Problem 5

Problem 10.

- (i) Characterize paving matroids in terms of their collection of independent sets and in terms of their collections of bases.
- (ii) Characterize uniform matroids in terms of their collections of circuits.

— . —

Proof.

- (i) Let M be a matroid. \mathfrak{B} its collection of bases and \mathcal{I} its collection of independent sets.

M is a paving matroid if and only if for every $A \subset E(M)$ with $|A| < r(M)$ there exists $B \in \mathfrak{B}$ such that $A \subset B$.

As M is matroid, by I2, this is exactly the same as saying:

M is a paving matroid if and only if for every $A \subset E(M)$ with $|A| < r(M)$, $A \in \mathcal{I}$.

Proof. If M is paving, by definition of paving matroid, any $A \in E(M)$ such that $|A| < r(M)$ then, $A \in \mathcal{I}$ (and therefore, there is some $B \in \mathfrak{B}$ such that $A \subset B$).

If M is a matroid such that for every $A \subset E(M)$ with $|A| \leq r(M)$ there exists $B \in \mathfrak{B}$ such that $A \subset B$ (and therefore, $A \in \mathcal{I}$). Then, suppose that C is a circuit of M . If $|C| < r(M)$, then there would be some $B \in \mathfrak{B}$ such that $C \subset B$, but then $C \in \mathcal{I}$, which is a contradiction and therefore $|C| \geq r(M)$. That is, M is paving.

□

- (ii) A matroid M is uniform if and only if it has no circuits of size less than $r(M) + 1$.

Proof. See (9)

□

□

2.8 Oxley, Section 1.4, Problem 2

Problem 11. *Show that a subset X of a matroid is a basis if and only if X is both independent and spanning.*

— . —

Proof.

Sufficiency

If X is a basis of a matroid M , then by definition of basis, it is independent. Also, as X is basis, then $r(X) = r(X \cup x)$ for any $x \in E(M)$ and, therefore, $cl(X) = E(M)$, that is, X is spanning.

Necessity

Suppose that X is both independent and spanning. As X is spanning, then $r(X) = r(M)$, that is, for any $B \in \mathfrak{B}(M)$, $|X| \geq r(X) = r(B) = |B|$. On the other hand, $X \in \mathcal{I}(M)$ so, for any $B \in \mathfrak{B}(M)$, $|X| \leq |B|$. We have then that $|B| \leq |X| \leq |B|$, that is $|B| = |X|$ for any $\mathfrak{B}(M)$ and therefore, X is a basis. \square

2.9 Oxley, Section 1.4, Problem 6

Problem 12. Prove that statements (a)-(g) below are equivalent for an element e of a matroid M :

- (a) e is in every basis.
- (b) e is in no circuits.
- (c) If $X \subseteq E(M)$ and $e \in cl(X)$, then $e \in X$.
- (d) $r(E(M) \setminus \{e\}) = r(E(M)) - 1$.
- (e) $E(M) \setminus \{e\}$ is a flat.
- (f) $E(M) \setminus \{e\}$ is a hyperplane.
- (g) If I is an independent set, then so is $I \cup \{e\}$.

— . —

Proof.

[(a) \Rightarrow (b)]

Suppose there is a circuit C such that $e \in C$, then $C \setminus \{e\} \in \mathcal{I}(M)$, then there must be $B \in \mathfrak{B}$ such that $C \setminus \{e\} \subseteq B$, but $e \notin B$, otherwise $C \subset B$. But this contradicts that e is in every basis and therefor there is not such circuit C .

[(b) \Rightarrow (c)]

Let $X \subseteq E(M)$ such that $e \in cl(X)$. Suppose that $e \notin X$ and let B_X be a basis of X . we have that $x \in cl(X) = cl(B_X)$, then $cl(B_X) = cl(B_X \cup \{e\})$, which means that $r(B_X \cup \{e\}) = r(B_X) = |B_X| < |B_X \cup \{e\}|$, and therefore $B_X \cup \{e\}$ is a dependent set, and it must contain a circuit C , as C can not be subset of B_X (because B_X is independent) then $e \in C$, which is a contradiction to (b).

[(c) \Rightarrow (d)]

As $E(M) \setminus \{e\}$ is missing only one element of $E(M)$, then $r(E(M) \setminus \{e\}) \geq r(E(M)) - 1$. If $r(E(M) \setminus \{e\}) = r(E(M))$ then $cl(E(M) \setminus \{e\}) = E(M)$ and $e \in E(M)$ but $e \notin E(M) \setminus \{e\}$, which contradicts (c).

[(d) \Rightarrow (e)]

By definition of closure, $E(M) \setminus \{e\} \subseteq cl(E(M) \setminus \{e\})$. And $E(M) \setminus \{e\}$ is contained in only two subsets of $E(M)$, there are $E(M)$ itself and $E(M) \setminus \{e\}$. So $cl(E(M) \setminus \{e\}) = E(M) \setminus \{e\}$ or $cl(E(M) \setminus \{e\}) = E(M)$. If the second occurs, then $r(E(M) \setminus \{e\}) = r(E(M))$, which contradicts (d), and therefore $cl(E(M) \setminus \{e\}) = E(M) \setminus \{e\}$, which means that $E(M) \setminus \{e\}$ is flat.

[(e) \Rightarrow (f)]

As $E(M) \setminus \{e\}$ is flat, then $r(E(M) \setminus \{e\}) < r(M)$, but $E(M) \setminus \{e\}$ is missing only e , so $r(E(M) \setminus \{e\}) = r(M) - 1$ and then $E(M) \setminus \{e\}$ is an hyperplane.

[(f) \Rightarrow (g)]

Let $I \in \mathcal{I}(M)$ such that $e \notin I$ and let $B \in \mathcal{I}$ be a basis for $E(M) \setminus \{e\}$ containing I . As $|B| = r(B) = r(E(M) \setminus \{e\}) = r(M) - 1$, then $r(B \cup \{e\}) = r(M) = |B| + |\{e\}| = |B \cup \{e\}|$ and therefore $B \cup \{e\}$ is a basis, and $I \cup \{e\} \subseteq B \cup \{e\}$ so $I \cup \{e\}$ is independent.

[(g) \Rightarrow (a)]

Let $B \in \mathfrak{B}(M)$, as $B \cup \{e\} \in \mathcal{I}(M)$ and B is maximal by contention over independent sets, so there must be the case that $B \cup \{e\} = B$. \square

2.10 Oxley, Section 1.6, Problem 1

Problem 13. *Show the following:*

- (i) *All uniform matroids are transversal.*
- (ii) *A transversal matroid need not be graphic.*
- (iii) *A paving matroid need not be transversal.*

— . —

Proof. (i) Let $U_{m,n}$ be a uniform matroid of rank m over n elements, define $\mathcal{A} = (A_j | j \in [1, m])$ and $A_j = E(U_{m,n})$ and $S = E(U_{m,n})$. Then, for every subset $J \subset [1, m]$ you can take $T = J$ and the bijection $\psi : J \rightarrow T$ such that $\psi(j) = j$, proving that every J is an independent set in the transversal matroid just defined. Just as every J is independent in $U_{m,n}$. By the definition of \mathcal{A} , there cannot be independent sets of size larger than m . Just as in $U_{m,n}$. These two observations determine $U_{m,n}$ proving that $U_{m,n}$ can be seen as a transversal matroid.

(ii) Let $U_{2,4}$ be a uniform matroid, as we proved above, $U_{m,n}$ is transversal. $\{1, 2, 3\}$ and $\{1, 2, 4\}$ are circuits (see 9), if they were triangles of a graph, then the edges 3 and 4 must be parallel. But $\{3, 4\}$ is independent in $U_{\{2,4\}}$, so $U_{2,4}$ is not graphic.

(iii) Take the uniform matroid $U_{2,3}$ again, and remove the set $\{1, 3\}$ and $\{2, 3\}$ from the collection of independent sets, what you get is a paving matroid that is not uniform. Now, we have only the independent sets $\{1, 2\}$, $\{1\}$, $\{2\}$, $\{3\}$, \emptyset . Let's suppose that this is a transversal matroid, as the largest independent set has two elements, then we can assume that $\{A\}$ has only two elements, let's call them a and b . As $\{1, 2\}$ is independent without loss of generality, there would be a relation $(a, 1), (b, 2)$ and there is someone linked to 3, let's think it is $(a, 3)$, then we can have the matching $(b, 2), (a, 3)$, which lead us to an independent set that was not supposed to be in our matroid.

The same will happen no matter how you try to pair a, b with 1, 2, 3 in a way that let you have the matching given by our collection of independent sets.

□

2.11 Oxley, Section 1.6, Problem 3

Problem 14. Let $S = \{1, 2, \dots, 6\}$ and $\mathcal{A} = \{A_1, A_2, A_3\}$ where $A_1 = \{1, 2, 3\}$, $A_2 = \{2, 3, 4\}$, and $A_3 = \{4, 5, 6\}$.

(i) Find $\Delta[\mathcal{A}]$.

(ii) Give a geometric representation for $M[\mathcal{A}]$.

— . —

Proof.

1. Graphic representation for $\Delta[\mathcal{A}]$



2. Geometric representation for $M[\mathcal{A}]$:



The dependent sets are $\{1, 2, 3\}$, $\{1, 5, 6\}$, $\{2, 5, 6\}$, $\{3, 5, 6\}$, $\{4, 5, 6\}$ and $\{5, 6\}$

□

2.12 Oxley, Section 1.6, Problem 4

Problem 15. Characterize the circuits of $M[\mathcal{A}]$ in terms of the bipartite graph $G = \Delta[\mathcal{A}]$.

— . —

Proof. Lets remember that $G = \Delta[\mathcal{A}]$ is a bipartite graph with partition (S, \mathcal{A}) and that in terms of transversal matroids, a subset $T \subset S$ is independent if there is an injection $\phi : T \rightarrow \mathcal{A}$ such that $(t, \phi(t))$ is an edge for every $t \in T$.

Halls theorem states that a subset $T \subset S$ is independent if and only if $|U| \leq |N_G(U)|$. For every $U \subset T$.

That is, a subset $T \subset S$ is dependent if and only if there is $U \subset T$ such that $|U| > |N_G(U)|$. We are looking for subsets T that satisfies this condition and that are minimal.

If there is $U \subsetneq T$ such that $|U| > |N_G(U)|$, then T can not be minimal.

Then T must have the property $|T| > |N_G(T)|$ (we will refer to this property as the $(*)$ -property) and its minimal.

Then, T is a circuit if and only if it has the $(*)$ -property and its minimal with such condition. (first approach). We can learn a little more about these T 's. Lets check what happens if the graph induced by T has at least 2 connected components. Lets call T_1 and T_2 such that T_1 induces one connected component and $T_2 = T \setminus T_1$. Notice that $N_G(T_1) \cap N_G(T_2) = \emptyset$, and then $|N_G(T)| = |N_G(T_1)| + |N_G(T_2)|$. As $|T| = |T_1| + |T_2| > |N_G(T_1)| + |N_G(T_2)| = |N_G(T)|$ it must be that either $|T_1| > |N_G(T_1)|$ or $|T_2| > |N_G(T_2)|$, without loss of generality, if the first one is true, then at least T_1 is smaller than T and then T is not minimal. Then the graph induced by T must be connected.

T is circuit if and only if the graph induced by T (lets call it $G[T]$) is isomorphic to $K_{|T|, |T|-1}$.

Necessity

Suppose the $G[T]$ is isomorphic to $K_{|T|, |T|-1}$, T is minimal with $(*)$ -property, if you remove any vertex from T , its induced graph will be $K_{|T|-1, |T|-1}$ which has perfect matches, if you remove more you will have a $K_{m, n}$ with $m < n$ which has $K_{m, m}$ as subgraph.

Sufficiency

Suppose you have a circuit T , as T is bipartite and connected, then it must contain a $K_{m, n}$ as subgraph induced by vertices in T . If $n < m$, for one of them the necessity proves that T must be as desired, or then it has a subset that is dependent. If it is not the case and $m \leq n$ for every such graph, then T doesn't have the $(*)$ -property. \square

2.13 Oxley, Section 1.7, Problem 5

MISSING

Problem 16. *Prove that a finite lattice \mathcal{L} is semimodular if, for all x, y in \mathcal{L} , the following condition holds:*

IF BOTH x AND y COVER $x \wedge y$, THEN $x \vee y$ COVERS BOTH x AND y .

— . —

Proof.

□

2.14 Oxley, Section 1.8, Problem 1


Problem 17. (i) Find a maximum-weight spanning tree of the graph in the next figure. Is this the unique such tree?



(ii) Find all maximum-weight spanning trees and all minimum-weight spanning trees of the graph in the next figure, where the edge labels are interpreted as weights.



Proof. 1. There are only 2 maximum-weight spanning trees of the graph. I have illustrated what will greedy do step by step, and with all possible decisions being made.






2. As the weight function is injective, the maximum-weight and minimum-weight spanning trees are unique.

Here is again an illustration of what greedy would do in both scenarios.

Maximum-weight:



Minimum-weight:



□

2.15 Oxley, Section 1.8, Problem 4

Problem 18. Let M be a matroid and $\omega : E(M) \rightarrow \mathbb{R}$ be a one-to-one function. Prove that M has a unique basis of maximum weight.

— . —

Proof. Let B be a maximum-weight basis found using greedy. Suppose that there is another basis B' with maximum-weight also and suppose that they are different.

Let $e \in B' \setminus B$ of minimum-weight (it exists because ω is one-to-one), and then, there exists $f \in B \setminus B'$ such that $(B' \setminus \{e\}) \cup \{f\}$ is basis. We can choose such f of maximum weight. If $\omega(e) < \omega(f)$ then we have increased the weight of B' , which is a contradiction. If $\omega(f) < \omega(e)$ we can change the roles of B and B' and get the same contradiction.

So, there is no other basis B' with maximum-weight and different from B . □

3

Second Test

3.1 Oxley, Section 2.1, Problem 1

Problem 19. Find each of the following:

- (i) all self-dual uniform matroids;
- (ii) all identically self-dual uniform matroids;
- (iii) all self-dual graphic matroids on six or fewer elements;
- (iv) all identically self-dual graphic matroids on six or fewer elements;
- (v) an infinite family of simple graphic self-dual matroids.

— . —

Proof.

- (i) Let $U_{n,m}$ be an uniform matroid.

Lets asume that it is self-dual. As any basis B for $U_{n,m}$ has size n , if it is going to be self-dual, then its complement must have size n as well. This means that $m = 2n$.

Now lets suppose that $m = 2n$. So, any subset of size n is a basis for $U_{n,m}$, but it is also true that any subset of size n is the complement of another subset of size n and then, any basis is the complement of another basis. Then $U_{n,m}$ is not only self-dual, but also identically self-dual.

- (ii) Just as we saw above, if an uniform matroid is self-dual, then it is identically self-dual, and an uniform matroid $U_{n,m}$ is self-dual if and only if $m = 2n$.
- (iii) It is clear that if a matroid is to be self-dual, then its size must be two times the size of any of its basis (any forest that is not contained in any other forest). This means that there are only self-dual graphic matroids with an even number of edges.

Lemma 2.3.7 from [Oxl92] stays that if G^* is a geometric dual of the planar graph G , then $M(G^*) \cong M^*(G)$. In particular for self-dual matroids we will have that $M(G^*) \cong M^*(G) \cong M(G)$.

From now on we will use this fact to look for the remaining self-dual matroids. As the maximum size of these matroids is 6 and K_4 is a complete planar graph with 6 edges, then every one of the matroids will have a planar representation. And the previously mentioned lemma will let us work with graph representations instead of a complicated structure of subsets and its intersections.

Also, notice that if G is a planar graph. And $H = G \cup G^*$, then $M(H)$ is self-dual. The only trick is to send $M(G)$ to $M(G)$ in the dual and $M(G^*)$ to $M(G^*)$ in the dual.

Considering this. We will try to find self-duality through blocks (2-connected maximal subgraphs) when it is possible. Also, the drawings will be done thinking of these, we are going to separate in as many blocks as possible, given that it will not affect the resultant matroid.

We will try to order our analysis by the size of the maximum cycle contained in a graph or its dual matroid. And for this remember that the dual of a n -cycle is a set of n edges all of them parallel to each other.

2 edges:

- **No cycles.** There is no such graph. If a graph has no cycles, then its dual contains only loops (which are cycles). This will happen in any of the other graphs so we are not going to list this case anymore.
- **Cycles of size at most 1.** If there has to be a loop, then it has to be an edge that is not a loop. The loop will be mapped to the dual of the edge that is not a loop and viceversa. This one is not identically self-dual.



Figure 3.1: self-dual graphic matroid representation

- **Cycles of size at most 2.** There is only one way to get a cycle of size two with two edges, and it is easy to see that it has not only a self-dual matroid but an identically auto-dual matroid.




Figure 3.2: identically self-dual graphic matroid representation

There cannot be larger cycles with only 3 edges.

4 edges:

- **Cycles of size at most 1.** Again. For every loop there must be an edge that is not a loop. And for every edge that is not a loop there must be a loop. So the only possibility left is to have two loops and two edges that are not loops.



- **Cycles of size at most 2.** If we have a block as in 3.2, then, under the matroid isomorphism there are two possibilities, that such block is mapped into itself, or that it is mapped into another block. If it is mapped into another block, then the other block must be essentially the same graph and then we have something like



Figure 3.3: identically self-dual graphic matroid representation

(Notice that this will result in an identically self-dual matroid also. Every forest is a pair of non-parallel edges, and its complement is also a pair of non-parallel edges.)

If the first block is mapped into itself, then the other block must be mapped into itself also. Then the other block must be a representation of a self-dual matroid on 2 edges. And then, there are two possibilities. If the other block is like in 3.2, then we will obtain again 3.3. If the other block is like 3.1 then we will obtain something like




Figure 3.4: self-dual graphic matroid representation

(Notice that this could not be identically self-dual. As it has loops, any loop will be in a cobasis, but no basis contains loops.)

- **Cycles of size at most 3.** A graph that will represent a self-dual matroid and that has a 3-cycle and at most 4 edges, must have at most one block. Otherwise, one block must be the triangle, but a triangle is not self-dual and could not be mapped in the other block that only could have 1 edge.

This gives us essentially only one opportunity. A triangle adding a parallel edge to one of its edges. An isomorphism would be sending the simple edges to the parallel edges and the parallel edges to the simple edges. The two simple edges form a set that intersects any of the basis of the matroid, so it is a cocircuit. The proposed isomorphism send that cocircuit to a circuit in the original matroid. The two parallel edges along with any of the simple edges form another set that intersects any of the basis of the original matroid, so it is another cocircuit in the dual matroid. Those edges are sent to a triangle in the original matroid. There are two of these cocircuits and each one is sent to one of the two possible triangles in the original matroid. Any basis formed for a simple edge and one of the two parallel edges has as complement the other simple edge and the other parallel edge, and under the proposed isomorphism any of such basis are sent to other of the same type. The only other basis that is left is the one that consists of the two simple edges, and they are sent to the pair of parallel edges, which is independent in the dual matroid. We have proved so far that this isomorphism sends any of the basis in the original matroid to a basis in the dual matroid and any cocircuit to a circuit in the original matroid. Then such isomorphism is a matroid isomorphism between the graphic matroid and its dual.



Figure 3.5: self-dual graphic matroid representation

This could not be identically self-dual as it has a cobasis that contains parallel edges.

- **Cycles of size at most 4.** There is none, as the only cycle of size 4 has 4 edges, and as the 4-cycle is not self-dual, there cannot be self-dual matroids with 4-elements and 4-cycles.

There cannot be larger cycles with only 4 edges.

6 edges:

MISSING!

- (iv) Any graph on six or fewer elements obtained from a forest adding a parallel edge to each edge in the forest. We have proved this in the last part, indicating which of the self-dual matroids are identically self-dual or non-identically self-dual.
- (v) Any wheel. Proof in [29]

□

3.2 Oxley, Section 2.1, Problem 2

Problem 20. Let M be a matroid. Show that M^* has two disjoint circuits if and only if M has two hyperplanes whose union is $E(M)$.

— . —

Proof.

Sufficiency

Let C^* and C'^* be two disjoint circuits in M^* . From Proposition 2.1.6 in [Oxl92] we know that $C = E(M) \setminus C^*$ and $C' = E(M) \setminus C'^*$ are hyperplanes in M . Moreover, as C^* and C'^* are disjoint, then $C^* \subset C'$ and $C'^* \subset C$. Then

$$\begin{aligned}
 C \cup C' &= (C \cup C'^*) \cup (C' \cup C^*) \\
 &\quad \text{(we have added nothing new to each subset)} \\
 &= (E(M) \setminus C^*) \cup C'^* \cup (E(M) \setminus C'^*) \cup C^* \\
 &\quad \text{(reordering terms)} \\
 &= (E(M) \setminus C^*) \cup C^* \cup (E(M) \setminus C'^*) \cup C'^* \\
 &= E(M) \cup E(M) \\
 &= E(M).
 \end{aligned}$$

Necessity

Let H and H' be two hyperplanes such that $H \cup H' = E(M)$. Again, from Proposition 2.1.6 in [Oxl92] we know that $H^* = E(M) \setminus H$ and $H'^* = E(M) \setminus H'$ are cocircuits. Suppose there exists $x \in H^* \cap H'^*$, that is $x \in H^*$ and $x \in H'^*$. But $x \in H^*$ means $x \notin H$ and $x \in H'^*$ means $x \notin H'$, then $x \notin H \cup H' = E(M)$ which is a contradiction. Then there is not such x and $H^* \cap H'^*$ are cocircuits such that $H^* \cap H'^* = \emptyset$. \square

3.3 Oxley, Section 2.1, Problem 6

Problem 21. *Let e and f be distinct elements of a matroid. Prove that every circuit containing e also contains f if and only if $\{e\}$ or $\{e, f\}$ is a cocircuit.*

— . —

Proof.

Sufficiency

Suppose every circuit containing e also contains f .

If every basis contains e , then, no cobasis contains e , which means $\{e\}$ is a cocircuit.

Suppose then that there is a basis B such that $e \notin B$. As $C(e, B)$ is a circuit that contains e , then $f \in B$. That is, any basis that doesn't contain e , contains f . That is, any cobasis that contains e , doesn't contain f .

Note that $B \setminus \{f\} \cup \{e\}$ is a basis given that it has the right size to be a basis and it is obtained from $B \cup \{e\}$ getting rid of the only circuit that it contains (that is, it is independent). Then, there is a basis that doesn't contain e and then $\{e\}$ is coindependent, and there is also a basis that doesn't contain f and then $\{f\}$ is also coindependent.

Now suppose there is a basis B' such that $f \notin B'$. If $e \notin B'$, then $C(e, B')$ is a circuit that contains e but not f which contradicts our hypothesis. Then any basis that doesn't contain f , contains e . That is, any cobasis containing f , doesn't contain e .

Then $\{e, f\}$ is a subset that is not contained in any cobasis, then it is codependent but any of its subsets is coindependent. Then it is a cocircuit as we wished to show.

Necessity

Suppose $\{e\}$ is a cocircuit. Then, again by Proposition 2.1.6 in [Ox192], $E(M) \setminus \{e\}$ is an hyperplane, and then e is contained in every basis, then e is in no circuits (by [12]). Then it is true that every circuit containing e also contains f , since there are not circuits containing e .

Now suppose that $\{e, f\}$ is a cocircuit. Then, again by Proposition 2.1.6 in [Ox192], $H = E(M) \setminus \{e, f\}$ is an hyperplane, so any basis must contain e or f .

Let C be a circuit such that $e \in C$. Then $C \setminus \{e\}$ is independent. Let B be a basis such that $C \setminus \{e\} \subset B$. As B is basis and doesn't contain e , then $f \in B$. Notice that $C(e, B) = C$

Now, $B^* = E(M) \setminus B$ is a cobasis that contains e but doesn't contain f , then $C * (f, B^*) = \{e, f\}$ is the only cocircuit in $B^* \cup \{f\}$, that means that $B^* \cup \{f\} \setminus \{e\}$ is a coindependent of maximum size, that is a cobasis. Taking its complement we will get a basis, and it is

$$E(M) \setminus (B^* \cup \{f\} \setminus \{e\}) = B \cup \{e\} \setminus \{f\}.$$

Which means that $f \in C$. □

3.4 Oxley, Section 2.1, Problem 10

Problem 22. Let B be a basis of a matroid M and B^* be $E(M) - B$. If $e \in B$, let $C^*(e, B^*)$ denote the fundamental cocircuit of e with respect to the cobasis B^* of M , that is, $C^*(e, B^*) = C_{M^*}(e, B^*)$.

- (i) Show that $C^*(e, B^*)$ is the unique cocircuit that is disjoint from $B - e$.
- (ii) If $f \in B^*$, prove that $f \in C^*(e, B^*)$ if and only if $e \in C(f, B)$.

— . —

Proof.

1. $C^*(e, B^*)$ is the only cocircuit contained in $B^* \cup e$, then any other circuit must intersect the complement of $B^* \cup e$, but the complement of $B^* \cup e$ is $B - e$. Then any other cocircuit must intersect $B - e$.
2. Suppose $e \in C(f, B)$. Now, $B \setminus e \cup f$ must be a basis, it has the same size as B and $B \cup f$ has an unique circuit which is $C(f, B)$ and which contains f , so $B \setminus e \cup f$ cannot contain a circuit and then it is independent with maximum size, that is, $B \setminus e \cup f$ is a basis.

Then, its complement must be a cobasis, but its complement is $B^* \setminus f \cup e$, we know that $B^* \setminus f \cup e \cup f = B^* \cup e$ has an unique cocircuit and then it must contain f .

We have already proved that $e \in C(f, B)$ implies $f \in C^*(e, B)$, the proof for the sufficiency is completely analog.

□

3.5 Oxley, Section 2.2, Problem 2

Problem 23. Let A be an $m \times n$ matrix over a field F .

- (i) For each of the row operations, specify an $m \times m$ matrix L such that multiplying A on the left by L has the same effect as the row operation.
- (ii) For each of the column operations, specify an $n \times n$ matrix R such that multiplying A on the right by R has the same effect as the column operation.

— . —

Proof. ,

1. Take the $m \times m$ identity matrix. Perform any of the row operations to it and let L be the result.

Such matrix is called elemental matrix, and if you multiply A on the left by L it will have the same effect as the row operation.

2. The proof is exactly the same, but working with columns. Perform any of the column operations to the $n \times n$ identity matrix, and let R be the result.

□

3.6 Oxley, Section 2.2, Problem 4

Problem 24. *Show that, in a binary matroid, a circuit and cocircuit cannot have an odd number of common elements.*

— . —

Proof. Let M be a binary matroid with rank n and such that $|E(M)| = n + r$.

Theorem 2.2.8 from [Oxl92] states that if a linear matroid is expressed as $[I_n|D]$, then its dual matroid can be represented with the matrix $[-D^T|I_r]$.

In the case of binary matroids is the same $[-D^T|I_r]$ than $[D^T|I_r]$ (In general it is always the same, multiplying columns by -1 doesn't change dependency/independency conditions).

Lets say that C is a circuit of M and C^* is a cocircuit of M . As C is a circuit in a binary matroid, the sum of their elements must be the zero vector.

The representation of M can always be arranged in such way that C consists of $|C| - 1$ columns from I_n (lets call them I) and one column of D (lets call it d_j assuming that d_j is the j -th column of D).

With such representation is easy to see that if $e_i \in I$ is the i -th colmun of I_n , then $(d_j[i] = 1$ (that is, the i -th entry of d_j). Conversely, if $d_j[i] = 1$ then $e_i \in I$.

In the j -th row of $[D^T|I_r]$ the first n entries are exactly the entries of d_j , in the next r entries are exactly one and it is in the j -th entry of I_r . This is, the j -th row of $[D^T|I_r]$ has a one in its k -entry if and only if the k column of $[I_n|D]$ was in C .

Given that C^* is a circuit in M^* , the sum of its columns must be the zero vector. In particular, in $[D^T|I_r]$ the sum of their j -th entry, must be 0. That is, the number of ones in their j -entry must be even. As we saw that the j -th row of $[D^T|I_r]$ have ones exactly in the entries corresponding of columns of C , then the intersection of C and C^* must be even. \square

3.7 Oxley, Section 2.2, Problem 8

Problem 25. Let T_8 and R_8 be the vector matroids of the following matrices over $GF(3)$:

$$\left[\begin{array}{cccc|cccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ & & & & 0 & 1 & 1 & 1 \\ & & & & 1 & 0 & 1 & 1 \\ & & & & 1 & 1 & 0 & 1 \\ & & & & 1 & 1 & 1 & 0 \end{array} \right], \quad \left[\begin{array}{cccc|cccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ & & & & -1 & 1 & 1 & 1 \\ & & & & 1 & -1 & 1 & 1 \\ & & & & 1 & 1 & -1 & 1 \\ & & & & 1 & 1 & 1 & -1 \end{array} \right]$$

- (i) Show that T_8 and R_8 are both self-dual.
- (ii) Show that R_8 is identically self-dual but T_8 is not.
- (iii) Give geometric representation for T_8 and R_8 .
- (iv) Show that if $M \in \{T_8, R_8\}$ and $X = E(M) \setminus \{8\}$, then $(M|X)^* \cong F_7^-$.
- (v) Consider the following matrices over $GF(3)$:

$$A_1 = \left[\begin{array}{cccc|cccc} & & & & 0 & 1 & 1 & 1 \\ & & & & 1 & 0 & 1 & 1 \\ & & & & 1 & 1 & 0 & -1 \\ & & & & 1 & 1 & 1 & 0 \end{array} \right], \quad A_2 = \left[\begin{array}{cccc|cccc} & & & & 0 & 1 & 1 & -1 \\ & & & & 1 & 0 & 1 & -1 \\ & & & & 1 & 1 & 0 & -1 \\ & & & & 1 & 1 & 1 & 1 \end{array} \right]$$

Show, by applying a sequence of the row and column operations to A_1 and A_2 that $M[A_1] \cong T_8$ and $M[A_2] \cong R_8$.

- (vi) Show that R_8 can be obtained from $AG(3, 2)$ by relaxing two disjoint circuit-hyperplanes.

— . —

Proof.

- (i) Theorem 2.2.8 from [Oxl92] states that if a linear matroid is expressed as $[I_n|D]$, then its dual matroid can be represented with the matrix $[-D^T|I_r]$.

As multiplying columns by -1 doesn't change dependency/independency conditions. The dual matroid can be represented as $[D^T|I_r]$.

For both, T_8 and R_8 , we have that $n = 4$, $r = 4$, and $D = D^T$. Then, in both cases, $[D^T|I_4] = [D^T|I_4] = [D|I_4]$. And with this representation is clear how we would build an isomorphism from T_8 to T_8^* and from R_8 to R_8^* .

- (ii) Let's write $T_8 = [I_4|D_1]$, it is easy to prove that T_8 is not identically self-dual by simply pointing out a cobasis that is not a basis.

All the columns of I_4 form a basis. Then all the columns of D form a cobasis, but such cobasis is dependent, if you sum up all of their columns you will get the zero vector.




Figure 3.6: Geometric representation of T8

In this representation, there are points in the center of each face of the tetrahedron and the sphere that is inside the tetrahedron tangent to all the faces, is a plane that contains all the centers of the faces (just as in Fano's plane the circle inside the triangle is a line).

Assuming that the tetrahedron is regular, all the other planes that have 4 vertices and that are not indicated are those that can be found in the euclidean space. For example $\{1, 2, 5, 6\}$.




Figure 3.7: Geometric representation of R8

This representation is simply a cube. All the other planes that contain 4 vertices and that are not indicated are those that can be found in the euclidean space. For Example $\{7, 8, 3, 4\}$.

(iv)

(v)

(vi)

□

3.8 Oxley, Section 2.3, Problem 1

Problem 26.

- (i) Show that the geometric dual of a plane graph is connected.
- (ii) Give an example of a plane graph G for which $(G^*)^* \neq G$.

— . —

Proof.

- (i) Take any embedding of a plane graph, take its dual, and take any pair of vertices of its dual. such vertices are faces in the original embedding. It is always possible to find a curve that joins that pair of vertices and that avoids any other vertex in the original embedding. Such curve defines a path bewteen the pair of vertices.
- (ii) Any disconnected graph will work, $(G^*)^*$ will be connected but G not. Look at the following example.




Figure 3.8: disconnected graph




Figure 3.9: disconnected graph and its dual




Figure 3.10: disconnected graph dual and its dual




Figure 3.11: disconnected graph double dual

□

3.9 Oxley, Section 2.3, Problem 2

Problem 27. Find geometric duals of the graphs obtained from K_5 and $K_{3,3}$ by deleting a single edge of each.

— . —

Proof.




Figure 3.12: K_5 without one edge




Figure 3.13: K_5 without one edge and its dual




Figure 3.14: Dual of K_5 without one edge




Figure 3.15: $K_{3,3}$ without one edge




Figure 3.16: $K_{3,3}$ without one edge and its dual






Figure 3.17: Dual of $K_{3,3}$ without one edge (K_4)



3.10 Oxley, Section 2.3, Problem 6

Problem 28. *Construct the geometric duals of the plane graphs in the next figures.*



— . —

Proof.




Figure 3.18: First graph and its dual




Figure 3.19: First graph's dual




Figure 3.20: Second graph with its dual




Figure 3.21: Second graph's dual

□

3.11 Oxley, Section 2.3, Problem 10

Problem 29. Let r be an integer exceeding one and W_r be the r -wheel. Show that:

- (i) $W_3 \cong K_4$.
- (ii) $W_r^* \cong W_r$.
- (iii) $M(W_4)$ is isomorphic to a restriction of $M^*(K_{3,3})$.

— . —

Proof.

1. It is sufficient to prove that W_3 is complete.

The central vertex is always adjacent to the rest of the vertices.

Any vertex in the exterior cycle is adjacent to exactly other two of its type, but as W_3 has only 3 vertices in its exterior cycle, then any vertex in the exterior cycle is adjacent to the other two, plus, it is adjacent to the center.

We have proved that any vertex has degree 3 and there are 4 vertices. Then the graph is complete and we are done.

2. Any triangle that is induced by the central vertex and two of the external vertices under the dual operator will end up as a vertex of degree tree, adjacent to two triangles of its type and to the exterior face.

There are exactly r triangles of this type in W_r , then the external face will be adjacent to every one of these triangles.

Then the dual operator will make a graph such that it has a vertex adjacent to the rest of the vertices and the rest of the vertices are adjacent in addition to exactly two of its type. This is, the dual of W_r is W_r itself.


.

- 3.

□

3.12 Oxley, Section 2.4, Problem 3

Problem 30. Let G be the directed graph shown in next figure.



- (i) Construct the corresponding bipartite graph \hat{G} .
- (ii) Let $V = \{1, 2, 3, 4, 5, 6, 7, 8\}$, $X = \{5, 7\}$, $Y = \{4, 2\}$, and the paths linking X to Y in G be 5134 and 7862 . Construct the corresponding matching of $V \setminus X$ to \hat{Y} in \hat{G} .
- (iii) Now take the matching $\{2\hat{1}, 3\hat{5}, 4\hat{3}, 5\hat{4}, 6\hat{8}, 8\hat{7}\}$ in \hat{G} and construct the corresponding disjoint paths in G as in the proof of Lemma 2.4.3.

— . —

Proof.

(i)




Figure 3.22: Bipartite graph

(ii)





Figure 3.23: Bipartite graph and matching

(iii) The corresponding paths are $(1, 2)$ and $(7, 8, 6)$. It is generated also a cycle

□

3.13 Oxley, Section 2.4, Problem 4

Problem 31. Let G be the directed graph shown in the next figure



and let $L(G, B_0)$ where $B_0 = \{1, 2, 3\}$.

- (i) Find geometric representations for M and M^* .
- (ii) Give a presentation for the transversal matroid M^* .
- (iii) Reverse the directions on the arcs $(5, 4)$ and $(7, 5)$ of G and repeat (i) and (ii).

— . —

Proof. 1.




Figure 3.24: Geometric representation for M

2.

3.

□

3.14 Oxley, Section 3.1, Problem 1

Problem 32. Show that if $T \subset E(M)$, then

- (i) $M \setminus T = (M^*/T)^*$;
- (ii) $(M/T)^* = M^* \setminus T$; and
- (iii) $M^*/T = (M \setminus T)^*$.

— . —

Proof. (i) Let B be a basis for $M \setminus T$. We would like to prove that B is also a basis for $(M^*/T)^*$, or, what is the same, that $E \setminus (B \cup T)$ is a base for M^*/T .

As B is a basis for $M \setminus T$, B is independent in M . And then there is a basis B' of M such that $B \subset B'$. Now $E \setminus B'$ is basis for M^* , and then $B_T^* = E \setminus B' \cap T$ is basis for T in M^* .

Now

$$\begin{aligned}
 E \setminus ((E \setminus (T \cup B)) \cup B_T^*) &= (T \cup U) \cap (E \setminus B_T^*) \\
 &= (T \cup U) \cap (E \setminus ((E \setminus B') \cap T)) \\
 &= (T \cup U) \cap (B' \cup (E \setminus T)) \\
 &= ((T \cup B) \cap B) \cup ((T \cup B) \cap (E \setminus T)) \\
 &= (T \cap B') \cup B \\
 &= (T \cap B') \cup ((E \setminus T) \cap B') \\
 &= B'
 \end{aligned}$$

Then, every basis for $M \setminus T$ is a basis for $(M^*/T)^*$.

Now, let B be a basis for $(M^*/T)^*$, then $(E \setminus T) \setminus B = E \setminus (B \cup T)$ is a basis for M^*/T , then, there is a basis B_T^* for T in M^* such that $(E \setminus (B \cup T)) \cup B_T^*$ is a basis for M^* .

Let $B' = (B \cup T) \cap (E \setminus B_T^*)$, B' is a basis for M . Now

$$\begin{aligned}
 (E \setminus T) \cap B' &= (E \setminus T) \cap (B \cup T) \cap (E \setminus B_T^*) \\
 &\quad \text{(remember that } B \cap T = \emptyset \text{)} \\
 &= B \cap (E \setminus B_T^*) \\
 &= B.
 \end{aligned}$$

This means that B is a basis for $M \setminus T$ also. And we have finished.

(ii) Applying the previous part,

$$\begin{aligned}
 (M/T)^* &= ((M^*)^*/T)^* \\
 &= M^* \setminus T.
 \end{aligned}$$

(iii) Applying the previous part (the first one),

$$\begin{aligned}
 M^*/T &= ((M^*/T)^*)^* \\
 &= (M \setminus T)^*.
 \end{aligned}$$

□

3.15 Oxley, Section 3.1, Problem 4

Problem 33. Suppose that, for all elements f of a loopless matroid M , $r(M \setminus \{f\}) = r(M)$, but that, for some elements e and g , $r(M \setminus e, g) = r(M \setminus e/g)$. Show that $\{e, g\}$ is a cocircuit of M .

— . —

Proof.

□

3.16 Oxley, Section 3.2, Problem 3

Problem 34. Consider the wheel graph W_r . Show that if $r \geq 3$, then $M(W_r)$ has $M(K_4)$ as a minor.

— . —

Proof. Thinking of graphs, we can obtain K_4 which is the same as W_3 from W_r by simply erasing $r - 3$ contiguous edges of the central vertex and contracting all those edges that contain vertices that were disconnected from the central vertex.

As in graph matroids it is the same to take the minor matroid or taking the matroid of the minor graph, what we have said is already sufficient. \square

3.17 Oxley, Section 3.2, Problem 5

Problem 35. Let T_8 and R_8 be matroids in [25]. Give geometric representations for each of $T_8/8$, $T_8/1$, $R_8/8$ and $R_8/1$.

— . —

Proof.

□

Bibliography

[Oxl92] *James G. Oxley*. Matroid Theory. *Oxford University Press*, 1992.