

This project is a small Python library, `geometry.py`, to use in drawing figures for my geometry textbook. The library is contained in a single file at top level: `geometry.py`.

I am running Python 3.13 obtained via Homebrew, so I put a symbolic link to `geometry.py` in:

```
/usr/local/lib/python3.13/site-packages/geometry.py
```


The `projects` directory contains sub-directories with various scripts using it. Other scripts written during development to test particular issues are in `tests`.

Here is a [list](#) of all the functions defined there.

Some examples of figures made using the library:

### midline reverse


Starting with triangle  $ABC$ , construct the parallelograms on each side, by bisecting the side and then doubling the length (here  $AD = \text{twice } AK$ ).



$ABK$  and  $DCK$  are congruent by SAS. It follows that  $CD$  is parallel to  $AB$ . The same argument applies to  $BD$  and  $AC$ . With both pairs of opposing sides parallel,  $ABDC$  is a parallelogram.


All four of the smaller triangles are congruent and similar to the large triangle DEF. This is like the midline theorem in reverse.

### Ptolemy by similar triangles



Divide one vertex angle of a cyclic quadrilateral to form a triangle with sides  $a$  and  $x_1$  as shown.

$$x_1/a = c/y$$




$$x_2/d = b/y$$

$$x_1 + x_2 = x = ac/y + bd/y$$


$$xy = ac + bd$$

### Euclid I.2




$$AF = BC.$$

### Euclid II.5



Let  $C$  bisect  $AB$  and  $D$  be an arbitrary point on  $CB$ . Draw the squares on  $DB$  and  $CB$ . It follows that  $CDHL$  equals  $MFGH$ , so  $AD \cdot BD$  is equal to the gnomon  $CBFGHL$ . The latter is equal to the difference of squares  $BC^2 - CD^2$ .

### Euclid II.11



Division of a line segment into the golden mean.

Draw the square on AB and then bisect AC at E. Draw the circle on center E with radius BE and find where it cuts the extension of CA at F. Draw the square on AF.

The idea of the proof is then to show that  $AB/AH = AH/HB$  and

$$AB \cdot HB = AH^2$$

The LHS is the area of the red rectangle and the RHS is the area of the small blue square. We show they are equal by subtracting the rectangle AHKC from two other equal areas: the square on AB, and the rectangle CFGK.

To obtain the last result, start from Euclid II.6


$$CF \cdot AF = EF^2 - AE^2$$

But  $EF = EB$  and  $EB^2 - AE^2 = AB^2$  by Euclid I.47 so

$$CF \cdot AF = AB^2$$

We have shown that the rectangle CFGK is equal in area to the square on AB. Subtract AHKC from both and we obtain the needed equality.

**Euclid IV.10**



By construction, we have that  $AC = BD$  and

$$AB/AC = AC/BC$$


$$AC^2 = AB \cdot BC = BD^2$$

By the converse of the tangent-secant theorem, it follows that  $BD$  is tangent to the dotted circle.  $\angle BDC$  cuts the same arc as  $\angle CAD$ , so they are equal. Since the triangles  $BAD$  and  $BDC$  have the same vertex angle and share the base angle at  $B$ , they are similar. They are also isosceles, since  $AB$  and  $AD$  are radii of the first circle.

Thus  $CD = BD = AC$ . So triangle  $ACD$  is also isosceles. From this and the external angle theorem, we can deduce that  $\angle B = \angle BCD$  is twice  $\angle A$ . This is the basic construct for the pentagon.

The vertex of the isosceles triangles measures  $1/5$  of a right angle.

### **Pentagon construction**




Scale the circle so the radius is 2.

The key step is to bisect  $\angle OPA$  with  $OQ$ . Then  $AQ/OQ = \sqrt{5}$ . But  $AQ = 2 - OQ$ . So  $OQ = 1/\phi$ .

Make  $QPR$  a right angle, then  $OR = \phi$ . [Bogolmony]

$\phi$



Let the diameter of the circle be 1. Then the length of the diagonal of the rectangle is  $\sqrt{5}$ . Let  $x$  be the distance from one corner to the circle.

$$2x + 1 = \sqrt{5}$$

$$x = (\sqrt{5} - 1)/2 = \phi - 1$$

But

$$\phi^2 = 1 + \phi$$


$$1/\phi = \phi - 1$$

The rest of the diagonal is  $1 + x = \phi$ .


[Bogolmony]

### Pizza theorem


[wikipedia](https://en.wikipedia.org/wiki/Pizza_theorem)



**Euclid I.47**



### Pappus



On two sides of triangle  $ABC$  draw arbitrary parallelograms. Extend the two new sides to meet at  $H$ .

Draw  $RAJ$  parallel to  $HCTU$  with  $RA = AJ$ , and do likewise for  $SBI$ .



From the properties of parallelograms, is easy to show that


$$(ACDE) = (ACHR) = (AJUT).$$

Thus the area of the parallelogram on side AB is the sum of the areas of the other two.


The Pythagorean theorem is a special case for a right triangle and squares on the sides.

[wikipedia](https://en.wikipedia.org/wiki/Pythagorean_theorem)

### triangle rotation




### nine point circle



Show that  $AB = CD$ . [Acheson]

## Heron's theorem



Draw the incircle for triangle ABC on center I. Draw IK perpendicular to CI and BL perpendicular to CB.

We have two pairs of similar triangles. (Show that  $\angle L$  is complementary to the half angle at A).  
Then:

$$r/x = BL/BC = BL/(y+z)$$

From the other pair:

$$BL/ID = BL/r = BK/DK$$

Combining the two:

$$(y+z)/x = BK/DK$$

Add 1 to both sides:

$$(x+y+z)/x = (BK+DK)/DK = y/DK$$

Let  $s = x + y + z$ :

$$s \cdot DK = xy$$

Since triangle CIK is right:


$$DK \cdot z = r^2$$

$$xyz = sr^2$$


$$r^s = x y z s$$

But  $r_s$  is the area of triangle ABC.

**excircle**




## similar triangles



A proof that extends similarity of right triangles to similarity for all triangles.

### $\phi$ in the hexagon



Show that  $AP/AB = \phi$ .

$MF$  is half the radius of the circle on  $Q$ .

The rest is algebra and an application of the Pythagorean theorem. [Bogolmony]

### Notes

There are also a couple of write-ups, including one about Archimedes' broken chord theorem and another about excircles.

This is functional programming. The only objects we define are members of the class **Point**, to allow access via  $P.x$  and  $P.y$ .

The variable name **pL** found in most function definitions stands for *point list*, i.e. a list of **Point** objects. This may be a line segment, a triangle or another polygon.

We pretend to implement some of Euclid's constructions, but intersections between lines and circles are computed by analytic geometry. Under the hood, it is algebra.

When there are two points in a result to be returned, the order in which they are returned is sometimes challenging to determine. In the latest version, for two points, say, perpendicular to a line segment, we return the point "above" the line segment first, if you visualize the line segment

as oriented left-to-right.

Some examples may differ. For circle-circle intersection, we return the point closer to the origin first.

For a perpendicular or angle bisector, the *length* of the perpendicular or bisector is arbitrary. It should be adjusted by the caller, using the following trick:

```
S,T = geo.get_perp_at_point_by_fractional_length([A,B],f=0.5)
X = geo.get_intersection_for_two_lines([A,B],[S,T])

d = 10    # or whatever the desired length is
f = d/geo.get_length([X,S])
geo.get_point_by_fractional_length([X,S],f)
```

As I fiddled with the code, inconsistency in the order of return of two points has messed up many a diagram. I believe that's all fixed now.

For most examples, output paths for figures are hard-coded so it will require a bit of configuration to get it to work on another machine. I have tried to automate that for newer code, but haven't fixed the old examples.

Here are a few of the functions we can call:

```
geo.get_intersection_for_two_lines([A,B],[C,D])
geo.get_point_perp_on_line_for_point(P,[A,B])
geo.get_perp_at_point_by_fractional_length([A,B],f=0.5)

get_intersection_line_segment_circle([A,B],[Q,r])
get_intersection_circle_circle([Q1,r1],[Q2,r2])
get_tangent_points_on_circle_for_point([Q,r],P)
```

These are from the caller's POV. In the library's function definition, you cannot have

`([A,B],[C,D])`, it is

```
geo.get_intersection_for_two_lines(pL1,pL2)
```

## Opaque error messages

`matplotlib` gives error messages that can be challenging to interpret. If you forget an argument in a function call, it simply reports the *last* one missing. So, in drawing functions like

```
geo.outline_polygon(ax,[A,B,E,D],ec='k')
geo.draw_line_segments(ax,[[D,F],[C,F]])
```

If you forget `ax` in the first one, the error is:

```
TypeError: outline_polygon() missing 1 required
positional argument: 'pL'
```

If you forget to make a list of line segments by adding a second pair of brackets in the second one:

```
TypeError: draw_line_segments() got multiple values
for argument 'ec'
```

Other mistakes with brackets may result in Python trying to access a coordinate like `P.x` and complaining that a list doesn't have one.

```
AttributeError: 'tuple' object has no attribute 'x'
```

Finally, the functions in the library have long, but I hope explicit, names. I'm experimenting with shortcut definitions ([here](#)).

The shortcuts live in the library (at the end).

For this we might do `from geometry import *` in spite of the fact that it's generally not good practice. Or, one might call them like

```
geo.gtr()
```