

VIENNA UNIVERSITY OF TECHNOLOGY

184.725 HIGH PERFORMANCE COMPUTING

TU WIEN INFORMATICS

---

## Exercise 1

---

*Authors:*

Camilo TELLO FACHIN  
12127084

*Supervisor:*

Prof. Dr. Jesper Larsson TRÄFF

December 4, 2022



TECHNISCHE  
UNIVERSITÄT  
WIEN

Vienna University of Technology

## Abstract

Here documented the results of exercise 1.

# Contents

<b>1</b>	<b>Exercise 1 - Closed Form Expressions</b>	<b>1</b>
1.1	$\sum_{i=0}^d k^i$ for $k > 0$ (Ex1.1) . . . . .	1
1.2	$\sum_{i=1}^d i k^i$ for $k > 0$ (Ex1.4) . . . . .	1
1.3	$\sum_{i=1}^d i 2^{d-i}$ (Ex1.3) . . . . .	1
1.4	$\sum_{i=1}^d i 2^i$ (Ex1.2) . . . . .	1
<b>2</b>	<b>Exercise 2 - Graph Tree's with Canonical Numbering</b>	<b>2</b>
2.1	$T_k^d$ with $k = 3$ and $d = 3$ . . . . .	2
2.2	$B_k^d$ with $k = 3$ and $d = 4$ . . . . .	2
<b>3</b>	<b>Exercise 3 - Planar Graph <math>H_d</math></b>	<b>2</b>
<b>4</b>	<b>Exercise 4 - Gray Code Embedding</b>	<b>2</b>
<b>5</b>	<b>Exercise 5 - Inverse Gray Code</b>	<b>2</b>
<b>6</b>	<b>Exercise 6 -</b>	<b>2</b>
<b>7</b>	<b>Exercise 7 -</b>	<b>2</b>

# 1 Exercise 1 - Closed Form Expressions

## 1.1 $\sum_{i=0}^d k^i$ for $k > 0$ (Ex1.1)

$$\begin{aligned}
 \sum_{i=0}^d k^i &= \sum_{i=0}^d k^i \\
 \sum_{i=0}^d k^i - k \sum_{i=0}^d k^i &= \sum_{i=0}^d k^i - k \sum_{i=0}^d k^i \\
 \sum_{i=0}^d k^i - k \sum_{i=0}^d k^i &= \sum_{i=0}^d k^i - \sum_{i=1}^{d+1} k^i \\
 \sum_{i=0}^d k^i (1 - k) &= 1 - k^{d+1} \\
 \sum_{i=0}^d k^i &= \frac{k^{d+1} - 1}{k - 1}
 \end{aligned} \tag{1}$$

## 1.2 $\sum_{i=1}^d i k^i$ for $k > 0$ (Ex1.4)

$$\begin{aligned}
 \sum_{i=1}^d i k^i &= \sum_{i=0}^d i k^i = \sum_{i=0}^d k \frac{d}{dk} k^i = k \frac{d}{dk} \sum_{i=0}^d k^i \quad \text{use (1)} \\
 \sum_{i=1}^d i k^i &= k \frac{d}{dk} \frac{1 - k^{d+1}}{1 - k} = \frac{d k^{d+2} - (d+1) k^{d+1} + k}{(1 - k)^2}
 \end{aligned} \tag{2}$$

## 1.3 $\sum_{i=1}^d i 2^{d-i}$ (Ex1.3)

$$\begin{aligned}
 \sum_{i=1}^d i 2^{d-i} &, \quad \text{use } k \text{ instead of } 2 \\
 \sum_{i=1}^d i k^{d-i} &= \sum_{i=0}^d d k^{d-i} - \sum_{i=0}^d (d-i) k^{d-i} \\
 &= d \sum_{j=0}^d k^j - \sum_{j=0}^d j k^j \quad \text{with } j := d - i \\
 &\quad \text{use (1)} \quad \text{use (2)} \\
 &= \frac{d(k^{d+1} - 1)}{k - 1} - \frac{d k^{d+2} - (d+1) k^{d+1} + k}{(1 - k)^2} \quad \text{set } k \text{ back to } 2 \\
 &= d 2^{d+1} - d - d 2^{d+2} + d 2^{d+1} + 2^{d+1} - 2 \\
 \sum_{i=1}^d i 2^{d-i} &= 2^{d+1} - 2 - d
 \end{aligned} \tag{3}$$

## 1.4 $\sum_{i=1}^d i 2^i$ (Ex1.2)

$$\sum_{i=1}^d i 2^i = d 2^{d+2} - (d+1) 2^{d+1} + 2 \quad \text{with use of (2)} \tag{4}$$

## 2 Exercise 2 - Graph Tree's with Canonical Numbering

2.1  $T_k^d$  with  $k = 3$  and  $d = 3$

2.2  $B_k^d$  with  $k = 3$  and  $d = 4$

## 3 Exercise 3 - Planar Graph $H_d$

## 4 Exercise 4 - Gray Code Embedding

## 5 Exercise 5 - Inverse Gray Code

## 6 Exercise 6 -

## 7 Exercise 7 -