



TECHNISCHE
UNIVERSITÄT
WIEN

184.754 Seminar on Algorithms

Paper: Coloring the Vertices of 9-pt and 27-pt
Stencils with Intervals

Camilo Tello Fachin
December 5, 2023

Introduction

Interval Vertex Coloring (IVC)

Special Case Analysis

Definitions

Special Cases

Heuristics

NP - Completeness


Algorithms

Application and Experiments


References

Still to do..

- Given $G(V, E)$




- Given $G(V,E)$
- Find a vertex coloring s.t. colors on adjacent vertices differ



- Given $G(V, E)$
- Find a vertex coloring s.t. colors on adjacent vertices differ

Formal Definition of VCP given $G(V, E)$

find $f(v)$:
 $\forall v \in V : \forall w \text{ in } \Gamma(v) : f(v) \neq f(w).$





- Given $G(V, E)$
- Find a vertex coloring s.t. colors on adjacent vertices differ

Formal Definition of VCP given $G(V, E)$


find $f(v)$:

$\forall v \in V : \forall w \text{ in } \Gamma(v) : f(v) \neq f(w).$




- Let $G(V,E)$ be an undirected graph and $w : V \mapsto \mathbb{Z}^+$ the weights.

$G(V,E)$




- Let $G(V,E)$ be an undirected graph and $w : V \mapsto \mathbb{Z}^+$ the weights.

$G(V,E)$




- A interval coloring of the vertices of G is a function $\text{start} : V \mapsto \mathbb{Z}^+$

$G(V, E)$



- Let $G(V, E)$ be an undirected graph and $w : V \mapsto \mathbb{Z}^+$ the weights.
- A interval coloring of the vertices of G is a function $\text{start} : V \mapsto \mathbb{Z}^+$
- Vertex v is colored with open interval: $[\text{start}(v), \text{start}(v) + w(v))$

$G(V, E)$ 

- Let $G(V, E)$ be an undirected graph and $w : V \mapsto \mathbb{Z}^+$ the weights.
- A interval coloring of the vertices of G is a function $\text{start} : V \mapsto \mathbb{Z}^+$
- Vertex v is colored with open interval: $[\text{start}(v), \text{start}(v) + w(v))$
- Neighboring Vertices must have disjoint color intervals: $\forall (a, b) \in E : [\text{start}(a), \text{start}(a) + w(a)) \cap [\text{start}(b), \text{start}(b) + w(b)) = \emptyset$.

- A interval coloring of the vertices of G is a function $\text{start} : V \mapsto \mathbb{Z}^+$


- A interval coloring of the vertices of G is a function $\text{start} : V \mapsto \mathbb{Z}^+$

- A particular coloring "start" of the vertices of G is said to use:

$$\text{maxcolor} = \max_{v \in V} \text{start}(v) + w(v) \text{ colors.}$$

- A interval coloring of the vertices of G is a function $\text{start} : V \mapsto \mathbb{Z}^+$

Interval Colored $G(V, E)$
with $\text{maxcolor} = 30$




- A particular coloring "start" of the vertices of G is said to use:

$$\text{maxcolor} = \max_{v \in V} \text{start}(v) + w(v) \text{ colors.}$$

- A interval coloring of the vertices of G is a function $\text{start} : V \mapsto \mathbb{Z}^+$

Interval Colored $G(V, E)$
with $\text{maxcolor} = 30$



- A particular coloring "start" of the vertices of G is said to use:

$$\text{maxcolor} = \max_{v \in V} \text{start}(v) + w(v) \text{ colors.}$$

Optimization Problem Instance:

Find a coloring $\text{start} : V \mapsto \mathbb{Z}^+$ that minimizes maxcolor .


- A maxcolor that is indeed minimal, is denoted with maxcolor^* .

Definition: 2D Stencil Interval Vertex Coloring (2DS-IVC)

A problem where G is a 9-pt 2D stencil, composed of $X \times Y$ vertices on a 2D grid such that two vertices (i,j) and (i',j') are connected iff $|i - i'| \leq 1$ and $|j - j'| \leq 1$.


Definition: 2D Stencil Interval Vertex Coloring (2DS-IVC)

A problem where G is a 9-pt 2D stencil, composed of $X \times Y$ vertices on a 2D grid such that two vertices (i,j) and (i',j') are connected iff $|i - i'| \leq 1$ and $|j - j'| \leq 1$.



Definition: 2D Stencil Interval Vertex Coloring (2DS-IVC)

A problem where G is a 9-pt 2D stencil, composed of $X \times Y$ vertices on a 2D grid such that two vertices (i, j) and (i', j') are connected iff $|i - i'| \leq 1$ and $|j - j'| \leq 1$.




Definition: 3D Stencil Interval Vertex Coloring (3DS-IVC)

A problem where G is a 27-point 3D stencil, composed of $X \times Y \times Z$ vertices on a 3D grid such that two vertices (i, j, k) and (i', j', k') are connected iff $|i - i'| \leq 1$, $|j - j'| \leq 1$, and $|k - k'| \leq 1$.

Definition: 2D Stencil Interval Vertex Coloring (2DS-IVC)

A problem where G is a 9-pt 2D stencil, composed of $X \times Y$ vertices on a 2D grid such that two vertices (i, j) and (i', j') are connected iff $|i - i'| \leq 1$ and $|j - j'| \leq 1$.



Definition: 3D Stencil Interval Vertex Coloring (3DS-IVC)

A problem where G is a 27-point 3D stencil, composed of $X \times Y \times Z$ vertices on a 3D grid such that two vertices (i, j, k) and (i', j', k') are connected iff $|i - i'| \leq 1$, $|j - j'| \leq 1$, and $|k - k'| \leq 1$.

Theorem

Let $G(V, E)$ be a clique K_n with n vertices, then

$$\text{maxcolor} = \sum_{v \in V} w(v)$$

is indeed an optimal coloring maxcolor^* .

Theorem

Let $G(V, E)$ be a clique K_n with n vertices, then

$$\text{maxcolor} = \sum_{v \in V} w(v)$$

is indeed an optimal coloring maxcolor^* .

- One can build such a coloring by listing vertices in descending order of weights and greedily allocating the color interval with the lowest available $\text{start}(v)$ with $\mathcal{O}(n)$ □.

Theorem

Let $G(V, E)$ be a clique K_n with n vertices, then

$$\text{maxcolor} = \sum_{v \in V} w(v)$$

is indeed an optimal coloring maxcolor^* .

- One can build such a coloring by listing vertices in descending order of weights and greedily allocating the color interval with the lowest available $\text{start}(v)$ with $\mathcal{O}(n)$ □.
- Particularly important result since the 2D-IVC's and 3D-IVC's are composed of K_4 's and K_8 's respectively.

Theorem

Let $G(V, E)$ be a clique K_n with n vertices, then

$$\text{maxcolor} = \sum_{v \in V} w(v)$$

is indeed an optimal coloring maxcolor^* .

- One can build such a coloring by listing vertices in descending order of weights and greedily allocating the color interval with the lowest available $\text{start}(v)$ with $\mathcal{O}(n)$ □.
- Particularly important result since the 2D-IVC's and 3D-IVC's are composed of K_4 's and K_8 's respectively.
- Therefore any identified K_4 in a 2D-IVC Problem and any K_8 in a 3D-IVC Problem are immediately (lose) lower bounds on maxcolor.

Definition

A graph $G(V,E)$ is considered Bipartite iff its vertices can be partitioned into two sets A and B , such that all edges have one endpoint in A and the other in B .

Definition

A graph $G(V,E)$ is considered Bipartite iff its vertices can be partitioned into two sets A and B , such that all edges have one endpoint in A and the other in B .

- All edges provide trivial lower bound:
 $\text{maxcolor}^* \geq w(i) + w(j), \forall (i,j) \in E.$

Definition

A graph $G(V, E)$ is considered Bipartite iff its vertices can be partitioned into two sets A and B , such that all edges have one endpoint in A and the other in B .

- All edges provide trivial lower bound:
 $\text{maxcolor}^* \geq w(i) + w(j), \forall (i, j) \in E.$
- One can find a coloring
 $\text{maxcolor}^* = \max_{(i,j) \in E} w(i) + w(j)$ in $\mathcal{O}(|E|)$. (Proof Omitted)

Definition

A graph $G(V, E)$ is considered Bipartite iff its vertices can be partitioned into two sets A and B , such that all edges have one endpoint in A and the other in B .

- All edges provide trivial lower bound:
 $\text{maxcolor}^* \geq w(i) + w(j), \forall (i, j) \in E.$
- One can find a coloring
 $\text{maxcolor}^* = \max_{(i,j) \in E} w(i) + w(j)$ in $\mathcal{O}(|E|)$. (Proof Omitted)
- Another important result, since the 9-pt stencil contains a Bipartite 5-pt stencil and the 27-pt stencil contains at least a Bipartite 7-pt stencil, and all paths are Bipartite.

Definition

A graph $G(V, E)$ is considered Bipartite iff its vertices can be partitioned into two sets A and B , such that all edges have one endpoint in A and the other in B .

- All edges provide trivial lower bound:
 $\text{maxcolor}^* \geq w(i) + w(j), \forall (i, j) \in E.$
- One can find a coloring
 $\text{maxcolor}^* = \max_{(i,j) \in E} w(i) + w(j)$ in $\mathcal{O}(|E|)$. (Proof Omitted)
- Another important result, since the 9-pt stencil contains a Bipartite 5-pt stencil and the 27-pt stencil contains at least a Bipartite 7-pt stencil, and all paths are Bipartite.
- Authors use this to construct approximation algorithms.

- Graphs that are not Bipartite, contain at least one cycle of odd length.

- Graphs that are not Bipartite, contain at least one cycle of odd length.
- Odd cycles have optimal interval colorings that are strictly greater than the largest weight of any clique in the cycle.

- Graphs that are not Bipartite, contain at least one cycle of odd length.
- Odd cycles have optimal interval colorings that are strictly greater than the largest weight of any clique in the cycle.
- The clique with the largest weight from the cycle shown earlier would be of interval size 25, while the optimal coloring of the entire graph (the odd cycle) has $\text{maxcolor}^* = 30$.

- Graphs that are not Bipartite, contain at least one cycle of odd length.
- Odd cycles have optimal interval colorings that are strictly greater than the largest weight of any clique in the cycle.
- The clique with the largest weight from the cycle shown earlier would be of interval size 25, while the optimal coloring of the entire graph (the odd cycle) has $\text{maxcolor}^* = 30$.
- Understanding of coloring of odd cycles yields new lower bounds.

Definition

Let maxpair be the maximal sum of any two consecutive vertices. ¹

$$\text{maxpair} = \max_i w(i, i + 1)$$

¹Here G is cycle, neighbors of vertex x are $x + 1$ and $x - 1$.

Definition

Let maxpair be the maximal sum of any two consecutive vertices. ¹

$$\text{maxpair} = \max_i w(i, i + 1)$$

Definition

Let minchain3 be the minimum sum of any 3 consecutive vertices:

$$\text{minchain3} = \min_i w(i, i + 1, i + 2)$$

¹Here G is cycle, neighbors of vertex x are $x + 1$ and $x - 1$.

Definition

Let maxpair be the maximal sum of any two consecutive vertices. ¹

$$\text{maxpair} = \max_i w(i, i + 1)$$

Definition

Let minchain3 be the minimum sum of any 3 consecutive vertices:

$$\text{minchain3} = \min_i w(i, i + 1, i + 2)$$

Theorem

Let G be an odd cycle, then there holds:

$$\text{maxcolor}^* = \max(\text{maxpair}, \text{minchain3})$$

¹Here G is cycle, neighbors of vertex x are $x + 1$ and $x - 1$.

Idea of Proof

Idea of Proof

- If G is an odd cycle, there is an algorithm that yields $\max(\text{maxpair}, \text{minchain3})$ colors, which means $\text{maxcolor}^* \leq \max(\text{maxpair}, \text{minchain3})$.

Idea of Proof


- If G is an odd cycle, there is an algorithm that yields $\max(\text{maxpair}, \text{minchain3})$ colors, which means $\text{maxcolor}^* \leq \max(\text{maxpair}, \text{minchain3})$.
- If G is an odd cycle, then $\text{maxcolor}^* \geq \max(\text{maxpair}, \text{minchain3})$.

Idea of Proof

- If G is an odd cycle, there is an algorithm that yields $\max(\text{maxpair}, \text{minchain3})$ colors, which means $\text{maxcolor}^* \leq \max(\text{maxpair}, \text{minchain3})$.
- If G is an odd cycle, then $\text{maxcolor}^* \geq \max(\text{maxpair}, \text{minchain3})$.
- Coinciding bounds conclude proof [1] \square .

Idea of Proof

- If G is an odd cycle, there is an algorithm that yields $\max(\text{maxpair}, \text{minchain3})$ colors, which means $\text{maxcolor}^* \leq \max(\text{maxpair}, \text{minchain3})$.
- If G is an odd cycle, then $\text{maxcolor}^* \geq \max(\text{maxpair}, \text{minchain3})$.
- Coinciding bounds conclude proof [1] \square .
- One can verify this for the previous example



Lemma

$3DS-IVC \in NP$

Lemma

$3DS-IVC \in NP$

Proof.

Given a solution for 3DS-IVC, encoded with pairs of integers which are bounded between 0 and $\sum_{v \in V} w(v_i)$ \mapsto a solution in polynomial space.

Lemma

$3DS\text{-}IVC \in NP$

Proof.

Given a solution for 3DS-IVC, encoded with pairs of integers which are bounded between 0 and $\sum_{v \in V} w(v_i) \mapsto$ a solution in polynomial space.

Verify that no adjacent edges have overlapping intervals.

$\forall (u, v) \in E :$

$$[\text{start}(u), \text{start}(u) + w(u)) \cap [\text{start}(v), \text{start}(v) + w(v)) = \emptyset.$$

Lemma

$3DS\text{-}IVC \in NP$

Proof.

Given a solution for 3DS-IVC, encoded with pairs of integers which are bounded between 0 and $\sum_{v \in V} w(v_i) \mapsto$ a solution in polynomial space.

Verify that no adjacent edges have overlapping intervals.

$\forall (u, v) \in E :$

$$[\text{start}(u), \text{start}(u) + w(u)) \cap [\text{start}(v), \text{start}(v) + w(v)) = \emptyset.$$

Since $|E| \leq \frac{n(n-1)}{2}$ and checking if two intervals overlap is in $\mathcal{O}(1)$, the verification can be done in $\mathcal{O}(n^2)$ □.

Idea of Proof

- Construct an instance 3DS-IVC from an instance of NAE-3SAT in polynomial time.

Idea of Proof

- Construct an instance 3DS-IVC from an instance of NAE-3SAT in polynomial time.
- Verify that a positive instance of NAE-3SAT results in a positive instance of 3DS-IVC.

Idea of Proof

- Construct an instance 3DS-IVC from an instance of NAE-3SAT in polynomial time.
- Verify that a positive instance of NAE-3SAT results in a positive instance of 3DS-IVC.
- Verify that if the created instance of 3DS-IVC is positive, then the instance of NAE3-SAT is also positive.

Since the 3DS-IVC problem is in NP and is harder than NAE-3SAT which is an NP-Complete problem, 3DS-IVC is NP-Complete □.

Idea of Proof

- Construct an instance 3DS-IVC from an instance of NAE-3SAT in polynomial time.
- Verify that a positive instance of NAE-3SAT results in a positive instance of 3DS-IVC.
- Verify that if the created instance of 3DS-IVC is positive, then the instance of NAE3-SAT is also positive.

Since the 3DS-IVC problem is in NP and is harder than NAE-3SAT which is an NP-Complete problem, 3DS-IVC is NP-Complete \square .

The proof yields:

Theorem

Deciding whether a 27-pt stencil can be colored with less than k colors is NP-Complete. [1]

Lemma

In the application section I will probably cite [2] and also [3].

-  **D. Durrman and E. Saule**, "Coloring the vertices of 9-pt and 27-pt stencils with intervals," in **2022 IEEE International Parallel and Distributed Processing Symposium (IPDPS)**, 2022, pp. 963–973. DOI: [10.1109/IPDPS53621.2022.00098](https://doi.org/10.1109/IPDPS53621.2022.00098).
-  **A. Hohl, E. Delmelle, W. Tang, and I. Casas**, "Accelerating the discovery of space-time patterns of infectious diseases using parallel computing," **Spatial and Spatio-temporal Epidemiology**, vol. 19, pp. 10–20, 2016, ISSN: 1877-5845. DOI: <https://doi.org/10.1016/j.sste.2016.05.002>. [Online]. Available: <https://www.sciencedirect.com/science/article/pii/S187758451530040X>.
-  **E. Saule, D. Panchananam, A. Hohl, W. Tang, and E. M. Delmelle**, "Parallel space-time kernel density estimation," **2017 46th International Conference on Parallel Processing (ICPP)**, pp. 483–492, 2017. [Online]. Available: <https://api.semanticscholar.org/CorpusID:6645797>.